



City Research Online

City St George's, University of London

Citation: Colombo, A., Xu, Y. & Dong, H. (2023). Postprandial glycaemic response to white and wholemeal bread consumption between normal weight and overweight/obese healthy adults. *Proceedings of the Nutrition Society*, 82(OCE1), doi: 10.1017/s0029665123000241 ISSN 0029-6651 doi: 10.1017/s0029665123000241

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/30059/>

Link to published version: <https://doi.org/10.1017/s0029665123000241>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

Postprandial glycaemic response to white and wholemeal bread consumption between normal weight and overweight/obese healthy adults.

A Colosimo¹, Y Xu¹, H Dong^{2*}

1. School of Life Sciences, Coventry University, Coventry, UK, 2. School of Health and Psychological Sciences, City, University of London, London, UK.

*Corresponding author, honglin.dong@city.ac.uk

Introduction

The impact of postprandial glucose response has been implicated in the development of chronic metabolic diseases, and obesity is an important risk factor⁽¹⁾. Few studies investigated the effect of body weight on postprandial glycaemic response. This study aimed to investigate the difference in postprandial glycaemic response to commonly consumed white and wholemeal bread in the UK⁽²⁾ between normal weight and overweight/obese adults, and to investigate the difference in postprandial glycaemic response between white and wholemeal bread consumption in adults. Wholemeal bread contains higher dietary fibre than white bread, therefore is regarded as a healthier choice⁽³⁾.

Methods

Twenty healthy adults were each given two slides of white or wholemeal bread alongside 150ml of pure orange juice and 10g butter on separate visits at random order after fasting for 12 hours. The white bread meal contains 369.5 kcal, 60g carbohydrates, 17.8g sugar and 3.6g fibre, while the wholemeal bread meal contains 355.5 kcal, 48g carbohydrates, 17.4g sugar, and 5.6g fibre. The blood glucose concentration was measured before taking meals, 30 min, 60 min, 90 min and 120 min postprandially by finger pricks using Biosen Blood Glucose/Lactate Analyser (EKF Diagnostics, Cardiff). Participants consumed the meal within 10 minutes and kept sedentary with only water consumption allowed during the study period. The participant information of age, sex, ethnicity, body weight and height, and fat composition measured by Tanita MC-980MA PLUS (Tanita Company, Tokyo) were collected on the first visit. The difference in the area under the curve and the peak value of the postprandial glycaemic response were analysed between different groups using the independent t-test or between different meals using paired t-test.

Results

There was no significant difference in the area under the curve of postprandial glucose response between normal weight (n=10) and overweight/obese (n=10), between females (n=12) and males (n=8) or between white Caucasians (n=13) and non-white (n=7) participants, regardless of white or wholemeal bread consumption. However, the peak value was significantly higher in non-white than white participants (6.11 nmol/L vs 5.15 nmol/L, P=0.015) after white bread consumption, while males showed a significantly higher peak value than females after the wholemeal bread consumption (6.22 nmol/L vs 4.78 nmol/L, P=0.021). There was no significant difference in the area under the curve or peak values between white and wholemeal bread consumption.

Conclusion

The results indicated that body weight did not play significant role in postprandial glycaemic response to white or wholemeal bread consumption, and the dietary fibre content in the wholemeal bread used in the current study may not high enough to deliver the health benefit of postprandial glycaemic response. Further research is needed to include a larger sample size with power calculation to investigate the postprandial glycaemic response between sexes and different ethnicities.

Reference

1. Blaak EE, Antoine JM, Benton D, et al. (2012) *Obes Rev* 13(10), 923–984
2. Lockyer S & Spiro A (2020) *Nutr Bull* 45(2), 133–164.
3. Lužnik IV I, Lužnik A, Mateja Lušnic Polak ML, et al. (2019) *J Nutr Intermed Metab* 16, 100097-100103