



City Research Online

City St George's, University of London

Citation: Velthorst, E., Socrates, A., Alizadeh, B. Z., van Amelsvoort, T., Bartels-Velthuis, A. A., Bruggeman, R., Cahn, W., de Haan, L., Schirmbeck, F., Simons, C. J. P., et al (2023). Age-Related Social Cognitive Performance in Individuals With Psychotic Disorders and Their First-Degree Relatives. *Schizophrenia Bulletin*, 49(6), pp. 1460-1469. doi: 10.1093/schbul/sbad069

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/30519/>

Link to published version: <https://doi.org/10.1093/schbul/sbad069>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).



Draft Manuscript for Review. Please review online at <http://mc.manuscriptcentral.com/oup/szbltn>

Age-related social cognitive performance in individuals with psychotic disorders and their first-degree relatives

| | |
|-------------------------------|---|
| Journal: | <i>Schizophrenia Bulletin</i> |
| Manuscript ID | SZBLTN-ART-22-0642.R1 |
| Manuscript Type: | Regular Article |
| Date Submitted by the Author: | 12-Apr-2023 |
| Complete List of Authors: | Velthorst, Eva; GGZ Noord-Holland-Noord Socrates, Adam; Icahn School of Medicine at Mount Sinai Alizadeh, Behrooz; University of Groningen, University Medical Center Groningen, Epidemiology, Unit of Genetic Epidemiology Amelsoort, Therese; maastricht university, Bartels-Velthuis, Agna; University Medical Center Groningen, University Center for Psychiatry Investigators, GROUP; UMCG, psychiatry Cahn, Wiepke; UMC Utrecht Hersencentrum, Psychiatry de Haan, Lieuwe; Amsterdam UMC -AMC Campus, Psychiatry Schirmbeck, Frederike; Amsterdam UMC Locatie AMC, Psychiatry Simons, Claudia; Maastricht University, Department of Psychiatry & Neuropsychology; GGzE, Department of Research, Development and Innovation van Os, Jim; UMC Utrecht Fett, Anne-Kathrin; City University, Department of Psychology; Vrije Universiteit Amsterdam, Department of Clinical, Neuro and Developmental Psychology; King's College London, CSI Lab, Institute of Psychiatry, Psychology and Neuroscience, Department of Psychosis Studies |
| Keywords: | Psychosis, Social cognition, Aging, Siblings, schizophrenia |
| | |

SCHOLARONE™
Manuscripts

Age-related social cognitive performance in individuals with psychotic disorders and their first-degree relatives

Running title: psychosis and age-related social cognitive deficit

Eva Velthorst¹, Adam Socrates^{2,3}, GROUP investigators (Behrooz Z. Alizadeh^{4,5}, Therese van Amelsvoort⁶, Agna A. Bartels-Velthuis⁴, Richard Bruggeman^{4,7}, Wiepke Cahn^{8,9}, Lieuwe de Haan^{3,10}, Frederike Schirmbeck^{3,10}, Claudia J.P. Simons^{6,11}, Jim van Os^{8,12}), Anne-Kathrin Fett^{13,14}

1. Department of Research, Mental Health Organization 'GGZ Noord-Holland-Noord', Heerhugowaard, The Netherlands

2. Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, US

3. Seaver Center for Autism Research and Treatment, Icahn School of Medicine at Mount Sinai, New York, US

4. University of Groningen, University Medical Center Groningen, University Center for Psychiatry, Rob Giel Research center, Groningen, The Netherlands

5. University Medical Center Groningen, Department of Epidemiology, Groningen, The Netherlands

6. Maastricht University, Department of Psychiatry and Neuropsychology, School for Mental Health and Neuroscience, Maastricht, The Netherlands

7. University of Groningen, Department of Clinical and Developmental Neuropsychology, Groningen, The Netherlands

8. University Medical Center Utrecht, Department of Psychiatry, Brain Centre Rudolf Magnus, Utrecht University, Utrecht, The Netherlands

9. Altrecht, General Mental Health Care, Utrecht, The Netherlands

10. Arkin, Institute for Mental Health, Amsterdam, The Netherlands

11. GGzE Institute for Mental Health Care, Eindhoven, The Netherlands

12. King's College London, King's Health Partners, Department of Psychosis Studies, Institute of Psychiatry, London, UK

13. Department of Psychology, City, University of London, London, UK

14. Department of Psychosis Studies, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK

Corresponding author:

Eva Velthorst, PhD

Mental Health Service Organization 'GGZ Noord-Holland-Noord', Department of Research

Stationsplein 138, 1703 WC Heerhugowaard, The Netherlands

E: e.velthorst@ggz-nhn.nl

1
2
3 **Abstract:** 245/250 words; **Word count:** 3998/4000 words

4 **Abstract**

5 **Background**

6 Social cognitive impairment is a recognized feature of psychotic disorders. However, potential age-related
7 differences in social cognitive impairment have rarely been studied.
8
9
10

11
12 **Study Design**

13 Data came from 905 individuals with a psychotic disorder, 966 unaffected siblings, and 544 never-
14 psychotic controls aged 18-55 who participated in the Genetic Risk and Outcome of Psychosis (GROUP)
15 study. Multilevel linear models were fitted to study group main effects and the interaction between group
16 and age on emotion perception and processing (EPP; Degraded Facial Affect Recognition) and theory of
17 mind (ToM; Hinting Task) performance. Age-related differences in the association between
18 sociodemographic and clinical factors, and EPP and ToM were also explored.
19
20
21
22
23
24

25 **Study Results**

26 Across groups, EPP performance was associated with age ($\beta = -.02$, $z = -7.60$, CI95%: $-.02$, $-.01$, $p <$
27 $.001$), with older participants performing worse than younger ones. A significant group-by-age interaction
28 on ToM (group-by-age interaction: $(X^2(2) = 13.15$, $p = .001$)) indicated that older patients performed
29 better than younger ones, while no age-related difference in performance was apparent among siblings
30 and controls. In patients, the association between negative symptoms and ToM was stronger for younger
31 than older patients ($z = 2.16$, $p = .03$).
32
33
34
35
36
37

38 **Conclusions**

39 The findings point to different age-related performance patterns on tests of two key social cognitive
40 domains. ToM performance was better in older individuals, although this effect was only observed for
41 patients. EPP was less accurate in older compared with younger individuals. These findings have
42 implications for when social cognitive training in patients should be offered.
43
44
45
46
47
48

49 **Keywords:** aging, schizophrenia, siblings, psychosis, cognition, functional impairment
50
51
52
53
54
55
56
57
58
59
60

Introduction

Individuals with schizophrenia and related psychotic disorders often experience impairments in social cognition.^{1,2} Social cognition may impact functional outcomes more strongly than non-social cognition and psychotic symptoms³. While social cognition, referring to the psychological processes that enable people to understand social contexts, spans multiple domains (e.g., mentalization or theory of mind [ToM]), emotion perception and processing (EPP), and social perception, the vast amount of existing psychosis research has focused on ToM and EPP. However, even within these domains, there are knowledge gaps.

One concerns the age-related development of ToM and EPP impairment. Some studies suggest that impairments in EPP and ToM are already present prior to the first psychotic episode^{4,5}, and that social cognitive performance in individuals with a psychotic disorder is relatively stable over time⁶⁻⁹ and across illness phases.^{2,10,11} However, the effect of age on social cognitive performance remains largely unexplored and it is unclear whether potential age-related differences in patients with a psychotic disorder are comparable to those of never-psychotic individuals.¹² One study showed comparable ToM impairment in young individuals in the early phase and older individuals in more chronic phase of the illness.¹⁰ However, the study did not include a control group, which limits conclusions relating to ‘typical’ age-related functioning.

Further, most, but not all¹³, studies suggest that the social cognitive performance of unaffected siblings of patients lies in between never-psychotic individuals and those diagnosed with a psychotic disorder (for a review see¹⁴). As siblings share genetic and/or environmental factors that might be related to an increased risk for psychosis but are not exposed to the effects that are related to having a psychotic disorder (e.g., medication, stigma), investigations of social cognition in siblings can help to understand the impact of risk versus disorder-related mechanisms. Similar social cognitive performance in siblings and controls while patients diverge as they age, would suggest age-related processes specific to the disorder. Similar social cognitive performance in siblings and patients, would suggest familial rather than disorder-related mechanisms.

It remains unknown whether illness-related factors, such as symptom severity, illness duration, and non-social cognitive impairment relate differentially to social cognition throughout adulthood¹⁵, or whether possible age-related cognitive processes in psychotic disorders differ by sex. While previous psychosis studies found only small sex differences (e.g.,^{16,17}), or no sex difference¹⁸, the potential interaction

1
2
3 between age and sex on social cognitive performance in adults with psychotic disorders has not yet been
4 studied. Finally, social cognition has been related to functional outcomes, but this relationship may vary
5 with age. Identifying age-related patterns can inform interventions that aim to improve or retain patients
6 independent functioning, social connections, and a better quality of life as they age.
7
8
9

10
11 We investigated these knowledge gaps in a large sample of patients with a non-affective psychotic
12 disorder, unaffected siblings, and controls from the Genetic Risk and Outcome of Psychosis (GROUP)
13 study. The cross-sectional data on ToM and EPP comprised 2,415 individuals aged 18 to 55. Specifically,
14 we examined (1) whether possible group differences in EPP and ToM varied by age, (2) age-related
15 differences in the cross-sectional associations between EPP, ToM and sociodemographic, and illness
16 related factors, and (3) explored associations between age related patterns of EPP, ToM at baseline and
17 everyday functioning of the patient group three years later. **Based on previous cross-sectional and**
18 **longitudinal work, we expected to find relatively similar ToM and EPP scores by age.**
19
20
21
22
23
24

25 **Methods**

26 **Study population**

27
28 Data on 2,415 study participants, including 905 individuals with a non-affective psychotic disorder, 966
29 unaffected siblings, and 544 never-psychotic controls aged 18-55 were collected in 35 mental health care
30 institutes across the Netherlands and Belgium. **The control group was selected through a random system**
31 **of mailings to addresses in the catchment area and were excluded if they had a (history of a) psychotic**
32 **disorder, of a first-degree family member with a lifetime psychotic disorder.** The GROUP study, a
33 multisite longitudinal observational study was carried out between April 2004 and December 2013, with
34 the overall goal to identify vulnerability and protective factors that influence the onset and course of
35 psychotic disorders.¹⁹ The study was approved by the Medical Ethics Committee of the University
36 Medical Center Utrecht and subsequently by local review boards of each participating institute.
37
38
39
40
41
42
43

44 **Procedure**

45 **Non-social and social cognitive tests were administered first to make sure participants were focused and**
46 **fully concentrated. These tests were followed by the PANSS interview and social functioning scale.¹⁹**
47
48
49

50 **Materials**

51
52 Demographic information (i.e., sex, age, ethnicity, study site) and information on illness duration was
53 captured by means of a general demographic questionnaire **at baseline assessment.**
54
55
56
57
58
59
60

Assessments of social cognitive functioning

All participants completed at least one social cognition measure in the domain of (i) EPP (i.e., the ability to infer emotional information from facial expressions), and/or (ii) ToM (i.e., the ability to comprehend the mental states of others) at baseline assessment:

(i) *Emotion perception and processing (EPP)*

The computerized Degraded Facial Affect Recognition (DFAR) task, presented participants with 64 photographs of four individuals (two males/two females) expressing happy, angry, fearful and neutral emotions (16 images for each).²⁰ To increase difficulty and enhance the contribution of perceptual expectancies and interpretation, photographs of the faces were passed through a filter resulting in a reduced visual contrast by 30%. The total scores were calculated by the percentage of total correct answers per domain, and across domains (DFAR).

(ii) *Theory of Mind (ToM)*

The Hinting Task²¹ measures the ability to infer real intentions behind indirect speech. Ten short passages read aloud by the interviewer, present an interaction between two characters that included dropping a hint. Participants had to indicate what the hint meant. If the first response was inaccurate, a second hint was given, allowing participants to earn partial credit. An example question from the task is: *Rebecca's birthday is approaching. She says to her Dad, 'I love animals, specially dogs.'* Question 1: *What does Rebecca really mean when she says this? Answer: She wants her dad to get her a dog for her birthday. If an incorrect answer given to question 1: Rebecca goes on to say, 'Will the pet shop be open on my birthday, Dad?'* Question 2: *What does Rebecca want her dad to do? Answer: She wants her dad to get her a dog for her birthday.* Total scores range from 0 to 20. The Hinting Task has adequate test-retest reliability, small practice effects, and limited potential for floor effects, although ceiling effects have been noted.^{22,23}

Face recognition

A measure of the ability to match unfamiliar faces, the short version of the Benton Facial Recognition Test²⁴ was used at baseline to assess whether potential deficits in facial affect recognition are accounted for by general facial-recognition ability.

Non-social cognitive functioning

An abbreviated WAIS-III²⁵, comprising the information, arithmetic, block design, and digit symbol coding subtests, was used to measure performance in the domains of verbal knowledge, working memory, visuospatial processing, and processing speed, respectively. Here we used baseline IQ estimates.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Symptom severity

For patients, clinical psychotic symptoms at baseline were assessed with the Positive And Negative Syndrome Scale (PANSS)²⁶, consisting of three subscales: positive syndrome scale (item P1–P7), a negative syndrome scale (items N1–N7) and general psychopathology scale (item G1–G16).

Social functioning

The Social Functioning Scale (SFS)²⁷ measures social functioning in people diagnosed with a psychotic disorder and includes scales for. Here, we focused on interpersonal behavior and prosocial activities subscale only (interpersonal behavior (e.g., number of friends), pro-social activities (e.g., sports, visiting friends). The SFS was assessed at the three-year follow-up.

Data analyses

Data release 8.0 was used for the analyses. Group differences in sample characteristics were examined using χ^2 , t- and Mann–Whitney-U-tests. Raw scores on the social cognitive tests were z transformed against the average scores of the entire study sample (i.e., combining patients, siblings, and controls). Thus, β values throughout represent standardized effect sizes (ES), with values 0.2, 0.5, and 0.8 indicating small, medium, and large ES, respectively. These z scores were used in all statistical analyses. To account for multiple comparisons, a p -value of $(0.05 / 3 \text{ (number of groups)}) \leq 0.017$ was considered statistically significant.

- (1) To examine whether group-differences in EPP and ToM varied by age, multilevel linear models (MLMs) were fitted to account for the hierarchical structure of the data (i.e., with random intercepts for family and study site). We examined age-by-group interactions, group, and age as predictors if interactions were non-significant. Statistically significant age-by-EPP or ToM interactions were followed up with simple effects analyses. Non-significant age-by-group interactions were removed from the models before interpretation of any main effects. Sex was entered as a covariate in all models that investigated between group effects. Separate models including IQ as additional covariate (for both EPP and ToM analyses) and including face recognition ability scores (for EPP analyses) were also conducted.
- (2) To explore whether the association between symptom severity (PANSS positive, negative, and general), illness duration, non-social cognitive impairment (IQ), and socio demographic

characteristics (sex, ethnicity (white/non-white)), in the patient group differed by age, the predictors were added as interaction (with age) and main effects to the MLM's separately. Statistically significant age-by-factor of interest interactions were followed up with simple effects analyses. Non-significant interactions were removed from the models before interpretation of any main effects.

- (3) To explore whether the association between social cognition and social functional outcome varied by age similar MLM models were run, this time with the social functioning score at three-year follow-up as dependent variable, and social cognition at baseline, age, and their interaction as predictors. In sensitivity analyses we separately explored age-related patterns in social functioning. In this MLM model age-by-group interactions on social functioning were run without adding the social cognition scores as predictor to the model.

For visualization purposes and ease of interpretation, figures in the main text are presented based on age categorizations (very early adulthood: 18-25; early adulthood: 26-30; mid-adulthood: 31-40; later-adulthood: 41-55). The distribution of age categorizations for each group can be found in **Supplementary Table 1**.

Results

-----Table 1-----

Group differences in social cognitive performance

EPP: DFAR performance ($X^2(2) = 91.45, p < .001$) and performance in subcategories neutral ($X^2(2) = 25.49, p < .001$), angry ($X^2(2) = 42.62, p < .001$), fearful ($X^2(2) = 49.33, p < .001$), and happy ($X^2(2) = 11.01, p < .01$) differed significantly between patients, siblings, and controls. Patients showed medium-sized deficits in the recognition of most emotions in comparison to controls and siblings (DFAR: $z = -8.29$ and -8.11 , respectively, $p < .001$; neutral $z = -4.51$ and $-3.95, p < .001$; angry: $z = -6.03$ and $-4.91, p < .001$; fearful: $z = -6.23$ and $-5.95, p < .001$), except for the recognition of happy faces, where patients only differed from siblings ($z = -3.29, p < .01$) but not controls ($z = -1.19, p = .23$). Siblings performed marginally worse than controls on the total DFAR ($z = -2.55, p = .011$). Specifically, they were less accurate in recognizing fearful emotions ($z = -2.35, p < .01$), but performed similar to controls for neutral ($z = -1.19, p = .32$), angry ($z = -2.07, p = .04$), and happy emotions ($z = 1.7, p = .08$).

ToM: The groups differed significantly on the Hinting Task ($X^2(2) = 240.85, p < .001$). Patients performed worse than controls ($z = -13.16, p < .001$) and siblings ($z = -13.32, p < .001$). Siblings performed in between patients ($z = -14.64, p < .001$) and controls \ controls ($z = -2.41, p = .016$).

See supplementary table 2 for an EPP and TOM score distribution by group.

-----Figure 1-----

(1) Age-related patterns for EPP and ToM skills differ by group

EPP: The group-by-age interactions for DFAR, neutral, happy, and angry were not significant (all $p > .16$), suggesting that age differences in EPP followed a similar pattern in all groups (see **Figure 1**). Across groups, age was associated with the DFAR ($\beta = -.02$, $z = -7.60$, CI95%: $-.02$, $-.01$, $p < .001$), and with the recognition of happy ($\beta = -.01$, $z = -4.05$, CI95%: $-.02$, $-.005$, $p < .001$), and angry faces ($\beta = -.006$, $z = -2.60$, CI95%: $-.01$, $-.002$, $p < .01$), such that older participants performed worse than younger ones. The age effects for DFAR and happy were comparable after controlling for IQ (DFAR: $\beta = -.02$, $z = -6.92$, CI95%: $-.02$, $-.01$, $p < .001$; happy: $\beta = -.009$, $z = -3.38$, CI95%: $-.01$, $-.004$, $p = .001$), but only marginally reduced for the recognition of angry faces ($\beta = -.005$, $z = -2.10$, CI95%: $-.01$, $-.0004$, $p = .04$). No age effect was observed for the recognition of neutral faces ($\beta = -.003$, $z = -1.16$, CI95%: $-.01$, $.002$, $p = .25$). For fearful faces a group-by-age interaction was observed ($X^2(2) = 7.63$, $p = 0.02$), which remained after controlling for IQ ($X^2(2) = 8.58$, $.01$); age effects were significant in all groups, although effects were stronger in siblings ($\beta = -.03$, $z = -7.25$, CI95%: $-.04$, $-.02$, $p < .001$) than controls ($\beta = -.02$, $z = -3.83$, CI95%: $-.04$, $-.001$, $p < .001$) and patients ($\beta = -.02$, $z = -3.87$, CI95%: $-.03$, $-.01$, $p < .001$).

No age-related differences in face recognition ability, i.e., Benton Facial Recognition Test scores were observed ($\beta = -.003$, $z = -1.44$, CI95%: $-.009$, $.001$, $p = .15$), and all observed DFAR effects were comparable after controlling for Benton scores. **Supplementary Figure 1** shows the individuals graphs by emotion.

ToM: There was a significant group-by-age interaction on Hinting Task performance ($X^2(2) = 13.15$, $p = .001$), where older patients performed better than younger ones ($\beta = .02$, $z = 4.13$, CI95% $.01$, $.03$, $p < .001$) while no such age-related pattern emerged for controls (see **Figure 2**; $\beta = -.003$, $z = -.89$, CI95%: $-.01$, $.004$, $p = .38$) or siblings ($\beta = .01$, $z = 1.78$, CI95% $-.001$, $.01$, $p = .08$). These findings were not accounted for by IQ differences (group-by-age interaction after controlling for IQ: $X^2(2) = 10.35$, $p = .006$).

-----Figure 2-----

(2) *The effect of age on the association between illness factors and EPP and ToM performance in patients*

We conducted exploratory analyses to investigate whether the effect of predictors of social cognitive performance differed by age. Sociodemographic and clinical correlates (r) of performance on DFAR and Hinting Task in the patient group are displayed in **Supplementary Figure 2**.

Patient characteristics

There were no *age-by-sex* interactions for DFAR ($\beta = -.007$, $z = -.66$, CI95%: $-.03$, $.01$, $p = .51$), and Hinting Task performance ($\beta = -.003$, $z = -.28$, CI95%: $-.03$, $.02$, $p = .78$). Removing the interaction from the models, females outperformed males on **the DFAR task** (DFAR: $\beta = .24$, $z = 2.63$, CI95%: $.06$, $.41$, $p = .009$).

Similarly, no *age-by-ethnicity* interactions were found (DFAR: $\beta = .02$, $z = 1.21$, CI95%: $-.001$, $.04$, $p = .23$; Hinting Task: $\beta = .01$, $z = 0.97$, CI95%: $-.01$, $.04$, $p = .33$). Non-white did not differ from white participants on the DFAR ($\beta = -.02$, $z = -.27$, CI95%: $-.20$, $.14$, $p = .79$), but had lower scores on the Hinting Task ($\beta = -.32$, $z = -3.14$, CI95%: $-.52$, $-.26$, $p = .002$).

Finally, there were no *age-by-illness duration* interactions (DFAR: $\beta = .0005$, $z = -.64$, CI95%: $-.002$, $.001$, $p = .53$; Hinting: $\beta = .0007$, $z = 0.76$, CI95%: $-.001$, $.002$, $p = .44$). No main effect of illness duration was detected for DFAR or Hinting Task performance (DFAR: $\beta = .005$, $z = .55$, CI95%: $-.01$, $.02$, $p = .58$; Hinting: $\beta = -.006$, $z = -.63$, CI95%: $-.03$, $.01$, $p = .53$).

Symptom severity

We observed a **marginal** *age-by-negative symptom* interaction for the Hinting Task, suggesting **a slightly** stronger association between negative symptoms and ToM performance in younger participants ($\beta = .01$, $z = 2.16$, CI95%: $.001$, $.03$, $p = .031$). There was no interaction for the DFAR ($\beta = .004$, $z = .68$, CI95%: $-.007$, $.02$, $p = .50$). Regardless of age, higher negative symptom severity was associated with poorer DFAR performance (DFAR: $\beta = -.18$, $z = -4.24$, CI95%: $-.26$, $-.10$, $p = .009$).

There were no significant *age-by-positive symptom* interactions on social cognitive performance (DFAR: $\beta = .006$, $z = .88$, CI95%: $-.007$, $.02$, $p = .38$; Hinting: $\beta = .005$, $z = .62$, CI95%: $-.01$, $.02$, $p = .54$).

Participants with higher positive symptoms performed worse on DFAR and Hinting task (DFAR: $\beta = -.13$, $z = -2.72$, CI95%: $-.23$, $-.04$, $p = .007$; Hinting: $\beta = -.23$, $z = -4.14$, CI95%: $-.35$, $-.12$, $p < .001$).

No *age-by-general symptom* interactions were observed (DFAR: $\beta = .003$, $z = 0.35$, CI95%: $-.01$, $.02$, $p = .72$; Hinting: $\beta = .009$, $z = .93$, CI95%: $-.01$, $.03$, $p = .35$), but general symptoms had a negative association with performance of both social cognitive tests (DFAR: $\beta = -.21$, $z = -2.95$, CI95%: $-.35$, $-.07$, $p = .003$; Hinting: $\beta = -.46$, $z = -5.60$, CI95%: $-.61$, $-.30$, $p < .001$).

Non-social cognition

There were no *age-by-IQ* interactions for DFAR ($\beta < .0001$, $z = -.05$, CI95%: $-.0006$, $.0005$, $p = .96$) or Hinting Task ($\beta = -.0002$, $z = -.91$, CI95%: $-.0009$, $.0003$, $p = .36$). IQ predicted DFAR ($\beta = .01$, $z = 5.97$, CI95%: $.009$, $.02$, $p < .0001$) and Hinting Task performance ($\beta = .02$, $z = 10.95$, CI95%: $.02$, $.03$, $p < .0001$), regardless of age.

(3) EPP and ToM as a predictor of social functioning across ages

We explored the association between DFAR and Hinting Task performance, age and two indices of social functioning three years later (see **Supplementary Table 3** for a partial correlation matrix). There was no interaction between DFAR and age at baseline on interpersonal functioning or prosocial activities at follow-up (both $p > .15$). Regardless of age, DFAR performance was significantly associated with interpersonal functioning ($\beta = 1.91$, $z = 2.51$, CI95%: $.42$, 3.40 , $p < .01$), but not with pro-social activities ($p = .30$). There was no interaction between Hinting Task performance and age at baseline on interpersonal functioning or prosocial activities at follow-up (both $p > .22$). Regardless of age, Hinting Task performance was **marginally** associated with interpersonal functioning ($\beta = 1.59$, $z = 2.35$, CI95%: $.26$, 2.91 , $p = .02$), but not with pro-social activities ($p = .76$).

Sensitivity analyses: age-related patterns in social functioning among the patient group

MLM models exploring age-related differences in social functioning revealed an age effect for prosocial activities. In this model, participation in pro-social activities was significantly lower among older patients ($\beta = -.23$, $z = -2.96$, CI95%: $-.39$, -0.08 , $p = .003$). No age-related differences were observed for interpersonal functioning ($\beta = -.19$, $z = -.12$, CI95%: $-.41$, $.03$, $p = .09$).

Discussion

This large cross-sectional study provides the first comprehensive analysis of age-related differences in EPP and ToM impairment in individuals with a non-affective psychotic disorder, unaffected siblings, and never-psychotic controls. Group differences in EPP were unrelated to age. Across groups, EPP was worse in older compared with younger individuals. In contrast, in patients ToM was better in older than in younger individuals, while ToM was not significantly related to age in never-psychotic controls or siblings. In patients, associations between EPP and ToM and most sociodemographic and clinical factors of interest did not vary with age. Interestingly, however, the association between negative symptoms and ToM appeared somewhat stronger for younger than for older patients. Finally, both ToM and EPP were positively associated with interpersonal functioning, but not with prosocial activities across all ages. Findings from this study add to the literature in several important ways.

First, our results suggest that EPP impairment in patients diagnosed with a non-affective psychotic disorder is stable across ages. Regardless of group, EPP was worse in older individuals, with group differences between patients, unaffected siblings, and never-psychotic controls that were of similar magnitude across the adult lifespan. The observed age-related worsening of EPP is in congruence with studies of normative aging that show declining EPP from mid adulthood²⁸, and previous work in samples with schizophrenia.¹⁷ Several, non-exclusive, theories have been proposed to explain this decline. From an evolutionary perspective it might be more important for younger people to accurately recognize others' emotions to enable learning from others, to promote partner selection, or to avoid risk in social situations.²⁹ This 'survival' theory is also supported by neuroimaging research, which showed that viewing negative emotional expressions is associated with less amygdala activation in older than younger adults.^{30, 31} Other work suggests that age-related changes in EPP might be due to changes in the visual scan paths of faces; changes that may be related to normative age-related changes in executive functioning.³² We also cannot rule out that task characteristics contributed to the age-related differences in EPP, as suggested previously³³, for example, bias may arise due to participants being presented with mostly young faces in the DFAR. However, studies that test whether own-age faces are recognized more accurately than other-age faces are limited in number and report inconsistent results.³⁴

Second, our results suggest that differences in ToM between patients, unaffected siblings and never-psychotic controls are smaller in individuals in their early to mid-50's, in line with previous work.³⁵ Age-related improvements on the Hinting Task in patients with schizophrenia, but not controls have previously been reported.¹⁷ There are various potential explanations. First, the smaller difference may suggest that patients 'regain some function' during the later phase of the illness when they might have adjusted to

1
2
3 some of the negative consequences, such as negative symptoms. The weaker association between Hinting
4 Task performance and negative symptoms in older individuals supports this idea. In addition, initial
5 ceiling effects in controls and siblings may have prevented us from noticing any further age-related
6 improvement. Ceiling effects on the Hinting Task have been reported previously.³⁶ Studies making use of
7 different ToM tasks could shed light on this. However, possibly patients may be catching up with controls
8 in ToM performance as they age because of gained experience and/or practice with social situations.
9 Alternatively, around the age of 50 controls and siblings (to a lesser degree) may start to show decline
10 that already occurred earlier in patients. Supportive of this hypothesis are studies which show age-related
11 declines in ToM in healthy individuals that are associated with declines in other cognitive function, such
12 as attention and working memory.^{28,37} It is also possible that the larger deficits in our younger patient
13 group are due to their earlier onset of the disorder. An early illness onset may disrupt the development of
14 ToM and/or may be related to a distinct neural pathology where disease mechanisms interact with neural
15 development.³⁸ As recently suggested by Armando, Hutsebaut, Debbané³⁹ the arrested development in the
16 specialization of social cognition during adolescence and early adulthood may account for residual social
17 functioning impairment in psychosis. Thus, individuals with an earlier illness onset may follow different
18 cognitive and social functioning trajectories over time. While meta-analyses suggest that age of onset
19 might not relate to the magnitude of ToM difficulties, whereas illness duration does⁴⁰, longitudinal
20 studies including large social cognitive test batteries are needed to formally investigate this. If our
21 findings were corroborated in such studies, this would suggest that young adulthood may offer a very
22 good window of opportunity for social cognitive interventions that can promote the development of social
23 cognitive skills⁴¹ and that may positively impact on patients social functioning further down the line.^{3,42}
24 Vice versa, a potential beneficial downstream effect on social functioning and the additional early
25 application of interventions that address social skills and deficits may foster ToM across the life span.

26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41 Finally, associations between EPP and ToM and most sociodemographic and clinical factors of interest
42 did not vary with age and ToM and EPP significantly predicted interpersonal functioning, but not
43 prosocial activities across all ages. These results tentatively suggest that interventions that aim to improve
44 or retain patients interpersonal functioning are useful across the entire adult lifespan.

45
46
47
48
49 Our results need to be interpreted considering several limitations. First, we cannot rule out age-associated
50 effects in EPP or ToM in early life or late adulthood since our youngest and oldest participants were 18
51 and 55, respectively. For instance, non-social cognition and functional outcome psychosis studies, a
52 second period of decline beyond the age of 65 has been suggested.^{43,44} Future research is needed to
53 further explore whether cognitive processes show different patterns in individuals with a psychotic
54
55
56
57
58
59
60

1
2
3 disorder from healthy controls after the age of 55.

4
5 Second, individuals with better EPP/ToM may have been more likely to participate in this study,
6 particularly in certain age groups. For example, the older age group, which was substantially smaller may
7 have presented individuals with overall better functioning in various domains. In a similar vein, it is
8 possible that young patients with more severe EPP/ToM impairments may have been overrepresented in
9 the study because they were more likely to be pushed to participate by family members or staff, in
10 comparison to older patients.
11
12
13

14 Third, our findings regarding the two social cognitive domains require replication using larger
15 test batteries that cover each domain in greater detail and that tap into other domains as well. Future
16 studies should consider the reliability of tests in older age groups. For example, DFAR faces are all young
17 to mid adulthood and it may be that participants are particularly skilled in recognizing emotions of similar
18 aged peers. Relatedly, and regarding social cognition, ceiling effects, cultural differences and cultural
19 sensitivity of the stimulus and test materials should be considered.
20
21
22

23
24 Fourth, it is important to note that some individuals in the control and sibling group had
25 diagnoses that might have impacted on their social cognitive functioning. As such the current findings
26 represent a conservative estimate of differences between the included participant groups.
27
28

29 Finally, any age-related differences in EPP and ToM need to be investigated longitudinally; the
30 cross-sectional nature of our study cannot speak to life-time trajectories of ToM and EPP within person
31 and may be subject to cohort effects.
32
33

34
35 To conclude, our findings suggest that, at least without intervention, EPP and ToM impairments are
36 present in individuals with a non-affective psychosis across adulthood. Our results show that EPP and
37 ToM impairment is associated with poorer interpersonal functioning both in younger and older patient
38 groups but suggest that younger individuals might benefit particularly from support with ToM skills,
39 whereas older individuals might be more likely to benefit from support with EPP skills. Longitudinal
40 research is needed to gain much needed insights into age-related trajectories of social cognition.
41
42
43
44
45
46
47
48

49 **Acknowledgements:**

50 We are grateful for the generosity of time and effort by the patients, their families and healthy subjects.
51 Furthermore, we would like to thank all research personnel involved in the GROUP project, in particular:
52 Joyce van Baaren, Erwin Veermans, Ger Driessen, Truda Driessen, Erna van 't Hag.
53
54
55
56
57
58
59
60

Funding:

The infrastructure for the GROUP study was funded through the Geestkracht programme of the Dutch Health Research Council (Zon-Mw, grant number 10-000-1001), and matching funds from participating pharmaceutical companies (Lundbeck, AstraZeneca, Eli Lilly, Janssen Cilag) and universities and mental health care organizations (Amsterdam: Academic Psychiatric Centre of the Academic Medical Center and the mental health institutions: GGZ Ingeest, Arkin, Dijk en Duin, GGZ Rivierduinen, Erasmus Medical Centre, GGZ Noord Holland Noord. Groningen: University Medical Center Groningen and the mental health institutions: Lentis, GGZ Friesland, GGZ Drenthe, Dimence, Mediant, GGNet Warnsveld, Yulius Dordrecht and Parnassia psycho-medical center The Hague. Maastricht: Maastricht University Medical Centre and the mental health institutions: GGzE, GGZ Breburg, GGZ Oost-Brabant, Vincent van Gogh voor Geestelijke Gezondheid, Mondriaan, Virenze riagg, Zuyderland GGZ, MET ggz, Universitair Centrum Sint-Jozef Kortenberg, CAPRI University of Antwerp, PC Ziekeren Sint-Truiden, PZ Sancta Maria Sint-Truiden, GGZ Overpelt, OPZ Rekem. Utrecht: University Medical Center Utrecht and the mental health institutions Altrecht, GGZ Centraal and Delta). EV is supported by the Seaver Foundation.

References:

1. Vaskinn A, Horan WP. Social Cognition and Schizophrenia: Unresolved Issues and New Challenges in a Maturing Field of Research. *Schizophr Bull* Apr 10 2020;46(3):464-470.
2. Green MF, Bearden CE, Cannon TD, et al. Social cognition in schizophrenia, Part 1: performance across phase of illness. *Schizophr Bull* Jun 2012;38(4):854-864.
3. Fett AK, Viechtbauer W, Dominguez MD, Penn DL, van Os J, Krabbendam L. The relationship between neurocognition and social cognition with functional outcomes in schizophrenia: a meta-analysis. *Neurosci Biobehav Rev* Jan 2011;35(3):573-588.
4. Lee TY, Hong SB, Shin NY, Kwon JS. Social cognitive functioning in prodromal psychosis: A meta-analysis. *Schizophr Res* May 2015;164(1-3):28-34.
5. van Donkersgoed RJ, Wunderink L, Nieboer R, Aleman A, Pijnenborg GH. Social Cognition in Individuals at Ultra-High Risk for Psychosis: A Meta-Analysis. *PLoS One* 2015;10(10):e0141075.
6. Horan WP, Green MF, DeGroot M, et al. Social cognition in schizophrenia, Part 2: 12-month stability and prediction of functional outcome in first-episode patients. *Schizophr Bull* Jun 2012;38(4):865-872.
7. McCleery A, Lee J, Fiske AP, et al. Longitudinal stability of social cognition in schizophrenia: A 5-year follow-up of social perception and emotion processing. *Schizophr Res* Oct 2016;176(2-3):467-472.
8. Lysaker PH, Olesek KL, Warman DM, Martin JM, Salzman AK, Nicolo G, Salvatore G, Dimaggio G. Metacognition in schizophrenia: correlates and stability of deficits in theory of mind and self-reflectivity. *Psychiatry Res* Nov 30 2011;190(1):18-22.
9. Hamm JA, Renard SB, Fogley RL, Leonhardt BL, Dimaggio G, Buck KD, Lysaker PH. Metacognition and social cognition in schizophrenia: stability and relationship to concurrent and prospective symptom assessments. *J Clin Psychol* Dec 2012;68(12):1303-1312.
10. Vohs JL, Lysaker PH, Francis MM, et al. Metacognition, social cognition, and symptoms in patients with first episode and prolonged psychoses. *Schizophr Res* Mar 2014;153(1-3):54-59.
11. Sprong M, Schothorst P, Vos E, Hox J, van Engeland H. Theory of mind in schizophrenia: meta-analysis. *Br J Psychiatry* Jul 2007;191:5-13.
12. Moran JM, Jolly E, Mitchell JP. Social-cognitive deficits in normal aging. *J Neurosci* Apr 18 2012;32(16):5553-5561.
13. Mucci A, Galderisi S, Green MF, et al. Familial aggregation of MATRICS Consensus Cognitive Battery scores in a large sample of outpatients with schizophrenia and their unaffected relatives. *Psychol Med* Jun 2018;48(8):1359-1366.
14. Lavoie MA, Plana I, Bedard Lacroix J, Godmaire-Duhaime F, Jackson PL, Achim AM. Social cognition in first-degree relatives of people with schizophrenia: a meta-analysis. *Psychiatry Res* Sep 30 2013;209(2):129-135.
15. Balogh N, Egerhazi A, Berecz R, Csukly G. Investigating the state-like and trait-like characters of social cognition in schizophrenia: a short term follow-up study. *Schizophr Res* Nov 2014;159(2-3):499-505.
16. Ferrer-Quintero M, Green MF, Horan WP, Penn DL, Kern RS, Lee J. The effect of sex on social cognition and functioning in schizophrenia. *NPJ Schizophr* Dec 1 2021;7(1):57.

17. Pinkham AE, Kelsven S, Kouros C, Harvey PD, Penn DL. The Effect of Age, Race, and Sex on Social Cognitive Performance in Individuals With Schizophrenia. *J Nerv Ment Dis* May 2017;205(5):346-352.
18. Navarra-Ventura G, Vicent-Gil M, Serra-Blasco M, et al. Group and sex differences in social cognition in bipolar disorder, schizophrenia/schizoaffective disorder and healthy people. *Compr Psychiatry* Aug 2021;109:152258.
19. Korver N, Quee PJ, Boos HB, Simons CJ, de Haan L, investigators G. Genetic Risk and Outcome of Psychosis (GROUP), a multi-site longitudinal cohort study focused on gene-environment interaction: objectives, sample characteristics, recruitment and assessment methods. *Int J Methods Psychiatr Res* Sep 2012;21(3):205-221.
20. van't Wout M, Aleman A, Kessels RP, Larøi F, Kahn RS. Emotional processing in a non-clinical psychosis-prone sample. *Schizophrenia research* 2004;68(2-3):271-281.
21. Corcoran R, Mercer G, Frith CD. Schizophrenia, symptomatology and social inference: investigating “theory of mind” in people with schizophrenia. *Schizophrenia research* 1995;17(1):5-13.
22. Ludwig KA, Pinkham AE, Harvey PD, Kelsven S, Penn DL. Social cognition psychometric evaluation (SCOPE) in people with early psychosis: A preliminary study. *Schizophrenia research* 2017;190:136-143.
23. Fett A-KJ, Maat A, Investigators G. Social cognitive impairments and psychotic symptoms: what is the nature of their association? *Schizophrenia bulletin* 2013;39(1):77-85.
24. Benton A, Van Allen M, Hamsher Kd, Levin H. Test of facial recognition. *Manual Neurosurgery Center Publication* 1973(287).
25. Wechsler D. Wechsler adult intelligence scale. *Archives of Clinical Neuropsychology* 1955.
26. Kay SR, Fiszbein A, Opler LA. The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophr Bull* 1987;13(2):261-276.
27. Birchwood M, Smith J, Cochrane R, Wetton S, Copestake S. The Social Functioning Scale. The development and validation of a new scale of social adjustment for use in family intervention programmes with schizophrenic patients. *Br J Psychiatry* Dec 1990;157:853-859.
28. Kemp J, Després O, Sellal F, Dufour A. Theory of Mind in normal ageing and neurodegenerative pathologies. *Ageing Research Reviews* 2012/04/01/ 2012;11(2):199-219.
29. Frith CD, Frith U. Mechanisms of Social Cognition. *Annual Review of Psychology* 2012;63(1):287-313.
30. Gunning-Dixon FM, Gur RC, Perkins AC, et al. Age-related differences in brain activation during emotional face processing. *Neurobiology of Aging* 2003/03/01/ 2003;24(2):285-295.
31. Fischer H, Nyberg L, Bäckman L. Age-related differences in brain regions supporting successful encoding of emotional faces. *Cortex* 2010/04/01/ 2010;46(4):490-497.
32. Circelli KS, Clark US, Cronin-Golomb A. Visual scanning patterns and executive function in relation to facial emotion recognition in aging. *Neuropsychol Dev Cogn B Aging Neuropsychol Cogn* 2013;20(2):148-173.
33. Sze JA, Goodkind MS, Gyurak A, Levenson RW. Aging and emotion recognition: not just a losing matter. *Psychol Aging* 2012;27(4):940-950.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
34. Short LA, Mondloch CJ. Aging faces and aging perceivers: young and older adults are less sensitive to deviations from normality in older than in young adult faces. *Perception* 2013;42(8):795-812.
 35. Smeets-Janssen MMJ, Meesters PD, Comijs HC, Eikelenboom P, Smit JH, de Haan L, Beekman ATF, Stek ML. Theory of Mind differences in older patients with early-onset and late-onset paranoid schizophrenia. *International Journal of Geriatric Psychiatry* 2013;28(11):1141-1146.
 36. Klein HS, Springfield CR, Bass E, Ludwig K, Penn DL, Harvey PD, Pinkham AE. Measuring mentalizing: A comparison of scoring methods for the hinting task. *International Journal of Methods in Psychiatric Research* 2020;29(2):e1827.
 37. Johansson Nolaker E, Murray K, Happé F, Charlton RA. Cognitive and affective associations with an ecologically valid test of theory of mind across the lifespan. *Neuropsychology* 2018;32(6):754.
 38. Burke L, Androustos C, Jogia J, Byrne P, Frangou S. The Maudsley Early Onset Schizophrenia Study: The effect of age of onset and illness duration on fronto-parietal gray matter. *European Psychiatry* 2008/06/01/ 2008;23(4):233-236.
 39. Armando M, Hutsebaut J, Debbané M. A mentalization-informed staging approach to clinical high risk for psychosis. *Frontiers in psychiatry* 2019;10:385.
 40. Bora E, Yucel M, Pantelis C. Theory of mind impairment in schizophrenia: meta-analysis. *Schizophrenia research* 2009;109(1-3):1-9.
 41. Yamada Y, Inagawa T, Sueyoshi K, et al. Social cognition deficits as a target of early intervention for psychoses: a systematic review. *Frontiers in psychiatry* 2019;10:333.
 42. Halverson, F. T, Orleans-Pobee M, Merritt C, Sheeran P, Fett AK, Penn DL. Pathways to functional outcomes in schizophrenia spectrum disorders: Meta-analysis of social cognitive and neurocognitive predictors. *Neurosci Biobehav Rev* Oct 2019;105:212-219.
 43. Friedman JI, Harvey PD, Coleman T, et al. Six-year follow-up study of cognitive and functional status across the lifespan in schizophrenia: a comparison with Alzheimer's disease and normal aging. *Am J Psychiatry* Sep 2001;158(9):1441-1448.
 44. Moran JM. Lifespan development: The effects of typical aging on theory of mind. *Behavioural Brain Research* 2013/01/15/ 2013;237:32-40.

Table 1 displays the sociodemographic and clinical characteristics of the GROUP study sample.

| Table 1. Study sample characteristics | | | | |
|---|-----------------------------|-----------------------------|-----------------------------|--------------------------------|
| | Patients^P | Siblings^S | Controls^C | Statistics^{**} |
| | (n = 905) | (n = 966) | (n = 544) | |
| Female (n, (%)) | 198 (21.88) | 526 (54.45) | 301 (55.33) | P<S,C |
| Age (mean, (sd)) | 27.74 (7.47) | 28.61 (7.77) | 31.26 (10.21) | P<S<C |
| IQ (mean, (sd)) | 94.74 (16.26) | 102.88 (15.60) | 110.11 (15.14) | P<S<C |
| White (n, (%)) | 673 (78.81) | 780 (83.60) | 492 (92.83) | P<S<C |
| Antipsychotics yes (n, %) | 874 (99.5) | | | |
| Benton face recognition (mean, (sd)) | 22.75 (2.32) | 23.22 (2.15) | 23.19 (2.02) | P<S,C |
| Social Functioning (mean, (sd))* | 112.43 (9.42) | 122.20 (6.70) | 123.94 (5.63) | P<S<C |
| Interpersonal Score | 124.47 (19.53) | 138.48 (12.69) | 140.74 (10.38) | P<S<C |
| Prosocial activities | 113.1 (13.88) | 120.95 (11.58) | 123.39 (10.12) | P<S<C |
| DFAR (mean, (sd)) | 68.30 (10.75) | 72.47 (9.27) | 73.35 (9.18) | P<S,C |
| DFAR neutral | 77.92 (17.29) | 81.71 (14.81) | 82.41 (14.42) | P<S,C |
| DFAR happy | 86.85 (12.69) | 88.11 (10.63) | 87.45 (11.02) | P<S |
| DFAR fearful | 48.27 (20.24) | 53.04 (19.20) | 54.78 (18.38) | P<S,C |
| DFAR angry | 63.95 (21.02) | 70.34 (19.27) | 72.05 (18.48) | P<S,C |
| Hinting Task (mean, (sd)) | 17.54 (2.78) | 18.87 (1.62) | 19.13 (1.26) | P<S<C |
| PANSS positive (weight. mean, (sd)) | 1.81 (.76) | | | |
| PANSS negative (weight. mean, (sd)) | 2.03 (.87) | | | |
| PANSS general (weight. mean, (sd)) | 1.75 (.52) | | | |
| Age of 1 st psychosis (mean, (sd)) | 23.31 (6.95) | | | |
| Illness duration in yrs. (mean, (sd)) | 4.62 (4.50) | | | |

* measured at three-year follow-up, ** differences significant at 0.05. DFAR = Degraded Facial Affect Recognition; IQ = Intelligence Quotient; PANSS = Positive and Negative Syndrome Scale




Figure 1. Age-related performance on the Degraded Facial Affect Recognition (DFAR) task across groups, adjusting for sex, family ID (in case multiple individuals from one family participated in the study) and study site.




Figure 2. Age-related performance on the Hinting Task across groups, adjusting for sex, family ID (in case multiple individuals from one family participated in the study) and study site.

Supplementary Table 1. Age distribution by age categorizations

| | Patients^a n (%), mean (SD) | Siblings^b n (%), mean (SD) | Controls^c n (%), mean (SD) | Statistics |
|-----------|---|---|---|-------------------|
| Age 18-25 | 434 (47.96), 21.89 (2.17) | 408 (42.24), 21.78 (2.28) | 213 (39.15), 20.93 (2.34) | C≠P, S |
| Age 26-30 | 206 (22.76), 27.70 (1.38) | 233 (24.12), 28.03 (1.37) | 72 (13.24), 27.76 (1.51) | |
| Age 31-40 | 197 (21.77), 34.57 (2.72) | 238 (24.64), 34.76 (2.70) | 120 (22.06), 35.45 (2.91) | |
| Age 41-55 | 68 (7.51), 45.47 (4.20) | 87 (9.01), 45.40 (3.68) | 139 (25.55), 45.27 (3.08) | |

Supplementary Table 2. Z-distribution of social cognition scores by group

| Z score | Patients^a | | Siblings^b | | Controls^c | |
|---------------------|-----------------------------|--------------------|-----------------------------|--------------------|-----------------------------|--------------------|
| | DFAR n (%) | Hints n (%) | DFAR n (%) | Hints n (%) | DFAR n (%) | Hints n (%) |
| Z < -3 | 15 (1.8) | 41 (4.7) | 2 (0.2) | 5 (0.5) | 1 (0.2) | |
| Z = -3 to -2 | 34 (4.1) | 39 (4.5) | 7 (0.8) | 7 (0.7) | 6 (1.2) | 3 (0.6) |
| Z = -2 to -1 | 123 (14.7) | 168 (19.3) | 81 (9.0) | 59 (6.2) | 35 (6.8) | 18 (3.4) |
| Z = -1 to 0 | 331 (39.7) | 237 (27.3) | 291 (32.4) | 224 (23.6) | 151 (29.5) | 111 (10.8) |
| Z = 0 to 1 | 244 (29.3) | 384 (44.2) | 343 (38.2) | 654 (68.9) | 199 (38.9) | 403 (75.3) |
| Z = 1 to 2 | 84 (10.1) | ‡ | 160 (17.8) | ‡ | 113 (22.1) | ‡ |
| Z > 2 | 3 (0.4) | ‡ | 15 (1.7) | ‡ | 6 (1.2) | ‡ |

Supplementary Table 3. Z-distribution of social cognition scores by age group

| Z score | Age 18-25 | | Age 26-30 | | Age 31-40 | | Age 41-55 | |
|--------------|------------|-------------|------------|-------------|------------|-------------|------------|-------------|
| | DFAR n (%) | Hints n (%) | DFAR n (%) | Hints n (%) | DFAR n (%) | Hints n (%) | DFAR n (%) | Hints n (%) |
| Z < -3 | 9 (0.9) | 28 (2.7) | 4 (0.8) | 12 (2.4) | 1 (0.2) | 4 (0.7) | 4 (1.5) | 2 (0.7) |
| Z = -2 to -3 | 18 (1.8) | 23 (2.2) | 11 (2.3) | 10 (2.0) | 9 (1.8) | 13 (2.4) | 9 (3.3) | 3 (1.0) |
| Z = -2 to -1 | 80 (8.1) | 133 (13.0) | 58 (12.3) | 49 (9.9) | 68 (13.2) | 42 (7.8) | 33 (12.0) | 21 (7.2) |
| Z = -1 to 0 | 336 (34.2) | 259 (25.2) | 147 (31.1) | 121 (24.3) | 214 (41.6) | 131 (24.3) | 114 (41.5) | 61 (21.0) |
| Z = 0 to 1 | 346 (35.2) | 583 (56.8) | 183 (38.8) | 305 (61.4) | 181 (35.2) | 350 (64.8) | 76 (27.6) | 203 (70.0) |
| Z = 1 to 2 | 180 (18.3) | ‡ | 66 (14.0) | ‡ | 72 (14.0) | - | 39 (14.2) | ‡ |
| Z > 2 | 14 (1.4) | ‡ | 3 (0.6) | ‡ | 7 (1.4) | - | ‡ | ‡ |


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38

| Supplementary Table 4. Correlations between study measures, unadjusted and adjusted for age | | | | | | | | | | | | | | | | | | | | |
|--|----------------|--------------|----------------|--------------|-------------------|--------------|----------------|--------------|----------------|--------------|----------------|--------------|----------------|--------------|------------------|--------------|----------------|--------------|----------------|--------------|
| | DFAR total | | Hinting task | | SFS interpersonal | | SFS prosocial | | Ethnicity | | Sex | | IQ | | Illness duration | | PANSS pos | | Panss neg | |
| | Unadj. for age | Adj. for age | Unadj. for age | Adj. for age | Unadj. for age | Adj. for age | Unadj. for age | Adj. for age | Unadj. for age | Adj. for age | Unadj. for age | Adj. for age | Unadj. for age | Adj. for age | Unadj. for age | Adj. for age | Unadj. for age | Adj. for age | Unadj. for age | Adj. for age |
| Hinting task | 0.23 | 0.24 | | | | | | | | | | | | | | | | | | |
| SFS interpers. | 0.18 | 0.19 | 0.24 | 0.24 | | | | | | | | | | | | | | | | |
| SFS pros | 0.14 | 0.13 | 0.13 | 0.13 | 0.50 | 0.49 | | | | | | | | | | | | | | |
| Ethnicity | -0.01 | -0.02 | -0.05 | -0.04 | -0.09 | -0.08 | -0.02 | -0.02 | | | | | | | | | | | | |
| Sex | 0.16 | 0.18 | 0.19 | 0.18 | 0.22 | 0.21 | 0.11 | 0.12 | -0.03 | -0.02 | | | | | | | | | | |
| IQ | 0.20 | 0.21 | 0.35 | 0.35 | 0.27 | 0.27 | 0.16 | 0.17 | -0.12 | -0.12 | 0.006 | -0.003 | | | | | | | | |
| Patient group only: | | | | | | | | | | | | | | | | | | | | |
| Illness duration | -0.009 | 0.03 | 0.03 | -0.02 | -0.06 | -0.05 | -0.10 | -0.06 | 0.02 | 0.04 | 0.05 | -0.04 | 0.05 | 0.04 | | | | | | |
| Age of 1 st psychosis | -0.07 | -0.01 | 0.11 | 0.03 | -0.08 | -0.06 | -0.06 | 0.04 | -0.04 | -0.04 | 0.20 | 0.09 | 0.006 | -0.04 | -0.17 | - | | | | |
| PANSS positive | -0.14 | -0.15 | -0.25 | -0.24 | -0.32 | -0.32 | -0.24 | -0.25 | 0.07 | 0.07 | -0.22 | -0.21 | -0.25 | -0.24 | 0.09 | 0.13 | | | | |
| PANSS negative | -0.19 | -0.20 | -0.38 | -0.37 | -0.42 | -0.42 | -0.34 | -0.34 | 0.09 | 0.09 | -0.27 | -0.25 | -0.37 | -0.36 | -0.07 | -0.03 | 0.47 | 0.47 | | |
| PANSS general | -0.15 | -0.16 | -0.30 | -0.29 | -0.39 | -0.39 | -0.30 | -0.31 | 0.06 | 0.06 | -0.25 | -0.23 | -0.31 | -0.30 | -0.002 | 0.06 | 0.74 | 0.74 | 0.70 | 0.69 |


39 DFAR = Degraded Facial Affect Recognition, SFS = social functioning scale (measured at three-year follow up), PANSS = Positive and Negative Syndrome Scale

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Supplementary Figure 1. Age-related performance on the degraded facial affect recognition (DFAR) task by emotion type across groups, adjusting for sex and accounting for family ID and recruitment centre.



Supplementary Figure 2. Socio-demographic and clinical correlates of performance on the DFAR and Hinting Task in the patient group.



DFAR = degraded facial affect recognition, PANSS_pos = Positive and Negative Syndrome Scale, positive symptoms subscale, PANSS_neg = PANSS negative symptoms subscale, PANSS_gen = general symptoms subscale SFS = Social Functioning Scale