



City Research Online

City, University of London Institutional Repository

Citation: Kam, J. H., Shinhmar, H., Powner, M. B., Hayes, M. J., Aboelnour, A. & Jeffery, G. (2023). Mitochondrial decline in the ageing old world primate retina: Little evidence for difference between the centre and periphery. PLoS One, 18(5), e0273882. doi: 10.1371/journal.pone.0273882

This is the published version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/30592/>


Link to published version: <https://doi.org/10.1371/journal.pone.0273882>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

RESEARCH ARTICLE

Mitochondrial decline in the ageing old world primate retina: Little evidence for difference between the centre and periphery

Jaimie Hoh Kam¹, Harpreet Shinhmar¹, Michael Barry Powner², Matthew John Hayes¹, Asmaa Aboelnour³, Glen Jeffery^{1*}¹ Department of Visual Science, Institute of Ophthalmology, University College London, London, United Kingdom, ² Centre for Applied Vision Research, City University, London, United Kingdom, ³ Department of Histology, Damanhour University, Damanhour, Egypt* g.jeffery@ucl.ac.uk

OPEN ACCESS

Citation: Kam JH, Shinhmar H, Powner MB, Hayes MJ, Aboelnour A, Jeffery G (2023) Mitochondrial decline in the ageing old world primate retina: Little evidence for difference between the centre and periphery. PLoS ONE 18(5): e0273882. <https://doi.org/10.1371/journal.pone.0273882>

Editor: Michele Madigan, Save Sight Institute, AUSTRALIA

Received: August 15, 2022

Accepted: April 10, 2023

Published: May 2, 2023

Copyright: © 2023 Kam et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Data are all within the paper.

Funding: GJ and MBP. BB/N000250/1 Biotechnology and Biological Sciences Research Council. <https://www.ukri.org/councils/bbsrc/> The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Abstract

Mitochondrial decline is a key feature of ageing. The retina has more mitochondria than any other tissue and ages rapidly. To understand human retinal ageing it is critical to examine old world primates that have similar visual systems to humans, and do so across central and peripheral regions, as there is evidence for early central decline. Hence, we examine mitochondrial metrics in young and ageing *Macaca fascicularis* retinae. In spite of reduced ATP with age, primate mitochondrial complex activity did not decline. But mitochondrial membrane potentials were reduced significantly, and concomitantly, mitochondrial membrane permeability increased. The mitochondrial marker Tom20 declined significantly, consistent with reduced mitochondria number, while VDAC, a voltage dependent anion channel and diffusion pore associated with apoptosis increased significantly. In spite of these clear age-related changes, there was almost no evidence for regional differences between the centre and the periphery in these mitochondrial metrics. Primate cones do not die with age, but many showed marked structural decline with vacuous spaces in proximal inner segments normally occupied by endoplasmic reticulum (ER), that regulate mitochondrial autophagy. In many peripheral cones, ER was displaced by the nucleus that transposed across the outer limiting membrane and could become embedded in mitochondrial populations. These data are consistent with significant changes in retinal mitochondria in old world primate ageing but provide little if any evidence that aged central mitochondria suffer more than those in the periphery.

Introduction

Mitochondria regulate metabolism and ageing [1, 2]. The significance of these features come together in the outer retina that has the highest metabolic rate in the body, the greatest concentration of mitochondria in any tissue and also a very rapid rate of ageing [3, 4]. Further, as vision is the primary sensory modality for primates, mitochondrial function in the retina

underpins our lives. However, the rapid pace of retinal ageing is problematic in relation to progressive extensions to human lifespan. For much of our evolution, lifespan was probably under 40 years, but is now more than double that in the Western world and parts of Asia. This extension exposes the vulnerability of the retina [5, 6].

By the age of approximately 70 years, around 30% of human central rod photoreceptors have died, and cones that do not show age related loss have significantly reduced ability [7, 8]. Mitochondrial function in the ageing primate retina declines significantly with over a 70% reduction in adenosine triphosphate (ATP) production, which is a key energy source needed for metabolic function. This is in spite of a significant increase in glycolysis [9]. These factors likely have an impact on aged human vision as improving aged mitochondrial function is associated with improved colour contrast [10] and scotopic thresholds [11].

Studies that examine patterns of retinal ageing have commonly used rodents. However, vision is not their primary sensory modality, and the animals prefer darkness. They have not evolved a high degree of central visual acuity mediated by a region of high cell density that in humans can be a focus of aged disease. But importantly, murine patterns of outer retinal cell loss are profoundly different from those in primates. In mice, unlike primates, cone photoreceptor loss is not only an ageing feature, but is initiated before rod loss [12], and rodent age related patterns of cytokine expression are completely different from that found in primates in both the retina and serum [9].

For these reasons it is important that we examine retinal ageing in the closest living relative to humans, old world primates, and that this should have a focus on mitochondria that are critical in the ageing process [1] and take account of central and peripheral region, as there is evidence for preferential and progressive accumulation of mitochondrial DNA (mtDNA) mutations in the central human retina compared to the periphery [13]. Here we examine mitochondrial decline in the old world primate (*Macaca fascicularis*), describing key features and also asking if similar processes are found in the central retina, which mediates the majority of our visual function as in the periphery. We also describe key aged changes in cone photoreceptors that do not die with age and on which, because of artificial lighting, we have become increasingly dependent.

Materials and methods

Animals

Primate ocular tissues were acquired from a large, long established colony of *Macaca fascicularis* maintained by UK Public Health laboratory regulated under local and U.K. Home Office regulation. No experiments were undertaken on these primates as the main concern was to reduce colony size. All the animals were healthy and were sacrificed by Public Health England. The primary purpose of animal usage was different from the aims of this study and eyes were only retrieved for this study after death. The primates are of Mauritian origin.

Macaca fascicularis have a highly diverse geographic distribution extending from Vietnam and Myanmar down to Java and Timor. Local populations can show extreme genetic, physiological and behavioural variations. This is particularly true comparing mainland with island populations and extends to mtDNA sequencing [14, 15].

Mauritian macaques have been established as an isolated colony for approximately 400 years and have distinctive mtDNA sequencing with nucleotide variations in mtDNA 10 fold lower than in the Indonesian population, consistent with their isolation. As a distant island population, Mauritian animals show little group variation compared to their distant cousins who can mate with rhesus macaques. They likely originated from Indonesia but show morphological and genomic distinction [14–16]. While there are examples to the contrary, it is

strongly argued that ageing and lifespan is partly driven by genetic diversity in a population, which Mauritian animals lack relative to their Indonesian ancestors [17, 18]. For these reasons Mauritian animals were used in this study.

The normal maximum lifespan of Mauritian animals in this colony is around 18 years compared to approximately up to a maximum of 30 years in a similar wild Indonesian population. However, lifespan in the wild varies considerably across subspecies and in relation to habitat. However, this makes the Mauritian animals used here an ideal model for old world primate ageing.

Each of the animals used was bred in the unit and had a full life long medical history. None had any history of disease. They were each subject to monthly health checks by a designated veterinary officer. Every 3 months this included sedation and blood samples that were run for standard clinical markers of disease and organ function. Eyes of all animals were normal without sign of disease or drusen deposition. Old animals had normal noticeable browning of the lens.

Eye retrieval followed termination after sedation with ketamine (200 mg/kg) and overdose of intravenous sodium pentobarbital (50 mg/kg). The eyes were removed rapidly. One eye from each animal was rapidly dissected on ice. Following removal of the anterior eye and lens, four radial cuts were made to open the eye cup. Regions of tissue spanning from the retina down to the choroid were separated from different regions of the eye cup. Sample sizes varied from approximately 2mm x 2mm in central regions to approximately 3mm x 3mm in other regions. These were gently dissected with fine forceps and micro scissors on a chilled Petri dish and separated into retina, retinal pigment epithelial (RPE) and choroid. These were from the macular and defined region of the periphery. Samples were taken from the central macular just beyond the fovea by approximately 2mm. Those from the periphery were outside the macular area with tissue pooled from the nasal, ventral and dorsal regions approximately 4mm distant from the optic nerve head. These samples were snap frozen on dry ice and stored at -80°C for different assays. The fellow eyes were fixed in 4% paraformaldehyde in PBS for 24h before their transfer into PBS in 1% sodium azide. Samples were taken from 10 young animals (average age 4 years. SD \pm 4 months) and 10 older primates (average age 13 years and 10 months. SD \pm 4 months) in this study.

Mitochondrial extraction and isolation

Mitochondria from 5 old (all females) and 5 young adults (3 females and 2 males) from the central macular and peripheral regions were homogenised in pre-chilled mitochondrial storage buffer consisting of 230mM Mannitol, 70mM sucrose, 1mM EGTA, 10mM HEPES and protease inhibitor cocktail (Sigma-Aldrich, UK). The homogenate was centrifuged at 800 X g for 30 mins at 4°C and then the supernatant was centrifuged at 11,000 X g for 10 mins to precipitate the mitochondria. Mitochondrial pellet was resuspended in fresh mitochondrial storage buffer and protein concentration of each sample was measured using the BCA assay (ThermoFisher Scientific, UK). Here the right eye was used.

Mitochondrial membrane potentials

The membrane potential of isolated mitochondria from above was investigated using a coloured dye described by Figueira et al. [19] in which they follow changes in fluorescence of the membrane potential probe, Safranin O is a lipophilic dye commonly used to measure membrane potential with a spectrofluorometer. Safranin O undergoes an optical shift upon its potential-dependent distribution between the external medium and intramitochondrial compartment, and on its stacking to mitochondrial membrane ionic sites. There is a spectral shift

in safranine produced by the energy state of mitochondria with a linear relationship between its absorbance and mitochondrial membrane potential. 20 μ g of mitochondrial protein was added to an incubation medium containing 220mM Mannitol, 100mM Sucrose, 1mM EGTA, 4mM K₂PO₄, 0.2mg/ml BSA, 20mM HEPES, 10 μ M Safranin O and 15mM of D-glyceraldehyde-3-phosphate, pH 7.4 and the changes were read at excitation wavelength 495nm and emission wavelength 586nm for 1 minute.

Mitochondrial membrane permeability and light scattering

Isolated mitochondrial suspensions from above containing about 50 μ g protein/ml were subjected to swelling using a hypotonic solution to examine their membrane permeability. The hypotonic solution consisting of 90mM KCl, 20mM MOPS-KOH, pH 7.2 and the absorbance changes were measured spectrophotometrically at 540nm for 1 minute. The absorption rate (ΔA) and the initial rate (1 min) of mitochondrial swelling ($\Delta A/\text{min}$) were recorded. Light scattering measurements of the mitochondria were determined from absorption values prior to the addition of the hypotonic solution.

Immunohistochemistry

The right eye of 5 older (4 females and 1 male) and 5 young (4 female and 1 male) primates were fixed in 4% paraformaldehyde in PBS for approximately 24h. Eyes were then washed in 1XPBS, dissected and the anterior tissues removed. A horizontal strip containing the retina, RPE and choroid was removed running from the temporal periphery, through the macular and fovea into the nasal retina. These were cryoprotected in 30% sucrose and then embedded in OCT (Optimal Cutting Temperature Tissue-Tek compound, Leica). Frozen sections were cryosectioned at 10 μ m and processed for immunohistochemistry using the streptavidin horseradish peroxidase complex. The sections of the eye were incubated for 1 hour in a 5% Normal Donkey serum in 0.3% Triton X-100 in PBS, pH 7.4, followed by an overnight incubation with primary antibodies in the [Table 1](#) below which was made in 1% Normal Donkey Serum in 0.3% Triton X-100 in PBS. After the primary antibody incubation, the sections were washed three times in 0.1 M PBS and then treated with 0.3% hydrogen peroxide in PBS to quench endogenous peroxidase activity. After several washes, the tissues were incubated with the appropriate biotin-SP conjugated secondary antibody (Jackson ImmunoResearch Laboratories, 11000) which were made up in 2% Normal Donkey Serum in 0.3% Triton X-100 in PBS, were added to the sections and incubated for 1 hour at room temperature. Negative controls were undertaken omitting the primary antibody. After the secondary antibody incubation, sections were washed several times and incubated in a ready to use horseradish peroxidase Streptavidin solution (VECTOR Laboratories, UK) for 30 minutes, followed by a peroxidase substrate solution, 3,3-diaminobenzidine (DAB) for 1 minute. Slides were mounted in glycerol and coverslipped after several washes in PBS and TBS. Sections were viewed and images captured using an Epi-fluorescence bright-field microscope (Olympus BX50F4, Olympus, Japan), where data were captured as 24-bit colour images at 3840 \times 3072 pixel resolution using Nikon DXM1200 (Nikon, Tokyo, Japan).

Western blotting

Snap frozen retinal samples from the left eye 5 older (4 females and 1 male) and 5 young (4 female and 1 male) primates from both central macular and peripheral regions were homogenized in 2% sodium dodecyl sulfate (SDS, Sigma Aldrich, UK) with protease inhibitor cocktail (Sigma Aldrich, UK), and centrifuged at 13,000 X g. The supernatant was transferred to a new microcentrifuge tube and for Western blots of the proteins in [Table 1](#); translocase of outer

Table 1.

Primary antibodies used	Catalogue number	Concentration		Company
		Western blot	Immunohistochemistry	
Mouse monoclonal Tom20 (F-10)	SC-17764	0.667 µg/ml		Santa Cruz Biotechnology Inc.
Rabbit polyclonal to VDAC	NB100-695	0.05 µg/ml		Novus Biologicals
Mouse monoclonal to β -actin (2D4H5)	66009-1g	0.1 µg/ml		Proteintech
Mouse monoclonal to Succinate Dehydrogenase (Complex II)	Ab14715	2 µg/ml	5 µg/ml	Abcam
Goat polyclonal to cytochrome C oxidase subunit III (Complex III) (N-20)	Sc-23986	0.2 µg/ml	0.4 µg/ml	Santa Cruz Biotechnology Inc.
Mouse monoclonal to COXIV (Complex IV)	Ab33985	1 µg/ml	10 µg/ml	Abcam
Mouse monoclonal to α -tubulin	T8203	0.4 µg/ml		Sigma-Aldrich
Rabbit polyclonal to GAPDH	ABS16	0.245 µg/ml		Milipore

<https://doi.org/10.1371/journal.pone.0273882.t001>

membrane 20 (Tom20), a marker of mitochondria and the mitochondrial porins voltage-dependent anion channel (VDAC), proteins of mitochondrial function: complex II, III and IV. The protein concentration of each sample was measured using the BCA Protein Assay (ThermoFischer Scientific, UK). Equal amounts of protein (50 µg/ml) were separated by a 10% SDS gel electrophoresis and electrophoretically transferred onto nylon membranes. Then the nylon membranes were pre-treated with 5% non-fat dried milk in 1M phosphate buffered saline (PBS), pH7.4 for 1 h and incubated overnight at room temperature with the primary antibodies in the table below followed by several washes in 0.05% Tween-20 in 1M PBS. The membranes were then incubated with the respective secondary antibodies; goat anti-mouse HRP conjugated (1/5000, ThermoFisher Scientific UK), rabbit anti goat HRP conjugated HRP (1/5000, Dako UK) and goat anti-rabbit HRP conjugated (1/5000, Dako UK) for 1 h. Immunoreactivities were visualised by exposing x-ray films to blots incubated with enhanced chemiluminescence (ECL) reagent (SuperSignal West Dura, ThermoFisher Scientific UK). Total protein profile was determined by staining blots with Ponceau S solution to check the transfer efficiency and quantification. Protein bands were then photographed and scanned. After each protein detected, the nylon membranes were then stripped with a stripping buffer containing 6M guanidine hydrochloride, 0.2% Nonidet P-40, 0.1M β -mercaptoethanol and 20mM Tris-HCL and washed several times with the washing buffer made up of 0.14M NaCl, 10mM Tris-HCL and 0.05% Nonidet P-40 before being reprobed by another primary antibody of interest (Table 1). The absolute intensity of each band was measured using Adobe Photoshop CS5 extended.

Resin embedded histology

Primate eyes of 5 older (5 females) and 5 young (3 females and 2 males) were fixed in 4% paraformaldehyde in PBS, and then transferred to 0.1M PBS for dissection. Horizontal strip of the neural retina together with the RPE/choroidal tissues were dissected from the periphery to the macular and were processed for resin embedded plastic. They were post-fixed with 2% paraformaldehyde and 2% glutaraldehyde in PBS for 24h, followed by repeated PBS washing and dehydrated through a series of ethanol. The samples were infiltrated, polymerised and embedded in Technovit 7100 historesin (Taab Laboratories equipment, UK). Resin sections were cut at 2.5 µm using a microtome, histologically stained with Acid Fuchsin (mitochondrial stain) and mounted in Depex, cover slipped.

Results

Mitochondrial complex activities

Complex activities were examined by western blot and immunostaining for complexes II, III and IV. Patterns in immunostaining for the complexes were similar across retinal layers. As expected, the most prominent feature of the immunostaining was the heavy label in photoreceptor inner segments. Staining was also present in the inner retina but largely absent from the outer nuclear layer. There were no obvious differences with age or location in the extent of the label (Figs 1–3). In each case the retinal sections used come from similar locations with similar proportions of cones that can be identified in the higher power images for each complex

A similar analysis was undertaken with Western blot to quantify complex protein expression. Complexes II and III were consistently more abundant in the periphery than the centre in both young and older animals. In complex IV greater concentrations were found in the centre than the periphery in both young and older, but these differences were not statistically significant. These data are similar to those presented previously by us [9] where we measured the enzyme kinetic of complex IV activities, however previously the difference between young and older in the periphery did just reach statistical significance. The overall result for complex measurements was that there were no significant changes with age in spite of the 70% reduction in ATP documented in the same population of primates [9] (Figs 1–3).




Fig 1. A–D. Immunohistochemistry of succinate dehydrogenase (complex II) activities on young and old primate central (C) and peripheral (P) retina. Staining is present mostly in the ganglion cell layer (GCL), Inner plexiform layer (IPL) and strongly in inner segment of the photoreceptor layer (PR). E–H. Inner segments at higher magnification. Staining appears stronger in peripheral sections particularly in the older animals. I. Western blot analysis for complex II confirms the immune staining with higher levels of complex II found in the periphery particularly in older animals. However, there is no evidence of consistent age related changes in any retinal layer. N = 5 per group. Abbreviations: GCL, ganglion cell layer. IPL, inner plexiform layer. INL, inner nuclear layer. ILM, inner limiting membrane. OPL, outer plexiform layer. ONL, outer nuclear layer. ILM, outer limiting membrane. Photoreceptors PR. Young centre, Y (C). Young periphery Y(P). Old centre, O(C). Old periphery, O (P). Statistical significance, * $p < 0.05$. See supplementary for full uncropped images.

<https://doi.org/10.1371/journal.pone.0273882.g001>




Fig 2. A–D. Immunohistochemistry of cytochrome C oxidase subunit III in young and old primate central (C) and peripheral (P) retina. Positive staining is present in the ganglion cell layer (GCL), Inner plexiform layer (IPL), inner nuclear layer (INL) and inner segment of the photoreceptor layer (PR). In both age groups, staining in the periphery appeared heavier than in the central retina. E–H. Images of photoreceptor inner segments at higher magnification. The peripheral images in both aged groups appeared to show heavier staining than central region. I. Western blot analysis confirming higher levels of expression of Complex III in the periphery of both age groups, but also revealing no consistent age related changes in expression in any retinal layer. N = 5 per group. Abbreviations: GCL, ganglion cell layer. IPL, inner plexiform layer. INL, inner nuclear layer. ILM, inner limiting membrane. OPL, outer plexiform layer. ONL, outer nuclear layer. ILM, outer limiting membrane. Photoreceptors, PR. Young centre, Y(C). Young periphery Y (P). Old centre, O(C). Old periphery, O(P). Statistical significance, * $p < 0.05$. Uncropped images were not available for these data.

<https://doi.org/10.1371/journal.pone.0273882.g002>

Mitochondrial membrane potential, permeability and size

A measure of mitochondrial viability can be obtained by examining membrane potential, membrane permeability and mitochondrial size that were derived from isolated mitochondrial preparations. Generally, with age there are reductions in mitochondrial membrane potential, an increase in their membrane permeability and an overall increase in mitochondrial size [20–23]. These features were seen here in the primate retinal tissues. Membrane potentials were significantly lower in older than younger animals although there were not significant regional differences. Overall reductions across age were approximately 30–40% (Fig 4A). Reduced membrane potentials were associated with increased membrane permeability. Again, there were no significant regional differences within either age group but there was a significant increase in permeability with age of approximately 30–40% (Fig 4B).

Reductions in light scatter are associated with an increase in mitochondrial size. In both young and old primates, mitochondrial size was greater in the centre than the periphery and in both regions, there was an increase in mitochondrial size with age, but this was only statistically significant for the central retina (Fig 4C).

Mitochondrial number and voltage-dependent anion channels (VDAC)

Estimating mitochondrial number in whole retinae is challenging and not realistically possible in electron microscopy (EM) or using Mito trackers that require fresh tissue. Hence, here we have used Tom20 a mitochondrial membrane marker to estimate their number. Western blot analysis shows a significant approximate 50% reduction in Tom20 between young and older




Fig 3. A-D. Immunohistochemistry of cytochrome C oxidase subunit IV in young and old primate central (C) and peripheral (P) retina. Staining is present mostly in the ganglion cell layer (GCL), Inner plexiform layer (IPL) and outer plexiform layer (OPL), also in inner segments of the photoreceptor layer (PR). Central expression appeared heavier than that in the periphery. E-H. Images of inner segment of photoreceptors at higher magnification. Central inner segments of both age groups appear to show higher levels of expression than those in the periphery. I. Immunostaining results were confirmed with Western blot showing higher levels of complex IV in central regions. However, there were no consistent age related differences in any retinal layer. N = 5 per group. Abbreviations: GCL, ganglion cell layer. IPL, inner plexiform layer. INL, inner nuclear layer. ILM, inner limiting membrane. OPL, outer plexiform layer. ONL, outer nuclear layer. ILM, outer limiting membrane. Photoreceptors, PR. Young centre, Y(C). Young periphery Y(P). Old centre, O(C). Old periphery, O(P). See supplementary for full uncropped images.

<https://doi.org/10.1371/journal.pone.0273882.g003>

retinae with no regional difference between centre and periphery in either age group, consistent with there being significant mitochondrial decline with age (Fig 5A). VDAC is a general diffusion pore for small hydrophilic molecules that opens when mitochondrial membrane




Fig 4. Three mitochondrial metrics were assessed from the same retinal samples across age and location. **A.** Retinal mitochondrial membrane potentials from young and old primates from central and peripheral regions. An overall significant age-related decrease of between 30–40% was revealed in membrane potentials for both central and peripheral regions. **B.** Mitochondrial membrane permeability was assessed revealing an inverse relationship with mitochondrial membrane potential with permeability increasing with age across the retina associated with declining potentials. Increased aged permeability was approximately 30–40% independent of location. **C.** Mitochondrial size was determined via light scatter. Larger mitochondria scatter less light due to increased absorbance by their larger surface area. In both age groups, mitochondria were larger in the periphery than the centre. However, with age mitochondria increased in size, however, this only reached statistical significance in the central retina. Hence, with age mitochondria have reduced membrane potential, increased permeability and in the centre increased size. N = 5 per groups. Abbreviations: Central, C. Peripheral, P. Statistical significance, * p < 0.05, ** p < 0.01.

<https://doi.org/10.1371/journal.pone.0273882.g004>




Fig 5. A. Quantitative assessment of the mitochondrial marker Tom20 normalised with β -actin using western blots. Tom20 is localised in the outer mitochondrial membrane and it recognises precursors to facilitate protein across the outer mitochondrial membrane. The graph shows that young primates have more mitochondria than the old primates do. Peripheral retina in young have significantly higher mitochondria than peripheral retina of older primates ($P = 0.0143$). B. Western blot analysis of the protein voltage-dependent anion channel (VDAC) which is localised in the outer membrane of the mitochondria and it controls hydrophilic solutes between the mitochondria and the other cell compartments. The overall results show that younger primates have more VDAC than older primates when normalised with β -actin but central retina of both young and old primates have more VDAC than the peripheral retina. The central retina of young primates have significantly more VDAC than the young peripheral retina ($P = 0.0278$). There was no statistical significant decrease between old primate peripheral retina compared to the centre but there is a trend that central retina has less VDAC than peripheral when normalised with beta-actin. C. Western blot results of VDAC normalised with the amount of mitochondria in retina of young and old primates. It shows that older primates both in the central and periphery have significantly more VDAC/Tom20 ratio than young primates do ($P = 0.0179$ and $P = 0.0079$ respectively). $N = 5$ in each groups * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$; C- central and P-peripheral; Y-young and O- old. See supplementary for full uncropped images.

<https://doi.org/10.1371/journal.pone.0273882.g005>

potentials decline and plays a role in metabolite transport and can release cytochrome c that triggers apoptosis. When this is normalised against beta actin less is found in the periphery in young and older, and overall less in older animals (Fig 5B). But when it is normalised against Tom20 there is a significant increase with age across both retinal regions (Fig 5C). This is consistent with the notion that there is an increase in VDAC pore in age that could be indicative of increased apoptosis.

Outer retinal structure

Mitochondria are tightly packed in the apical region of photoreceptor inner segments [24] and their density here accounts for a significant proportion of those found in the retina. With age there is significant rod loss in the primate but there is no evidence for cone loss [7, 25], although cone function declines [26]. Hence, we ask what mitochondrial features of ageing are present in cones that may explain their loss of function.

A key feature of the ageing outer retina in the primate was the loss of content of the proximal inner segment in many cone photoreceptors. This region is normally occupied by the endoplasmic reticulum [24] that is critical for the maintenance of fission (OPA1) and fusion (Fis1) processes in mitochondrial dynamics. This feature of the aged inner segment was twice as prevalent in the peripheral compared to the centre (Fig 6).

It has been noted that in human retinae some cones have a displacement of their nuclei through the outer limiting membrane [27, 28]. Fig 7A shows a series of adjacent cones that display progressive translocations of their nuclei through the outer limiting membrane into the inner segment. Star (*) in this figure is a rod that also appears to have a nucleus breaking through the outer limiting membrane. In Fig 7B, a cone nucleus has fully transitioned into the




Fig 6. Representative photomicrographs showing age related changes in primate retina. Retina from young and old primates were processed and embedded in plastic, sectioned and stained with Acid Fuchsin (mitochondrial stain). **A.** Photomicrograph of a young primate taken at the centre of the retina showing compact and elongated photoreceptors. The mitochondria (arrowheads) are thin, tubular and elongated. **B.** Photomicrograph from old primate central retina showing the photoreceptors are swollen and some of them there seem to have less Golgi and endoplasmic reticulum in the inner segment (* asterix). The mitochondria are sparse, fragmented and rounder in shape. **C.** Photomicrograph of young primate retina from the peripheral. The mitochondria have thin tubular and elongated shape. **D.** Photomicrograph of old retinae from the peripheral area. The photoreceptors are swollen and some have a void in them at the Golgi and endoplasmic reticulum area of the inner segments (* asterix). The mitochondria appear fragmented and rounder in shape.

<https://doi.org/10.1371/journal.pone.0273882.g006>

inner segment. In Fig 7C, the cone indicated by the arrow has not only transitioned into the inner segment but is directly adjacent to the mitochondrial population and the vacuous region is present in the outer nuclear layer. On the right hand side other vacuous regions can be seen crossing into the outer nuclear layer.

Discussion

The results of this study show that in spite of a 70% reduction in retinal ATP in the same colony of old world primates as we have used here [9], complex activity is not reduced. However, other mitochondrial changes are marked with age with reduced membrane potential and increased mitochondrial membrane permeability, along with evidence of mitochondrial swelling as shown by reduced light scatter. Further, aged changes in Tom20 are consistent with an aged decline in mitochondrial number, and when mapped against VDAC the data are consistent with opening of membrane channels that can be associated with apoptosis. This study also shows significant structural changes in regions that would normally be occupied by endoplasmic reticulum in cones. Here the nucleus in some cells crosses the outer retinal margin and enters the inner segments and can become embed in the mitochondrial population. The endoplasmic reticulum is a key player in regulating mitochondrial dynamics and these structural changes may undermine this process.

As there is little evidence for primate cone loss with age, but photopic function declines significantly [7, 25, 26], it is likely that the mitochondrial decline documented here, along with structural changes in the inner segment are associated with visual decline. The structural changes in the inner segment appear to be unique to primates, and have not to our knowledge,




Fig 7. Displacement of nuclei from the outer nuclear layer to the inner segment. Representative photomicrographs of old primate peripheral retina showing displacement of the nuclei from the outer nuclear layer (ONL) to the inner segment (IS) of the photoreceptors (black arrows). The retinae were sectioned at 2.5 μ m and stained with Acid Fuchsin. **A.** Nuclei displacements seem to occur mainly in cones (black arrows) and rarely in rods (*asterix). It is observed that nuclei displacement occurs more in the old primate periphery than in young peripheral retina. **B.** Micrograph shows that the nucleus has completely crossed the outer limiting membrane into the inner segment, in the area of endoplasmic reticulum and Golgi (the black arrow on the right). **C** shows that the nuclei have migrated and is sitting on top of the mitochondria (black arrow). Scale bar = 10 μ m.

<https://doi.org/10.1371/journal.pone.0273882.g007>

been reported in other animals. This reinforces the need to examine retinal ageing in a model appropriate for humans.

Our primates were of Mauritian origin and most likely have a restricted gene pool because of a higher rate of inbreeding within and closed island population. While the Indonesian macaques have longer life spans with a maximum around 30 years the average age of death in our long established population is close to 18 years. By this time the primates display a wide range of hallmarks of ageing including reduced fertility, grey hair, arthritic joints and an advanced browning of the lens. As such they are ideal models for ageing research.

A challenging question arising from the data in this study is the apparent discrepancy between the 70% reduction in retinal ATP found with age by us in a previous study on animals from the same colony as used here [9], compared to the failure to find any significant age related reduction in mitochondrial complex activity. Here there are a number of resolving possibilities. First, it is possible that there are the same number of electron transport chains, but fewer ATPase molecules to convert energy held within H ions, into energy trapped within ATP. Second, ATPase pumps work both ways, as such there could be a continual breakdown of ATP alongside production. Third, a component of the electron transport chain could be lost with age, as such even though most proteins are there, the chain is disrupted. At this stage it is

unclear which of these mechanisms, or any alternative, is responsible for the discrepancy between ATP production and complex activity.

In our previous study [9], we showed that there is a significant drop in the enzyme activities of complex IV with age. Here the results showed that the same amount of enzyme complex IV is present in both young and older primates. This indicates that even the enzymes are present in the same amount, but their activity is significantly reduced. There is a further complication. It is important that the data presented here for mitochondrial function is seen as a snapshot of aged differences. Mitochondria change their function over the day and this includes complex activity, ATP production and ADP/ATP ratios [29]. The primates in this study were all culled between 9am and 10.30am. However, sacrifice at different times may result in different patterns between the age groups, as we have little idea how the mitochondrial metrics in the primate retina change over the day or how any daily patterns in these in young animals are impacted by age.

The reductions in mitochondrial number, reduced membrane potential, increased membrane permeability and mitochondrial size are all consistent with age related changes [30–32]. These metrics are also consistent with the increase in VDAC expression. Taken together they demonstrate declines in mitochondria with age. However, the one striking factor is that this study offers almost no evidence for the central retina suffering a greater degree of age related decline than found in the periphery as has been proposed [13]. But there are a number of differences between the Barron et al study finding regional differences and our own. Barron et al examined the fovea directly, which we did not, our analysis was confined to the central macular. Further, their analysis was of mtDNA deletions and cytochrome c oxidase deficiency. We made no attempt to examine mtDNA deletions, although our inability to find any significant reductions in cytochrome c oxidase (complex IV) is perplexing. But there are a number of different factors. Barron et al. studied human post-mortem human tissue spanning ages, collected a number of hours after death, and likely at different times of the day, while our study was taken at the point of death from animals in only two defined categories of age. Being human tissue, it would have suffered from varied environmental factors over life such as diet, different patterns of exercise and potentially smoking. Our primates experienced almost no environmental variability and were maintained on a standard vegetable-based diet over life. Taken together, this array of differences makes the two studies hard to compare.

Overall, the symmetry of changes between the two retinal regions used in this study in the analysis of mitochondrial extracts is marked in each of the data sets we present with the exception of changes in mitochondrial size. Here there was evidence of an increase in size in central mitochondria compared to the periphery. Such changes are associated with pathology [32, 33]. But this was the only instance of differential mitochondrial behaviour between the two regions examined.

Interactions between the endoplasmic reticulum and mitochondria are key in regulating mitochondrial biogenesis [34, 35] and their disruption may lead to neurodegeneration and aged disease [36–38]. The endoplasmic reticulum is located between the nucleus and the mitochondrial population in the inner segment, although elements of the endoplasmic reticulum project up through mitochondrial population, particularly those located proximally to the nucleus [24]. The apparent loss of the endoplasmic reticulum in many cones, particularly those in the periphery is likely to be associated with disruption to mitochondrial dynamics that play an important role in autophagy and the maintenance of mitochondrial health.

Apparent loss of the endoplasmic reticulum was associated with movement of the nucleus across the outer limiting membrane to fill this void. This was obvious in cones, but also present in some rods. Nuclear translation across the outer limiting membrane has been noted before in aged human retinae [27, 28], and as in this study, it is primarily a feature of the peripheral

retina. In some cases, the nucleus embeds in the mitochondrial population and the vacuous region is left behind and crosses the outer limiting membrane into the ONL. These data fit with others who have shown redistribution of cone opsin and swelling of cone distal axons with age in human retinae [39]. Cone nuclear displacement has also been noted as a feature in aged human retina suffering from macular degeneration [40]. These structural changes have not been noted in other aged animals, particularly mice that are commonly used as aged models.

There is little evidence for cone loss in the ageing primate retina including humans [7, 25], although Reumueller et al. found a weak reduction in number, but this was from imaging with adaptive optics [41]. Panda-Jones et al. also noted a small decline in cone numbers in human ageing but the scatter in the data was very large [42]. Counting primate cones is often undertaken from whole mounted retinae where the large cone inner segment distinguishes them from rods [7], or they can be identified when stained for a cone opsin and the outer segments are counted [7]. In neither preparation would the structural features we have described be readily identified. However, given the marked changes that occur to the internal structure of many aged cones it would be surprising if these cells functioned normally. Such changes along with those in mitochondrial integrity including reduced membrane potential and their increased permeability are likely to impact on ageing human photopic vision and may be related to aged reductions in colour sensitivity [26]. An additional factor that is relevant to this study is the finding that the spatial organisation of mitochondria in mammalian cones has the ability to act as light guide, focussing the light path onto the outer segments. It is likely that the changes we describe here on aged mitochondria impact upon this ability [43].

Supporting information

S1 File.
(DOCX)

Author Contributions

Conceptualization: Michael Barry Powner, Matthew John Hayes, Glen Jeffery.

Investigation: Jaimie Hoh Kam, Harpreet Shinhmar, Asmaa Aboelnour.

Methodology: Jaimie Hoh Kam, Harpreet Shinhmar.

Project administration: Jaimie Hoh Kam.

Writing – original draft: Glen Jeffery.

References

1. López-Otín C, Blasco MA, Partridge L, Serrano M, Kroemer G. The hallmarks of aging. *Cell*. 2013; 153(6):1194. <https://doi.org/10.1016/j.cell.2013.05.039> PMID: 23746838
2. Giacomello M, Pyakurel A, Glytsou C, Scorrano L. The cell biology of mitochondrial membrane dynamics. *Nat Rev Mol Cell Biol*. 2020; 21(4):204–24. <https://doi.org/10.1038/s41580-020-0210-7> PMID: 32071438
3. Country MW. Retinal metabolism: A comparative look at energetics in the retina. *Brain Res*. 2017; 1672:50–7. <https://doi.org/10.1016/j.brainres.2017.07.025> PMID: 28760441
4. Viegas FO, Neuhauss SCF. A Metabolic Landscape for Maintaining Retina Integrity and Function. Vol. 14, *Frontiers in Molecular Neuroscience*. 2021;p.60. <https://doi.org/10.3389/fnmol.2021.656000> PMID: 33935647
5. Finch CE. Evolution of the human lifespan and diseases of aging: Roles of infection, inflammation, and nutrition. *Proc Natl Acad Sci*. 2010 Jan; 107(suppl 1):1718 LP– 1724.

6. Rose MR, Mueller LD. Evolution of human lifespan: Past, future, and present. *Am J Hum Biol.* 1998 Jan; 10(4):409–20. [https://doi.org/10.1002/\(SICI\)1520-6300\(1998\)10:4<409::AID-AJHB2>3.0.CO;2-B](https://doi.org/10.1002/(SICI)1520-6300(1998)10:4<409::AID-AJHB2>3.0.CO;2-B) PMID: 28561475
7. Curcio CA. Photoreceptor topography in ageing and age-related maculopathy. *Eye.* 2001; 15(3):376–83. <https://doi.org/10.1038/eye.2001.140> PMID: 11450761
8. Wuergler S. Colour Constancy Across the Life Span: Evidence for Compensatory Mechanisms. *PLoS One.* 2013 May; 8(5):e63921. <https://doi.org/10.1371/journal.pone.0063921> PMID: 23667689
9. Hoh Kam J, Weinrich TW, Shinhmar H, Powner MB, Roberts NW, Aboelnour A, et al. Fundamental differences in patterns of retinal ageing between primates and mice. *Sci Rep.* 2019; 9(1):1–14.
10. Shinhmar H, Hogg C, Neveu M, Jeffery G. Weeklong improved colour contrasts sensitivity after single 670 nm exposures associated with enhanced mitochondrial function. *Sci Rep.* 2021 Dec; 11(1):1–9.
11. Grewal MK, Sivapathasuntharam C, Chandra S, Gurudas S, Chong V, Bird A, et al. A Pilot Study Evaluating the Effects of 670 nm Photobiomodulation in Healthy Ageing and Age-Related Macular Degeneration. *J Clin Med.* 2020; 9(4):1001. <https://doi.org/10.3390/jcm9041001> PMID: 32252424
12. Cuneo A, Powner MB, Jeffery G. Death by color: Differential cone loss in the aging mouse retina. *Neurobiol Aging.* 2014; 35(11):2584–91. <https://doi.org/10.1016/j.neurobiolaging.2014.05.012> PMID: 24929970
13. Barron MJ, Johnson MA, Andrews RM, Clarke MP, Griffiths PG, Bristow E, et al. Mitochondrial Abnormalities in Ageing Macular Photoreceptors. *Invest Ophthalmol Vis Sci.* 2001 Nov; 42(12):3016–22. PMID: 11687550
14. Liedigk R, Kolleck J, Böker KO, Meijaard E, Md-Zain BM, Abdul-Latiff MAB, et al. Mitogenomic phylogeny of the common long-tailed macaque (*Macaca fascicularis fascicularis*). *BMC Genomics.* 2015; 16(1):222. <https://doi.org/10.1186/s12864-015-1437-0> PMID: 25887664
15. Haus T, Ferguson B, Rogers J, Doxiadis G, Certa U, Rose NJ, et al. Genome typing of nonhuman primate models: implications for biomedical research. *Trends Genet.* 2014 Nov 1; 30(11):482–7. <https://doi.org/10.1016/j.tig.2014.05.004> PMID: 24954183
16. Kanthaswamy S, Ng J, Satkoski Trask J, George DA, Kou AJ, Hoffman LN, et al. The genetic composition of populations of cynomolgus macaques (*Macaca fascicularis*) used in biomedical research. *J Med Primatol.* 2013 Jun 1; 42(3):120–31. <https://doi.org/10.1111/jmp.12043> PMID: 23480663
17. Lohr JN, David P, Haag CR. Reduced lifespan and increased ageing driven by genetic drift in small populations. *Evolution (NY).* 2014 Sep 1; 68(9):2494–508. <https://doi.org/10.1111/evo.12464> PMID: 24897994
18. Kraus C, Snyder-Mackler N, Promislow DEL. How size and genetic diversity shape lifespan across breeds of purebred dogs. *GeroScience.* 2022. <https://doi.org/10.1007/s11357-022-00653-w> PMID: 36066765
19. Figueira TR, Melo DR, Vercesi AE, Castilho RF. Safranin as a Fluorescent Probe for the Evaluation of Mitochondrial Membrane Potential in Isolated Organelles and Permeabilized Cells. In: Palmeira CM, Moreno AJ, editors. *Mitochondrial Bioenergetics: Methods and Protocols.* Totowa, NJ: Humana Press; 2012. p. 103–17.
20. Navarro A, Boveris A. Rat brain and liver mitochondria develop oxidative stress and lose enzymatic activities on aging. *Am J Physiol—Regul Integr Comp Physiol.* 2004; 287(5 56–5):1244–9. <https://doi.org/10.1152/ajpregu.00226.2004> PMID: 15271654
21. Paradies G, Paradies V, Ruggiero FM, Petrosillo G. Changes in the mitochondrial permeability transition pore in aging and age-associated diseases. *Mech Ageing Dev.* 2013; 134(1–2):1–9. <https://doi.org/10.1016/j.mad.2012.12.006> PMID: 23287740
22. Crompton M. Mitochondria and aging: a role for the permeability transition? *Aging Cell.* 2004 Feb; 3(1):3–6. <https://doi.org/10.1046/j.1474-9728.2003.00073.x> PMID: 14965348
23. Chalmers S, Saunter CD, Girkin JM, McCarron JG. Age decreases mitochondrial motility and increases mitochondrial size in vascular smooth muscle. *J Physiol.* 2016 Aug; 594(15):4283–95. <https://doi.org/10.1113/JP271942> PMID: 26959407
24. Hayes MJ, Jeffery G, Tracey-White D, Kam JH, Powner MB, Jeffery G. The 3D Organisation of Mitochondria in Primate Photoreceptors. *Sci Rep.* 2021; 11(1):18863. <https://doi.org/10.1038/s41598-021-98409-7> PMID: 34552195
25. Weinrich TW, Powner MB, Lynch A, Jonnal RS, Werner JS, Jeffery G, et al. No evidence for loss of short-wavelength sensitive cone photoreceptors in normal ageing of the primate retina. *Sci Rep.* 2017; 7:1–7.
26. Werner JS, Steele VG. Sensitivity of human foveal color mechanisms throughout the life span. *J Opt Soc Am A.* 1988; 5(12):2122–30. <https://doi.org/10.1364/josaa.5.002122> PMID: 3230481

27. Gartner S, Henkind P. Aging and degeneration of the human macula. 1. Outer nuclear layer and photoreceptors. *Br J Ophthalmol*. 1981 Jan; 65(1):23–8. <https://doi.org/10.1136/bjo.65.1.23> PMID: 7448155
28. Lai YL, Masuda K, Mangum MD, Lug R, Macrae DW, Fletcher G, et al. Subretinal displacement of photoreceptor nuclei in human retina. *Exp Eye Res*. 1982; 34(2):219–28. [https://doi.org/10.1016/0014-4835\(82\)90056-2](https://doi.org/10.1016/0014-4835(82)90056-2) PMID: 7060649
29. Weinrich TW, Kam JH, Ferrara BT, Thompson EP, Mitrofanis J, Jeffery G. A day in the life of mitochondria reveals shifting workloads. *Sci Rep*. 2019; 9(1):1–8.
30. Haas RH. Mitochondrial dysfunction in aging and diseases of aging. Vol. 8, *Biology*. 2019. <https://doi.org/10.3390/biology8020048> PMID: 31213034
31. Nicholls DG. Mitochondrial membrane potential and aging. *Aging Cell*. 2004; 3(1):35–40. <https://doi.org/10.1111/j.1474-9728.2003.00079.x> PMID: 14965354
32. Chapa-Dubocq X, Makarov V, Javadov S. Simple kinetic model of mitochondrial swelling in cardiac cells. *J Cell Physiol*. 2018/01/23. 2018 Jul; 233(7):5310–21. <https://doi.org/10.1002/jcp.26335> PMID: 29215716
33. Sivapathasuntharam C, Hayes MJ, Shinhmar H, Kam JH, Sivaprasad S, Jeffery G. Complement factor H regulates retinal development and its absence may establish a footprint for age related macular degeneration. *Sci Rep*. 2019; 9(1):1–13.
34. Rowland AA, Voeltz GK. Endoplasmic reticulum–mitochondria contacts: function of the junction. *Nat Rev Mol Cell Biol*. 2012; 13(10):607–15. <https://doi.org/10.1038/nrm3440> PMID: 22992592
35. Marchi S, Patergnani S, Pinton P. The endoplasmic reticulum–mitochondria connection: One touch, multiple functions. *Biochim Biophys Acta—Bioenerg*. 2014; 1837(4):461–9. <https://doi.org/10.1016/j.bbabi.2013.10.015> PMID: 24211533
36. Lee K-S, Huh S, Lee S, Wu Z, Kim A-K, Kang H-Y, et al. Altered ER-mitochondria contact impacts mitochondria calcium homeostasis and contributes to neurodegeneration in vivo in disease models. *Proc Natl Acad Sci U S A*. 2018 Sep; 115(38):E8844–53. <https://doi.org/10.1073/pnas.1721136115> PMID: 30185553
37. Martinvalet D. The role of the mitochondria and the endoplasmic reticulum contact sites in the development of the immune responses. *Cell Death Dis*. 2018; 9(3):336. <https://doi.org/10.1038/s41419-017-0237-7> PMID: 29491398
38. Molledo O, Remondelli P, Amodio G. The Mitochondria-Endoplasmic Reticulum Contacts and Their Critical Role in Aging and Age-Associated Diseases. *Front cell Dev Biol*. 2019; 7:172. <https://doi.org/10.3389/fcell.2019.00172> PMID: 31497601
39. Shelley EJ, Madigan MC, Natoli R, Penfold PL, Provis JM. Cone Degeneration in Aging and Age-Related Macular Degeneration. *Arch Ophthalmol*. 2009 Apr 1; 127(4):483–92. <https://doi.org/10.1001/archophthalmol.2008.622> PMID: 19365029
40. Pow D V, Sullivan RKP. Nuclear kinesis, neurite sprouting and abnormal axonal projections of cone photoreceptors in the aged and AMD-afflicted human retina. *Exp Eye Res*. 2007 May; 84(5):850–7. <https://doi.org/10.1016/j.exer.2007.01.005> PMID: 17343852
41. Reumueller A, Wassermann L, Salas M, Schranz M, Told R, Kostolna K, et al. Three-dimensional assessment of para- and perifoveal photoreceptor densities and the impact of meridians and age in healthy eyes with adaptive-optics optical coherence tomography (AO-OCT). *Opt Express*. 2020 Nov; 28(24):36723–39. <https://doi.org/10.1364/OE.409076> PMID: 33379760
42. Panda-Jonas S, Jonas JB, Jakobczyk-Zmija M. Retinal Photoreceptor Density Decreases with Age. *Ophthalmology*. 1995; 102(12):1853–9. [https://doi.org/10.1016/s0161-6420\(95\)30784-1](https://doi.org/10.1016/s0161-6420(95)30784-1) PMID: 9098287
43. Ball JM, Chen S, Li W. Mitochondria in cone photoreceptors act as microlenses to enhance photon delivery and confer directional sensitivity to light. *Sci Adv*. 2022; 8(9):1–13. <https://doi.org/10.1126/sciadv.abn2070> PMID: 35235359