



City Research Online

City, University of London Institutional Repository

Citation: Rodgers, L., Botting, N., Cartwright, M., Harding, S. & Herman, R. (2023). Shared characteristics of intervention techniques for oral vocabulary and speech comprehensibility in preschool children with co-occurring features of developmental language disorder and a phonological speech sound disorder: protocol for a systematic review with narrative synthesis. *BMJ Open*, 13(6), e071262. doi: 10.1136/bmjopen-2022-071262

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/30680/>

Link to published version: <https://doi.org/10.1136/bmjopen-2022-071262>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

TITLE PAGE

TITLE:

The shared characteristics of intervention techniques for oral vocabulary and speech comprehensibility in pre-school children with co-occurring features of developmental language disorder and a phonological speech sound disorder: protocol for a systematic review with narrative synthesis

AUTHORS:

Corresponding author: Lucy Rodgers. Department of Language and Communication Science, Myddelton Street Building, City University of London, EC1V 0HB. Lucy.rodgers@city.ac.uk
<https://orcid.org/0000-0001-6585-7206>

Co-author: Prof. Nicola Botting. Department of Language and Communication Science, City University, London, UK.

Co-author: Dr Martin Cartwright. Department of Health Services Research and Management, City University, London, UK.

Co-author: Dr Sam Harding. Bristol Speech and Language Therapy Research Unit, North Bristol NHS Trust, Bristol, UK. <https://orcid.org/0000-0002-5870-2094>

Co-author: Prof. Rosalind Herman. Department of Language and Communication Science, City University, London, UK. <http://orcid.org/0000-0001-5732-9999>

Word count (excluding title page/abstract/references): 4,527

Key words: Language Development Disorders; Speech Sound Disorder; Child Language; Speech-Language Pathology

ABSTRACT

Introduction:

Evidence suggests that over a third of young children with Developmental Language Disorder (DLD) or Speech Sound Disorder (SSD) have co-occurring features of both. A co-occurring DLD and SSD profile is associated with negative long-term outcomes relating to communication, literacy and emotional wellbeing. However, the best treatment approach for young children with this profile is not understood. The aim of the proposed review is to identify intervention techniques for both DLD and SSD, along with their shared characteristics. The findings will then be analysed in the context of relevant theory. This will inform the content for a new or adapted intervention for these children.

Methods and analysis:

This search will build on a previous systematic review by Roulstone et al. (2015) but with a specific focus on oral vocabulary (DLD outcome) and speech comprehensibility (SSD outcome). These outcomes were identified by parents and Speech and Language Therapists within the pre-study stakeholder engagement work. The following databases will be searched for articles from January 2012 onwards: Ovid EMCare, Medline Complete, CINAHL, APA PsychINFO, Communication Source and ERIC. Two reviewers will independently perform the title/abstract screening and the full text screening with the exclusion criteria document being revised in an iterative process. Articles written in languages other than English will be excluded. Data will be extracted regarding key participant and intervention criteria, including technique dosage and delivery details. This information will then be pooled into a structured narrative synthesis.

Ethics and dissemination: Ethical approval is not needed for a systematic review protocol. Dissemination of findings will be through peer-reviewed publications, social media, and project steering group networks.

Study registration number: PROSPERO, CRD4202237393

Strengths and limitations of this study

- This protocol follows the referred Reporting Items for Systematic Review and Meta-Analysis Protocols (PRISMA-P) guidelines.
- Electronic databases spanning medicine, education, and psychology will be searched.
- Electronic databases in languages other than English will not be searched.
- Meta-bias(es) within the literature cannot be fully controlled.
- The level of detail within intervention reporting, as per the TIDieR guidelines, has the potential to vary amongst studies.

INTRODUCTION

Within the field of child language disorders, there are often overlapping or co-occurring difficulties which create unique patient experiences. Yet, while there is ample literature on treatment for singly occurring difficulties, there is a notable gap in evidence for treating children with co-occurring disorders. This review focusses on intervention for children who have co-occurring features of both Developmental Language Disorder (DLD) and Speech Sound disorder (SSD).

Co-occurring Developmental Language Disorder (DLD) and Speech Sound Disorder (SSD)

An estimated 7.58% of 4 year olds present with features of a Developmental Language Disorder (DLD)(1). DLD is characterised by idiopathic difficulties in using and understanding spoken language (2). One feature is limited vocabulary development (2), which has a known association with childhood temper tantrums/mental health, and later language and literacy skills (3,4). Such features of DLD may co-occur with a Speech Sound Disorder (SSD); that is, difficulties in producing speech sounds (5). An estimated 3.4% of 4- year olds have SSD (6). One of the most devastating impacts SSD is the impact on a child's ability to make themselves understood to others in everyday life (7). The term for this is speech comprehensibility (8). A related term, speech intelligibility, refers to the acoustic-phonetic decoding of utterances, and is very closely related to speech comprehensibility as both are linked to the functional use of speech. As with limited vocabulary, poor speech comprehensibility/intelligibility within the early years have also been associated with negative longer-term outcomes, including persisting speech difficulties (9) and poor literacy skills (10,11). Although it is typical for very young children not to be fully understood to those around them as their speech develops, by 4 years of age a child would typically be at least 50% intelligible (12).

Thirty six percent of 4-year olds with idiopathic SSD also have oral (i.e. expressive-spoken) language features of DLD (6). This high rate of co-occurrence is in keeping with historical research in the area (13), as well as study data from clinical caseloads (14). The combined impact of co-occurring features of DLD/SSD is twofold; for example, for a child with limited oral vocabulary and speech comprehensibility, not only are they unable to use many words, but the limited words they do have will not be understood to others within their daily lives. It therefore may be unsurprising that co-occurring phonological DLD/SSD features in early childhood are associated with negative long-term outcomes relating to literacy (15,16) and communication (17,18), with downstream consequences for quality of life (18,19) and emotional wellbeing (20). Consequently, access to effective and appropriately targeted intervention for children with this profile is crucial.

Phonological SSDs are the most frequently presenting SSD subtype (5), and occur when a child has difficulties with manipulating the different sound contrasts (phonemes) which are needed to form words (21). There are different types of phonological SSDs, including consistent phonological disorder (where the child makes consistent sound omissions or substitutions) and inconsistent phonological disorder (where these errors have no consistent pattern) (21). Research highlights a known link between DLD and phonological SSDs, as both disorders are underpinned by shared linguistic deficits (2). This overlap is

represented in the seminal CATALISE DLD consensus paper (2). In contrast to phonological SSDs, the CATALISE authors' speech, language and communication needs diagram highlights how other SSD subtypes, such as motor based SSDs like dysarthria, have a less marked overlap with DLD. They also often have a known cause (i.e. are non-idiopathic). Due to their significant overlap with DLD which has no known causation, this review will focus on phonological SSDs which are also idiopathic in nature.

Figure 1: DLD within the broader category of Speech, Language and Communication Needs (SLCN) (2)

The overlap between language and phonological SSDs is further supported by studies on the speech and language development of young children, where complex and bi-directional relationships between the development of individual sounds (phonology) and words (the lexicon) have been identified (22,23). For example, the first words of young children primarily consist of the speech sounds already established within their emerging phonological inventory (23). This relationship between phonology and the lexicon may have implications for intervention with children with co-occurring features of DLD and a phonological SSD. For example, growth in vocabulary and/or the strengthening of phonological representations has the potential to impact speech and vocabulary development concurrently through a process known as 'lexical restructuring' (24). A further psycholinguistic theory of potential relevance is the speech processing model (25), which suggests that individual children with co-occurring features of DLD/a phonological SSD may have difficulties at one or more levels of speech processing, rather than just with phonological representations alone. Such theories are important within interventions for co-occurring DLD/phonological SSD as they can be used to inform intervention content and delivery.

Current interventions for pre-school co-occurring DLD/SSD

Although this overlap exists between DLD and phonological SSDs, there is currently a paucity of theoretically informed interventions which have been specifically developed for this group (26). Additionally, intervention studies within existence primarily target morphosyntactic aspects of expressive language, alongside accuracy of speech sound production (26,27). However, for younger children with this profile, and children whose features of DLD are more severe, building vocabulary is typically targeted in speech and language therapy prior to morphosyntax (28).

'Child Talk' (29) was a large National Institute of Health Research (NIHR) funded mixed methods programme of work, including a systematic review. This involved investigating the use of early years' speech and language therapy interventions. The findings led to the specification of a) a typology of early years' speech and language therapy (SLT) intervention, b) key intervention ingredients for each typology theme. A technique can be described as "the specific teaching behaviours/actions thought to effect change" (30). The findings highlighted that for children with co-occurring features of DLD/SSD, clinicians often adapt existing interventions by selecting and combining different techniques. This enables them to use their knowledge and experience to provide the best treatment that they can (29,31).

Although our knowledge of what techniques work best for children with this profile is limited, techniques identified might be related to underlying theories of potential relevance. For example, language modelling is typically linked to growth in expressive language (32). However, based on the lexical restructuring hypothesis, it is hypothesised that the subsequent impact of this language growth on the accuracy and segmentation of the child's phonological representations could also influence their phonological speech sound production (24). 'Building things into play' was also a key technique highlighted in Child Talk. Theoretically, this is supported by the latest research on the brain basis of speech and language learning, which indicates that learning best takes place within interactions which are meaningful for the child (33,34). Considering this, it is hypothesised that this technique supports speech and language learning through capitalising on the child's heightened attention and motivation during the play activity.

These considerations highlight a valuable opportunity for an intervention specific to this clinical group to be developed, utilising techniques which can be supported by relevant theory. Due to the current paucity of evidence, the associated negative impact of this co-occurring profile on long term outcomes, and the high level of presentation on clinical caseloads, there is an urgent need for such intervention development to take place. The first stage in this development would be to conduct a systematic review to identify potential techniques of relevance.

Broader context: an intervention development study

The proposed review updates the systematic review findings from 'Child Talk' (29) whilst refining the focus to techniques within interventions for children with features of DLD or a phonological SSD. Techniques will be extracted from included studies and then analysed in relation to shared characteristics and underpinning theory. The synthesis will then be used to inform the content of a new intervention which is being developed for young children with co-occurring features of DLD/phonological SSD.

Both DLD and SSD are heterogenous disorders (2,21), and therefore have a range of associated outcomes. This review, and body of intervention development work it is a part of, will focus exclusively on the outcomes of oral vocabulary (DLD outcome) and speech comprehensibility (SSD outcome). This is due to the afore mentioned impact of such difficulties on the everyday lives of young children; this decision is also elaborated on in the 'patient and public involvement' section of this paper.

Based on the dose form framework (30), shared characteristics for DLD/phonological SSD intervention techniques may include similarities in:

- 1) *who* delivers the technique; for example, is it the parent, clinician, or both?
- 2) *where* the technique is delivered; for example, at home, nursery, clinic or a combination of these?
- 3) the nature of *technique delivery*; for example, is it during an adult led structured game, child led play, everyday routines, or a combination of these?

Underpinning theory may relate to:

- 1)The lexical restructuring hypothesis (24)
- 2) Psycholinguistic models of speech and language development; such as the speech processing model (25)
- 3)The neural basis for speech and language development; for example, the role of meaningful interactions within language learning (33)

Objectives

The overarching aim of the review is to bring together intervention techniques from DLD and phonological SSD interventions. The objectives within this are to:

1. Identify the shared core characteristics of the techniques; this includes the deliverer, place of delivery, format of delivery and nature of delivery (*e.g. child or adult led*)
2. Compare and synthesise the shared core characteristics of the techniques in relation to underlying theory
3. Establish the best available evidence for interventions that incorporate these core characteristics of the intervention techniques

Research questions:

- 1)What are the shared core characteristics of intervention techniques in preschool interventions targeting speech comprehensibility and/or oral vocabulary?
- 2)How do these shared core characteristics relate to underlying theory?
- 3)What evidence is there for the effect of interventions that incorporate these core characteristics of intervention techniques?

METHODS AND ANALYSIS

In accordance with the guidelines, our systematic review protocol has been registered with the International Register of Systematic Reviews (PROSPERO) on 16/12/2022 (registration number CRD42022373931). In the event of any amendments to methodology set out below, the date of each amendment will be accompanied by a description of the change and the rationale in either the Prospero register and/or the final results paper.

Eligibility criteria

The eligibility criteria stated below are in line with the criteria from the original 'Child Talk' systematic review (29) with amendments according to the objectives of the current review. Most importantly, this review will focus specifically on the 'expressive language' and 'speech' themes generated from their initial typology of early years' Speech and Language

Therapy (SLT) interventions, as these themes encompass the two outcomes for which we are seeking to identify techniques.

Study designs

Included studies must report on an empirical evaluation of the effectiveness of an intervention. To ensure we identify all relevant literature, a range of study designs will be included. These include randomised control trials (RCTs), experimental and quasi-experimental studies, within subjects designs (e.g. pre-post studies) and case studies (which may include multiple baseline or other systematic manipulation of the intervention). Studies which report on single timepoint (e.g. cross-sectional studies) will be excluded. Studies focusing on efficacy, including lab-based training, will not be excluded if all other inclusion criteria are met. This is because information on the efficacy of speech/language learning techniques can be gleaned from these studies, although careful consideration will be given to how these results are integrated into the narrative analysis (further information on this is provided under 'data synthesis').

Population

To capture the age group most typically seen within clinical services, 80% of children within included studies must have been aged between 2:0 and 5:11 years. Additionally, although this review is part of a wider intervention development study for children aged 3 and 4 years, an expanded age range within this review will help to ensure all that techniques of potential relevance will be captured. The children within included studies must have presented with phonological speech production difficulties and/or difficulties relating to oral vocabulary, with all subtypes of phonological SSD included (e.g. consistent and inconsistent phonological disorder, phonological delay). These difficulties may be identified by standardised assessments such as the Preschool Language Scale (35), parental and/or professional observation reports such as the intelligibility in context scale (36) and/or probes. Probes may also be used to assess progress through the repeated measurement of the dependent variable before, during and after the intervention. As already observed in the literature, common probes within speech and language therapy interventions may include a selection of words containing the child's targeted speech sound/s or vocabulary (37,38). In keeping with the afore mentioned diagnostic description within CATALISE (2), included papers must state that the participants' needs had no obvious cause, i.e., excluding children with neurodevelopmental differences that have a known association with speech and/or language development, such as Autism or Cerebral Palsy. Due to the challenges in diagnosing DLD in very young children (2), and in order to maximise the identification of potentially relevant intervention techniques, studies will be included where the child does not have a formal diagnosis of DLD but is a late talker.

Intervention

We will include studies reporting on interventions delivered in any setting (e.g., home based, clinic) or format (e.g., face to face, online). The deliverer may be a speech and language therapist, speech and language therapy assistant, or equivalent professional (including education staff), and the intervention may involve professionals training up others (e.g., parents) to deliver some or all of the intervention.

Comparator

Comparators for included studies may be a control without an intervention (including multiple baseline and within subjects designs) or an alternative experimental group (i.e., intervention comparison).

Outcomes

Included papers must measure the effectiveness of the intervention on a) oral vocabulary, and/or b) speech comprehensibility. These outcomes must be evaluated via standardised assessment, probes, and/or observational ratings or scales.

If composite speech and language assessments are used, studies must report on the separate sub test results for oral vocabulary and/or speech comprehensibility to be included.

Studies with only syntactic measures of language change will be excluded; this includes mean length of utterance in morphemes (MLUm). However, they will be included if a proximal measure of vocabulary change is used alongside syntactic measures, such as the number of different words (NDW).

Speech comprehensibility is the SSD outcome in focus. As previously mentioned, comprehensibility and intelligibility are overlapping but differing constructs, with a shared focus on functional human communication (8). Therefore, we will also include studies with an outcome of improved speech intelligibility as a proxy for comprehensibility. This was deemed more suitable than using measures such as PCC as a proxy for comprehensibility, with their focus being on speech accuracy. Due to the very recent consensus in terminology, measures for comprehensibility might include measures with 'intelligibility' within their title, such as the 'intelligibility in context' scale, which is becoming increasingly utilised in SSD intervention research (36).

Information sources

The search will be conducted in Ovid EMCare, Medline Complete, CINAHL, APA PsychINFO, ERIC and Communication Source. These sources have been selected as they encompass the fields of health (medical, nursing and allied health professions), speech and language therapy, education and psychology and have been successfully utilised in previous reviews in the field (26,39). To support with literature saturation, supplementary search methods will be employed; this includes screening the reference lists from prominent reviews in the field post 2012 (40,41). We have selected reviews from 2012 onwards due to the original search going up to this date (29). Reference lists from included papers within the current search will also be screened for potential study eligibility. Forward citation searches in Web of Science (using the core collection) will also be carried out, with additional searches in Scopus if the titles are not available in Web of Science.

Due to resource constraints, articles written in languages other than English will be excluded. However, articles written in English where the participants speak languages other than English will be included. Additionally, grey literature searching will be confined to the inclusion of theses/dissertations, via the databases stated above. Thesis/dissertations have

been selected as although the original review (29) included a range of grey literature, thesis/dissertations were the only grey literature sources which contributed studies within the final included papers. In keeping with the original review, thesis/dissertations will only be included when a corresponding journal article cannot be found for the study.

Search strategy

Together with support from a specialist librarian, we will conduct an update of the original 'Child Talk' systematic review (29), searching articles from January 2012 to the present day. One of the researchers (SH) undertaking the current search also led on the original review. Relevant studies from the original 'speech' and 'expressive language' typology themes within the original 'Child Talk' review have already been located by reviewing the recorded outcomes for each study as stated on the original data extraction spreadsheet. Out of 41 papers from the 'speech' theme, two were found to address the outcome of comprehensibility/intelligibility. From the 30 papers within the 'expressive language' theme, 12 were found to include oral vocabulary as an outcome. These 14 papers will be further screened at stage 2 of the screening process (full text stage, outlined below).

The original review search strategy (29) has been updated for the current review, accounting for advances in terminology, e.g., consensus on the term 'Developmental Language Disorder' (2). The original 'Child Talk' search encompassed a broader range of speech and language outcomes, therefore the search terms for the current review have been adjusted to focus on our two specific outcomes of interest; oral vocabulary and speech comprehensibility. The updated search strategy was initially reviewed by two independent post-doctoral researchers in the field and adjusted as needed, for example, adding in the term 'specific language impairment', which may be relevant to older papers in the search. For the revised search strategy draft for each database, please see supplementary material 1.

Study records, Selection and data collection process

Search results will initially be imported into RefWorks, where duplicates will be removed by the first author (LR). The remaining articles will then be uploaded to the Covidence systematic review management database.

Initially, the first author (LR) will trial the exclusion guidance criteria document on 30 papers. For the initial draft of this exclusion criteria guidance document, please see supplementary material 2. The 30 papers will be randomly selected using a random number generator. These 30 papers will then be reviewed by a second reviewer. The reviewers will then meet to discuss discrepancies and make amendments to the exclusion guidance document if needed.

The screening and data extraction will be carried out as follows:

1) *Title/abstract screening*

The full set of titles/abstracts will be screened by the first author (LR). If uncertainty arises about how to apply to eligibility criteria to a specific paper, these articles will

be discussed with a member of the review team (who is not involved in the formal screening process). This may then lead to further revisions to the exclusion criteria document. Following this, a second independent reviewer (SH) will independently screen the titles/abstracts. Any disagreements, and how these may relate to the exclusion guidance document, will be discussed in consensus meetings. Any disputed articles will then be re-screened should alterations have been made to the exclusion criteria document. If disagreement is not caused by confusion over the exclusion criteria document, and consensus cannot be reached through discussion, a third reviewer will be consulted.

2) *Full paper screening*

At the full text screening stage, two reviewers (LR, SH) will independently appraise all of the remaining articles for inclusion, following the iterative process as outlined for stage 1.

To enable transparency of the reliability of screening at stages 1 and 2, Cohen's K for these stages will be reported in the final paper.

3) *Risk of bias/internal validity*

Retained studies will then undergo assessment of internal validity by two independent reviewers (LR, SH). The reviewers will have regular consensus meetings, after independently assessing up to 4 papers at a time, to resolve potential conflicts. If disagreements persist, a third reviewer will be involved. Disagreements that arise (including those that have been resolved) will be recorded and reported in the final paper.

For the PEDro-P (42), papers with a rating of six and over will be retained for data extraction. This aligns with the original review (29). On the Risk of Bias in N-of-1 Trials (RoBiNT) scale, included studies will be rated as *fair* or above (43).

4) *Data extraction*

The first author (LR) will extract data from the first 25% of studies. These will be randomly selected using random number generation. A second extractor (SH) will then independently extract data from the same studies. The two extractors will then meet to discuss potential discrepancies, and to update the data extraction form if needed. Following this, the first author (LR) will extract the remainder of the data.

Data items

Data will be sought regarding general study information (e.g., date; study type; location; participant numbers), population characteristics (e.g., male/female; age; speech/language profile-including phonological SSD subtype), intervention techniques (e.g., dosage; underpinning theory and justification given by the authors) and reported impact on the outcome of interest. Data on reported participant SES background will also be obtained, due to this being a known risk factor within developmental speech and language disorders (44). We will also collate information on the number of languages spoken by the participants, as

well as reported ethnicities, with this being a potential factor for the external validity of findings (i.e., relevance to 'everyday' clinical practice).

The Template for Intervention Description and Replication (TIDieR) will be used as a framework to guide the extraction process (45), combined with the speech and language therapy specific 'Dose form Framework and Definitions', based on the work of Warren and colleagues (46); this has been applied in other reviews specific to paediatric speech and language therapy intervention (40). Details of techniques will be extracted regarding intervention contexts (e.g. the overarching activity the technique is presented in), method of instruction (e.g. who delivers the technique, where and when), and technique dosage (dose frequency and dose duration). All reported dosage information will be extracted in order to allow for variation in study design; most notably, studies which target both oral vocabulary and speech comprehensibility concurrently.

Outcomes and prioritisation

The two outcomes (oral vocabulary, speech comprehensibility) are of equal interest within this review, regardless of whether they are primary or secondary outcomes within the included studies.

Risk of bias in individual studies

Individual studies will be assessed for internal validity. To encompass the range of study designs included within this review, we will use the PEDro-P (42). Specifically, for single case experimental designs, the RoBiNT scale will be used (43).

Data synthesis

Quantitative data

Overarching details for each included study, including the individual internal validity ratings, will be given in the first table. Two summary graphs will also be presented to convey the percentage of overall ratings from the PEDro-P and RoBiNT Scales. The frequency of techniques within the included papers will be presented either numerically within a table, or within a graph or chart if this deemed more suited to the data collected. We will be guided by the synthesis without meta-analysis (SWiM) in systematic review guidelines (47) and will report on the direction of effect of the interventions, using vote counting with a sign test if appropriate.

Qualitative data

A description of the identified techniques will be presented in a table, including details regarding how they were operationalised, based on TIDieR (45) and the dose form framework (23).

The narrative synthesis will include sections on:

- 1) Similarities and differences (including shared core characteristics) between techniques used for the different outcomes
- 2) Patterns of technique dosage and delivery across the interventions

3)How the similarities and differences (including shared core characteristics) in techniques relate to underlying theory. Depending on findings, this section will be broken down into sub sections focusing on each theory of interest, potentially including (but not necessarily limited to):

- the lexical restructuring hypothesis
- the speech processing model
- the neural basis for speech and language development

4)The effectiveness of interventions which contain these techniques/shared core characteristics of techniques

If relevant, any observed differences between interventions for different phonological SSD sub types will be incorporated into the narrative synthesis, or given in an additional section if deemed to be more appropriate to the data found.

In the event of lab-based training studies meeting the final inclusion criteria, this data will be presented on a separate table. Additionally, within the narrative synthesis itself they will not be directly compared to the effectiveness studies. Instead, they will be used to support any potential theory building arising from the synthesis.

If challenges are identified regarding gaps and quality in the knowledge base, this will also be explored within the results and discussion section of the corresponding results paper.

Patient and Public Involvement

According to the James Lind Alliance, knowing how to best select communication strategies according to a child's individual profile is the 2nd most important recommendation for research (48). This is strongly in keeping with the aims of this review, and highlights the broader relevance of this work.

For our wider intervention development study, outcomes were prioritised by clinicians and parents of pre-school children with DLD/SSD within pre-study PPIE work (49,50). They identified the outcomes of increasing 1) oral vocabulary (*DLD outcome*), and 2) speech comprehensibility (*SSD outcome*). This provides further support focusing on techniques that directly target oral vocabulary and speech comprehensibility.

In keeping with the integral role of PPIE throughout, a newly formed project PPIE steering group will provide input at key points in the review process. This is a diverse group consisting of parents, speech and language therapists, a person with DLD, a specialist early years educator, a bi/multi-lingual educational family support worker and a clinical equality, diversity and inclusion (EDI) expert. During the review, they will be involved with:

- 1) Reviewing the content for the data extraction form (supplementary document 3), prior to the data extraction phase
- 2) Identifying what data has the most relevance in the 'real world', with these potentially informing recommendations within the final paper

- 3) Defining and agreeing key messages to take from the review, and dissemination through the steering group networks.

Steering group input will be recorded and reported in the final article, in accordance with the GRIPP 2 reporting criteria short form (51).

Meta-bias (es)

It is important to acknowledge that meta-bias, including reporting and publication bias, is present within all aspects of health research. Although it is not possible to completely control for such bias, we will:

- 1) Establish if the protocol for each study was published before recruitment for participants commenced (where possible)
- 2) Compare the outcomes and results sections of the published report when a protocol is available (for when considering selective reporting bias)
- 3) Assess potential publication bias through the inclusion of prioritised grey literature (thesis/dissertations)

Confidence in Cumulative evidence

Confidence within the evidence as a whole will be based on the summary of the internal validity, as presented in the two summary tables (see data synthesis section). We will also acknowledge and discuss key factors relating to meta bias, and how the review findings should be interpreted based on this.

Figure 1: DLD within the broader category of Speech, Language and Communication Needs (SLCN) (2)

Ethics and dissemination:

As a systematic review this study does not warrant ethics board approval. Findings will be disseminated through peer reviewed publications, social media, and project steering group networks.

Registration details:

Study registration number: PROSPERO, CRD42022373931

Authors' contributions

LR led on this work and independently developed an initial draft of the manuscript (and appended documents). RH, NB, MC and SH suggested amendments after reviewing this and 3 subsequent re-drafts by LR. LR completed the fourth and final version of the manuscript, which was reviewed and agreed by all of the authors. LR led on the manuscript revisions following peer review, in collaboration with NB and SH. All authors agreed to the final submitted version.

The authors would also like to acknowledge and thank Dr Pauline Frizelle (University College Cork) and Dr Helen Stringer (Newcastle University) for providing expert independent reviews of the initial terms for the search strategy.

Competing interests' statement

There are no competing interests to declare.

Funding

This research was funded in whole, or in part, by the Wellcome Trust (Grant number 223500/Z/21/Z). For the purpose of open access, the author has applied a CC BY public copyright licence to any Author Accepted manuscript version arising from this submission.

FULL REFERENCES

- 1)Norbury, C.F., Gooch, D., Wray, C., Baird, G., Charman, T., Simonoff, E., Vamvakas, G. and Pickles, A., 2016. The impact of nonverbal ability on prevalence and clinical presentation of language disorder: Evidence from a population study. <https://doi.org/10.1111/jcpp.12573>
- 2)Bishop, D.V., Snowling, M.J., Thompson, P.A., Greenhalgh, T., Catalise-2 Consortium, Adams, C., Archibald, L., Baird, G., Bauer, A., Bellair, J. and Boyle, C., 2017. Phase 2 of CATALISE: A multinational and multidisciplinary Delphi consensus study of problems with language development: Terminology. <https://doi.org/10.1111/jcpp.12721>
- 3)Duff, F.J., Reen, G., Plunkett, K. and Nation, K., 2015. Do infant vocabulary skills predict school-age language and literacy outcomes?. *Journal of Child Psychology and Psychiatry*, 56(8), pp.848-856. <https://doi.org/10.1111/jcpp.12378>
- 4)Manning BL, Roberts MY, Estabrook R, Petitclerc A, Burns JL, Briggs-Gowan M, Wakschlag LS, Norton ES., 2019. Relations between toddler expressive language and temper tantrums in a community sample. *Journal of Applied Developmental Psychology*, volume 65, article 101070. <https://doi.org/10.1016/j.appdev.2019.101070>
- 5)Dodd, B., Reilly, S., Ttofari Eecen, K. and Morgan, A.T., 2018. Articulation or phonology? Evidence from longitudinal error data. <https://doi.org/10.1080/02699206.2018.1488994>
- 6)Eadie, P., Morgan, A., Ukoumunne, O.C., Ttofari Eecen, K., Wake, M. and Reilly, S., 2015. Speech sound disorder at 4 years: Prevalence, comorbidities, and predictors in a community cohort of children. <https://doi.org/10.1111/dmcn.12635>
- 7)McCormack J, McLeod S, McAllister L, Harrison LJ. My speech problem, your listening problem, and my frustration: The experience of living with childhood speech impairment. *Speech, Language and hearing Services in Schools*, 41(4), pp.379-392. [https://doi.org/10.1044/0161-1461\(2009/08-0129\)](https://doi.org/10.1044/0161-1461(2009/08-0129))
- 8)Pommée, T., Balaguer, M., Maclair, J., Pinquier, J. and Woisard, V., 2022. Intelligibility and comprehensibility: A Delphi consensus study. <https://doi.org/10.1111/1460-6984.12672>
- 9)To, C.K.S., McLeod, S., Sam, K.L. and Law, T., 2022. Predicting Which Children Will Normalize Without Intervention for Speech Sound Disorders. *Journal of Speech, Language, and Hearing Research*, 65(5), pp.1724-1741. https://doi.org/10.1044/2022_JSLHR-21-00444
- 10) Burgoyne K, Lervag A, Malone S, Hulme C. 2019. Speech difficulties at school entry are a significant risk factor for later reading difficulties. *Early Childhood Research Quarterly*. 1 (49), pp. 40-8. <https://doi.org/10.1016/j.ecresq.2019.06.005>
- 11)Jin F, Schjølberg S, Wang MV, Eadie P, Nes RB, Røysamb E, Tambs K. 2020. Predicting literacy skills at 8 years from preschool language trajectories: A population-based cohort study. *Journal of Speech, Language, and Hearing Research*, 63(8), pp. 2752-62. https://doi.org/10.1044/2020_JSLHR-19-00286

- 12)Hustad KC, Mahr TJ, Natzke P, Rathouz PJ. 2021. Speech development between 30 and 119 months in typical children I: Intelligibility growth curves for single-word and multiword productions. *Journal of Speech, Language, and Hearing Research*, 64(10), pp.3707-19.
https://doi.org/10.1044/2021_JSLHR-21-00142
- 13)Shriberg, L.D. and Kwiatkowski, J., 1994. Developmental phonological disorders I: A clinical profile. <https://doi.org/10.1044/jshr.3705.1100>
- 14)Broomfield, J. and Dodd, B., 2004. Children with speech and language disability: caseload characteristics. <https://doi.org/10.1080/13682820310001625589>
- 15)Hayiou-Thomas, M.E., Carroll, J.M., Leavett, R., Hulme, C. and Snowling, M.J., 2017. When does speech sound disorder matter for literacy? The role of disordered speech errors, co-occurring language impairment and family risk of dyslexia. <https://doi.org/10.1111/jcpp.12648>
- 16)Lewis, B.A., Freebairn, L., Tag, J., Ciesla, A.A., Iyengar, S.K., Stein, C.M. and Taylor, H.G., 2015. Adolescent outcomes of children with early speech sound disorders with and without language impairment. https://doi.org/10.1044/2014_AJSLP-14-0075
- 17)Wren, Y., Miller, L.L., Peters, T.J., Emond, A. and Roulstone, S., 2016. Prevalence and predictors of persistent speech sound disorder at eight years old: Findings from a population cohort study. https://doi.org/10.1044/2015_JSLHR-S-14-0282
- 18)Johnson, C.J., Beitchman, J.H. and Brownlie, E.B., 2010. Twenty-year follow-up of children with and without speech-language impairments: Family, educational, occupational, and quality of life outcomes. [https://doi.org/10.1044/1058-0360\(2009/08-0083\)](https://doi.org/10.1044/1058-0360(2009/08-0083))
- 19)Le, H.N., Mensah, F., Eadie, P., McKean, C., Sciberras, E., Bavin, E.L., Reilly, S. and Gold, L., 2021. Health-related quality of life of children with low language from early childhood to adolescence: Results from an Australian longitudinal population-based study. <https://doi.org/10.1111/jcpp.13277>
- 20)St Clair, M.C., Forrest, C.L., Yew, S.G.K. and Gibson, J.L., 2019. Early risk factors and emotional difficulties in children at risk of developmental language disorder: A population cohort study. https://doi.org/10.1044/2018_JSLHR-L-18-0061
- 21)Dodd, B., 2014. Differential diagnosis of pediatric speech sound disorder. <https://doi.org/10.1007/s40474-014-0017-3>
- 22)Petinou, K., Taxitari, L., Phinikettos, I. and Theodorou, E., 2021. Dynamic linguistic interconnectedness and variability in toddlers. <https://doi.org/10.1007/s10936-020-09747-y>
- 23)Stoel-Gammon, C., 2011. Relationships between lexical and phonological development in young children. <https://doi.org/10.1017/S0305000910000425>

- 24) Ainsworth, S., Welbourne, S. and Hesketh, A., 2016. Lexical restructuring in preliterate children: Evidence from novel measures of phonological representation. <https://doi.org/10.1017/S0142716415000338>
- 25) Pascoe, M., Stackhouse, J., and Wells, B. *Persisting Speech Difficulties in Children: Children's Speech and Literacy Difficulties*. Chichester, UK: Whurr Publishers limited 2006: page 383
- 26) Rodgers, L., Harding, S., Rees, R. and Clarke, M.T., 2022. Interventions for pre-school children with co-occurring phonological speech sound disorder and expressive language difficulties: A scoping review. <https://doi.org/10.1111/1460-6984.12719>
- 27) Tyler AA, Lewis KE, Haskill A, Tolbert LC. Efficacy and cross-domain effects of a morphosyntax and a phonology intervention. 2002. *Language, Speech, and Hearing Services in Schools*. [https://doi.org/10.1044/0161-1461\(2002/005\)](https://doi.org/10.1044/0161-1461(2002/005))
- 28) Department for Education. 2012. Exploring interventions for children and young people with speech, language and communication needs: A study of practice. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/219627/DFE-RR247-BCRP13.pdf
- 29) Roulstone, S.E., Marshall, J.E., Powell, G.G., Goldbart, J., Wren, Y.E., Coad, J., Daykin, N., Powell, J.E., Lascelles, L., Hollingworth, W. and Emond, A., 2015. Evidence-based intervention for preschool children with primary speech and language impairments: Child Talk—an exploratory mixed-methods study. <https://doi.org/10.3310/pgfar03050>
- 30) Frizelle, P. and McKean, C., 2022. Using Theory to Drive Intervention Efficacy: The Role of Dose Form in Interventions for Children with DLD. <https://doi.org/10.3390/children9060859>
- 31) Morgan, L., Marshall, J., Harding, S., Powell, G., Wren, Y., Coad, J. and Roulstone, S., 2019. 'It depends': Characterizing speech and language therapy for preschool children with developmental speech and language disorders. <https://doi.org/10.1111/1460-6984.12498>
- 32) Girolametto L, Pearce PS, Weitzman E. 1995. The effects of focused stimulation for promoting vocabulary in young children with delays: A pilot study. *Journal of Children's Communication Development*, 17(2), pp.39-49. <https://doi.org/10.1177/152574019501700205>
- 33) Romeo, R.R., Leonard, J.A., Robinson, S.T., West, M.R., Mackey, A.P., Rowe, M.L. and Gabrieli, J.D., 2018. Beyond the 30-million-word gap: Children's conversational exposure is associated with language-related brain function. <https://doi.org/10.1177/0956797617742725>
- 34) Wasik, B. A., & Jacobi-Vessels, J. L. (2017). Word play: Scaffolding language development through child-directed play. *Early Childhood Education Journal*, 45(6), pp. 769-776. <https://doi.org/10.1007/s10643-016-0827-5>

- 35) Zimmerman, I. L., Steiner, V. G., & Pond, R. E. (2002). *Preschool Language Scale, Fourth Edition (PLS-4)*. <https://doi.org/10.1037/t15140-000>
- 36) McLeod, S., Harrison, L.J. and McCormack, J., 2012. The intelligibility in context scale: Validity and reliability of a subjective rating measure. [https://doi.org/10.1044/1092-4388\(2011/10-0130\)](https://doi.org/10.1044/1092-4388(2011/10-0130))
- 37) Sugden, E., Baker, E., Williams, A.L., Munro, N. and Trivette, C.M., 2020. Evaluation of parent-and speech-language pathologist–delivered multiple oppositions intervention for children with phonological impairment: A multiple-baseline design study. https://doi.org/10.1044/2019_AJSLP-18-0248
- 38) Curran, M. & Owen Van Horne, A., 2019. Use of Recast Intervention to Teach Causal Adverbials to Young Children With Developmental Language Disorder Within a Science Curriculum: A Single Case Design Study. https://doi.org/10.1044/2018_AJSLP-17-0164
- 39) Wren, Y., Harding, S., Goldbart, J. and Roulstone, S., 2018. A systematic review and classification of interventions for speech-sound disorder in preschool children. <https://doi.org/10.1111/1460-6984.12371>
- 40) Frizelle, P., Tolonen, A.K., Tulip, J., Murphy, C.A., Saldana, D. and McKean, C., 2021. The influence of quantitative intervention dosage on oral language outcomes for children with developmental language disorder: A systematic review and narrative synthesis. https://doi.org/10.1044/2020_LSHSS-20-00058
- 41) Carson, L., Baker, E. and Munro, N., 2022. A Systematic Review of Interventions for Late Talkers: Intervention Approaches, Elements, and Vocabulary Outcomes. https://doi.org/10.1044/2022_AJSLP-21-00168
- 42) Murray, E., Power, E., Togher, L., McCabe, P., Munro, N. and Smith, K., 2013. The reliability of methodological ratings for speechBITE using the PEDro-P scale. <https://doi.org/10.1111/1460-6984.12007>
- 43) Tate, R.L., Perdices, M., Rosenkoetter, U., Wakim, D., Godbee, K., Togher, L. and McDonald, S., 2013. Revision of a method quality rating scale for single-case experimental designs and n-of-1 trials: The 15-item Risk of Bias in N-of-1 Trials (RoBiNT) Scale. <https://doi.org/10.1080/09602011.2013.824383>
- 44) Law, J., Charlton, J., Dockrell, J., Gascoigne, M., McKean, C. and Theakston, A., 2017. Early Language Development: Needs, provision and intervention for pre-school children from socio-economically disadvantaged backgrounds. London: Education Endowment Foundation. https://educationendowmentfoundation.org.uk/public/files/Law_et_al_Early_Language_Development_final.pdf
- 45) Hoffmann, T.C., Glasziou, P.P., Boutron, I., Milne, R., Perera, R., Moher, D., Altman, D.G., Barbour, V., Macdonald, H., Johnston, M. and Lamb, S.E., 2014. Better reporting of

interventions: template for intervention description and replication (TIDieR) checklist and guide. <https://doi.org/10.1136/bmj.g1687>

46)Warren, S.F., Fey, M.E. & Yoder, P.J. 2007. Differential treatment intensity research: A missing link to creating optimally effective communication interventions. <https://doi.org/10.1002/mrdd.20139>

47)Campbell, M., McKenzie, J.E., Sowden, A., Katikireddi, S.V., Brennan, S.E., Ellis, S., Hartmann-Boyce, J., Ryan, R., Shepperd, S., Thomas, J. and Welch, V., 2020. Synthesis without meta-analysis (SWiM) in systematic reviews: reporting guideline. <https://doi.org/10.1136/bmj.l6890>

48)Morris C, Simkiss D, Busk M, *et al.*Setting research priorities to improve the health of children and young people with neurodisability: a British Academy of Childhood Disability-James Lind Alliance Research Priority Setting Partnership. [doi: 10.1136/bmjopen-2014-006233](https://doi.org/10.1136/bmjopen-2014-006233)

49)Rodgers, L., and Harford., D. Collaborating with parents to identify outcomes for a preschool speech and language intervention. ESLA Congress 2022: New frontiers in speech and language therapy: advancing practice, research and education. DOI:[10.13140/RG.2.2.34626.45763](https://doi.org/10.13140/RG.2.2.34626.45763)
https://www.researchgate.net/publication/362910019_Collaborating_with_parents_to_identify_outcomes_for_a_pre-school_speech_and_language_intervention

50)Rodgers, L., and Harford., D. 2022. Part of the family. RCSLT Bulletin. Issue 832, autumn 2022.

51)Staniszewska, S., Brett, J., Simera, I., Seers, K., Mockford, C., Goodlad, S., Altman, D.G., Moher, D., Barber, R., Denegri, S. and Entwistle, A., 2017. GRIPP2 reporting checklists: tools to improve reporting of patient and public involvement in research. <https://doi.org/10.1136/bmj.j3453>