



City Research Online

City St George's, University of London

Citation: Magee, L., Goldsmith, L., Chaudhry, U. A. R., Donin, A. S., Wahlich, C., Stovold, E., Nightingale, C. M., Rudnicka, A. R. & Owen, C. G. (2022). Nonpharmacological Interventions to Lengthen Sleep Duration in Healthy Children: A Systematic Review and Meta-analysis. *JAMA Pediatrics*, 176(11), pp. 1084-1097. doi: 10.1001/jamapediatrics.2022.3172

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/31244/>

Link to published version: <https://doi.org/10.1001/jamapediatrics.2022.3172>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

Title: Non-Pharmacological Interventions to Lengthen Sleep Duration in Healthy Children

A Systematic Review and Meta-Analysis

Sub-title: Interventions to Lengthen Sleep Duration in Healthy Children

Dr Lucia Magee, MBChB ¹

Dr Lucy P Goldsmith, PhD ^{1,2}

Dr Umar A R Chaudhry, MBBS ¹

Dr Angela S Donin, PhD ¹

Dr Charlotte Wahlich, DPsych ¹

Ms Elizabeth Stovold, MA ¹

Dr Claire M Nightingale, PhD ¹

Professor Alicja R Rudnicka, PhD ¹

Professor Christopher G Owen, PhD ¹

1. Population Health Research Institute, St George's University of London, Cranmer Terrace, Tooting, London, SW17 0RE
2. Institute for Infection and Immunity, St George's University of London, Cranmer Terrace, Tooting, London, SW17 0RE

Date of revision: 8th June 2022

Word count: Abstract 317 words, manuscript (text only) 3250 words

Corresponding Author:

Professor Christopher G Owen

Address: Population Health Research Institute, St George's University of London, Cranmer Terrace, Tooting, London, SW17 0RE

Tel: +44 (0)20 87252796

Email: cowen@sgul.ac.uk

Key Points

QUESTION Are non-pharmacological interventions effective in lengthening sleep duration in healthy children? If so, what are the key elements?

FINDINGS In this systematic review and meta-analysis of 45 trials, interventions to improve sleep in healthy children were associated with a small increase in sleep duration (by 10 minutes per night). Face-to-face delivery was an important component of interventions, but interventions that included earlier bedtimes were associated with a substantial 47 minutes longer sleep duration per night.

MEANING Supporting children to go to bed earlier can increase sleep duration in healthy children, and interventions to encourage earlier bedtimes should be included.

Abstract

IMPORTANCE Adequate sleep duration is necessary for many aspects of child health, development and wellbeing yet sleep durations for children are declining, and effective strategies to increase sleep in healthy children remain to be elucidated.

OBJECTIVE To determine whether non-pharmaceutical interventions to improve sleep duration in healthy children are effective, and to identify the key components of these interventions.

DATA SOURCES CENTRAL, MEDLINE, Embase, PsycINFO, Web of Science Core collection, ClinicalTrials.gov, and WHO trials databases were searched from inception to 15th November 2021.

STUDY SELECTION Randomised controlled trials of interventions to improve sleep duration in healthy children were independently screened by 2 researchers. A total of 28,478 studies were identified.

DATA EXTRACTION AND SYNTHESIS Data were processed according to the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA). Random effects meta-analytic models were used to estimate pooled effect sizes.

MAIN OUTCOME AND MEASURE The outcome was difference in sleep duration measured in minutes.

RESULTS Pooled results indicate that sleep interventions (45 trials, n=13,539) were associated with 10.5 minutes (95%CI, 5.6 to 15.4) longer nocturnal sleep. There was substantial variation between trials. Sources of variation that were not associated with the study effect size included age group, whether the population was identified as having a sleep problem or being disadvantaged, method of assessment of sleep duration (objective versus subjective), location of intervention delivery (home versus school), whether interventions were delivered in person, used parental involvement, behavioural theory, environmental change, or had greater or lower intensity. Interventions that included earlier bedtimes were associated with a 47-minute sleep extension (3 trials, 95%CI 18.9 to 75.0), compared with remaining studies (42 trials, 7.4 minutes 95%CI 2.9 to 11.8; p=0.006 for group difference). Trials of shorter duration (≤ 6 month) had larger effects.

CONCLUSIONS AND RELEVANCE Interventions focussed on earlier bedtimes may offer a simple, pragmatic, effective way to meaningfully increase sleep duration that could have important benefits for child health.

Introduction

Early childhood experiences and development can have a significant impact on subsequent health, and adequate sleep during childhood is a critical component of wellbeing. Healthy sleep involves a combination of appropriate sleep timing, quality and duration^{1,2}. Sleep in children is vital for appropriate growth, cognitive processing, mental wellbeing, effective social interactions and metabolic health³⁻⁶.

Marked differences in sleep patterns are observed through childhood, with declines in sleep duration with increasing age. Guidance from the US National Sleep Foundation recommends that toddlers aged 1-2 years should sleep 11 to 14 hours per night, pre-school aged children 3-5 years 10 to 13 hours, school aged children 6-13 years 9 to 11 hours, and teenagers 14-17 years 8 to 10 hours,¹ with similar decreasing sleep durations with increasing child age in the UK,⁷ reflecting the changes in circadian rhythm and sleep homeostatic sleep processes during child development. While these recommendations are not derived from systematic reviews and meta-analysis of the evidence, they represent the best available consensus from a multidisciplinary expert panel.¹ Unfortunately, fewer children are achieving these recommendations, with a trend towards shorter sleep durations,⁸⁻¹¹ where currently a third or less of younger children and adolescents are achieving adequate sleep durations.⁸⁻¹⁰ These short sleep durations have been attributed to increased screen time, electronic media use, stimulant intake (particularly caffeine)¹²⁻¹⁹, and increasing emergence of unhealthy sleep behaviours, including poor bedtime routines (such as irregular sleep timings and unsupportive environments)^{20,21}.

Effective strategies are therefore needed to encourage healthy sleep and optimize sleep behaviour in children. Recent systematic reviews and meta-analyses largely of intervention studies have focussed on those with sleep problems and/or medical issues²²⁻²⁴, restricted to narrow age groups^{21,25-28} including infants and pre-schoolers where circadian sleep patterns are not yet fully established²⁶⁻²⁸, have also considered suboptimal non-experimental studies²⁶, and have lacked sufficient evidence to unravel the role of single component versus multi-component interventions as part of broader health improvement agendas²¹. Most reviews to date have also been narrative, which may reflect a lack of standardization in reporting sleep-related outcomes, particularly sleep duration²⁵. Of the few meta-analyses that have quantified overall intervention effects, pooled estimates have been modest at best²⁶.

Increasingly, objective measurements are being used, which limits comparisons with previously reported self- or parentally-reported sleep²⁶.

Interventions in school-aged children may offer an ideal age group to alter nocturnal sleep behaviour, given circadian sleep patterns are well established, and scope to deliver interventions within both school and home settings. Hence, we have carried out an up-to-date systematic review of intervention studies to promote optimal sleep patterns in all children, including those of school age (aged 1 to 18 years).

Aim

The main aim of this systematic review and meta-analysis was to estimate the effectiveness of sleep-related interventions in healthy children. Secondary aims included subgroup exploration to identify which sleep-related intervention components are most or least effective, and to estimate the effect size of these different elements.

Methods

This systematic review and meta-analysis was reported according to PRISMA guidelines²⁹, aligned with the EQUATOR reporting guidelines. The protocol was registered on the international prospective register of systematic reviews (PROSPERO registration CRD42019160089).

Eligibility Criteria

Eligible studies were both individual and cluster RCTs (at both class and school level) investigating interventions to affect sleep duration in children, aged between 1-18 years, without a known medical condition or disability. Studies were excluded if they included: infants under 1 year of age, those with a formal diagnosis of a sleep disorder or use of pharmacological interventions. Pharmacological interventions are not recommended first-line for sleep improvement and differ from behavioural interventions³⁰.

The sleep-related intervention would be stand-alone or part of a multicomponent health promotion strategy, e.g., addressing obesity. The comparator group would be delayed intervention, alternative child health or safety advice, or usual care. The primary outcome was change in mean self-reported, parental-reported, or objectively measured daily total sleep-time (in minutes) in children receiving the intervention compared with controls. Non-English language studies were excluded at the full-text screening stage.

Additional proposed outcomes were change in measures of sleep quality (based on study specific subjective scales) and sleep hygiene (based on study specific sleep hygiene scales). However, due to the heterogeneity of these measures and lack of consistency between studies, it was not possible to meaningfully synthesise these outcomes.

Search Strategy

We carried out searches using both text words and MeSH/index terms in 7 databases: CENTRAL, MEDLINE (Ovid), Embase (Ovid), PsycINFO (Ovid), Web of Science Core Collection, ClinicalTrials.gov, and WHO trials databases.

Searches were carried out from database inception to 15th November 2021. The search strategy used is provided in eTable 1.

We used the Cochrane RCT Classifier³¹ for an initial assessment of the results. Following this, all remaining titles and abstracts were screened independently by two reviewers using Covidence version 1.0³², with discrepancies resolved by a third reviewer. Full text screening was carried out independently by two reviewers in the research group and disagreements were resolved with a third reviewer.

Data Extraction

Study information was extracted using the Covidence data extraction form, edited to include relevant fields specific to this review, including details about the study population, intervention design and delivery. Use of theory in the interventions was assessed using a framework³³ which has been utilised in previous systematic reviews^{34,35}, assessed against 4 levels: informed by theory, applied theory, testing theory and building theory. Data were extracted by two reviewers independently with discrepancies discussed and resolved with a third reviewer (LM,AD,CW), as necessary.

Quantitative data were extracted using a structured data extraction sheet in Microsoft Excel. The data extraction sheet was piloted on 6 trials by (UC) and (CO). Extraction included data in different forms (e.g., means at end-point or change-on-change), using different measurement methods (e.g. actigraphy or self-/parental-report) related to sleep at different times of the week (e.g., weekend or weekday) and at different follow-up periods. Quantitative data extraction was undertaken independently by two reviewers with discrepancies discussed and resolved with a third reviewer where necessary (UC,LG,AR). Due to reporting inconsistency of intervention length and follow-up period, these were combined under study period.

Study Quality

Cochrane's 'Risk of Bias' tool³⁶ was chosen over alternative instruments as it is sensitive to variation in the quality of randomised controlled trials. Two reviewers independently scored the domains (selection bias, performance bias, detection bias, attrition bias, reporting bias, and other biases) prior to data analysis, with discrepancies resolved by a third reviewer (LM,CW,AD).

Data Analysis

We used Cochrane's Review Manager (version 5.4)³⁷ to calculate effect estimates (in minutes) and associated 95% confidence intervals (CI), using a random effects mean difference model. Trials were excluded if the number of participants at follow up could not be established. The comparator group was split to avoid double counting in three-arm trials, and where a study did not report pooled results for the entire sample, effects reported were combined using a fixed effects meta-analysis. Adjusted effect estimates were preferred over unadjusted estimates, and the longest follow-up period was selected for inclusion where multiple follow-up periods were reported. Data for weekday sleep were used in preference to weekend sleep.

Further analyses were performed in STATA version 17 (using *metan* and *metareg* commands)³⁸. Subgroup analyses, examining differences in effect estimates for different populations (e.g., by age group, continent, level of disadvantage, self-identified sleep problem), and study design features (e.g., method of measuring of sleep duration, length of intervention, location and mode of delivery) were performed (using Z-tests). Statistical heterogeneity was assessed using the I^2 statistic, and p-values. An I^2 value of 50-75% suggested moderate heterogeneity, and 75% or above high heterogeneity³⁹. We assessed publication bias by visual examination of funnel plots, and using Begg and Egger tests, overall and by subgroups.

Results

A total of 28,478 studies were identified, of which 8,854 were duplicates. The RCT classifier excluded 5,460 records and a further 436 duplicates were identified resulting in 13,728 studies to be screened. After screening of titles and abstracts, 411 potentially relevant full-text articles were screened. Figure 1 shows the article selection process. In total, 45 RCTs with 47 estimates of mean differences in sleep duration between intervention and control groups were included in the analyses (with 13,539 child participants, mean age range 18 months to 19 years; 6,897 in the

intervention group, 6,642 controls). The key characteristics of selected trials and interventions are shown in Table 1 (more detailed characteristics in eTable 2, study estimates in eTable 3). Of 45 trials, 10 were in pre-school age, 11 primary school age and 26 in secondary school age.

The primary analysis found that sleep interventions were associated with 10.5 minutes (95% CI 5.6 to 15.4 mins, $p < 0.0001$) longer nocturnal sleep duration compared with the control condition. Figure 2 shows the primary meta-analysis forest plot, with quality assessments. There was marked heterogeneity across study estimates ($I^2 = 87.2\%$). All subgroup comparisons, by population characteristics, study design features and intervention features are summarised in Table 2.

Differences in sleep duration by study population characteristics

There were no significant differences in sleep duration estimates across different pre-school, primary and secondary school age groups, presence of social disadvantage, or whether the study population included those reporting sleep problems. Most trials were carried out in high income countries in children of largely white European ancestry (i.e., 15 trials were carried out in North America, 12 trials Australasia, 9 in Europe); the remaining trials were carried out in Asia (7 trials) and South America (4 trials). There were significant differences in the effect size estimates by study geography ($p = 0.015$); pooled estimates from North American and Asian trials appeared to show the largest difference in sleep duration, 15.6, 95%CI 3.8 to 27.5 mins and 16.9 95% CI 4.5 to 29.4 respectively), compared with pooled trials from Australasia and South America that showed no change (-0.6, 95% CI -5.8 to 4.7 and 5.0, 95% CI -1.2 to 11.0 mins, Table 2) with wide confidence intervals.

Differences in sleep duration by study design features

There were no overall differences in sleep duration when measured using objective (accelerometer or electroencephalography) vs subjective (sleep diaries, child or parental report) methods. However, there were considerable differences in sleep duration among the trials relying on subjective reports ($p < 0.001$), where those using parental-report were associated with a very small change in sleep (0.4, 95%CI -3.4 to 4.3 mins), those relying on child report were associated with a non-significant difference in sleep duration (7.8, 95%CI -2.3 to 17.7 mins), whereas trials using sleep diaries were associated with a large difference in sleep duration (26.4, 95% CI 17.7 to 35.2). Estimates from trials recording data on 7 nights per week (all week nights) were similar to those recording data 5 nights per week (weekday nights); 11.9 minutes (95%CI 4.0 to 19.7) vs 10.3 minutes (95%CI 2.5 to 18.2),

respectively. Two trials only reported data for weekend nights, reporting a lower effect estimate (2.9, 95%CI -0.1 to 5.8); the difference between subgroups was significant ($p=0.035$). Trials that reported final follow-up data within 6 months of the baseline reported significantly greater effect sizes than those reporting final follow-up beyond 6 months.

Differences in sleep duration by intervention features

There were no statistically significant differences in sleep duration estimates by the location of intervention delivery (e.g., home or school), with parental involvement in the intervention, or in whether the intervention was delivered face-to-face. Trials which focussed on lengthening sleep alone showed significantly more change in sleep duration (15.8, 95%CI 8.9 to 22.6 minutes) than interventions with broader aims (1.2, 95%CI -2.5 to 5.0 minutes; $p<0.0001$). There were no statistically significant differences in the effects of trials by intensity of intervention delivery, use of environmental change to support longer sleep, or whether behavioural theory was used to inform intervention design.

Trials which focussed on obesity had a smaller effect; 3.3 minutes, (95%CI -4.8 to 11.3), compared with other trials 12.8 minutes, (95%CI 7.0 to 18.7); ($p=0.06$). Similarly, trials with a physical activity component had a significantly smaller effect size compared with those with no physical activity component; 1.6 minutes (95%CI -1.2 to 4.5) vs 13.4 minutes (95%CI 7.1 to 19.6), respectively ($p=0.001$). Trials which involved an earlier bedtime had a significant and substantially greater effect estimate (47.0 minutes, 95%CI 18.9 to 75) compared with the effect of trials which did not encourage participants to go to bed earlier (7.4 minutes, 95% CI 2.9 to 11.8; $p=0.006$). The effect of early bedtimes was in fewer studies (3 trials), but the effect appeared similar in studies carried out in primary school aged children, and 2 studies solely in secondary school aged children ($p=0.15$).

Heterogeneity in subgroup analyses varied between 0% and 95.5%.

Possible effect of bias and other factors on differences in estimates of sleep duration

Visual inspection did not reveal any clear pattern of small study / publication bias overall (Egger test=0.85); There was no clear pattern of small study bias by sub-groups. Equally, risk of bias assessments, including selection bias, performance bias, detection bias, attrition bias, reporting bias, and other biases (shown in Figure 2), did not provide notable patterns in the data. There was no evidence that differences in sleep duration between intervention and

control differed by quartile groups of baseline levels of sleep duration ($p=0.57$). Moreover, there was no evidence of a systematic difference in unadjusted versus adjusted sleep duration analyses, both at a study ($n=45$) or estimate ($n=58$) level ($p=0.34, 0.48$, respectively).

Discussion

This systematic review found that sleep interventions in children aged 1 to 18 years were associated with 10.5 minutes longer sleep duration per night. Many intervention components have small effects on sleep duration. Substantial increases in sleep duration were observed in a subgroup of trials encouraging earlier bedtimes, with objectively measured sleep durations increasing by 47 minutes. Interventions which have a broader focus (e.g., interventions which focus on obesity) were ineffective in increasing sleep duration, extending sleep by 1.2 minutes, compared with 16 minutes in interventions focussed on sleep only.

The pooled effect estimate of 10.5 minutes longer sleep duration per night is similar to previous reviews which included fewer trials with correspondingly fewer participants²⁶, or narrower age groups (e.g., infants, or infants and pre-school children only²⁶⁻²⁸). While a recent meta-analysis of older children and young adults (12 to 25 years) found a larger intervention effect (35 mins), the pooled estimate was associated with wide confidence intervals (95% CI 8.70, 61.14) inclusive of our overall finding⁴⁰. Notably, the pooled effect estimate from that review is similar to the estimate reported in the present review for trials that encouraged earlier bed-times, and importantly employed objective actigraphy measurement of sleep duration (with 2 trials being used in both reviews^{41,42}), which limits well-known biases associated with self-or parentally-reported sleep times²⁶. While prior qualitative reviews have suggested that multi-component interventions, underpinned by behavioural theory might be more effective²⁵, our findings (in agreement with another previous review²⁶) demonstrate that simple sleep-focused interventions are more effective.

We included a large number of RCTs (45 trials with 13,539 participants) which provide the best available evidence of potential intervention. This review spans the entirety of childhood (excluding infants <1 year) and finds no clear evidence of differences in overall intervention effects by age. Most trials included in this review were carried out in primary- and secondary-school aged children, where circadian sleep patterns and nocturnal sleep behaviour are well established. The trials included delivered interventions in both school and home settings, and no difference was found in the effect sizes of interventions in either setting. These findings suggest that reinforcement of interventions

in both settings could be used for future implementation. Differences in individual study results were reflected by high levels of heterogeneity in both the main and sub-group analyses.

There were inconsistencies measuring sleep related outcomes across trials. Sleep duration was the most commonly reported outcome, but there was variation in the analytical approach, with some trials reporting mean sleep durations by intervention and control groups, others mean differences in sleep duration between groups at follow-up, or differences-in-differences (with differing levels of adjustment); although, there was no evidence of systematic differences across these groups. Other important sleep-related outcomes beyond duration, such as sleep quality and latency (not reported here), were inconsistently and less frequently reported, which presents challenges for appropriately pooling results. Similarly, other intervention characteristics were not routinely reported and while interventions underpinned by theory did not appear to be associated with longer sleep durations, more detailed reporting of the use of theory (including behavioural models underpinning proposed interventions) and study fidelity (including uptake and compliance with a given intervention) could usefully contribute to the evidence by allowing effective components to be more readily identified, which would support the development of novel interventions.

While an overall 10-minute increase in sleep duration would have a modest effect in addressing declining sleep durations in children⁸⁻¹¹, a 47-minute increase associated with earlier bedtimes would substantially impact on this population-level problem. However, while the difference in sleep duration among trials employing earlier bedtimes was much larger, this was only based on 3 studies (1 trial in primary school aged⁴³ and 2 trials in secondary school aged children^{42,44}), largely for the USA^{42,43}, and these findings need to be replicated in further studies and settings. If confirmed, earlier bedtimes are needed given wake-up times are less changeable, especially on schooldays. While one randomised controlled trial from Israel included in our review showed 55 minutes longer sleep duration by delaying school start times by 1 hour within the first week, this was not sustained.⁴⁵ Further high quality experimental evidence is needed,⁴⁶ especially as interventions to shift wake-up times by delaying school start times in a UK context have been shown to be impractical.⁴⁷ However, recent observational evidence from the USA has shown that increased sleep duration (by as much as 40 mins) could be achieved by delaying school start times, suggesting that such an approach might be feasible^{48,49}. In our study, control group children achieved only 8.6 hours sleep overall, with 10.4 hours in 10 trials of pre-school children, 9.2 hours in 9 trials of primary school-aged children, and as little as 7.6 hours in 22 trials of secondary school-aged children. These durations are much shorter than

current recommended levels of 10-11 hours in 4 to 12 year olds, and 8-10 hours in 13 to 18 year olds¹. Lengthening time asleep by 47-minutes among those with inadequate sleep duration could not only address these shortfalls but could also have important benefits for child health. Observational evidence in 8 to 10-year olds suggests that such increases could have an appreciable impact on cardiometabolic health (e.g., 0.1 kg/m² lower BMI, and 2% reduction in insulin resistance)⁵⁰, as well as having potential benefits for mental health and well-being⁶. Patterning sleep behaviour from early age could also track into adulthood, with potential benefits for health in later life. Currently the only studies encouraging earlier bedtimes within the age ranges of this study were conducted largely in adolescents. Further research is urgently needed to establish whether similarly large effect sizes are observed in sleep interventions encouraging earlier bedtimes in younger children, and, moving beyond this, to establish whether the effects can be sustained. Additional research is also needed to clarify the long-term benefits of such interventions on health outcomes.

Conclusion

Typically, non-pharmacological interventions increase sleep by a small amount (10.5 minutes). Intervention elements, such as a narrow focus on sleep including earlier bed times should be taken forward in terms of study design to help children achieve adequate sleep to support growth, cognitive development, mental wellbeing and metabolic health. Based on the best available evidence to date, earlier bedtimes offer a simple, pragmatic strategy to meaningfully increase sleep duration.

Author Contributions

LM was involved in data collection, data interpretation, project administration, preparation of figures for publication, writing the original draft, and ongoing review & editing. LG contributed to data collection, data analysis, statistical analysis, data interpretation, preparation of figures for publication, writing the original draft, and ongoing review & editing. UC contributed to data collection, data analysis, statistical analysis, data interpretation, manuscript writing, review & editing. AD contributed to data collection, data interpretation, manuscript review & editing. CW contributed to, data collection, data interpretation, manuscript review & editing. ES contributed to data searches, manuscript review & editing. CMN contributed to data collection, data interpretation, statistical analysis, manuscript review & editing. AR conceptualized and supervised the study, and led data collection, data interpretation, study design, validation of results, manuscript review & editing. CGO conceptualized and supervised the study, and led

data collection, data interpretation, study design, validation of results, writing the original draft, and ongoing review & editing. The corresponding author had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. All authors shared the final responsibility for the decision to submit for publication.

Declaration of Interests

All authors declare no competing interests.

Funding/Support Statement

No funding. The sponsor had no role in design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

References

1. Hirshkowitz M, Whiton K, Albert SM, et al. National sleep foundation's sleep time duration recommendations: Methodology and results summary. *Sleep Health*. 2015;1(1):40-43.
2. Ohayon M, Wickwire EM, Hirshkowitz M, et al. National Sleep Foundation's sleep quality recommendations: first report. *Sleep Health*. 2017;3(1):6-19.
3. Cappuccio FP, D'Elia L, Strazzullo P, Miller MA. Quantity and quality of sleep and incidence of type 2 diabetes: a systematic review and meta-analysis. *Diabetes Care*. 2010;33(2):414-420.
4. Fallone G, Owens JA, Deane J. Sleepiness in children and adolescents: clinical implications. *Sleep medicine reviews*. 2002;6(4):287-306.
5. Gozal D, Kheirandish-Gozal L. Neurocognitive and behavioral morbidity in children with sleep disorders. *Current opinion in pulmonary medicine*. 2007;13(6):505-509.
6. Short MA, Bartel K, Carskadon MA. Chapter 32 - Sleep and mental health in children and adolescents. In: Grandner MA, ed. *Sleep and Health*. Academic Press; 2019:435-445.
7. NHS. How much sleep do children need? <https://www.nhs.uk/live-well/sleep-and-tiredness/how-much-sleep-do-kids-need/>. Published 2020. Accessed.
8. Guan H, Zhang Z, Wang B, et al. Proportion of kindergarten children meeting the WHO guidelines on physical activity, sedentary behaviour and sleep and associations with adiposity in urban Beijing. *BMC Pediatrics*. 2020;20(1):70.
9. Keyes KM, Maslowsky J, Hamilton A, Schulenberg J. The great sleep recession: changes in sleep duration among US adolescents, 1991-2012. *Pediatrics*. 2015;135(3):460-468.
10. Chaput J-P, Janssen I. Sleep duration estimates of Canadian children and adolescents. *Journal of Sleep Research*. 2016;25(5):541-548.
11. Singh GK, Kenney MK. Rising Prevalence and Neighborhood, Social, and Behavioral Determinants of Sleep Problems in US Children and Adolescents, 2003-2012. *Sleep disorders*. 2013;2013:394320-394320.
12. Hysing M, Pallesen S, Stormark KM, Jakobsen R, Lundervold AJ, Sivertsen B. Sleep and use of electronic devices in adolescence: results from a large population-based study. *BMJ Open*. 2015;5(1):e006748.
13. Dube N, Khan K, Loehr S, Chu Y, Veugelers P. The use of entertainment and communication technologies before sleep could affect sleep and weight status: a population-based study among children. *The international journal of behavioral nutrition and physical activity*. 2017;14(1):97.
14. Hale L, Guan S. Screen time and sleep among school-aged children and adolescents: a systematic literature review. *Sleep medicine reviews*. 2015;21:50-58.

15. Cain N, Gradisar M. Electronic media use and sleep in school-aged children and adolescents: A review. *Sleep medicine*. 2010;11(8):735-742.
16. Bruni O, Sette S, Fontanesi L, Baiocco R, Laghi F, Baumgartner E. Technology Use and Sleep Quality in Preadolescence and Adolescence. *Journal of Clinical Sleep Medicine*. 2015;11(12):1433-1441.
17. Gamble AL, D'Rozario AL, Bartlett DJ, et al. Adolescent Sleep Patterns and Night-Time Technology Use: Results of the Australian Broadcasting Corporation's Big Sleep Survey. *PLOS ONE*. 2014;9(11):e111700.
18. Ahluwalia N, Herrick K, Moshfegh A, Rybak M. Caffeine intake in children in the United States and 10-y trends: 2001-2010. *Am J Clin Nutr*. 2014;100(4):1124-1132.
19. Temple JL. Review: Trends, Safety, and Recommendations for Caffeine Use in Children and Adolescents. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2019;58(1):36-45.
20. Allen SL, Howlett MD, Coulombe JA, Corkum PV. ABCs of SLEEPING: A review of the evidence behind pediatric sleep practice recommendations. *Sleep medicine reviews*. 2016;29:1-14.
21. Belmon LS, van Stralen MM, Busch V, Harmsen IA, Chinapaw MJM. What are the determinants of children's sleep behavior? A systematic review of longitudinal studies. *Sleep medicine reviews*. 2019;43:60-70.
22. Meltzer LJ, Mindell JA. Systematic Review and Meta-Analysis of Behavioral Interventions for Pediatric Insomnia. *Journal of Pediatric Psychology*. 2014;39(8):932-948.
23. Nikles J, Mitchell GK, de Miranda Araújo R, et al. A systematic review of the effectiveness of sleep hygiene in children with ADHD. *Psychology, health & medicine*. 2020;25(4):497-518.
24. Keogh S, Bridle C, Siriwardena NA, et al. Effectiveness of non-pharmacological interventions for insomnia in children with Autism Spectrum Disorder: A systematic review and meta-analysis. *PloS one*. 2019;14(8):e0221428-e0221428.
25. Busch V, Altenburg TM, Harmsen IA, Chinapaw MJ. Interventions that stimulate healthy sleep in school-aged children: a systematic literature review. *European journal of public health*. 2017;27(1):53-65.
26. Fangupo LJ, Haszard JJ, Reynolds AN, et al. Do sleep interventions change sleep duration in children aged 0-5 years? A systematic review and meta-analysis of randomised controlled trials. *Sleep medicine reviews*. 2021;59:101498.
27. Kempner L, Sharpe L, Miller CB, Bartlett DJ. Do psychosocial sleep interventions improve infant sleep or maternal mood in the postnatal period? A systematic review and meta-analysis of randomised controlled trials. *Sleep medicine reviews*. 2016;29:15-22.
28. Bryanton J, Beck CT, Montelpare W. Postnatal parental education for optimizing infant general health and parent-infant relationships. *The Cochrane database of systematic reviews*. 2013(11):Cd004068.
29. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Bmj*. 2021;372:n71.
30. Wilson S, Anderson K, Baldwin D, et al. British Association for Psychopharmacology consensus statement on evidence-based treatment of insomnia, parasomnias and circadian rhythm disorders: An update. *Journal of Psychopharmacology*. 2019;33(8):923-947.
31. Wallace BC, Noel-Storr A, Marshall IJ, Cohen AM, Smalheiser NR, Thomas J. Identifying reports of randomized controlled trials (RCTs) via a hybrid machine learning and crowdsourcing approach. *Journal of the American Medical Informatics Association*. 2017;24(6):1165-1168.
32. Covidence systematic review software VHI. Covidence systematic review software. www.covidence.org. Published 2021. Accessed.
33. Painter JE, Borba CP, Hynes M, Mays D, Glanz K. The use of theory in health behavior research from 2000 to 2005: a systematic review. *Annals of behavioral medicine : a publication of the Society of Behavioral Medicine*. 2008;35(3):358-362.
34. Pang B, Kubacki K, Rundle-Thiele S. Promoting active travel to school: a systematic review (2010–2016). *BMC Public Health*. 2017;17(1):638.
35. Hurley E, Dietrich T, Rundle-Thiele S. A systematic review of parent based programs to prevent or reduce alcohol consumption in adolescents. *BMC Public Health*. 2019;19(1):1451.
36. Higgins JPT, Altman DG, Gøtzsche PC, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*. 2011;343:d5928.
37. *Review Manager (RevMan)* [computer program]. Version Version 5.42020.
38. *Stata Statistical Software* [computer program]. Version Release 15. College Station, TX: StataCorp LLC.2017.
39. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *Bmj*. 2003;327(7414):557-560.

40. Griggs S, Conley S, Batten J, Grey M. A systematic review and meta-analysis of behavioral sleep interventions for adolescents and emerging adults. *Sleep medicine reviews*. 2020;54:101356.
41. Dewald-Kaufmann JF, Oort FJ, Meijer AM. The effects of sleep extension on sleep and cognitive performance in adolescents with chronic sleep reduction: an experimental study. *Sleep medicine*. 2013;14(6):510-517.
42. Van Dyk TR, Zhang N, Catlin PA, et al. Feasibility and emotional impact of experimentally extending sleep in short-sleeping adolescents. *Sleep*. 2017;40(9).
43. Hart C, Hawley N, Egleston B, et al. 0995 BRIEF BEHAVIORAL INTERVENTION ENHANCES CHILDREN'S SLEEP AND IMPROVES WEIGHT STATUS. *Sleep*. 2017;40(suppl_1):A370-A370.
44. Dewald-Kaufmann JF, Oort FJ, Meijer AM. The effects of sleep extension on sleep and cognitive performance in adolescents with chronic sleep reduction: an experimental study. *Sleep medicine*. 2013;14(6):510-517.
45. Lufi D, Tzischinsky O, Hadar S. Delaying school starting time by one hour: some effects on attention levels in adolescents. *Journal of clinical sleep medicine*. 2011;7(2):137-143.
46. Minges KE, Redeker NS. Delayed school start times and adolescent sleep: a systematic review of the experimental evidence. *Sleep medicine reviews*. 2016;28:86-95.
47. Illingworth G, Sharman R, Jowett A, Harvey CJ, Foster RG, Espie CA. Challenges in implementing and assessing outcomes of school start time change in the UK: experience of the Oxford Teensleep study. *Sleep medicine*. 2019;60:89-95.
48. Widome R, Berger AT, Iber C, et al. Association of Delaying School Start Time With Sleep Duration, Timing, and Quality Among Adolescents. *JAMA Pediatrics*. 2020;174(7):697-704.
49. Bowers JM, Moyer A. Effects of school start time on students' sleep duration, daytime sleepiness, and attendance: a meta-analysis. *Sleep Health*. 2017;3(6):423-431.
50. Rudnicka AR, Nightingale CM, Donin AS, et al. Sleep Duration and Risk of Type 2 Diabetes. *Pediatrics*. 2017;140(3).
51. Beijamini F, Louzada FM. Are educational interventions able to prevent excessive daytime sleepiness in adolescents? *Biological Rhythm Research*. 2012;43(6):603-613.
52. Blake M, Waloszek JM, Schwartz O, et al. The SENSE Study: post Intervention Effects of a Randomized Controlled Trial of a Cognitive-Behavioral and Mindfulness-Based Group Sleep Improvement Intervention among At-Risk Adolescents. *Journal of consulting and clinical psychology*. 2016;84(12):1039-1051.
53. Delli Bovi AP, Manco Cesari G, Rocco MC, et al. Healthy Lifestyle Management of Pediatric Obesity with a Hybrid System of Customized Mobile Technology: The PediaFit Pilot Project. *Nutrients*. 2021;13(2).
54. Cain N, Gradisar M, Moseley L. A motivational school-based intervention for adolescent sleep problems. *Sleep medicine*. 2011;12(3):246-251.
55. Cepni AB, Taylor A, Crumbley C, et al. Feasibility and Efficacy of the "FUNPALs Playgroup" Intervention to Improve Toddler Dietary and Activity Behaviors: A Pilot Randomized Controlled Trial. *Int J Environ Res Public Health*. 2021;18(15).
56. Dong L, Dolsen MR, Martinez AJ, Notsu H, Harvey AG. A transdiagnostic sleep and circadian intervention for adolescents: six-month follow-up of a randomized controlled trial. *J Child Psychol Psychiatry*. 2020;61(6):653-661.
57. Haines J, McDonald J, O'Brien A, et al. Healthy Habits, Happy Homes: randomized trial to improve household routines for obesity prevention among preschool-aged children. *JAMA pediatrics*. 2013;167(11):1072-1079.
58. Hammersley ML, Okely AD, Batterham MJ, Jones RA. An Internet-Based Childhood Obesity Prevention Program (Time2bHealthy) for Parents of Preschool-Aged Children: randomized Controlled Trial. *Journal of medical Internet research*. 2019;21(2):e11964.
59. Hammersley ML, Wyse RJ, Jones RA, et al. Translation of Two Healthy Eating and Active Living Support Programs for Parents of 2-6-Year-Old Children: Outcomes of the 'Time for Healthy Habits' Parallel Partially Randomised Preference Trial. *Nutrients*. 2021;13(10).
60. Hart CN, Hawley NL, Wing RR. Development of a Behavioral Sleep Intervention as a Novel Approach for Pediatric Obesity in School-aged Children. *Pediatr Clin North Am*. 2016;63(3):511-523.
61. Hiscock H, Quach J, Paton K, et al. Impact of a Behavioral Sleep Intervention on New School Entrants? Social Emotional Functioning and Sleep: a Translational Randomized Trial. *Behavioral sleep medicine*. 2019;17(6):698-712.
62. Kalak N, Gerber M, Kirov R, et al. Daily morning running for 3 weeks improved sleep and psychological functioning in healthy adolescents compared with controls. *The Journal of adolescent health : official publication of the Society for Adolescent Medicine*. 2012;51(6):615-622.

63. Kira G, Maddison R, Hull M, Blunden S, Olds T. Sleep education improves the sleep duration of adolescents: a randomized controlled pilot study. *Journal of clinical sleep medicine*. 2014;10(7):787-792.
64. Knebel MTG, Borgatto AF, Lopes MVV, et al. Mediating role of screen media use on adolescents' total sleep time: A cluster-randomized controlled trial for physical activity and sedentary behaviour. *Child Care Health Dev*. 2020;46(3):381-389.
65. Lin CY, Strong C, Scott AJ, Brostrom A, Pakpour AH, Webb TL. A cluster randomized controlled trial of a theory-based sleep hygiene intervention for adolescents. *Sleep*. 2018;41(11) (no pagination).
66. Marsh S, Taylor R, Galland B, Gerritsen S, Parag V, Maddison R. Results of the 3 Pillars Study (3PS), a relationship-based programme targeting parent-child interactions, healthy lifestyle behaviours, and the home environment in parents of preschool-aged children: A pilot randomised controlled trial. *PLoS One*. 2020;15(9):e0238977.
67. Mindell JA, Telofski LS, Wiegand B, Kurtz ES. A nightly bedtime routine: impact on sleep in young children and maternal mood. *Sleep*. 2009;32(5):599-606.
68. Mindell JA, Sedmak R, Boyle JT, Butler R, Williamson AA. Sleep Well!: a pilot study of an education campaign to improve sleep of socioeconomically disadvantaged children. *Journal of clinical sleep medicine*. 2016;12(12):1593-1599.
69. Mitchell JA, Morales KH, Williamson AA, et al. Engineering a mobile platform to promote sleep in the pediatric primary care setting. *Sleep Adv*. 2021;2(1):zpab006.
70. Moore SM, Borawski EA, Love TE, et al. Two family interventions to reduce BMI in low-income urban youth: a randomized trial. *Pediatrics*. 2019;143 (6) (no pagination)(e20182185).
71. Morell-Azanza L, Ojeda-Rodriguez A, Ochotorena-EliceGUI A, et al. Changes in objectively measured physical activity after a multidisciplinary lifestyle intervention in children with abdominal obesity: a randomized control trial. *BMC pediatrics*. 2019;19 (1) (no pagination)(90).
72. Moseley L, Gradisar M. Evaluation of a school-based intervention for adolescent sleep problems. *Sleep*. 2009;32(3):334-341.
73. Moula Z, Powell J, Karkou V. An Investigation of the Effectiveness of Arts Therapies Interventions on Measures of Quality of Life and Wellbeing: A Pilot Randomized Controlled Study in Primary Schools. *Front Psychol*. 2020;11:586134.
74. Mousarrezaei Z, Valizadeh L, Alizadeh M, Aghajari P, Jafarabadi MA, Janani R. The effects of education through short message service for mothers on sleep duration among school-aged children: A randomized trial. *Nurs Midwifery Stud*. 2020;9:9-15.
75. Pablos A, Nebot V, Vano-Vicent V, Ceca D, Elvira L. Effectiveness of a school-based program focusing on diet and health habits taught through physical exercise. *Physiologie appliquee, nutrition et metabolisme [Applied physiology, nutrition, and metabolism]*. 2018;43(4):331-337.
76. Puder JJ, Marques-Vidal P, Schindler C, et al. Effect of multidimensional lifestyle intervention on fitness and adiposity in predominantly migrant preschool children (Ballabeina): cluster randomised controlled trial. *BMJ (Clinical research ed)*. 2011;343:d6195.
77. Quach J, Spencer-Smith M, Anderson PJ, Roberts G. Can working memory training improve children's sleep? *Sleep medicine*. 2018;47:113-116.
78. Rigney G, Blunden S, Maher C, et al. Can a school-based sleep education programme improve sleep knowledge, hygiene and behaviours using a randomised controlled trial. *Sleep medicine*. 2015;16(6):736-745.
79. Santiago LCS, Lyra MJ, Germano-Soares AH, et al. Effects of Strength Training on Sleep Parameters of Adolescents: A Randomized Controlled Trial. *J Strength Cond Res*. 2020.
80. Sousa IC, Souza JC, Louzada FM, Azevedo CVM. Changes in sleep habits and knowledge after an educational sleep program in 12th grade students. *Sleep and biological rhythms*. 2013;11:144-153.
81. Sundgot-Borgen C, Friberg O, Kolle E, et al. Does the Healthy Body Image program improve lifestyle habits among high school students? A randomized controlled trial with 12-month follow-up. *J Int Med Res*. 2020;48(3):300060519889453.
82. Tamura N, Tanaka H. Effects of sleep education with self-help treatment for elementary schoolchild with nocturnal lifestyle and irritability. *Sleep and biological rhythms*. 2014;12:169-179.
83. Tamura N, Tanaka H. Effects of a sleep education program with self-help treatment on sleeping patterns and daytime sleepiness in Japanese adolescents: a cluster randomized trial. *Chronobiology international*. 2016;33(8):1073-1085.
84. Taylor RW, Cox A, Knight L, et al. A Tailored Family-Based Obesity Intervention: A Randomized Trial. *Pediatrics*. 2015;136(2):281-289.

85. Tomayko EJ, Prince RJ, Cronin KA, Kim K, Parker T, Adams AK. The Healthy Children, Strong Families 2 (HCSF2) Randomized Controlled Trial Improved Healthy Behaviors in American Indian Families with Young Children. *Current developments in nutrition*. 2019;3(Suppl 2):53-62.
86. Uhlig S, Groot J, Jansen E, Scherder E. Rap & Sing Music Therapy and sleep in adolescents: a single-blind cluster randomized controlled trial. *Nordic journal of music therapy*. 2019;28(1):60-70.
87. van Rijn E, Koh SYJ, Ng ASC, et al. Evaluation of an interactive school-based sleep education program: a cluster-randomized controlled trial. *Sleep Health*. 2020;6(2):137-144.
88. Walton K, Filion AJ, Gross D, et al. Parents and tots together: pilot randomized controlled trial of a family-based obesity prevention intervention in canada. *Canadian journal of public health-revue canadienne de sante publique*. 2015;106(8):e555-e562.
89. Wilson KE, Miller AL, Bonuck K, Lumeng JC, Chervin RD. Evaluation of a sleep education program for low-income preschool children and their families. *Sleep*. 2014;37(6):1117-1125.
90. Wing YK, Chan NY, Yu MWM, et al. A school-based sleep education program for adolescents: A cluster randomized trial. *Pediatrics*. 2015;135(3):e635-e643.
91. Wolfson AR, Harkins E, Johnson M, Marco C. Effects of the Young Adolescent Sleep Smart Program on sleep hygiene practices, sleep health efficacy, and behavioral well-being. *Sleep health*. 2015;1(3):197-204.
92. Yoong SL, Grady A, Stacey F, et al. A pilot randomized controlled trial examining the impact of a sleep intervention targeting home routines on young children's (3-6 years) physical activity. *Pediatric obesity*. 2019;14(4):e12481.
93. Blake MJ, Blake LM, Schwartz O, et al. Who benefits from adolescent sleep interventions? Moderators of treatment efficacy in a randomized controlled trial of a cognitive-behavioral and mindfulness-based group sleep intervention for at-risk adolescents. *Journal of child psychology and psychiatry, and allied disciplines*. 2018;59(6):637-649.

Figure 1: Study selection process





Figure 2: Forest plot showing mean differences in sleep duration (minutes) in 45 studies, comparing sleep intervention groups with control



Key: Mean differences in sleep duration (minutes) for each study, comparing duration in those receiving a sleep intervention versus control. Point estimate shown by blue solid squares, with horizontal lines showing the 95% CI. The study authors are indicated on the y-axis in alphabetical order. The pooled estimate based on a random-effects model is shown with a blue diamond (95% CI). Risk of bias legend: (A) Random sequence generation (selection bias), (B) Allocation concealment (selection bias), (C) Blinding of participants and personnel (performance bias), (D) Blinding of outcome assessment (detection bias), (E) Incomplete outcome data (attrition bias), (F) Selective reporting (reporting bias), (G) Other bias. Risk of bias criteria graded from green '+' low risk, yellow '?' unclear, or red '-' high risk.

Table 1: Study Characteristics

Name	Study title	Age range/mean	Target population	Setting ^a	Inclusion criteria	Mode of delivery	Intervention overview ^{b,c}	Dose	Intervention duration	Study duration
Bejamini 2012 ⁵¹	Educational Programme for Sleep (EPS)	13-14 years	School students in Public High schools	School	General population	Face-to-face	Educational programme of 4 themed lessons on sleep ^b	4x50min lessons, over 5 days	<1week	3-weeks
Blake 2016 ⁵²	Study Sense/Sleep Sense	12-16 years	Students in year 7-10 at risk for major depressive disorder	Secondary schools	High levels of anxiety and sleeping difficulties but without a depressive disorder	Face-to-face	Sleep education programme, including 7 sessions & information book	7x90min sessions	7-weeks	9-weeks
Bovi 2021 ⁵³	PediaFit	6-14 years	Obese children	Pediatric Obesity Clinic in University Hospital	BMI >95th percentile for age and sex according to the CDC 2000 growth curves	Face-to-face group sessions and Whatsapp texts	Clinician led groups session - health nutrition/physical activity	Group session with 3 messages a week for 24 weeks Or as above + monthly in presence visits (4 over 6 months)	24-weeks	9-months
Cain 2011 ⁵⁴	Motivational sleep intervention	16-17 years	Psychology students	Secondary schools	Students with discrepant weekday-week-end out-of-bed times (>2 h) and insufficient sleep on school nights (<8 h)	Face-to-face	Sleep education classes & motivational interviewing framework ^b	4x50min sessions	4-weeks	6-weeks
Cepni 2021 ⁵⁵	Families Understanding Nutrition and Physically Active Lifestyles (FUNPALs)	12-36 months	Parent-toddler dyads	Home environment	Low-income and ethnically diverse general population	Face-to-face, text reminders	Trained facilitators and health workers delivered health information and positive parenting coaching	10x90 minute session, weekly	10-weeks	12-weeks
Dewald-Kaufmann 2013 ⁴¹	Gradual Sleep extension	12-19 years	Adolescents with chronic sleep reduction	School + home phone calls	Adolescents with a Chronic Sleep Reduction Questionnaire (CSRQ) score of ≥40	Face-to-face and telephone	Personal sleep schedule to gradually extend sleep by 10min per night	Daily sleep schedule provided for two weeks	2-weeks	3-weeks

Name	Study title	Age range/mean	Target population	Setting ^a	Inclusion criteria	Mode of delivery	Intervention overview ^{b,c}	Dose	Intervention duration	Study duration
Dong 2020 ⁵⁶	Transdiagnostic sleep and circadian intervention for adolescents	10-18 years	Adolescents with eveningness chronotype	University based clinic	Reported eveningness and 7-day sleep diary with late onset sleep time	Face-to-face	Promoting behaviour change via four cross-cutting modules (functional analysis, goal setting, motivational interviewing, education)	6 x 50 min sessions, delivered weekly	6-weeks	6-months
Haines 2013 ⁵⁷	Healthy Habits, Happy Homes	2-5 years	Ethnic minority and low-income families	Community Health Centres in deprived areas	Low-income and ethnically diverse general population with a television in their room	Face-to-face, phone calls, texts and printed information	Motivational coaching to adopt healthy routines to reduce obesity	4xhome visits, 4xphone calls, weekly text messages	24-weeks	24-weeks
Hammersley 2019 ⁵⁸	Time2bHealthy	2-5 years	Parent-child dyads	Home environment	General population (healthy weight and overweight)	Online	Online modules targeting multiple behaviours including sleep	6x30min modules to complete over 11 weeks, weekly email reminders	23-weeks	24-weeks
Hammersley 2021 ⁵⁹	Time for Healthy Habits	2-6 years	Children aged 2-6 years	Home environment	General population	Telephone and online	Modules targeting multiple behaviours including sleep via telephone calls or online modules	6x30min telephone calls or 6 online modules	3-months	9-months
Hart 2017 , ⁴³ (additional detail from ⁶⁰)	Behavioural intervention to enhance time in bed	8-11 years	Children aged 8-11 years	Not stated	General population	Face to face, telephone	Behavioural strategies (e.g., goal setting, stimulus control, positive reinforcement) to increase time in bed by 1.5 hours	4 sessions over 8 weeks (week 1 60 mins, week 2 30 mins, telephone week 4, week 6)	8-weeks	8-weeks
Hiscock 2019 ⁶¹	Sleep well, Be well	5-6 years	Children with sleep problems in the 1st year of schooling	Primary schools	Parental report of "moderate" or "severe" sleep problems	Face-to-face, follow-up phone call	Nurse led consultation, providing information on good sleep practice and new sleep routine	1x45min consultation, 10min follow-up phone call, optional 20min consultation	4-weeks	52-weeks
Kalak 2012 ⁶²	Daily running intervention	17-19 years	School students	Secondary school	General population	Face-to-face	Daily morning run at school in small groups	30min run weekdays for 3 weeks	3-weeks	3-weeks
Kira 2014 ⁶³	ACES; adaptation of the Australian Centre for Education in Sleep programme	13-16 years	School students	Secondary school	General population	Face-to-face plus workbook	A sleep educational programme, including small group discussions ^b	4x50min sessions	4-weeks	10-weeks
Knebel 2020 ⁶⁴	Movimente programme	11-15 years	School students	Secondary school	General population	Face-to-face	Programme to increase physical activity and reduce sedentary time teacher training, environmental improvements, and health education ^c	Structures created and equipment provided to encourage physical activity and teachers encouraged to use	39-weeks	39-weeks

Name	Study title	Age range/mean	Target population	Setting ^a	Inclusion criteria	Mode of delivery	Intervention overview ^{b,c}	Dose	Intervention duration	Study duration
Lin 2018 ⁶⁵	A theory-based sleep hygiene intervention	15years	School students	Secondary school	General population	Face-to-face	A sleep education programme, encouraging behaviour change with individual plan and parental involvement	4x1hr sessions every two weeks	8-weeks	24-weeks
Lufi 2011 ⁴⁵	Delayed school start time intervention	13-14 years	School students	Secondary school	General population	Restructure environment	School start time delayed for one hour ^b	Daily for one week	1-week	2-weeks
Marsh 2020 ⁶⁶	3 Pillars Study	2-4 years	Parent – child dyads	Clinical and home	Children whose daily screen use exceeded recommendations (≥1 hour per day)	Face-to-face and online	Workshop and online materials aimed to promote positive parent-child interactions at bedtime, mealtimes, and when restricting screen time	A half-day workshop and subsequent access to the study website over a 6-week period	6-weeks	12-weeks
Mindell 2009 ⁶⁷	Bedtime routine intervention	18-36 months	Mother-child dyads	Home	Mother reported “small” to “severe” sleep problem, but not a sleep disorder	Face-to-face	A nightly 3-step bedtime routine (bath, massage, quiet activities), lights out within 30 mins of end of bath	Everyday	2-weeks	3-weeks
Mindell 2016 ⁶⁸	Sleep Well! Sleep Hygiene Education	2-12 years	Socioeconomically disadvantaged children	Home	Children from low income household, without proper bedding	Face-to-face and restructured environment	Sleep education messages for parents	Messages provided twice: week 1 & week 4	4-weeks	4-weeks
Mitchell 2021 ⁶⁹	Mobile platform to promote sleep	10-12 years	Children aged 10-12 years	Home	Parent/guardian report of typically 8–9 hours in bed	Online	Mobile health platform used to send goal achievement messages around bedtimes	Daily messages over a 7 week period	7-weeks	9 weeks
Moore 2019 ⁷⁰	Healthy Change and Systems Change	11-12 years	Low-income, minority race, overweight or obese children	Middle school	Children with BMI ≥85th percentile, entering 6 th grade	Face-to-Face and telephone	2 active arms: Healthy Change: behavioural strategies used in CBT and motivational interviewing. System Change: emphasis on restructuring family daily routines	25 face-to-face (f-t-f) sessions in Year 1, alternating monthly f-t-f and phone sessions year 2, 4 f-t-f sessions and 8 phone sessions in year 3	156-weeks	3 years (annual data collection)
Morell-Azanza 2019 ⁷¹	IGENOI (Intervention Grupo Estudio Navarro de Obesidad Infantil) study	7-16 years	Children with abdominal obesity	Clinical	Waist circumference above 90 th percentile, according to national data	Face-to-face	Family based lifestyle programme. Education sessions around diet	6 x 30-min sessions	8-weeks	8-weeks

Name	Study title	Age range/mean	Target population	Setting ^a	Inclusion criteria	Mode of delivery	Intervention overview ^{b,c}	Dose	Intervention duration	Study duration
Moseley 2009 ⁷²	Improving Adolescent Well-Being: Day and Night	15.6 years	Year 11, psychology class students with delayed sleep	Secondary school	Adolescents with discrepant out of bedtimes (school v weekend mornings >2 h), and insufficient sleep on school nights	Face-to-face	Education sessions promoting and maintaining a healthy lifestyle using a cognitive-behaviour therapy framework ^b	4 x 50-min sessions	4-weeks	6-weeks
Moula 2020 ⁷³	Arts therapies intervention study	7-10 years	Primary school children	Primary school	Children with mild emotional and behavioral difficulties	Face-to-face	Art therapies	1 hour per week for 8 weeks	8-weeks	8-weeks
Mousarrezai 2020 ⁷⁴	Education through Short Message Service	7-12 years	Mother – child dyads	Home	Children with sleep inadequacy based on sleep duration	Mobile phone	Educational messages sent to mothers via phone about sleep	Daily messages for 4 weeks	4-weeks	13-weeks
Pablos 2018 ⁷⁵	Healthy Habits Program	11-12 years	Students attending Spanish primary schools in Eastern Valencia	Primary school	General population	Face-to-face	Education sessions around diet, physical activity, sleep and hygiene + a practical physical activity session ^b	Twice weekly sessions lasting 150 mins	32-weeks	32-weeks
Puder 2011 ⁷⁶	Ballabeina study	5.1 years	Children at school in high migrant population areas in Switzerland	Pre-school	General population with >40% migrant children	Face-to-face and take home resources and restructured environment	Multidimensional lifestyle intervention, focused on physical activity, nutrition, media use, sleep and adaptation of the built environment of the preschool class ^b	Children: 4 x 45min sessions a week + 22 additional sessions Teachers: x2 workshops Parents: x3 evenings	42-weeks	42-weeks
Quach 2018 ⁷⁷	Cogmed Working Memory	6-7 years	Students in the second year (Grade 1) with low working memory	Primary schools	Children with low verbal and/or visuo-spatial working memory scores relative to their peers	Face-to-face	The Cogmed Working Memory Training tasks on temporary storage and manipulation of verbal and/or visuo-spatial information	20-25 sessions, 25 mins each, over 5-7 weeks	5-7-weeks	26-weeks
Rigney 2015 ⁷⁸	Australian Centre for Education in Sleep (ACES)	11-13 years	Students in Year 6/7 from 12 south Australian schools	Secondary schools	General population	Face-to-face and take home resources	Education sessions on sleep physiology and sleep hygiene ^c	4x weekly lessons then a group project on sleep which students then presented to parents.	4-weeks	18-weeks

Name	Study title	Age range/mean	Target population	Setting ^a	Inclusion criteria	Mode of delivery	Intervention overview ^{b,c}	Dose	Intervention duration	Study duration
Santiago 2020 ⁷⁹	Strength training for sleep	14-19 years	Students aged 14-19 years	Secondary schools	Adolescents reporting poor sleep quality or sleep disturbance, and daytime sleepiness	Face-to-face	Strength training sessions	55 mins, 3 times a week, for 12-weeks	12-weeks	12-weeks
Sousa 2013 ⁸⁰	Sleep Education Program (SEP)	16.8 years	12 th grade students	Secondary schools	General population	Face-to-face	Education programme about physiological and behavioural processes of sleep and healthy lifestyle	5 x 50-minute classes	<1-week (5 days)	4-weeks
Sundgot-Borgen 2020 ⁸¹	Healthy Body Image program	16.8 years	12 th grade students	Secondary schools	General population	Face-to-face	Interactive workshops, with three themes related to body image, social media literacy and lifestyle ^c	3 x 90-minute interactive workshops	13-weeks	52-weeks
Tamura 2014 ⁸²	Sleep Education programme with self-help	9-11 years	Children in years 4 to 6	Elementary schools	General population	Face-to-face	Education session with quiz about sleep habits, then practicing target sleep behaviour for 2 weeks. Monitoring practice ^b	1x 40 min sessions, then performing sleep behaviour for two weeks	2-weeks	2-weeks
Tamura 2016 ⁸³	Sleep Education with self-help	12-13 years	Adolescents in grade 7 public junior high schools	Junior high schools	General population	Face to face	Education session, then practicing a target sleep behaviour for 2 weeks. Monitoring ^b	1x 45 min session, then the sleep behaviour for two weeks	2-weeks	2-weeks
Taylor 2015 ⁸⁴	Tailored Family Based Obesity Intervention	6.5 years	Families with children aged 4 to 8 who were overweight and obese	Clinical setting	Children with BMI ≥85 th percentile	Face to face, phone calls	Families attended one multidisciplinary session to develop specific goals then met with a mentor	Initial session (1-2 hours). Meet mentor monthly in year 1; three monthly in year 2	104-weeks	104-weeks
Tomayko 2019 ⁸⁵	Healthy Children Strong Families	2-5 years	American Indian parent-child dyads on tribal reservations	Home	General population from American Indian communities	Mail, text reminders, social media	12 healthy lifestyle lessons (one mailed each month)	12 lessons (one each month for a year)	52-weeks	52-weeks
Uhlig 2019 ⁸⁶	Rap & Sing Music Therapy	8-13 years	Adolescents in eighth grade	Public primary school	General population	Face-to-face	Rap and sing music therapy sessions ^b	16 sessions, 45 min each, weekly over 4 months	16-weeks	16-weeks
Van Dyk 2017 ⁴²	Sleep Extension	14-18 years	High school students who sleep 5 to 7 hours on school nights	Home	General population with reduced sleep	Face to face	Increase time in bed on school nights	Increase time in bed by 1.5 hours/ night for two weeks	5-weeks	5-weeks
Van Rijn 2020 ⁸⁷	Interactive school-based sleep education program	14.0 years	Eighth grade secondary school students from an all-boys school in Singapore	Secondary school	General population	Face-to-face	Education program about the importance of sleep, barriers to getting enough sleep, and improving their opportunities for sleep ^b	4 x 1 hour lessons delivered over 5-weeks	4-weeks	11-weeks
Walton 2015 ⁸⁸	Parents and Tots Together	2-5 years	Parents of 2-5 year olds attending early years centres	Ontario early years centres	General population	Face to face	Group discussions about weight related topics, sometimes stimulated by videos of vignettes	9x 120 min weekly group sessions	9-weeks	36-weeks

Name	Study title	Age range/mean	Target population	Setting ^a	Inclusion criteria	Mode of delivery	Intervention overview ^{b,c}	Dose	Intervention duration	Study duration
Wilson 2014 ⁸⁹	Sweet Dreamzzz Sleep Education Programme	4.3 years	Low-income preschool children and their families	Head Start preschools	General population	Face to face	Education sessions for children with a 20 min daily lesson followed by 20 min of related activity ^b	8x 40 min sessions for children; 1x 45 min presentation for parents.	2-weeks	4-weeks
Wing 2015 ⁹⁰	Sleep Education Programme for adolescents	14.86 years	Seventh- to 11th-grade students aged between 12 and 18 years	Secondary schools	General population	Face to face	Didactic teaching with sleep hygiene tips, followed by small group discussions with cases ^b	1x 60 min town hall seminar, 2x 40-min small workshops, one month apart	12.2 weeks	17.2 weeks
Wolfson 2015 ⁹¹	Sleep Smart Program	12.5 years	Seventh graders and their parents	Public middle schools	General population	Face to face	Didactic information on adolescent sleep and specific strategies for establishing healthy sleep hygiene practices ^c	8x 40 min sessions over 4 weeks; then 4x booster sessions at two months & twelve months post-intervention	52-weeks	12 months (full results at 4 weeks only)
Yoong 2019 ⁹²	Sleep Intervention targeting physical activity	3-6 years,	Parents of 3 to 6-year-olds at a convenience sample of childcare centres	Home	General population	Video, phone, text message	Video targeting key behavioural beliefs about sleep practices & a telephone call using motivational interview techniques	1x 12 min video, 1-3x phone calls 2-4 weeks later, x2 text messages one week after call.	12-weeks	12-weeks

Footnote: ^a Setting includes where participants were recruited and tested and not necessarily where the intervention has taken place. Studies that reported cluster randomisation at class^b or school level^c

Table 2: Results of sub-group comparisons

Sub-group comparisons		Number of studies	Pooled estimate for sub-group in minutes (95% CI), minutes	Heterogeneity I ² , p	Difference in subgroup estimates, p
Combined estimate; no sub-groups		47	10.49 (5.56, 15.42), p<0.001	87.2%, p<0.001	-
Population Characteristics					
Age group	pre-school (>1 to 5)	10	5.73 (-5.81, 17.26), p=0.33	69.1%, p=0.001	p=0.68
	primary school (aged 6 to 11)	11	11.56 (3.34, 19.78), p=0.006	77.9%, p<0.001	
	secondary school (>11)	26	11.36 (4.45, 18.27), p=0.001	90.5%, p<0.001	
Continent	North America	15	15.63 (3.76, 27.50), p=0.01	90.4%, p<0.001	P=0.015
	Australasia	12	-0.56 (-5.82, 4.70), p=0.83	28.9%, p=0.16	
	Europe	9	9.97 (2.06, 17.88), p=0.014	49.0%, p=0.047	
	Asia	7	16.93 (4.47, 29.38), p=0.008	95.5%, p<0.001	
	South America	4	4.96 (-1.15, 11.02), p=0.11	0.0%, p=0.76	
Disadvantage	Population reported to be disadvantaged	6	7.90 (-3.28, 19.09), p=0.17	69.7%, p=0.006	p=0.67
	Population not reported to be disadvantaged	41	10.64 (5.23, 16.04), p<0.001	87.9%, p<0.001	
Population has a sleep problem	No	37	9.73 (4.46, 15.00), p<0.001	85.5%, p<0.001	p=0.75
	Yes	10	12.26 (-2.20, 26.73), p=0.10	91.9%, p=0.097	
Study design features					
Measure of sleep duration	Objective (actigraphy, EEG)	19	11.00 (1.42, 20.58), p=0.024	88.1%, p<0.001	p=0.86
	Subjective (child/parental report/sleep diary)	28	9.95 (3.98, 15.91), p=0.001	86.9%, p<0.001	
Subjective measures	Child self-report	11	7.76 (-2.31, 17.65), p=0.13	92.8%, p<0.001	p<0.001
	Parental report	10	0.43 (-3.41, 4.27), p=0.83	9.3%, p=0.36	
	Sleep diary	7	26.44 (17.69, 35.19), p<0.001	13.5%, p=0.33	
Nights used for data collection	Weekend nights	2	2.86 (-0.10, 5.83), p=0.059	0.0%, p=0.56	p=0.035
	Weekday nights	19	10.34 (2.50, 18.19), p=0.010	88.1%, p<0.001	
	All week nights	26	11.86 (4.04, 19.68), p=0.003	87.2%, p<0.001	
Study period for participants (including intervention length and follow-up)	One month or less	12	15.22 (7.55, 22.89), p<0.001	39.6%, p=0.077	p<0.001
	Longer than a month and up to 6 months	24	14.61 (6.21, 23.01), p=0.001	91.6%, p<0.001	
	Longer than six months and up to 12 months	6	1.64 (-4.65, 7.93), p=0.61	41.0%, p=0.13	
	Longer than 12 months	5	0.65 (-2.80, 4.10), p=0.71	0.0%, p=0.90	
Intervention features					
Location of intervention delivery	Home	13	16.42 (-1.21, 34.05), p=0.068	89.9%, p<0.001	p=0.49
	School	23	9.85 (3.50, 16.20), p=0.002	88.1%, p<0.001	
	Combination of locations /other/unspecified	11	5.84 (-1.57, 13.25), p=0.12	72.9%, p<0.001	
Mode of delivery	Face-to-face	31	12.86 (6.16, 19.55), p<0.001	89.1%, p<0.001	p=0.046
	Telephone, text message and/or online	14	4.00 (-1.67, 9.59), p=0.17	66.9%, p<0.001	
Parental involvement in delivery	No	19	10.61 (5.20, 16.02), p<0.001	69.5%, p<0.001	p=0.96
	Yes	28	10.39 (2.71, 18.09), p<0.001	90.5%, p<0.001	
Physical activity component	No	36	13.39 (7.14, 19.64), p<0.001	88.7%, p<0.001	P=0.001
	Yes	11	1.64 (-1.21, 4.49), p=0.26	0.0%, p=0.56	
Intervention aims to improve sleep only	No	18	1.25 (-2.54, 5.03), p=0.52	30.3%, p=0.11	P<0.001
	Yes	29	15.76 (8.90, 22.62), p<0.001	90.2%, p<0.001	
Intervention focused on obesity	No	35	12.82 (6.97, 18.68), p<0.001	88.5%, p<0.001	p=0.061
	Yes	12	3.29 (-4.78, 11.35), p=0.43	73.6%, p<0.001	
Behaviour models used	No	29	11.33 (5.78, 16.88), p<0.001	83.8%, p<0.001	p=0.65
	Yes	18	8.79 (-0.62, 18.19), p=0.067	87.8%, p<0.001	
Environmental change	No	41	11.06 (5.55, 16.58), p<0.001	88.3%, p<0.001	p=0.22
	Yes	6	5.34 (-2.01, 12.68), p=0.15	36.8%, p=0.16	
Intervention involves an earlier bedtime	No	44	7.37 (2.93, 11.82), p=0.001	82.8%, p<0.001	p=0.006
	Yes	3	46.97 (18.94, 75.00), p=0.001	89.6%, p<0.001	
<i>Earlier bedtime in younger v older children</i>	primary school (aged 6 to 11)	1	44.00 (28.74, 59.26), p<0.001		p=0.88
	secondary school (>11)	2	48.01 (-0.48, 96.52), p=0.052	94.4%, p=<0.001	
Intensity	Low/unclear	23	11.13 (4.78, 17.48), p=0.001	87.9%, p<0.001	p=0.64
	Medium	11	11.83 (0.74, 22.93), p=0.037	77.3%, p<0.001	
	High	13	7.01 (-0.11, 14.12), p=0.054	54.4%, p=0.010	

Footnote: Blake 2016⁵² data was supplemented by Blake 2018⁹³ data as both report the same study.