



City Research Online

City, University of London Institutional Repository

Citation: Kokke, K. H., Morris, J. A. & Lawrenson, J. (2008). Oral omega-6 essential fatty acid treatment in contact lens associated dry eye.. *Contact Lens & Anterior Eye*, 31(3), pp. 141-146. doi: 10.1016/j.clae.2007.12.001

This is the unspecified version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/3264/>

Link to published version: <https://doi.org/10.1016/j.clae.2007.12.001>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk

**Oral Omega-6 Essential Fatty acid Treatment in Contact Lens Associated
Dry Eye**

Karolien, H. Kokke ^a, Judith, A. Morris ^b & John, G. Lawrenson ^{b,*}

^a Private practitioner and contact lens department Sussex Eye Hospital

^b Henry Wellcome Laboratories for Visual Sciences, Department of Optometry and Visual Science, City University, London. UK. EC1V OHB

*Corresponding author

Address for Correspondence:

Henry Wellcome Laboratories for Visual Sciences, Department of Optometry and Visual Science, City University, UK. EC1V OHB

Tel: 0207 040 4310

Fax: 0207 0408494

Email: J.G.Lawrenson@city.ac.uk

Conflict of interest:

None

ABSTRACT

Purpose: Symptoms of dry eye are commonly reported in contact lens wearers and are a frequent cause of non-tolerance. The purpose of the present study is to evaluate the effects of oral treatment with particular omega-6 fatty acids in the form of evening primrose oil (EPO) on subjective symptoms, ocular surface signs and tear film characteristic in contact lens wearers with contact lens associated dry eye. **Methods:** The study design was randomised, double-masked and placebo controlled. 76 female patients were treated for 6 months with either EPO or placebo (olive oil). Patients underwent 3 examinations (baseline, 3 months and 6 months). At each examination patients were given a questionnaire relating to lens comfort and dry eye symptoms and underwent a series of tests of tear film characteristics (tear meniscus height, break-up time), Meibomian gland function (lipid layer thickness and quality) and ocular surface parameters (hyperaemia and staining). **Results:** The EPO group showed a significant improvement in the specific symptom of 'dryness' at 3 and 6 months ($p < 0.01$) and also a significant improvement in overall lens comfort at 6 months ($p < 0.01$). Tear meniscus height was increased in the EPO group at 6 months relative to baseline ($p < 0.01$), although all other objective signs were unchanged. **Conclusion:** This study provides evidence for a beneficial effect of particular orally administered omega-6 fatty acids in alleviating dry eye symptoms and improving overall lens comfort in patients suffering from contact lens associated dry eye.

Keywords: contact lens; evening primrose oil; dry eye; therapy

INTRODUCTION

A sensation of 'dryness' is a commonly reported symptom in contact lens wearers [1, 2] and is a frequent cause of reduced wearing times or discontinuation of wear [3]. The presence of a contact lens on the eye can have an adverse affect on tear physiology by increasing the rate of evaporation and decreasing tear break-up time [4, 5, 6]. The magnitude of the effect is influenced by variables such as lens material, care regime and environmental conditions [7, 8, 9]. In the majority of cases of contact lens-induced dry eye objective signs are lacking and a diagnosis is based on symptoms alone. However, a recent study has demonstrated that intolerant contact lens wearers show significantly reduced tear stability and volume compared to successful wearers [10].

Conventionally, a variety of management strategies have been used to alleviate contact lens-induced dry eye including: tear supplements [11, 12] changing lens parameters [13] or tear preservation with punctal plugs [14, 15]. However, novel treatment strategies that advocate the use of anti-inflammatory or immunomodulatory agents may also be appropriate. This approach is based on evidence for a putative role of inflammation of the lacrimal functional unit (lacrimal gland and ocular surface) in the pathogenesis of many forms of dry eye [16, 17]. The immunomodulatory drug cyclosporine, that specifically inhibits T-lymphocyte proliferation, was approved by the FDA in 2002 as a treatment for dry eye. Randomized placebo-controlled trials have shown that cyclosporine

ophthalmic emulsion is superior to vehicle in stimulating aqueous tear production in patients with keratoconjunctivitis sicca (KCS) [18]. Moreover, a recent pilot study offers evidence that cyclosporine may also be beneficial in contact lens intolerant patients [19].

Dietary modification or supplementation with essential fatty acids (EFAs) represents an alternative therapeutic strategy for dry eye. EFAs act as precursors for the synthesis of eicosanoids (prostaglandins, thromboxanes and leukotrienes) that regulate many aspects of the inflammatory process. By altering the EFA content of the diet, or administering particular EFAs as supplements, it is possible to modify the balance between pro and anti-inflammatory mediators [20]. The synthesis of inflammatory eicosanoids from their precursor EFAs is shown in Figure 1. 2-series prostaglandins and 4-series leukotrienes, which derive from arachidonic acid (AA) via the enzymes cyclooxygenase (COX) and 5-lipoxygenase (5-LOX) respectively, have proinflammatory effects e.g. pain, vasodilation and leukocyte recruitment. Dietary omega-3 fatty acid supplementation e.g. eicosapentaenoic acid (EPA), can reduce the production of these AA-derived mediators via a process of competitive enzyme inhibition, thereby shifting the balance to a less inflammatory state [20].

An alternative therapeutic approach involves the use of specific omega 6 EFA supplements [21]. Several studies have shown that gamma-linoleic acid (GLA), which is found in high concentration in oils derived from evening primrose (*Oenothera biennis*), has anti-inflammatory properties. GLA elevates dihomo- γ -linolenic acid (DGLA) concentrations, leading to an increased synthesis of 1-series prostaglandins that have a negative feedback role in chronic inflammation [21]. DGLA can also suppress the production of 2-series prostaglandins and 4-series leukotrienes via enzyme inhibition. Clinical studies have demonstrated that dietary supplementation with GLA and its precursor linoleic acid (LA) produces symptomatic relief in systemic diseases that are characterised by chronic

inflammation such as rheumatoid arthritis [22]. Recently, therapy with GLA and LA has also been shown to reduce ocular surface inflammation and improve symptoms in patients with dry eye syndrome with an inflammatory component [23, 24].

The purpose of the present study is to evaluate the effects of oral treatment with particular omega-6 fatty acids in the form of evening primrose oil (EPO) on subjective symptoms, ocular surface signs and tear film parameters in contact lens wearers with contact lens associated dry eye.

MATERIALS AND METHODS

Subject recruitment

76 Female soft contact lens wearers (Table 1) were recruited into the trial based either on their responses to a McMonnies dry eye history questionnaire indicating that they were suffering from dry eye or borderline dry eye (Mc Monnies score ≥ 10) or that they were experiencing symptoms of contact lens-induced dry eye. All patients who took part in the trial were wearing monthly or daily soft contact lenses. The lens types worn were divided into 4 groups, of which group 1 were silicone hydrogel lenses and groups 2 to 4 were separated according to FDA categories. The average wearing time of the contact lenses was calculated as hours per week. All subjects gave informed signed consent and the study was approved by the City University Research and Ethical Committee. Guidelines of the Declaration of Helsinki were adhered to.

Experimental design

This trial was longitudinal, double masked and placebo controlled. Subjects were randomly allocated to one of two groups. Each group was treated for a 6 month period with six capsules per day containing either LA/GLA in the form of EPO or a placebo, which was olive oil. Subjects were instructed not to change their diet, nor take any additional dietary supplements, for the duration of the study.

The active and placebo capsules were both supplied by Equazen (Equazen UK Ltd, London). The contents of the EPO capsules (*Qarma*[™]) used in the trial were LA 72.6%, GLA 10.5%, palmitic acid 6.5%, oleic acid 6.4% and stearic acid 1.8%. Each capsule contained 50 mg of GLA. The contents of the olive oil were 78.0% C18:1 (mainly oleic acid), 11.2% C16:0 (palmitic acid) and 5.6% C18:2 (mainly linoleic acid). Before encapsulation 1.6 mg natural tocopheryl (vitamin E) was added to both active and placebo to protect the contents of the capsule from oxidative damage. The olive oil capsules were made to taste, smell and appear the same as the active capsules. The capsules were loaded into standard unmarked tablet jars with a four-digit identifier code. Codes were allocated to one of two treatment groups (1 or 2), using an online random sequence generator. Coding, labelling and randomisation were carried out by the supplier and codes were not broken until the analysis was complete.

Experimental procedures

Subjects were evaluated at baseline and at three and six months during which the following procedures were carried out.

Symptoms questionnaire

Two visual analogue scales were designed. Firstly, lens comfort was graded on a scale from 1 to 10 (1=uncomfortable, to 10=no discomfort). Secondly each of the following symptoms was graded individually on a scale from 0 to 10 (0=no

symptoms to 10=extreme symptoms): soreness, scratchiness, dryness, grittiness, burning.

Tear meniscus height

The inferior tear meniscus height (TMH) was assessed with the slit lamp and recorded in mm.

Ocular surface health

Bulbar and palpebral conjunctival hyperaemia was assessed, together with fluorescein staining of the cornea and conjunctiva. The CCLRU decimalised grading scale was used for grading (1=very slight, 2=slight, 3=moderate, 4=severe) [25]. The van Bijsterveld score was used to grade the rose bengal staining of the cornea and conjunctiva (0=no staining to 3=confluent staining) [26].

Meibomian gland assessment

The quality and quantity of Meibomian secretions were measured by manual expression. The quantity was graded using a 3-point scale: 0=normal, 1=delay 2= partially blocked and 3=blocked. The quality was similarly graded: 0=clear, 1=cloudy, 2=granular, 3=opaque solid.

Lipid layer evaluation

The lipid layer was assessed on its appearance under specular reflection using the Keeler Tearscope Plus®. (Keeler Ltd, UK) The lipid layer was graded according to the Guillon Scale into five different interference patterns (1= meshwork, 2=wave, 3=amorphous, 4=colour fringes (1st order), 5=colour fringes (2nd order) [27].

Tear break-up time

The non invasive break-up time (NIBUT) was assessed using the Tearscope and fluorescein break-up time (BUT) was measured using conventional methodology. An average of three readings was taken in each case and the time in seconds from the blink to the first break noted.

Statistical analysis

Within group objective and subjective data were analysed using a non-parametric 2-way analysis of variance (ANOVA) test (Friedman). The Mann Whitney test was used for between group comparisons. Due to multiple testing across many parameters and the possibility of finding effects by chance, a inflammation such as rheumatoid arthritis [22]. Recently, therapy with GLA and LA has also been shown to reduce ocular surface inflammation and improve symptoms in patients with dry eye syndrome with an inflammatory component [23, 24].

The purpose of the present study is to evaluate the effects of oral treatment with particular omega-6 fatty acids in the form of evening primrose oil (EPO) on

subjective symptoms, ocular surface signs and tear film parameters in contact lens wearers with contact lens associated dry eye p-value of less than 0.01 was considered as statistically significant.

RESULTS

Of the 76 contact lens wearers recruited, 52 completed the trial (Placebo N=24, EPO N=28). The principal reason for drop out was non-compliance. All subjects, including dropouts, were included in the statistical analysis. Missing data was incorporated using the 'last observation carried forward' [28].

Demographics for all patients are shown in Table 1.

The median McMonnies score for all the study subjects was 13. Scores indicative of borderline dry eye (10-20) were found in 69% of subjects with 6% recording a score indicative of a more severe dry eye (>20). Although all subjects subjectively complained of symptoms of dry eye, 25% of subjects had a McMonnies score of <10. There was no significant difference in dry eye severity between the EPO group and the placebo group.

The EPO group had an average wearing time of 71.9 hours per week (SD 21.6 hours) and the placebo group had an average wearing time of 75.6 hours (SD 26 hours). This difference was not significant. The lens types worn by the subjects during the trial also showed no significant difference between groups (Table 1).

The overall contact lens comfort at six months for the EPO group showed an approximately 20% improvement compared to baseline ($p < 0.01$) (Figure 2). No significant changes in lens comfort were found for the placebo group. In terms of individual symptom scores, the EPO group showed a significant reduction in the symptom of 'dryness' at both three and six months ($p < 0.01$) (Figure 3). Both

groups showed no significant differences in all other individual symptom scores after three and six months.

Compared to baseline, measures of tear quality and stability, together with indicators of ocular surface health (e.g. hyperaemia and staining) did not show any significant differences at three and six months. Similarly there were no differences in meibomian gland secretions or lipid layer morphology. By contrast, the EPO group showed a significant increase in TMH after six months of EPO compared to baseline ($p < 0.01$) (Figure 4).

EPO and placebo were well tolerated and no significant adverse reactions were reported.

DISCUSSION

The present study found that dietary supplementation with the EPO-derived omega-6 EFAs LA and GLA ameliorated symptoms and improved overall lens comfort in female patients with contact lens associated dry eye.

Supplementation also caused a significant increase in tear production, as defined by tear meniscus height. It is reasonable to assume that the observed clinical improvement is primarily due to the well documented anti-inflammatory effects of these essential fatty acids [21].

Sub-clinical inflammation has been demonstrated in contact lens wearers, based on numbers of polymorphonuclear leukocytes [29], tear cytokine profile [29] and increased expression of the inflammatory markers HLA -DR and ICAM-1 on ocular surface epithelia [30]. HLA-DR is a MHC class II antigen whose expression by conjunctival epithelial cells has been shown to correlate with the severity of dry eye, as characterised by vital staining and Schirmer test [31,32,33]. Contact lens wearers with dry eye show greater expression of HLA-DR compared to non-dry eye wearers [34].

Recent studies have shown the beneficial effect of oral omega-6 supplementation (LA and GLA) in Sjögrens and dry eye syndrome with an inflammatory component [23,24]. Reduced symptoms of dry eye were reported, as well as an improvement in objective signs such as corneal staining and reduced conjunctival expression of HLA-DR. Supplementation with LA and GLA

has also been shown to increase tear production and reduce dry eye symptoms after photorefractive keratectomy (PRK) [35].

The present study similarly showed a reduction in the specific symptom of 'dryness', which was reduced by 40%. Although tear production was increased, other objective assessments e.g. ocular surface integrity and tear film quality were unaltered. Our sample size is not large enough to completely rule out that some effects may have gone undetected, although the magnitude of the observed non-significant effects in this sample suggests that changes would not be beyond measurement variability or be clinically significant. Furthermore, patients with contact lens associated dry eye typically display few clinical signs, even in the presence of marked symptoms. The study patients likewise showed minimal conjunctival hyperaemia and good ocular surface integrity at baseline and there was little or no suggestion of poor tear quality or quantity.

The anti-inflammatory effects of EPO are thought to be mediated by two principal mechanisms. The first involves a direct action of its component EFAs on immune cells. In-vitro studies have shown that the administration of LA and GLA suppresses T-cell activation by interfering with signal transduction [36]. The second mechanism is via their effect on eicosanoid synthesis. GLA is metabolised to DGLA, which inhibits the synthesis of pro-inflammatory cytokines

e.g. LT_4 and PGE_2 by competing with AA for COX and inhibiting 5-LOX via the 15-hydroxyl intermediate 15-hydroxyeicosatrienoic acid (15-HETE) [21]. DGLA is also metabolised to 1-series prostaglandins such as PGE_1 which has demonstrable anti-inflammatory effects at the ocular surface [37]. Furthermore, PGE_1 has also been shown to increase aqueous tear secretion via a presumed action on lacrimal adrenergic receptors leading to cyclic nucleotide synthesis [38].

Dietary supplementation with LA and GLA in patients with primary Sjögrens syndrome, at a lower dose than in the present study, elevates both plasma DGLA [39] and leads to raised PGE_1 levels in tears [24] in line with an improvement in ocular surface signs and symptoms. A similar mechanism could therefore explain the observed symptomatic improvement in the EPO group. A significant increase in tear meniscus height was also seen in the treatment group at 6 months compared to pre-trial values and a smaller increase in the placebo group (although this did not reach the 1% level of significance). The placebo used in the current study was olive oil. The choice of a truly inactive placebo can be problematic in randomised placebo-controlled trials of this nature [21] and therefore the possibility that olive oil influences tear secretion cannot be completely ruled out. Although olive oil contains predominantly oleic acid, an omega-9 fatty acid, it also contains a small quantity of LA, which in the body can be converted into GLA via the enzyme delta-6 desaturase and thus influence tear secretion by the same mechanism. Furthermore, olive oil has been recently

shown to contain a natural anti-inflammatory compound which displays a potency and pharmacological profile similar to ibuprofen [40].

Whilst the present study has provided evidence for the beneficial effect of LA and GLA in dry eye, other omega 6 EFAs may be deleterious e.g. a diet rich in AA can lead to an excessive production of proinflammatory mediators. For optimal nutrition, an appropriate balance should be maintained between omega-6 and omega-3 EFA consumption [41]. Western diets tend to be deficient in omega-3, particularly those derived from fish, which may promote the development of several chronic diseases, including dry eye [41]. Significantly, Miljanovich and co-workers found that dietary ratios of omega-6 to omega-3 EFAs in excess of 15:1 were associated with an increased risk of dry eye syndrome [42]. Omega 3 EFAs reduce the levels of AA-derived inflammatory mediators as well as maintaining the conversion of DGLA into anti-inflammatory 1-series prostaglandins.

There is evidence that omega-3 supplementation may independently be of benefit in the management of dry eye. A recent large epidemiological study involving 39,876 women in the USA found that higher intake of omega-3 fatty acids in the diet is associated with a decreased incidence of dry eye syndrome. Women who consumed at least five servings of tuna per week were at a 68% reduced risk of dry eye syndrome, compared to women who had one serving per week [42].

In conclusion, this study provides evidence for a beneficial effect of particular orally administered omega-6 fatty acids in alleviating symptoms and improving overall lens comfort in patients suffering from contact lens associated dry eye. It is likely that this is via a reduction in ocular surface inflammation that has been previously documented in contact lens wearers. A secondary effect on tear secretion may also be a contributing factor. When advising patients regarding EPO supplementation, the importance of omega-3 EFA intake should also be discussed. Future studies will investigate the efficacy of a combined preparation, since it is likely that omega-3 fatty acids could act synergistically with omega-6 to enhance the therapeutic effect of each EFA.

ACKNOWLEDGEMENTS

The authors wish to thank Dr D.P Crabb (City University, London) for statistical advice. None of the investigators have financial or personal interests in the products used during the trial.

REFERENCES

- [1] Brennan NA, Efron N, Symptomology of HEMA contact lens wear. *Optom Vis Sci* 1989;66:834-838.
- [2] Begley CG, Caffery B, Nichols KK, Chalmers R, Responses of Contact Lens Wearers to a Dry Eye Survey. *Optom Vis Sci* 2000;77:40-46.
- [3] Young G, Veys J, Pritchard N, Coleman S, A multi-centre study of lapsed contact lens wearers. *Ophth Physiol Opt* 2002;22:516-27.
- [4] Farris RL, The dry eye: its mechanisms and therapy, with evidence that contact lens is a cause. *CLAO J* 1986;12:234-46.
- [5] Lowther GE, Comparison of hydrogel contact lens patients with and without the symptom of dryness. *Int Contact Lens Clin* 1993 ;20:191-94.
- [6] Thai LC, Tomlinson A, Doane MG, Effect of contact lens materials on tear physiology. *Optom Vis Sci* 2004; 81:194-204.
- [7] Maruyama K, Yokoi N, Takamata A, Kinoshita S, Effect of environmental conditions on tear dynamics in soft contact lens wearers. *Invest Ophthalmol Vis Sci* 2004;45:2563-8.
- [8] Nichols JJ, Mitchell GL, King-Smith PE, Thinning Rate of the Precorneal and Prelens Tear Films. *Invest Ophthalmol Vis Sci* 2005;46: 2353-61.

[9] Nichols JJ, Mitchell GL, King-Smith PE, The impact of contact lens care solutions on the thickness of the tear film and contact lens. *Cornea* 2005;24:825-32.

[10] Glasson MJ, Stapleton F, Keay L, Sweeney D, Willcox MD, Differences in clinical parameters and tear film of tolerant and intolerant contact lens wearers. *Invest Ophthalmol Vis Sci* 2003 ;44:5116-24.

[11] Malet F, Karsenti D, Pouliquen P, Preservative-free ocular hydrating agents in symptomatic contact lens wearers: saline versus PVP solution. *Eye Contact Lens* 2003;29:38-43.

[12] Guillon M, Maissa C, Pouliquen P, Delval L, Effect of povidone 2% preservative-free eyedrops on contact lens wearers with computer visual syndrome: pilot study. *Eye Contact Lens* 2004;30:34-9.

[13] Riley C, Chalmers RL, Pence N, The impact of lens choice in the relief of contact lens related symptoms and ocular surface findings. *Contact Lens Ant Eye* 2005;28:9-13.

[14] Giovagnoli D, Graham SJ, Inferior punctal occlusion with removable silicone punctal plugs in the treatment of dry-eye related contact lens discomfort. *J Am Optom Assoc* 1992;63:481-5.

[15] Virtenal T, Huotari K, Harkonen M, Tervo T, Lacrimal plugs as a therapy for contact lens intolerance. *Eye* 1996;10:727-31.

Formatted: German (Germany)

Formatted: German (Germany)

Field Code Changed

Formatted: German (Germany)

Field Code Changed

Formatted: German (Germany)

Formatted: German (Germany)

Formatted: German (Germany)

Field Code Changed

Formatted: German (Germany)

Formatted: German (Germany)

Field Code Changed

Field Code Changed

Formatted: German (Germany)

Formatted: German (Germany)

[16] Stern ME, Gao J, Siemasko KF, Beuerman RW, Pflugfelder SC, The role of the lacrimal functional unit in the pathophysiology of dry eye. *Exp Eye Res* 2004;78:409-16.

Formatted: German (Germany)

Field Code Changed

[17] Smith RE, The Tear Film Complex: Pathogenesis and Emerging Therapies for Dry Eyes. *Cornea* 2005;24:1-7.

[18] Pflugfelder SC, Antiinflammatory therapy for dry eye. *Am J Ophthalmol* 2004;137:337-42.

[19] Hom MM, Use of cyclosporine 0.05% ophthalmic emulsion for contact lens intolerant patients. *Eye Contact Lens* 2006;32:109-11

[20] James MJ, Gibson RA, and Cleland LG, Dietary polyunsaturated fatty acids and inflammatory mediator production. *Am J Clin Nutr* 2000; 71 (suppl): S343-8

[21] Belch JJF and Hill A, Evening primrose oil and borage oil n rheumatic conditions. *Am J Clin Nutr* 2000; 71 (suppl): S352-6

[22] Zurier RB, Rosetti RG, Jacobson EW, DeMarco DM, Liu NY, Temming JE, White BM, Laposata M, Gammalinolenic acid treatment of rheumatoid arthritis. A randomized, placebo controlled trial. *Arthritis Rheum* 1996;39:1808-17.

[23] Barabino M, Rolando, P, Camicione, G Ravera, S, Systemic linoleic and gamma-linolenic acid therapy in dry eye syndrome with an inflammatory component. *Cornea* 2003;22:97-101.

[24] Aragona P, Bucolo C, Spinella R, Giuffrida S, Ferreri G, Systemic omega-6 essential fatty acid treatment and PGE1 tear content in Sjogren's syndrome patients. *Invest Ophthalmol Vis Sci* 2005;46:4474-9.

[25] Terry RL, Schnider CM, Holden BA, Cornish R, Grant T, Sweeney D et al, CCLRU standards for success of daily and extended wear contact lenses. *Optom Vis Sci* 1993;70: 234-43

Formatted: Dutch (Netherlands)

[26] van Bijsterveld, OP, Diagnostic tests in the Sicca syndrome. *Arch Ophth* 1969;82:10-4

[27] Craig JP, Tomlinson A, The importance of the lipid layer in human tear film stability and evaporation. *Optom Vis Sci* 1997;74: 8-13

[28] Barnes SA, Lindborg SR, Seaman JW Jr, Multiple imputation techniques in small sample clinical trials. *Stat Med* 2006; 25:233-45

[29] Thakur A, Willcox MD, Contact lens wear alters the production of certain inflammatory mediators in tears. *Exp Eye Res* 2000;70:255-9.

[30] Pisella PJ, Malet F, Lejeune S, Brignole F, Debbasch C, Bara J, Rat P, Colin J, Baudouin C, Ocular surface changes induced by contact lens wear. *Cornea* 2001;20:820-5.

[31] Tsubota K, Fujihara T, Saito K, Takeuchi T, Conjunctival Epithelium Expression of HLA-DR in Dry Eye Patients. *Ophthalmologica* 1999;213:16-19.

[32] Brignole F, Pisella PJ, Goldschild M, De Saint Jean M, Goguel A, Baudouin C, Flow cytometric analysis of inflammatory markers in conjunctival epithelial cells of patients with dry eyes. *Invest Ophthalmol Vis Sci* 2000;41:1356-63.

[33] Rolando M, Barabino S, Mingari C, Moretti S, Giuffrida S, Calabria G, Distribution of conjunctival HLA-DR expression and the pathogenesis of damage in early dry eyes. *Cornea* 2005;24:951-4.

[34] Albietsz JM, Conjunctival histologic findings of dry eye and non-dry eye contact lens wearing subjects. *CLAO J* 2001;27:35-40

[35] Macri A, Giuffrida S, Amico V, Lester M, Traverso CE, Effect of linoleic acid and gamma-linolenic acid on tear production, tear clearance and on the ocular surface after photorefractive keratectomy. *Graefes Arch Clin Exp Ophthalmol* 2003;41:561-6.

[36] Vassilopoulos D, Zurier RB, Rossetti RG, Tsokos GC, Gammalinolenic acid and dihomogammalinolenic acid suppress the CD3-mediated signal transduction pathway in human T cells. *Clin Immunol Immunopathol* 1997;83:237-44.

[37] Hoyng PF, Verbey N, Thong L, van Haeringen NJ, Topical prostaglandins inhibit trauma induced inflammation in the rabbit eye. *Invest Ophthalmol Vis Sci* 1986;27:217-25.

[38] Pholpramool C, Secretory effect of prostaglandins on the rabbit lacrimal gland in vivo. *Prostaglandins Med* 1979;3:185-92.

Formatted: English (U.K.)

[39] Oxholm P, Mathorpe R, Prause JU and Horrobin D, Patients with primary Sjögrens syndrome treated for two months with evening primrose oil. *Scand J Reumatol* 1996;15:103-8.

[40] Beauchamp GK, Keast RS, Morel D, Lin J, Pika J, Han Q, Lee CH, Smith AB, Breslin PA, Phytochemistry: ibuprofen-like activity in extra-virgin olive oil. *Nature* 2005;437:45-6.

[41] Simopoulos AP Essential fatty acids in health and chronic disease. *Am J Clin Nutr* 1999; 70 (suppl): S560-9.

[42] Miljanovic B, Trivedi KA, Dana MR, Gilbard JP, Buring JE, Schaumberg DA, Relation between dietary n-3 and n-6 fatty acids and clinically diagnosed dry eye syndrome in women. *Am J Clin Nutr* 2005;82:887-93.

TABLES

Patient demographics		EPO	Placebo
Wearing time (mean (SD) in hours)		71.9 (21.6)	75.6 (26.0)
Age (mean (SD) in years)		46.4 (12.6)	37.3 (10.7)
Lens type (proportions in each FDA category as %)	Group 1	0	3
	Group 2	74	64
	Group 3	0	0
	Group 4	26	33
McMonnies score (proportions in each category as %)	< 10	20	30
	10-20	74	64
	>20	6	6

Table 1. Patient demographics

FIGURES

Figure 1-Metabolism of EFAs to prostaglandins and leukotrienes. Dietary omega-6 EFAs linoleic acid and gamma linolenic acid are metabolised to dihomo-gamma linolenic acid (DGLA), resulting in an increased synthesis of anti-inflammatory 1-series prostaglandins. DGLA can be converted at a slow rate to arachidonic acid. The omega-3 amino acid eicosapentaenoic acid (EPA) competitively inhibits pathways leading to AA-derived proinflammatory. Solid lines represent synthetic pathways, broken lines represent enzyme inhibition.




Figure 2 - Overall contact lens comfort for the EPO and placebo group based on a 10-point visual analogue scale (1= uncomfortable, 10= no discomfort). Values represent means \pm SD. *P<0.01 (Friedman 1-way ANOVA).




Figure 3 – Mean symptom score for ‘dryness’ for the EPO and placebo group based on a 10-point visual analogue scale (1= no symptoms 10= extreme symptoms). Values represent means \pm SD. *P<0.01 (Friedman 1-way ANOVA) using a 10-point visual analogue scale




Figure 4 – Tear meniscus height for the EPO and placebo group (means ± SD).

*P<0.01 (Friedman 1-way ANOVA)

