



City Research Online

City, University of London Institutional Repository

Citation: Nketiah, L., Dzefi-Tettey, K., Mayeden, R. N., Agborli, A., Ohene-Botwe, B. & Mensah, Y. B. (2024). Correlation of sonographically-determined residual urine volume with lower urinary tract symptoms in adult males at a tertiary hospital. *Ghana Medical Journal*,

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/32773/>

Link to published version:

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk

Correlation of sonographically-determined residual urine volume with lower urinary tract symptoms in adult males at a tertiary hospital.

Linda Nketiah¹, Klenam Dzefi-Tettey², Raphael N Mayeden³, Ambrose Agborli⁴, Benard Ohene-Botwe⁵ Yaw B. Mensah⁶.

¹ Department of Diagnostics and Imaging, University of Ghana Medical Centre, Legon, Ghana.

² Department of Radiology, Korle Bu Teaching Hospital, Accra, Ghana.

³ Department of Radiology, School of Medicine, University of Health and Allied Sciences PMB 31, Ho, Ghana.

⁴ Department of Statistics and Actuarial Sciences, University of Ghana. Box LG 115 Legon, Ghana.

⁵ Department of Midwifery & Radiotherapy, School of Health and Psychological Sciences, City, University of London, London, UK.

⁶ Department of Radiology, University of Ghana Medical School, Legon, Ghana.

Corresponding Author: Linda Nketiah

Email address: lindabky@yahoo.com

Conflict of interest: None declared

Summary

Objective: To determine the correlation between the severity of LUTS as measured by the International Prostate Symptom Score (IPSS) and PVR urine volume measured by transabdominal ultrasound in patients with LUTS, and to determine the correlation between 'feeling of incomplete bladder emptying and sonographically measured PVR urine volume.

Design: Correlational cross-sectional study

Setting: Ultrasound Unit of the Radiology Department and Urology Clinic of Korle Bu Teaching Hospital

Participants: Male patients (n=256) aged 40 years or more and who presented to the urology department of Korle Bu Teaching Hospital with LUTS and gave their written consent were enrolled. The presence and severity of LUTS were evaluated using the IPSS. The PVR urine was measured using real-time transabdominal ultrasound scan.

Main outcome measure: Severity of LUTS and Residual urine volume

Results: The mean PVR urine volume was 84.5ml. Majority of respondents (57.3%, n=146) had PVR urine volume below 50ml with 27% (70 patients) having PVR urine volume above 100ml. PVR urine volume and total IPSS showed no statistically significant correlation. All age groups of respondents scored above 4 for quality of life (QoL). 'Intermittency' is the IPSS symptom which showed statistically significant correlation with PVR urine volume. PVR urine volume did not show a statistically significant correlation with the 'feeling of incomplete emptying'.

Conclusions: There was no statistically significant correlation between the total IPSS and PVR urine volume, thus residual urine volume does not correlate with severity of LUTS. The 'feeling of incomplete emptying' does not correlate with PVR urine volume.

Keywords: post-void residual urine, lower urinary tract symptoms, feeling of incomplete emptying, international prostate symptom score, quality of life.

Introduction

The International Continence Society (2015) defines LUTS as symptoms that result from conditions affecting the bladder and the urethra. LUTS have multifactorial causes, thus presenting a challenge to the management of this problem. Despite various options of medical treatments, lower urinary tract obstruction with its associated symptoms remains a significant economic burden and public health problem.¹

A review by Zhang et al² found that LUTS were highly prevalent in the world and estimated to affect 2.3 billion people in 2018, with 44.7% being men. In Ghana, Chokkalingam et al in 2011³ reported the prevalence of moderate-to-severe LUTS (IPSS \geq 8) to be 19.9%.

These symptoms negatively affect the quality of life (QoL) of affected individuals.⁴ The prevalence of LUTS generally increases with increasing age in men. Compared to younger men, men over 80 are more likely to have LUTS and more likely to complain of incomplete emptying, frequency, urgency, and weak stream.⁵

LUTS can be distressing for affected patients. Some individuals with LUTS organize their daily activities based on the availability of a toilet facility/restroom. LUTS can be evaluated using a questionnaire such as the International Prostate Symptom Score (IPSS) to establish the impact of these symptoms on a patient's routine. The IPSS is recognized globally as a tool for assessing LUTS severity.⁶ Although initially adopted for use in the clinical diagnosis of benign prostatic hyperplasia (BPH), the IPSS is now commonly used as a more specific measure of LUTS irrespective of the pathology in men and women⁷. It has been shown to be intrinsically consistent and show sensitivity to changes in symptomatology as well as having a good test-retest reliability.⁸ It is designed to be self-administered by patients; however, the inability to read and write can be a major drawback for the administration of the questionnaire.^{9,10}

The IPSS consists of seven questions encompassing storage, voiding, and post-micturition symptoms and an eighth question on QoL.⁸ The feeling of incomplete emptying assessed by the first question of the IPSS can be defined as "complaint that the bladder does not feel empty after micturition".¹¹ This symptom has been shown to be associated with lower QoL¹² and worsening of voiding and storage symptoms.¹³

A feeling of incomplete emptying does not always correlate strongly with the measured PVR urine volume.¹³ A study by Sountoulides et al¹⁴ reported that the feeling of incomplete bladder emptying does not always correlate with the finding of significant residual urine volume and can even be felt with an empty bladder. Cayetano-Alcaraz et al in 2016¹⁵ documented similar findings.

Measurement of PVR urine, the amount of residual urine in the bladder after a voluntary void¹⁶, is a screening test for evaluating voiding dysfunction. Like uroflowmetry, measurement of PVR is an effective tool that helps to identify patients in need of further evaluation and treatment follow-up. Threshold values for an abnormal PVR are poorly defined but most urologists agree that volumes of 50 mL to 100 mL constitute the lower threshold defining abnormal residual urine volume.¹⁷

There are 2 methods of measuring PVR urine: sterile catheterization and bladder ultrasound. Urethral catheterization has been accepted as the gold standard for PVR urine measurements, but this is a source of discomfort to the patient and carries a risk of urinary tract infection and urethral trauma.¹⁸ The bladder ultrasound on the other hand is a non-invasive, timesaving, widely available and well-tolerated method for measuring PVR urine.¹⁷ Due to its non-invasiveness, there is no risk of urethral injury and can be repeated as often as needed to follow up patient's response to management, thus the preferred method.

Traditionally, ultrasound estimation of bladder volume can be performed in two ways; either by using real-time ultrasound to directly visualize the bladder^{8,19,20} or by using a portable bladder scanner to calculate the volume automatically without directly visualizing the bladder.²¹ Most ultrasound machines have a function to automatically calculate volumes from the direct measurements of the antero-posterior, transverse and cranio-caudal dimensions of the bladder.

Referring urologists and general practitioners in Ghana often request for PVR urine volume but there has not been documented data on how this volume correlates with the patient's LUTS severity.

Significant chronic residual urine has been shown to lead to elevated intravesical pressures which may lead to hydronephrosis and eventually, renal failure.²² Patients with significant PVR urine will therefore require close follow-up to know when to initiate medical or surgical therapy. Very importantly, bothersome LUTS in spite of insignificant PVR urine also requires attention due to the great impact on QoL. For these reasons, urologists often would want to know the PVR urine volume in patients with LUTS. Unfortunately, there is limited data in Ghana regarding the relationship between PVR urine and the severity of LUTS, as well as the connection between the 'feeling of incomplete bladder emptying' and the measured PVR urine. This study aims to determine the correlation between the severity of LUTS, as assessed by the IPSS, and PVR urine volume, measured by transabdominal ultrasound in patients with LUTS. Additionally, it aims to establish the correlation between the feeling of incomplete bladder emptying and sonographically measured PVR urine volume.

METHODS

Study Design and Setting

The study was a correlational cross-sectional study which was conducted in the ultrasound unit of the Radiology Department with subjects from the Urology Clinic of the Korle Bu Teaching Hospital from August 2020 and June 2021. The Korle Bu Teaching Hospital is the largest tertiary hospital in Ghana, having the highest concentration of urologists in the country thus receiving referrals from most parts of the country. With an annual Urology outpatient average attendance of about 12500 patients, the urology clinic gets a fair number of patients with LUTS (average of 7180 cases of LUTS per annum). The Radiology Department of the Korle Bu Teaching Hospital has a busy ultrasound unit with individual scanning suites and built-in toilet facility.

Data collection

Sample size

Using a sample size formula proposed by Charan and Biswas for cross-sectional studies,²³

$$N = \frac{Z_{(1-\alpha/2)}^2 p(1-p)}{d^2}$$

Where N= sample size estimate of patients with lower urinary tract symptoms

$Z_{(1-\alpha/2)}$ = Standard normal variate (at 5% type 1 error ($p < 0.05$), it is 1.96.

Using a prevalence of 19.9 % per the study by Chokkalingam et al in 2011, the calculated minimum sample size was 245.

Adjusting for 10% participant attrition, the estimated sample size came to 270 respondents.

However, there were 256 respondents in the study after discarding incompletely filled questionnaires.

Study participants

Male patients aged 40 years or more who presented to the urology out-patient department of the Korle Bu Teaching Hospital with lower urinary tract symptoms and gave their written consent were enrolled in the study.

Patients with LUTS who had a history of previous urological (bladder/prostate) surgery or urethral /suprapubic catheter in-situ at the time of study, those with diabetes mellitus, history of urinary tract infection three months before presentation, known neurological disease which can affect sensation, or on diuretic medications were excluded.

Data collection

Using convenience sampling, patients who met the eligibility criteria and consented to participate in the study were consecutively recruited according to their availability, until the required sample size was obtained.

The investigator obtained written informed consent, after which a pre-tested questionnaire which consists of demographic characteristics and the IPPS survey questions was used to collect the data.

All PVR urine scans were performed by the principal investigator using a curvilinear transducer frequency of 3.5 MHz and volume estimator of Edan U60 ultrasound machine.

Procedure: Technique of sonographic determination of post-void residual urine

Just before the ultrasound scan was performed, subjects were instructed to empty their bladder as completely as possible in a private room.

The participant was positioned supine, and the area between each participant's umbilicus and pubic hairline was uncovered. It was prepped with acoustic/ultrasound gel and the transducer was placed at approximately 3 cm superior to the symphysis pubis so that the beam points toward the expected bladder location. The bladder was scanned in both axial (transverse) and sagittal planes, with

sweeps through the bladder in an arc. In the axial plane, the transducer was swept from cephalad to caudad, angling to include the whole bladder. In the sagittal plane, the transducer was initially placed in the midline and then swept in an arc to both sides to ensure that the whole bladder was visualised. The greatest transverse dimension on transverse view, and anteroposterior and superior-inferior dimensions on sagittal view were determined with the ultrasound calliper. The ultrasound calliper was placed in the inner wall of the bladder during the measurement of the urine volume, thus excluding the bladder wall.

The residual urine volume in this study was determined using the ultrasound machine's internal volume calculations. Two separate bladder ultrasound volume measurements were obtained. The average of the two measurements obtained was then recorded as the final PVR urine volume for that individual (this was done to minimize intra observer variability).

International Prostate Symptom Score (IPSS)

The IPSS questionnaire consists of eight items, which include seven 6-point scale questions on the LUTS (feeling of incomplete emptying, urinary frequency, interrupted stream, urinary urgency, weak urinary stream, urinary hesitancy/straining, and nocturia) and one 7-point scale question on

patients' satisfaction with their urinary condition. Based on the criteria, symptom severity was divided into three groups: mild (a symptom score of 0–7), moderate (8–19), and severe (20–35)⁸.

The QoL or level of satisfaction of LUTS patients is represented by seven grades : “No problem” (0 point = Very satisfied/Delighted), “I’m all right” (1 point= pleased), “Somewhat satisfied” (2 points= mostly satisfied), “Half-satisfied, half-dissatisfied” (3 points= mixed), “Somewhat dissatisfied” (4 points=mostly dissatisfied), “Distressed” (5 points= unhappy), and “I can’t stand it” (6 points =Terrible/Very dissatisfied).

Statistical analysis

Data was analysed using Microsoft Excel 2016 and IBM Statistical Package for the Social Sciences (SPSS) software version 20. The statistical significance level was set at $P < 0.05$.

A non-parametric Spearman’s Correlation test was used to assess the level of association between the data variables since the data was found not to follow a normal distribution. As a result, median values were reported alongside the mean values. Also, one-way non-parametric ANOVA (Kruskal-Wallis test) with post hoc analysis in SPSS was used to compare variables.

Ethical Considerations

Ethical approval was obtained from the Scientific and Technical Committee and Institutional Review Board of the Korle Bu Teaching Hospital (Ref: KBTH-STC 000022/2020 and KBTH-IRB/00022/2020 respectively).

RESULTS

Demographic information

A total of 256 males aged 40 years and above participated in this study. The ages of the subjects enrolled in this study ranged from 40-83 years with a mean of 65.1 ± 8.4 years old. The median age range was 60-69 years (with a median age of 65.0 years). The majority of the subjects were in the 60-69 years range. Table 1 shows the demographic data of respondents.

Table 1. Demographic characteristics of subjects

Variable	
Age	Years
Minimum	40
Maximum	83
Mean \pm SD	65.1 ± 8.4

Occupational status	Count (%)
Retired	161 (62.9)
Active Service	95 (37.1)
Educational level	Count (%)
None	7 (2.7)
Primary	76 (29.7)
Secondary	111 (43.4)
Tertiary	39 (15.2)
Postgraduate	23 (9.0)

As shown in Table 1 - the majority (n=161, 62.9%) of respondents recruited for this study had retired from active service and 37.1% (n=95) were in active service. Secondary/middle school leavers (n=111, 43.4%) formed the majority of the respondents, with those having no formal education (n=7, 2.7%) being the least.

The mean duration of symptoms was 3.8 ± 4.6 years (median -2 years), with the shortest duration being 0.1years (1-month 2weeks) and the longest duration being 20 years. Younger respondents had a shorter symptom duration.

Residual urine volume

The data for the post-void residual urine volume was skewed. The mean, standard deviation and median values were 84.5 ml, 112.6 and 49 ml, respectively with the least recorded residual urine being 0 (complete bladder emptying) and the largest volume being 669 ml. Most respondents (57.3%, n=146) had PVR urine volume below 50 ml, 15.7% (n=40) between 51 and 100 ml and 27.0 % (n=70) had volumes above 100 ml

IPSS and QoL

The observed statistics (mean, standard deviation, median and range values) across all the patients were: Total IPSS score (Mean \pm SD = 16.3 ± 6.4 ; median = 17; range = 4-31); QoL on a scale of 1-6 (Mean \pm SD = 4.4 ± 1.2 ; median = 5; range = 1-6), shown in Table 2.

Of the 7 LUTS assessed by the IPSS, the symptom with the highest mean score was nocturia, with a mean score of 3.2 ± 1.4 and median of 3.

Table 2: IPSS and the observed characteristics

Parameters	Mean± SD	Range		Median
		min	max	
Incomplete emptying	2.2±1.7	0	5	2.0
Frequency	2.6±1.5	0	5	2.0
Intermittency	2.2±1.7	0	5	2.0
Urgency	1.9±1.7	0	5	2.0
Weak stream	2.4±1.7	0	5	2.5
Straining	1.8±1.8	0	5	1.0
Nocturia	3.2±1.4	0	5	3.0
Total IPSS score	16.4±6.5	4	31	17.0
QoL	4.4±1.3	1	6	5.0
PVR	84.5±112.6	0	669	49.0

Min=Minimum; Max = Maximum; SD = Standard deviation

The mean QoL score across all age groups was above 4 irrespective of their total IPSS score or residual urine volume; the age group with the highest mean QoL score of 5 was the 40-49 age

range. The age group with the highest PVR was the 70-79 years group, with a mean PVR urine of 96.5ml (median = 45.5 ml). The 70-79 years group also recorded the highest mean duration of symptoms of 4.9 ± 4.8 years (Table 3).

Table 3: Age groupings and Duration of symptoms, IPSS and PVR urine volume

Age range (years)	Mean duration of symptoms score ± SD (range) in years/Median	Mean Total IPSS Score ± SD (range)/median	Mean QoL Score ± SD (range)/median	Mean PVR urine volume Score ± SD (range)/median
40-49	0.1± 0.0 (0.1-0.1)/ 0.1	7.25 ± 0.5 (7-8)/ 7.0	5 ± 0.0 (5-5)/ 5.0	54 ± 2.7 (50-56)/ 55.0
50-59	1.8 ± 1.7(0.1-5)/ 0.9	17.7 ± 7.5 (4-29)/ 20.0	4.2 ± 1.4 (1-6)/ 5.0	74.0 ± 125.0 (0-669)/ 45.0
60-69	4.1 ± 4.7 (0.1-20.0)/ 2.0	16.0 ± 5.9 (5-30)/ 16.0	4.4 ± 1.4 (1-6)/ 5.0	84.1 ± 98.1 (0-526)/ 51.0
70-79	4.9 ± 4.8 (0.1-20)/ 5.0	17.0 ± 6.3 (5-31)/ 16	4.5 ± 1.1 (2-6)/ 5.0	96.5 ± 133.7 (7-560)/ 45.5
80-83	4.8 ± 7.9 (0.2-20)/ 0.5	13.6 ± 6.4 (8-24)/ 9.0	4.7 ± 0.4 (4-5)/ 5.0	78.8 ± 65.3 (22-200)/ 50.0

Key: SD= standard deviation; QoL = Quality of life on a scale of 1-6 (1-Delighted, 6-Terrible); IPSS=International Prostate Symptom Score; PVR = postvoid residual.

Table 4: Groupings of IPSS severity with QoL and PVR

Total IPSS score category	n (%)	Mean QoL ± SD/(median)	Mean PVR± SD /(Median)
Mild IPSS score (0-7)	21(8.2)	2.52±1.3/ (2)	52.38±24.9/ (50)
Moderate IPSS score (8-19)	89(34.8)	4.30±1.2/ (5)	75.27±91.7/ (48.5)
Severe IPSS score (20-35)	146(57.0)	4.96±0.9/ (5)	107.06 ±147.8/ (49)

n =number of respondents, % =percentage

Majority of respondents (57 %, n=146) had severe LUTS. This category also scored high for QoL (poor quality of life). Those with severe symptoms recorded higher PVR urine than those with mild and moderate symptoms. Mild IPSS showed a statistically significant correlation with PVR urine

volume [r(254)=.90,p=.029], and the QoL r(254)=.64,p=.002]. However, moderate r(254)=.37,p=.001], and severe r(254)=.50,p=.001] IPSS showed a statistically significant correlation with QoL but not with PVR urine { moderate r(254)=.15, p=.070]; severe r(254)=.05, p=.652]}

Table 5: Correlation of PVR urine volume with individual IPSS, total IPSS, QoL and other variables

Test components	Correlation (r_s -value)	<i>p</i> -value
Frequency of micturition vs PVR urine volume	0.015	0.805
Incomplete emptying vs PVR urine volume	0.079	0.209
Urgency vs PVR urine volume	0.057	0.361
Straining Vs PVR urine volume	-0.098	0.118
Weak stream vs PVR urine volume	0.006	0.928
Intermittency vs PVR urine volume	0.212	0.001*
Nocturia vs PVR urine volume	-0.111	0.076
Duration of symptoms	0.130	0.038*
Age vs PVR urine volume	0.085	0.177
QoL score vs Total IPSS	0.541	0.001*
QoL vs PVR urine volume	0.038	0.549
Total IPSS vs PVR urine volume	0.085	0.176

*=*statistically significant*

There was a statistically significant positive correlation between PVR urine volume and duration of symptoms ($r_s = 0.130$, $p = 0.038$) as well as intermittency ($r_s = 0.212$, $p = 0.001$). Also, the QoL score and total IPSS demonstrated a statistically significant positive correlation ($r_s = 0.541$, $p = 0.001$). However, PVR urine volume did not show a statistically significant correlation with frequency of micturition, incomplete emptying, voiding urgency, straining, weak stream, nocturia, age, QoL and total IPSS (Table 8).

Discussion

A total of 256 respondents aged 40 years and above were evaluated in this study. The mean age of respondents was 65.1 ± 8.4 years and the 60-69 years' age range was the most represented in this study (45.7% , n=117). These findings support the fact that LUTS is common among the elderly.⁵

The mean duration of symptoms was 3.8 (median = 2 years). The reported duration of symptoms was wide, between a few months and 20 years. The disparity could be attributed many factors such as the insidious way LUTS present as well as the general perception that voiding symptoms are part of the ageing process. Studies elsewhere have shown

that many elderly males seek professional help only when their inconvenience exceeds tolerance level.^{25,26}

Younger respondents had shorter duration of symptoms compared to older respondents, with the shortest mean symptom duration (6 weeks) noted in the 40-49 years range, followed by the 50-59 years range. These age ranges comprised individuals who were in active service, thus the impact of LUTS on their daily activities /work schedule was deemed significant interference (impaired quality of life), resulting in their relatively earlier presentation to hospital. Ojewola et al²⁷ reported that one of the most common reasons for men with LUTS seeking help was impaired quality of life, supporting the fact that impaired quality of life is a push factor for early presentation to the hospital.

Majority of respondents (57.3%, n=146) had PVR urine below 50ml with 27% having PVR more than 100ml. Lammers et al²⁸ in a similar study also reported that 27% of respondents recorded PVR urine greater than 100ml. Although there is no numerical value of PVR that has been universally accepted into current clinical practice¹⁶, a study by Kelly¹⁷ considered PVR volume of less than 50 ml adequate bladder emptying in young adults and

etween 50 and 100 ml in the elderly. In the Lammers et al²⁸ study, residual urine volume of more than 100 mL was considered abnormal. Thus, most of the respondents in our study can be considered to have adequate bladder emptying.

A statistically significant positive correlation was seen between PVR urine and duration of symptoms. Thus, patients who have had LUTS for longer periods recorded higher PVR urine, supporting the assertion that LUTS if left untreated over a period, can result in urinary retention²⁹ with its attendant increased residual urine volume.

In the management of patients with LUTS, careful attention to the severity of symptoms as well as the impact of the symptoms on the quality of life of the affected individual is of utmost importance. The mean IPSS in the index study was 16.3 with nocturia being the symptom with the highest score. This compared favourably to what was documented by Lammers et al²⁸, the mean IPSS was similar (16.5) but weak stream was the symptom which most participants mentioned “almost always occur”

Significant LUTS is usually defined as a total IPSS of 8 and above.³⁰ This comprises the group with moderate and severe IPSS scores. In this study, majority of respondents (57%, n=146) had severe LUTS and a good proportion (34.8%, n=89) had moderate LUTS. A smaller percentage of respondents (8%, n= 21) had only mild LUTS. In a study by Hamid et al³¹ in men with LUTS, most respondents (50%) had severe symptom score, similar to the findings in the index study. Awaisu et al³² also reported that majority of respondents (55%) had moderate symptom scores. The low numbers for those with mild symptoms may be because most people generally do not seek medical attention when they have mild disease. It can therefore be inferred that most of the patients who present to the hospital for evaluation of LUTS have clinically significant symptoms and must be given prompt attention. Ojewola et al²⁷ corroborated this finding in their study which reported moderate-to-severe symptoms as one of the most common reasons why men with LUTS seek help.

In this study, the mean PVR urine for those with mild, moderate and severe IPSS was 52ml, 75.3ml and 107ml respectively. Among these three categories of severity grading, only mild IPSS showed statistically significant correlation with the measured PVR urine volume. This was slightly different from the findings by Aldamanhori³³ in Saudi Arabia. In that study, the PVR urine measurement showed no statistical difference in all three categories of symptom severity. The findings in both studies support the point that factors other than symptom severity, such as urinary effort and

total score for voiding symptoms do influence the residual urine volume.¹⁵

Although respondents with moderate and severe IPSS recorded relatively higher mean PVR urine, this was not found to be statistically significant. Overall, there was no statistically significant correlation between the total IPSS and PVR urine volume. There have been inconsistent results in previous studies on the correlation of IPSS with residual urine volume. Although a few studies^{34,35} have reported a correlation between IPSS and post-void residual urine volume, others like that by Lammers et al²⁸ did not find a relation between the IPSS score and an abnormal residual urine volume in men aged over 50 with LUTS consulting primary care. A possible reason for the differing findings between various studies could be due to the varying study settings, with some studies being community-based and others being hospital-based or primary care setting.

Out of the seven symptoms assessed by the IPSS questionnaire in this study, ‘intermittency’ showed statistically significant correlation with PVR. The other 6 LUTS (including ‘feeling of incomplete emptying’) assessed by the IPSS questionnaire in this study showed no statistically significant correlation with the measured PVR urine. It is however worth noting this study found that the feeling of incomplete emptying also impaired QoL. Cayetano-Alcaraz et al¹⁵ and Aldamanhori³³ reported similarly that the feeling of incomplete emptying showed no statistically significant correlation with the measured PVR urine

Ozlulderden et al³⁶ as part of their study investigated the association between PVR and the feeling of incomplete emptying in both men and women. It was interestingly noted that they found a positive correlation between the feeling of incomplete bladder emptying and PVR volume in all age groups of women but in men, such a relationship was found only in those above 60 years. The findings in the Ozlulderden et al study could be attributed to the fact that the mean post void residual urine in those with and without the complaint of incomplete emptying did not differ much in the age group below 60 years in men.

Regarding quality of life (QoL)/border score and IPSS, there was a statistically significant positive correlation between total IPSS and QoL. In a study on LUTS among middle-aged and elderly Nigerian men, Adegun et al³⁷ concluded that there is no level of severity of LUTS in which patients’ QoL is not impaired. In our study, those with moderate and severe IPSS recorded high bother scores, thus, it can be said that most patients with LUTS had their quality of life affected to some degree.

Limitations

IPSS questionnaire was designed in English language, thus patients who could not read English had to be assisted by translation into a comprehensible language. This has the tendency to create room for interpreter bias, thus influence the LUTS severity assessment.

Conclusion

The severity of LUTS as evaluated by the IPSS did not correlate with the measured PVR urine volume. Instead, the symptom severity correlated well with the degree of QoL impairment.

This study also showed that the feeling of incomplete emptying does not show a statistically significant correlation with the measured post-void residual urine volume. The IPSS symptom which correlated best with PVR urine volume was 'intermittency' in this study.

The severity of LUTS and PVR urine although do not show statistically significant correlation, independently influence choice of management, underscoring the importance of measuring and documenting the PVR urine in the abdominopelvic ultrasound report of all patients who present with LUTS. Those found to have significant PVR urine can be treated irrespective of the symptom severity and those found to have bothersome symptom despite insignificant residual urine will also benefit from medical therapy to improve their quality of life.

References

1. International Continence Society. (2015). ICS fact sheets: A background to urinary and faecal incontinence. Available from: <https://www.ics.org/documents/documentsdownload.aspx?documentid=2172> Accessed January 31, 2020.
2. Zhang AY, Xu X. Prevalence, Burden, and Treatment of Lower Urinary Tract Symptoms in Men Aged 50 and Older: A Systematic Review of the Literature. *SAGE Open Nursing*. 2018; 4:2377960818811773. doi:10.1177/2377960818811773
3. Truzzi JCI, Almeida FMR, Nunes EC, Sadi M v. Residual Urinary Volume and Urinary Tract Infection-When are They Linked? *J Urol*. 2008;180(1):182-185. doi:10.1016/j.juro.2008.03.044
4. Coyne KS, Sexton CC, Thompson CL, et al. The prevalence of lower urinary tract symptoms (LUTS) in the USA, the UK and Sweden: Results from the epidemiology of LUTS (EpiLUTS) study. *BJU Int*. 2009;104(3):352-360. doi:10.1111/j.1464-410X.2009.08427.x

5. Parsons JK, Bergstrom J, Silberstein J, Barrett-Connor E. Prevalence and Characteristics of Lower Urinary Tract Symptoms in Men Aged ≥ 80 Years. *Urol*. 2008;72(2):318-321. doi:10.1016/j.urology.2008.03.057
6. Kok ET, Groeneveld FPMJ, Gouweloos J, et al. Determinants of Seeking of Primary Care for Lower Urinary Tract Symptoms: The Krimpen Study in Community-Dwelling Men. *Eur Urol*. 2006;50(4):811-817. doi:10.1016/j.eururo.2006.03.024
7. McConnell JD, Barry MJ, Bruskewitz RC, Bueschen AJ, Denton SE, Holtgrewe HL, Lange JL, McClellan BL, Mebust WK, Reilly NJ, Roberts RG. Benign prostatic hyperplasia: Diagnosis and treatment. Clinical Practice Guideline, No. 8. AHCPR publication No. 94-0582, Rockville, Md. Agency for Health Care Policy and Research. Public Health Service, US Department of Health and Human Services. 1994;225.
8. Barry MJ, Fowler Jr FJ, O'Leary MP, Bruskewitz RC, Holtgrewe HL, Mebust WK, et al. The Measurement Committee of the American Urological Association. The American Urological Association symptom index for benign prostatic hyperplasia. *J Urol* 1992;148(5):1549-57.
9. Oztürk Mİ, Koca O, Keleş MO, Güneş M, Kaya C, Karaman Mİ. International prostate symptom score: really appreciated by all patients or not? *Urol J*. 2011;8(3):227-230. Accessed February 5, 2020. <http://www.ncbi.nlm.nih.gov/pubmed/21910103>
10. Badia X, Rodríguez F, Carballido J, et al. Influence of sociodemographic and health status variables on the American Urological Association Symptom Scores in patients with lower urinary tract symptoms. *Urol*. 2001;57(1):71-77. doi:10.1016/S0090-4295(00)00894-3
11. Haylen BT, de Ridder D, Freeman RM, et al. An International Urogynecological Association (IUGA)/International Continence Society (ICS) joint report on the terminology for female pelvic floor dysfunction. *Int Urogynecol J*. 2010;21(1):5-26. doi:10.1007/s00192-009-0976-9
12. Fujimura T, Kume H, Nishimatsu H, et al. Assessment of lower urinary tract symptoms in men by international prostate symptom score and core lower urinary tract symptom score. *BJU Int*. 2012;109(10):1512-1516. doi:10.1111/j.1464-410X.2011.10445.x
13. Lee JY, Lee DH, Lee H, Bang WJ, Hah YS, Cho KS. Clinical implications of a feeling of incomplete emptying with little post-void residue in men with lower urinary tract symptoms. *Neurourol Urodyn*. 2014;33(7):1123-1127. doi:10.1002/nau.22473

14. Sountoulides P, Mutomba WF. Feeling of incomplete bladder emptying: A definition with clinical implications. LUTS: Lower Urinary Tract Symptoms. 2019;11(1):85. doi:10.1111/luts.12251
15. Cayetano-Alcaraz AA, Herrera-Cáceres JO, García-Mora A. The relationship between the sensation of incomplete voiding and a high postvoid residual volume. *Spanish Urological Records (English Edition)*, Volume 40, Issue 5, June 2016, Pages 309-316.
16. Kaplan SA, Wein AJ, Staskin DR, Roehrborn CG, Steers WD. Urinary Retention and Post-Void Residual Urine in Men: Separating Truth From Tradition. *J Urol.* 2008;180(1):47-54. doi:10.1016/j.juro.2008.03.027
17. Kelly CE. Evaluation of voiding dysfunction and measurement of bladder volume. *Rev Urol.* 2004;6 Suppl 1:S32-7. Accessed February 5, 2020. <http://www.ncbi.nlm.nih.gov/pubmed/16985853>
18. Schaeffer AJ, Chmiel J. Urethral meatal colonization in the pathogenesis of catheter-associated bacteriuria. *J Urol.* 1983;130(6):1096-1099. doi:10.1016/S0022-5347(17)51701-2
19. Griffiths CJ, Murray A, Ramsden PD. Accuracy and repeatability of bladder volume measurement using ultrasonic imaging. *J Urol.* 1986;136(4):808-812. doi:10.1016/S0022-5347(17)45086-5
20. Abrams P, Cardozo L, Fall M, et al. The standardisation of terminology in lower urinary tract function: Report from the standardisation sub-committee of the International Continence Society. *Urol.* 2003;61(1):37-49. doi:10.1016/S0090-4295(02)02243-4
21. Hartnell GG, Kiely EA, Williams G, Gibson RN. Real-time ultrasound measurement of bladder volume: A comparative study of three methods. *BJR.* 1987;60(719):1063-1065. doi:10.1259/0007-1285-60-719-1063
22. Negro CLA, Muir GH. Chronic urinary retention in men: How we define it, and how does it affect treatment outcome. *BJU Int.* 2012;110(11):1590-1594. doi:10.1111/j.1464-410X.2012.11101.x
23. Charan J, Biswas T. How to Calculate Sample Size for Different Study Designs in Medical Research? *Indian J Psychol Med.* 2013;35(2):121. doi:10.4103/0253-7176.116232
24. Chokkalingam AP, Yeboah ED, DeMarzo A, et al. Prevalence of BPH and lower urinary tract symptoms in West Africans. *Prostate Cancer Prostatic Dis.* 2012;15(2):170-176. doi:10.1038/pcan.2011.43
25. Ward J, Sladden M. Urinary symptoms in older men, their investigation and management: is there an epidemic of undetected morbidity in the waiting room? *Fam Pract.* 1994;11(3):251-259. doi:10.1093/FAMPRA/11.3.251
26. Shim SR, Kim JH, Choi H, et al. Association between self-perception period of lower urinary tract symptoms and International Prostate Symptom Score: a propensity score matching study. *BMC Urol.* 2015;15(1). doi:10.1186/S12894-015-0021-X
27. Ojewola RW, Oridota ES, Balogun OS, Ogundare EO, Alabi TO. Lower Urinary Tract Symptoms: Prevalence, Perceptions, and Healthcare-Seeking Behavior amongst Nigerian Men. *World J Mens Health.* 2016;34(3):200. doi:10.5534/wjmh.2016.34.3.200
28. Lammers H, Teunissen T, Bor H, Smid I, Lagro-Janssen A. No Relationship Between the International Prostate Symptom Score and Post-Void Residual Volume in Primary Care. *Res Rep Urol.* 2020; 12:167-174. doi:10.2147/RRU.S241961
29. Stroup SP, Palazzi-Churas K, Kopp RP, Parsons JK. Trends in adverse events of benign prostatic hyperplasia (BPH) in the USA, 1998 to 2008. *BJU Int.* 2012;109(1):84-87. doi:10.1111/J.1464-410X.2011.10250.X
30. Homma Y, Araki I, Igawa Y, Ozono S, Gotoh M, Yamanishi T, et al. Clinical guideline for male lower urinary tract symptoms. *Int J Urol.* 2009;16(10):775-790. doi:10.1111/J.1442-2042.2009.02369.X
31. Hamid SHI, Mohammedahmed AY, Sharfi AR. Correlation between the Size of the Prostate, Post Void Residual Volume, PSA Level and IPSS in Men with LUTS in Three Major Urology Centers in Khartoum. *Int J Surg Res.* 2020;9(1):1-8. Accessed August 8, 2021. <http://article.sapub.org/10.5923.j.surgery.20200901.01.html>
32. Awaisu M, Ahmed M, Lawal AT, et al. Correlation of prostate volume with severity of lower urinary tract symptoms as measured by international prostate symptoms score and maximum urine flow rate among patients with benign prostatic hyperplasia. *Afr J Urol* 2021 27:1. 2021;27(1):1-7. doi:10.1186/S12301-021-00122-4
33. Aldamanhori R. Lower urinary tract symptoms and feeling of incomplete emptying in Saudi Arabian men and its correlation with postvoid residual urine. *Urol Ann.* 2019;11(2):132-134. doi:10.4103/UA.UA_133_18
34. Singla S, Garg R, Singla A, Sharma S, Singh J, Sethi P. Experience with Uroflowmetry in Evaluation of Lower Urinary Tract Symptoms in

Patients with Benign Prostatic Hyperplasia. *J Clin Diagn Res.* 2014;8(4):NC01. doi:10.7860/JCDR/2014/7909.4271

and elderly Nigerian men: Impact on quality of life. *Adv Urol.* 2016;2016. doi:10.1155/2016/1015796

35. Ezz El Din K, Kiemeny LALM, de Wildt MJAM, Debruyne FMJ, de La Rosette JJMCH. Correlation between uroflowmetry, prostate volume, postvoid residue, and lower urinary tract symptoms as measured by the international prostate symptom score. *Urol.* 1996;48(3):393-397. doi:10.1016/S0090-4295(96)00206-3

36. Özlülerden Y, Toktaş C, Zümrütbaş AE, et al. Can feeling of incomplete bladder emptying reflect significant postvoid residual urine? Is it reliable as a symptom solely? *Investig Clin Urol.* 2018;59(1):38. doi:10.4111/ICU.2018.59.1.38

37. Adegun PT, Adebayo PB, Areo PO. Severity of lower urinary tract symptoms among middle aged