



City Research Online

City St George's, University of London

Citation: Goozee, R., Barrable, A., Lubenko, J., Papadatou-Pastou, M., Haddad, M., McKeown, E., Hirani, S. P., Martin, M. & Tzotzoli, P. (2024). Investigating the feasibility of MePlusMe, an online intervention to support mental health, well-being, and study skills in higher education students. *Journal of Mental Health*, 33(5), pp. 576-586. doi: 10.1080/09638237.2022.2069699

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/32854/>

Link to published version: <https://doi.org/10.1080/09638237.2022.2069699>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

Investigating the feasibility of MePlusMe, an online intervention to support mental health, wellbeing, and study skills in higher education students

Rhianna Goozee¹, Alexia Barrable², Jelena Lubenko³, Marietta Papadatou-Pastou⁴, Mark Haddad⁵, Eamonn McKeown⁵, Shashivadan P Hirani⁵, Maryanne Martin⁶, and Patapia Tzotzoli⁷

¹Independent scholar, UK

²University of Dundee, School of Education and Social Work, UK

³Department of Health Psychology and Paedagogy, Rīga Stradiņš University, Latvia

⁴School of Education, National and Kapodistrian University of Athens, Greece

⁵School of Health Sciences, City, University of London, UK

⁶Department of Experimental Psychology, University of Oxford, UK

⁷My Psychology Clinic & iConcipio Ltd, UK

Corresponding author:

Ass. Prof. Marietta Papadatou-Pastou

National and Kapodistrian University of Athens, Greece

Tel: +30 6932398693;

Email: marietta.papadatou-pastou@seh.oxon.org

Acknowledgements

This article received writing support from Dr Rhianna Goozee, a freelance writer and editor. All authors have contributed to the writing of the paper and authorize the submission of the final manuscript.

Abstract

Introduction: While there are several web-based mental health interventions, few target higher education (HE) students. Importantly, more research is needed to establish their effectiveness. Here, we provide a pragmatic evaluation of an online intervention (MePlusMe) specifically designed to improve the mental health, wellbeing, and study skills of HE students.

Methods: In accordance with the published protocol for a feasibility study, we recruited a convenience sample of 137 HE students to participate in an 8-week intervention, with 26 participants retained at week 8. Validated measures of mood (depression and anxiety), wellbeing, and self-efficacy were collected at baseline, 2, 4 and 8 weeks, alongside two feedback forms assessing design and functionality (baseline), and engagement (week 4 and 8).

Results: We observed significant reductions in levels of anxiety and depression as well as increases in wellbeing, but no changes in self-efficacy. Participants rated the system design and functionality positively, and qualitative findings indicated high levels of satisfaction with MePlusMe.

Discussion: Findings support both the acceptability and the effectiveness of MePlusMe. Nonetheless, modest retention rates limit the precision and generalisability of these findings. Further investigation should ascertain optimal duration of engagement, most acceptable means of outcome assessment, and further detail about obstacles to utilisation.

Keywords

Students, online support, study skills, anxiety, depression, MePlusMe, mental health, wellbeing, digital intervention, higher education institutions, student support services, COVID-19

Introduction

Around three quarters of individuals with a lifetime mental health disorder experience their first episode by 25 years old.¹ Moreover, becoming a student at a higher education institution (HEI) constitutes a major transition and entails several challenges that can further exacerbate or even trigger the onset of mental health issues.²⁻⁴ It is therefore unsurprising that many higher education (HE) students worldwide experience mental health difficulties, including depression, anxiety, and other mental health and substance abuse disorders.⁴⁻⁸ Anxiety and depression are common among HE students in the US,² UK⁷ and further afield.⁵ A recent large-scale international survey of 14,000 HE students conducted by the World Health Organisation (WHO) reported 12-month prevalence rates of 17% (generalised anxiety disorder) and 19% (major depressive episode).⁹

Mental health difficulties can have a negative effect on several aspects of university and personal life, including reduced engagement with campus life, impaired academic performance, and lower graduation rates.^{8,10,11} Importantly, students may experience reduced quality of life.¹² Elevated prevalence of suicide ideation in HE students is also evident,^{10,13} and could lead to higher levels of suicide attempts and death by suicide.¹⁴ Difficulties with study skills are also related to student mental health. For example, examination stress and procrastination have been associated with mental health difficulties,^{15,16} suggesting that support to overcome these study challenges could have a positive impact on mental health in students.

Despite reported levels of both clinical and subclinical psychological disorders, few students receive treatment. According to a US study, only 15% of HE students experiencing moderate or severe depression received treatment, and less than 16% of those with suicide ideation were receiving help.¹⁷ There may be many reasons students do not seek help, with US students reporting that they view stress as a normal part of studying in HE, they wish to

deal with stress alone, and they do not have time for treatment.¹⁸ Stigma also likely plays a role. Students may be subject to personal or public stigma, with certain characteristics, such as race or religiosity, determining the level of stigma experienced.¹⁸ Higher personal stigma has been significantly associated with lower levels of help-seeking in a sample of over 5,500 college students.¹⁸ Moreover, male and minority students may be among those least likely to seek help for mental health problems due to stigma.^{18,19}

Student support services (SSS) exist onsite to address and manage students' mental health difficulties. There is increasing demand for such services, which SSS may struggle to meet due to lack of funding. Moreover, several further barriers that prevent students accessing SSS have been identified, including stigma (as discussed above), lack of time, and concerns about confidentiality.²⁰ Finally, the recent COVID-19 pandemic has highlighted the need for uninterrupted support in unprecedented times. Internet-based support systems could provide a solution to many of these difficulties by offering cost-effective support that can be accessed 24/7.²¹ Moreover, such interventions may be particularly well-suited to HE students who are unlikely to seek help elsewhere, and could offer a proactive approach focused on mental health promotion, including for distance learners.^{22,23} Internet-based systems can also provide support for mild to moderate difficulties allowing SSS to focus their limited resources on severe cases.

There has been a recent proliferation of online support systems, with increasing evidence supporting their use for mental health difficulties in various populations.²⁴⁻³¹ According to a recent review, among the 16 online support systems available to students, six were specifically designed for HE students.³² Most systems use CBT or mindfulness to support students and target depression, anxiety, stress, and study skills. Some also tackle other specific difficulties, such as relationships, substance abuse, eating disorders, and suicide. There is also a mix of formats, with some systems using multimedia (video, audio

and text), while others are text-based. Importantly, just four of these six systems have been subjected to empirical research to determine their effectiveness.³² Therefore, further research is necessary to establish the efficacy and effectiveness of online support systems in student populations.

Here, we report the results of a feasibility study of a proposed online support system, called MePlusMe, which is being developed specifically for HE students. Similar to other systems, CBT interventions presented in multimedia form are used to support students with symptoms of anxiety, depression, and stress, as well as helping with study skills. Unlike other systems, MePlusMe also looks to improve general wellbeing, including in those without psychological symptoms. More details on the system are provided in the Methods. In this study, we tested a simulation of the system to assess effects of its use and inform the next stage of its development.

Methods

Study design, participants, and procedure

The current study was based on methods outlined in a previously published protocol, with some necessary deviations due to sample size.³³ A within-subjects repeated measures study was conducted between October 2018 and February 2019, with students at Rīga Stradiņš University, Latvia. The study received ethical approval from the research ethics committee of Rīga Stradiņš University (Nr.6-3/1).

Participants were recruited via an advertisement sent out through the university's formal information channels (email, posters, and the official social media account). No incentives for participation were provided. Interested students completed initial self-report questions to identify severe difficulties and risk factors such as self-harm, substance abuse, and unusual sensory experiences or beliefs. Any students reporting these indicators were excluded from the study and immediately provided with clear details of HEI services and other sources of support (see protocol for full risk screening details³³). Students reporting no difficulties, or mild to moderate psychological and/or academic-related difficulties were eligible for inclusion.

Included participants completed demographic and baseline measures online before attending a group study day at which the MePlusMe system and one example video technique were presented, followed by an offline simulation of the system. Students completed the initial Questionnaire or Library form (see *MePlusMe* below) to inform the selection of appropriate video techniques, which were then provided digitally for further practice at home in the subsequent weeks.

Following the initial presentation, students provided feedback on the design and functionality of the system, and video format of the techniques. Outcome measures were

subsequently obtained at weeks 2, 4, and 8. At weeks 4 and 8, students also provided further feedback regarding their engagement with the video techniques and potential effects of use.

MePlusMe

MePlusMe is an English-language, web-based support system being developed to support HE students experiencing mild to moderate psychological difficulties or academic difficulties, and students without difficulties wanting to learn how to take care of their mental health and improve their study skills. It can be used alone or to complement traditional support services (for example, while waiting to receive support or following intervention). The final system will assess the risk of severe mental health difficulties and where relevant, signpost students to more appropriate sources of support.

Following risk assessment, eligible students will access videos with psycho-education and techniques via two routes, a symptoms-route ('Questionnaire'; in which identified symptoms are linked to recommended techniques) or a techniques-route ('Library'; where users choose which techniques they wish to learn). As they practise techniques over time, users will be able to monitor their progress with responses to a single item global function question – 'How are you getting on now in your daily life?'.

Motivational reminders and facilitated social networking will also be built into the system to encourage engagement and mutual support ('MyPlace'). Moreover, embedded filters will identify any users who develop more significant difficulties and refer them to seek further support from SSS or primary care. MePlusMe has been rigorously developed thus far, founded on current best practice and evidence-based techniques to ensure the system is fit for purpose.³⁴ Further details regarding the system and its development have been published elsewhere.³²⁻³⁶ A screenshot of the system is included in Appendix A.

Clinical measures

At baseline and at weeks 2, 4 and 8, students completed the Patient Health Questionnaire (PHQ-9),³⁷ Generalized Anxiety Disorder Scale (GAD-7),³⁸ Warwick-Edinburgh Mental Wellbeing scale (WEMWBS),³⁹ and Academic Self-Efficacy Scale (SEF).⁴⁰ All measures can be seen in Appendix B.

The PHQ-9 is a validated measure of depressive symptoms, widely used in primary care and general populations including HE students. It has nine items scored on four-point Likert scales ranging from ‘*not at all*’ (scoring “0”) to ‘*nearly every day*’ (scoring “3”). Total scores range from 0 to 27, with higher scores indicating greater severity of depressive symptoms. ... We examined PHQ-9 reliability among the study participants at baseline and identified a high level of internal consistency: Cronbach’s alpha value was 0.821 and item-total correlations (ITCs) ranged from 0.353 to 0.761.

The GAD-7 is a well-validated measure of generalized anxiety. It has seven items scored on four-point Likert scales ranging from ‘*not at all*’ (scoring “0”) to ‘*nearly every day*’ (scoring “3”). Total scores range from 0 to 21, with higher scores indicating greater severity of anxiety symptoms. The GAD-7 was found to be highly reliable in this study population: Cronbach’s alpha value was 0.851 and ITCs were between 0.498 and 0.735.

The WEMWBS was developed to measure mental wellbeing among the general population with an initial validation sample comprising university students together with general survey participants. It has 14 items scored on five-point Likert scales ranging from ‘*none of the time*’ (scoring “1”) to ‘*all of the time*’ (scoring “5”). Total scores range from 14 to 70, with higher scores indicating better mental wellbeing. For the WEMWBS, internal consistency in this sample was 0.833 and ITCs 0.440 to 0.708.

The original SEF developed with US students comprises 27 statements describing positive study-related behaviours scored on a 10-point scale (‘*not at all confident*’ scoring

“0” to ‘*extremely confident*’ scoring “10”). This scale was adapted (with permission) to: (i) adapt wording to the European setting (e.g., “term papers” replaced by “coursework”) and (ii) remove items with limited relevance (“Having more tests in the same week” and “Getting along with family members”). Total self-efficacy scores ranged from 0 to 250, with higher scores indicating greater self-efficacy. For the 25 SEF items used, the study responses indicated high reliability - Cronbach’s alpha was 0.949 and ITCs were between 0.366 and 0.799.

Open-ended responses and satisfaction ratings

In addition to clinical measures, two feedback forms were administered, with open-ended questions, scaled satisfaction items (‘liked’ ... ‘disliked’; ‘poor’ ... ‘very good’), and a question about how participants are getting on in their daily life (everyday functionality), rated from 1 (‘*extremely well*’) to 7 (‘*not well at all*’). They were also asked whether they had experienced any changes in their study skills and whether they had contacted anyone for help with them.

At the initial assessment, questions were asked on system design, general functionality, and video format. Further questions were administered at weeks 4 and 8 on system use, perceived benefits and user satisfaction. See Appendix B for the questions asked at each time point.

Data analysis

Descriptive statistics (means and standard deviations or standard errors) are reported for sample characteristics and outcomes. Linear mixed model (LMM) analyses were conducted with SPSS v25⁴¹ to examine differences in scores between time points for each outcome. LMM analyses were subsequently conducted to explore the main effects of time and

demographic/academic factors (e.g., gender, qualification level), and their potential interactions. p values were compared against α set at 0.05. Where interaction effects were significant, post hoc tests were conducted within time points and within each level of a factor to decompose effects (with Sidak's correction for multiple comparisons⁴²). Satisfaction ratings were examined with descriptive analyses. The textual data collected through the open-ended questions in the questionnaires was analysed using thematic analysis.⁴³ Coding was carried out independently by two members of the research team, and the agreed categories and patterns derived from the written comments are presented in tabular form in the Results. Data are available on request.

Results

Sample characteristics

Following recruitment, 137 students were enrolled in the study (mean age: 22.6 (\pm 3.38) years; male: $n=39$; female: $n=98$). The majority of participants were EU students ($n=76$, 55.5%), while 32.1% ($n=44$) were home students, and 12.4% ($n=17$) were international students. Most participants (60%) were in their first ($n=33$) or second ($n=36$) year of study, and most (83%) were studying medicine.

Following initial recruitment, 68 participants were retained at week 2, 45 at week 4, and 26 at week 8. Given this high dropout rate, LMM analyses were conducted to retain as much data as possible (Table 1).⁴⁴

[Insert Table 1]

Clinical measures

There was a significant reduction in anxiety across the eight weeks ($F_{(3,53.26)}=9.57$, $p<.001$), with baseline anxiety levels significantly higher than at all follow-up time points. There was also a significant reduction in depression across the intervention period ($F_{(3,55.82)}=4.94$, $p=.004$), with analysis at individual timepoints showing the greatest reduction by week 2, with non-significant changes in later weeks. Wellbeing scores steadily increased across time ($F_{(3,46.99)}=3.04$, $p=.038$), and there were significantly greater scores at week 8 than at baseline. The only measure not showing significant change was academic self-efficacy.

[Insert Figure 1]

LMM analyses were also conducted to determine the effect on the results of gender, qualification level, study program, accommodation, and whether participants opted for reminders.

Only the provision of reminders had a significant interaction effect with time on mental wellbeing and self-efficacy scores (see Table 1). While there were no significant differences between mental wellbeing scores at each time point for those who received reminders and those who did not, there were significantly greater scores at week 8 ($M = 54.95$, $SE = 3.86$) than at week 4 ($M = 44.45$, $SE = 3.15$) in the no-reminder group. Despite a significant interaction effect of the provision of reminders over time on self-efficacy scores, no significant differences were indicated by post hoc tests (either within time-point, or within-groups).

[Insert Figure 2]

[Insert Figure 3]

Use and impressions of the system

Reasons for participation. Participants gave various reasons for taking part in the study, with 39% ($n = 54$) reporting study difficulties and 33% ($n = 45$) reporting emotional or psychological difficulties. However, the most common reason for participating in the study was simply wanting to find out more about MePlusMe (61%, $n = 83$).

System design. Participants rated various design features of the ‘Questionnaire’, ‘Library’, ‘MyPlace’, and video techniques on a five-point scale from ‘very poor’ to ‘very good’. The ‘Questionnaire’, ‘Library’, and ‘MyPlace’ were all found to be easy to use and useful, with

clear wording (rated by over 83% of participants as ‘good’ or ‘very good’). Responses to open-ended questions supported this, however areas identified for improvement included a lack of relevance to their personal situation and limited or over-simplified content.

Both the illustrations and narrator’s voice in the videos were considered ‘good’ or ‘very good’ by the majority of participants (84% and 89%, respectively). The storyboard and ability to maintain interest were also positively rated (‘good’ or ‘very good’ by 95% and 85.5% of participants, respectively). Open-ended questions suggested that participants thought the videos and animations were high quality and the voices were soothing but negative aspects included that videos were not high definition, and the introduction could not be skipped.

Table 2 provides an overview of some of the strengths and areas for potential improvement in terms of system design reported by participants. As a whole, the system was reported to be clear, with an attractive colour scheme. Positive aspects of functionality included the ability to personalise content. The archiving and reminders features were seen as useful, and several participants commented that being an online system enhanced the user experience. A few areas for further development and improvement were identified. Some participants commented that the colour scheme was unengaging, and the interface was plain, while some illustrations were poorly produced. In terms of functionality, negative comments related to an overly complicated design and low resolution of videos.

The written and video content was found by participants to be positive, relevant, and motivational. They found that content helped them deal with their emotions and that goals set were realistic. The main concern regarding content was that it was too generic, with the questionnaire being singled out as not identifying certain feelings and having somewhat irrelevant questions, as well as being too long-winded.

[Insert Table 2]

Use of the system. At week 4, the largest proportion of students reported having only watched the videos once. However, the frequency increased by week 8, with 69% watching them 1–3 times each month. Frequency of practising techniques reduced over time, with most participants reporting a frequency of 1–3 times a week at week 4 (46%) and 1–3 times a month at week 8 (39%). Various reasons were reported for low frequency of use, including lack of time or not finding the videos or techniques personally relevant or useful. However, some participants found they could use the techniques as needed and did not need to revisit the videos as they had already internalised the content.

Most participants (70%) reported they did not access other types of support (for either feelings or study skills support) during the study, with reasons including being too busy or fear of being judged by others. However, at week 8, all those who did not seek help reported this was because it was not needed. Most participants (57% at week 4; 72% at week 8) reported that if they needed support in the future they would seek help from university resources, as well as MePlusMe.

Perceived benefits. In addition to clinical measures, participants were asked if they had experienced any change in their feelings or study skills across the study. The majority reported a perceived change in how they felt at both week 4 (74%) and 8 (76%), with improvements such as feeling more confident, optimistic, relaxed and rational, as well as having fewer breakdowns. However, for some there was an increased pressure to study and they felt more stressed. At week 4, only half of the participants reported a change in study skills, but by week 8 the majority (64%) reported a change. Improvements included increased concentration and motivation, as well as better time management.

Over the duration of the study, there were improvements in the participant's reports of their overall function, as identified by responses to the question about how they were getting on with their daily life. At baseline, a minority of participants (14%) reported function at the highest levels ('extremely well' or 'very well'), rising to a third by week 4, and half of the participants at week 8.

Satisfaction. The majority of participants were 'satisfied' or 'very satisfied' with the system at all timepoints (62% at baseline; 52% at week 4; 56% at week 8). Reasons for satisfaction included the system being straightforward to use, having learned some good techniques via the videos, and improved time management. Dissatisfied participants reported that the system was too simplistic for complex problems or not relevant to their current situation. At 4 and 8 weeks, the majority of participants also said they were either 'likely' or 'very likely' to recommend the system to a friend (59% at week 4; 76% at week 8) and were planning to continue using it (61% at week 4; 64% at week 8). Reasons included the benefits they had experienced and that it was likely to help others. For those who were not sure whether they would continue use, lack of time, forgetfulness and already knowing the techniques were cited as reasons.

Suggested improvements. Finally, participants were asked to suggest any possible improvements for the overall system. Overarching improvements included more specific information on study techniques and managing stress/mood, as well as targeting students early in their university career and including more techniques. In terms of functionality, improvements included better design, better animations, more interactivity, and increased personalisation. Further difficulties that participants suggested the system could address included homesickness, social skills, and dealing with family expectations.

Discussion

The present study investigated an online mental health and study skills intervention, MePlusMe, designed specifically for students attending HEI. The 8-week intervention was well-received, with results indicating a significant positive effect of its continued use. Across the intervention period, there were significant improvements in validated measures of anxiety, depression, and mental wellbeing. These clinical measures exhibited a high degree of reliability in our study population, and changes in these factors were not influenced by participant characteristics such as gender, qualification level, etc. In addition, participants reported a range of positive outcomes in response to open-ended questions, with initial views of the intervention being generally positive. However (and as noted below), these changes need to be interpreted in light of the high rates of participant loss to follow-up.

At baseline, mean scores for anxiety and depression were within the ‘mild’ range (5 – mild; 10 – moderate; 15 – severe). There was a small but significant decrease in depression scores over time, the mean anxiety score at final follow up was below the ‘mild’ cut off (GAD-7 score at week 8, 4.80 ± 0.71). As well as improvement in anxiety and depression symptoms, we identified positive changes in participants’ wellbeing. MePlusMe targets common psychological and emotional problems at lower levels of severity, such as mild to moderate depression and mixed anxiety and depressive symptoms. In contrast, other systems designed for HE students have been designed for more severe cases, which makes it difficult to compare results directly. Nonetheless, our finding that the intervention reduced anxiety and depression is similar to studies of these systems.^{45,46}

Despite improvements in mood and wellbeing, we did not find any statistically significant improvements in self-efficacy associated with system use. Self-efficacy is associated with both motivation and learning effectiveness. Prior research has indicated that digital intervention can lead to improvements in self-efficacy, where targeted

psychoeducational tools are used to increase self-efficacy through study skills training.⁴⁷ It is possible that the interventions in MePlusMe are not effective in targeting and improving self-efficacy. There may also be further unmeasured confounding or moderating variables (such as academic performance and feedback) that are relevant here. The extent of changes that we identified indicate that our study was under-powered to identify change in this variable. Alternatively, changes in self-efficacy may occur over the longer term, such that they were not captured in the current study period.

This study not only addressed potential effectiveness of the MePlusMe simulation in improving symptoms of psychological problems and their impact, but also investigated the acceptability of MePlusMe in terms of layout, functionality and content through open-ended questions. Overall, responses were positive for all these areas, while students also indicated some areas for development. Students' answers indicated that the system was easy to use, well laid out, and relevant. These responses were further supported by ratings of the layout, content, and usefulness of system features, content including the library of techniques and the video techniques. High ratings (of 'good' or 'very good') were given by the majority of users for the layout and ease of navigation of the 'Library', the video techniques, and the clarity of phrasing. Moreover, 87% rated the overall layout and ease of navigation of the system as 'very good' or 'good'. These ratings suggest that students were highly satisfied with the system in terms of functionality, content, and layout.

While users completing the current study indicated high satisfaction with MePlusMe, there was, unfortunately, a high attrition rate. This reflects a recognised difficulty in engaging university students in mental health interventions.⁴⁸ There are several possible explanations for dropout rates in the current study. Possibly, the length or format of the intervention did not appeal to all students. In addition, all participants had access to the video techniques regardless of their continued participation in the study. Therefore, it cannot be known

whether or not participants who failed to complete follow up measures were still using the videos but were not motivated to provide follow up feedback.

While high, the attrition rate seems comparable with other studies investigating digital mental health interventions. A recent systematic overview of studies of such interventions in young people reported that high dropout is common, with completion rates ranging from as low as 10% up to 94%, while other studies do not report attrition rates.⁴⁹ Some elements of design can ameliorate this dropout, including having some in-person elements,⁴⁹ something that could be considered in future versions of the system. Regardless, the high attrition rate is a limitation of this study, which compromises the validity of the study findings

Future directions

Given the increasing burden of psychological difficulties among HE students, there is a need to develop solutions that can overcome some of the barriers to access for those seeking support. Online interventions are one such solution and our data supports the potential for MePlusMe, an online intervention designed specifically for HE students, to lead to positive improvements in outcomes among a diverse HE student population. An improved study design, with a control group and initial randomisation would yield more robust evidence for efficacy. Moreover, addressing loss to follow-up is a key consideration for further research, and the use of incentives may be an important way to manage this potential sources of bias.⁵⁰

Conclusion

In this study, we observed significant positive changes in clinical measures associated with use of a simulation of MePlusMe, an online intervention specifically designed for students. Moreover, qualitative data suggest that initial views of the intervention were generally positive. Despite the study's limitations, these results provide qualified but positive support

for the feasibility and effectiveness of MePlusMe and can be used to further inform its future development.

References

1. Kessler RC, Amminger GP, Aguilar-Gaxiola S, et al. Age of onset of mental disorders: a review of recent literature. *Curr Opin Psychiatry* 2007; 20(4): 359–364. Available from: <https://doi.org/10.1097/YCO.0b013e32816ebc8c>.
2. Pedrelli P, Nyer M, Yeung A, et al. College students: mental health problems and treatment considerations. *Acad Psychiatry* 2015; 39(5): 503–511. Available from: <http://dx.doi.org/10.1007/s40596-014-0205-9>.
3. Ibrahim AK, Kelly SJ, Adams C, et al. A systematic review of studies of depression prevalence in university students. *J Psychiatr Res* 2013; 47(3): 391–400. Available from: <https://doi.org/10.1016/j.jpsychires.2012.11.015>.
4. Blanco C, Okuda M, Wright C, et al. Mental health of college students and their non-college-attending peers: results from the national epidemiologic study on alcohol and related conditions. *Arch Gen Psychiatry* 2008; 65(12): 1429–1437. Available from: <https://doi.org/10.1001/archpsyc.65.12.1429>.
5. Yi S, Peltzer K, Pengpid S, et al. Prevalence and associated factors of illicit drug use among university students in the association of southeast Asian nations (ASEAN). *Subst Abuse Treat Prev Policy* 2017; 12: 9. Available from: <https://doi.org/10.1186/s13011-017-0096-3>.
6. Bennett TH, Holloway KR. Drug use among college and university students: findings from a national survey. *J Subst Use* 2015; 20: 50–55. Available from: <https://doi.org/10.3109/14659891.2013.878762>.
7. Andrews B, Wilding JM. The relation of depression and anxiety to life-stress and achievement in students. *Br J Psychol* 2004; 95: 509–521. Available from: <https://doi.org/10.1348/0007126042369802>.
8. Auerbach RP, Mortier P, Bruffaerts R, et al. Mental disorder comorbidity and suicidal thoughts and behaviors in the World Health Organization World Mental Health Surveys International College Student initiative. *Int J Methods Psych Res* 2019; 28(2): e1752. Available from: <https://dx.doi.org/10.1002%2Fmpr.1752>.
9. Auerbach RP, Mortier P, Bruffaerts R, et al. The WHO World Mental Health Surveys International College Student Project: Prevalence and Distribution of Mental Disorders. *J Abnorm Psychol* 2018; 127(7): 623–638. Available from: <https://doi.org/10.1037/abn0000362>.
10. Garlow SJ, Rosenberg J, Moore JD, et al. Depression, desperation, and suicidal ideation in college students: results from the American Foundation for Suicide Prevention College Screening Project at Emory University. *Depress Anxiety* 2008; 25(6): 482–488. Available from: <https://doi.org/10.1002/da.20321>.
11. Downs MF, Eisenberg D. Help seeking and treatment use among suicidal college students. *J Am Coll Health* 2012; 60(2): 104–114. Available from: <https://doi.org/10.1080/07448481.2011.619611>.
12. Pillay N, Ramlall S, Burns JK. Spirituality, depression and quality of life in medical students in KwaZulu-Natal. *South Afr J Psych* 2016; 22(1): 731. Available from: <https://dx.doi.org/10.4102%2Fsajpsychiatry.v22i1.731>.
13. Cukrowicz KC, Schlegel EF, Smith PN, et al. Suicide Ideation Among College Students Evidencing Subclinical Depression. *J Am Coll Health* 2011; 59: 575–581. Available from: <https://doi.org/10.1080/07448481.2010.483710>.
14. Garlow SJ, Rosenberg J, Moore JD, et al. Depression, desperation, and suicidal ideation in college students: results from the American Foundation for Suicide Prevention College Screening

- Project at Emory University. *Depress Anxiety* 2008; 25: 482–488. Available from: <https://doi.org/10.1002/da.20321>.
15. Hailikari T, Katajavuori N, Asikainen H. Understanding procrastination: A case of a study skills course. *Soc Psychol Edu* 2021; 24(2): 589–606.
 16. Jones E, Priestly M, Brewster L et al. Student wellbeing and assessment in higher education: the balancing act. *Assessment & Evaluation in Higher Education* 2021; 46(3): 438–450.
 17. Eisenberg D, Hunt J, Speer N et al. Mental health service utilization among college students in the United States. *J Nerv Ment Dis* 2011; 199: 301–308. Available from: <https://doi.org/10.1097/nmd.0b013e3182175123>.
 18. Eisenberg D, Downs MF, Golberstein E, et al. Stigma and help seeking for mental health among college students. *Med Care Res Rev* 2009; 66: 522–541. Available from: <https://doi.org/10.1177/1077558709335173>.
 19. Loya F, Reddy R, Hinshaw SP. Mental illness stigma as a mediator of differences in Caucasian and South Asian college students' attitudes toward psychological counseling. *J Couns Psychol* 2010; 57: 484–490. Available from: <https://psycnet.apa.org/doi/10.1037/a0021113>.
 20. Givens JL, Tjia J. Depressed medical students' use of mental health services and barriers to use. *Acad Med* 2002; 77: 918–921. Available from: <https://doi.org/10.1097/00001888-200209000-00024>.
 21. Andersson G, Titov N. Advantages and limitations of Internet-based interventions for common mental disorders. *World Psychiatry* 2014; 13: 4–11. Available from: <https://doi.org/10.1002/wps.20083>.
 22. Ryan ML, Shochet IM, Stallman HM. Universal online interventions might engage psychologically distressed university students who are unlikely to seek formal help. *Adv Ment Health* 2010; 9: 73–83. Available from: <https://doi.org/10.5172/jamh.9.1.73>
 23. Lister K, Seale J, Douce C. Mental health in distance learning: a taxonomy of barriers and enablers to student mental wellbeing. *Open Learning: The Journal of Open, Distance and e-learning* 2021. Available from: <http://oro.open.ac.uk/75544/>.
 24. Stratton E, Lampit A, Choi I, et al. Effectiveness of eHealth interventions for reducing mental health conditions in employees: a systematic review and meta-analysis. *PLoS ONE* 2017; 12: e0189904. Available from: <https://doi.org/10.1371/journal.pone.0189904>.
 25. Schaub MP, Blankers M, Lehr D, et al. Efficacy of an internet-based self-help intervention to reduce co-occurring alcohol misuse and depression symptoms in adults: study protocol of a three-arm randomised controlled trial. *BMJ Open* 2016; 6: e011457. Available from: <https://doi.org/10.1136/bmjopen-2016-011457>.
 26. Ebert DD, Cuijpers P, Muñoz RF, et al. Prevention of mental health disorders using internet- and mobile-based interventions: a narrative review and recommendations for future research. *Front Psychiatry* 2017; 8: 116. Available from: <https://doi.org/10.3389/fpsy.2017.00116>.
 27. Andersson G, Carlbring P, Ljótsson B, et al. Guided internet-based CBT for common mental disorders. *J Contemp Psychother* 2013; 43: 223–233. Available from: <https://doi.org/10.1007/s10879-013-9237-9>.
 28. Christensen H, Batterham P, Mackinnon A, et al. Prevention of generalized anxiety disorder using a web intervention, iChill: randomized controlled trial. *J Med Internet Res* 2014; 16: e199. Available from: <https://doi.org/10.2196/jmir.3507>.

29. Jeong S, Alghowinem S, Aymerich-Franch L et al. A robotic positive psychology coach to improve college students' wellbeing. 29th IEEE International Conference on Robot and Human Interactive Communication (RO-MAN). 2020. doi: 10.1109/RO-MAN47096.2020.9223588.
30. Lattie E, Cohen KA, Winkquist N et al. Examining an app-based mental health self-care program, IntelliCare for College Students: single-arm pilot study. *JMIR Ment Health* 2020; 7(10): e21075. doi: 10.2196/21075.
31. Levin ME, Krafft J, Davis CH. et al. Evaluating the effects of guided coaching calls on engagement and outcomes for online acceptance and commitment therapy. 2021. *Cogn Behav Ther* 2021; 50(5): 395–408.
32. Papadatou-Pastou M, Goozee R, Payne E, et al. A review of web-based support systems for students in higher education. *Int J Ment Health Syst* 2017; 11: 59. Available from: <https://doi.org/10.1186/s13033-017-0165-z>.
33. Papadatou-Pastou M, Goozee R, Barley EA, et al. Online intervention, 'MePlusMe', supporting mood, wellbeing, study skills, and everyday functioning in students in higher education: a protocol for a feasibility study. *Pilot Feasibility Stud* 2015; 1: 34. Available from: <http://dx.doi.org/10.1186/s40814-015-0029-8>.
34. Barrable A, Papadatou-Pastou M, Tzotzoli P. Supporting mental health, wellbeing and study skills in Higher Education: an online intervention system. *Int J Ment Health Syst* 2018; 12: 54. Available from: <https://doi.org/10.1186/s13033-018-0233-z>.
35. Papadatou-Pastou M, Campbell-Thomson L, Barley E, et al. Exploring the feasibility and acceptability of the contents, design, and functionalities of an online intervention promoting mental health, wellbeing, and study skills in Higher Education students. *Int J Ment Health Syst* 2019; 13: 51. Available from: <https://doi.org/10.1186/s13033-019-0308-5>.
36. Touloumakos AK, Goozee RM, Papadatou-Pastou M, et al. Online support system for students in higher education: proof of concept study. *Digit Health* 2016; 2: 2055207616655012. Available from: <https://dx.doi.org/10.1177%2F2055207616655012>.
37. Spitzer RL, Williams JBW, Kroenke K, et al. Validity and utility of the PRIME-MD Patient Health Questionnaire in assessment of 3000 obstetric-gynecologic patients: the PRIME-MD Patient Health Questionnaire Obstetrics-Gynecology Study. *Am J Obstet Gynecol* 2000; 183: 759–769. Available from: <https://doi.org/10.1067/mob.2000.106580>.
38. Spitzer RL, Kroenke K, Williams JBW, et al. A brief measure for assessing Generalized Anxiety Disorder. The GAD-7. *Arch Intern Med* 2006; 166(10): 1092–1097. Available from: <https://doi.org/10.1001/archinte.166.10.1092>.
39. Tennant R, Hiller L, Fishwick R, et al. The Warwick-Edinburgh Mental Well-Being Scale (WEMWBS): development and UK validations. *Health Qual Life Outcomes* 2007; 5: 63. Available from: <https://dx.doi.org/10.1186%2F1477-7525-5-63>.
40. Zajacova A, Lynch SM, Espenshade TJ. Self-efficacy, stress, and academic success in college. *Res High Educ* 2005;46(6):677–706. Available from: <https://doi.org/10.1007/s11162-004-4139-z>.
41. IBM Corp. Released 2020. IBM SPSS Statistics for Windows, Version 25.0. Armonk, NY: IBM Corp
42. Šidák ZK. Rectangular confidence regions for the means of multivariate normal distributions. *J Am Stat Assoc* 1967; 62(318): 626–633. Available from: <https://doi.org/10.1080/01621459.1967.10482935>.
43. Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol* 2006; 3(2): 77–101.
44. Heck RH, Thomas SL, Tabata LN. Multilevel and longitudinal modeling with IBM SPSS, second edition. New York: Routledge, 2014.

45. Lynch S, Gander ML, Kohls N, et al. Mindfulness-based coping with university life: a non-randomized wait-list-controlled pilot evaluation. *Stress Heal* 2011; 27: 365–375. Available from: <https://doi.org/10.1177%2F2158244018758379>.
46. Musiat P, Conrod P, Treasure J, et al. Targeted prevention of common mental health disorders in university students: randomised controlled trial of a transdiagnostic trait-focused web-based intervention. *PLoS ONE* 2014; 9(4): 1–10. Available from: <https://doi.org/10.1371/journal.pone.0093621>.
47. Padilla Rodriguez BC, Armellini A. Developing self-efficacy through a massive open online course on study skills. *Open Praxis* 2017; 9(3): 335–343.
48. Melville KM, Casey LM, Kavanagh DJ. Dropout from Internet-based treatment for psychological disorders. *Br J Clin Psychol.* 2010; 49(Pt 4): 455–471. Available from: <https://doi.org/10.1348/014466509x472138>.
49. Lehtimaki S, Martic J, Wahl B, et al. Evidence on digital mental health interventions for adolescents and young people: systematic overview. *JMIR Ment Health* 2021; 8(4): e25847. doi: 10.2196/25847.
50. Wong E. Incentives in longitudinal studies, CLS Working Paper 2020/1. 2020. London: UCL Centre for Longitudinal Studies. Available from: <https://cls.ucl.ac.uk/wp-content/uploads/2020/04/CLS-working-paper-2020-1-Incentives-in-longitudinal-studies.pdf>

Tables and Figures

Table 1. Linear mixed models analyses showing changes in each outcome over time

Outcome ($M \pm SE$)	Time (weeks)				Effect of time			
	0	2	4	8	DF ₁	DF ₂	F	<i>P</i>
Anxiety	8.23 ^a ± 0.37	6.28 ^b ± 0.44	6.29 ^b ± 0.52	4.80 ^b ± 0.71	3	53.26	9.57	<0.001
Depression	8.49 ^a ± 0.42	6.92 ^b ± 0.46	7.45 ^{a,b} ± 0.73	6.55 ^{a,b} ± 0.82	3	55.82	4.94	0.004
Wellbeing	45.13 ^a ± 0.73	46.49 ^{a,b} ± 0.98	47.24 ^{a,b} ± 1.29	49.97 ^b ± 1.62	3	46.99	3.04	0.038
Academic self-efficacy	178.54 ± 3.49	182.38 ± 4.44	185.43 ± 6.05	196.57 ± 7.43	3	45.96	2.11	0.112

For each analysis, superscript letters indicate scores at each time point that are significantly different.

Table 2. Perceived strengths and areas of improvement identified by participants

Strengths	Areas for improvement
<p><i>Layout</i></p> <p>I thought it was designed in a clear way to find out exactly what my main problems are and how to help me with them (<i>Participant 29</i>)</p> <p>Well thought out concept, which includes every aspect of a student's life (<i>Participant 119</i>)</p> <p>Simple layout, ability to tailor needs (<i>Participant 15</i>)</p> <p>It is made so that it feel as if you are in your personal space, your little room (<i>Participant 35</i>)</p>	<p>The website is a little old fashioned in its design and could benefit from being perhaps more modern, particularly as your target group is students (<i>Participant 105</i>)</p> <p>The layout seemed a bit stiff and kind of too much emphasized studying (<i>Participant 4</i>)</p> <p>A more modern, clearer and minimalistic website design (<i>Participant 68</i>)</p> <p>It is fair and one would find it useful [...], but user experience really matters, so the design and layout shows room for improvement. More user friendly, more images instead of words as well. People rather have expressions than words (<i>Participant 83</i>)</p>
<p><i>Functionality</i></p> <p>It is quite easy to navigate through the portal. I like that it is kept simple without much distraction (<i>Participant 41</i>)</p> <p>This is a very good way to personalize help for students and their needs for a support (<i>Participant 72</i>)</p> <p>I like how you can personalize it according to your liking (<i>Participant 59</i>)</p> <p>It seems very straightforward (<i>Participant 44</i>)</p>	<p>Make it more interactive. If you're giving students study techniques, follow through and have them do things on the site to keep them engaged (<i>Participant 43</i>)</p>
<p><i>Content</i></p> <p>Just pure and easily understandable information that actually matches perfectly what I was looking for to cope with stress (<i>Participant 1</i>)</p> <p>Clarity of the information make[s] it useful for anyone (<i>Participant 94</i>)</p> <p>Specific features have names that makes it easy to understand what it is about so the clarity of the phrasing is understandable (<i>Participant 34</i>)</p> <p>I like that they have videos with solutions to almost anything that one might experience during university (<i>Participant 81</i>)</p>	<p>Terminology is kept rather general, which makes it quite hard to specify what is actually meant. More selection possibilities should be possible (<i>Participant 57</i>)</p> <p>Had a good selection but I feel like it could have more alternatives (<i>Participant 63</i>)</p> <p>Options can be elaborated a bit more, more relaxation techniques can be added, more user-initiated program structure that pushes the participant to record and build progress, and feel like he/she is improving using the system (<i>Participant 19</i>)</p> <p>Making it more interesting – making real situations, not animations. Looks sometimes too childish (<i>Participant 54</i>)</p>




Figure 1. Changes in anxiety (A), depression (B), mental wellbeing (C) and self-efficacy (D) over 8 weeks of treatment.




Figure 2. Changes in mental wellbeing scores over time for participants receiving or not receiving reminders.




Figure 3. Changes in self-efficacy scores over time for participants receiving or not receiving reminders.