



City Research Online

City St George's, University of London

Citation: King, N., Lewis, E. G., Kinnison, T., Langridge, A., Civai, C., May, S. A. & Cardwell, J. M. (2024). Mental health of veterinary nurses and student veterinary nurses: A scoping review. *Veterinary Record*, 194(9), no-. doi: 10.1002/vetr.4091

This is the published version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/33232/>

Link to published version: <https://doi.org/10.1002/vetr.4091>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

REVIEW

Mental health of veterinary nurses and student veterinary nurses: A scoping review

Naomi King¹  | Elisa G. Lewis²  | Tierney Kinnison³  | Alison Langridge⁴  |
Claudia Civali²  | Stephen Anthony May³  | Jacqueline M. Cardwell¹ 

¹Department of Pathobiology and Population Sciences, Royal Veterinary College, Hatfield, UK

²Division of Psychology, School of Applied Sciences, London South Bank University, London, UK

³Department of Clinical Science and Services, Royal Veterinary College, Hatfield, UK

⁴The College of Animal Welfare & CAW Business School, Godmanchester, UK

Correspondence

Naomi King, Department of Pathobiology and Population Sciences, Royal Veterinary College, Hawkshead Lane, North Mymms, Hatfield, Hertfordshire, AL9 7TA, UK.
Email: nking@rvc.ac.uk

Funding information

IDEXX Foundation

Abstract

Background: Research has shown that veterinarians around the world are at high risk of mental health problems, but far less research has examined the mental health of veterinary nurses (VNs) and student veterinary nurses (SVNs). This scoping review aimed to map existing evidence on this topic and identify knowledge gaps.

Methods: Literature searches of MEDLINE, PsycINFO, PubMed, Web of Science and Google Scholar were conducted, and a structured screening and selection procedure was applied. To be included, studies had to be peer reviewed, report relevant results specific to VNs and/or SVNs, and provide descriptive statistics if using quantitative methods.

Results: Of the 2118 publications identified, only 13 journal articles met the inclusion criteria. The findings were summarised in five categories: mental health and wellbeing, burnout, stress, compassion fatigue and moral distress. While the findings of five of the studies suggested that some VNs and SVNs experienced some form of poor mental health, these studies lacked generalisability or transferability for multiple reasons. There was also inconsistency and ambiguity in the interpretation of findings, as well as incompatible or oversimplified definitions of mental health problems.

Limitations: Our review excluded grey literature, such as reports, theses and conference presentations, as a preliminary search found very little empirical research on VNs' and SVNs' mental health in this type of publication.

Conclusions: More research is needed to address the gaps in the existing evidence supporting our understanding of VN and SVN mental health. This should establish baseline measures and include comparisons with other occupational and national populations.

INTRODUCTION

Research on mental health in veterinary professionals around the world has focused predominantly on veterinarians, finding that they are at high risk of stress, anxiety, depression, burnout, compassion fatigue and suicidal ideation.¹ Potential risk factors include heavy workloads, long working hours, poor work–life balance, unreasonable client expectations and demands, ethical and moral challenges, and the performance of euthanasia.^{1,2} Research using validated psychometric scales, in the UK and United States, also suggests that veterinary students experience poorer wellbeing and mental health than approximately age-

matched general population groups^{3,4} or some other student groups.^{5,6} Far less research has focused on veterinary nurses (VNs) or student veterinary nurses (SVNs), despite their vital role in effective veterinary teamwork.⁷

Some anecdotal evidence^{8,9} suggests that VNs and SVNs experience a range of mental health problems, but it is unclear whether they differ from the general population. The Royal College of Veterinary Surgeons reported that a 2019 survey of 4993 UK VNs¹⁰ found lower mean Warwick–Edinburgh Mental Wellbeing Scale (WEMWBS) scores, suggesting poorer wellbeing, in VNs than in the general population of England in 2016.¹¹ However, the difference was not significant

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Authors. *Veterinary Record* published by John Wiley & Sons Ltd on behalf of British Veterinary Association.

TABLE 1 Search terms used.

Database or search engine	Search terms
MEDLINE, PsycINFO, Web of Science (advanced search)	'veterinary nurse' OR 'veterinary nursing' OR 'veterinary professional' OR 'veterinary practice' (first search box) AND 'mental health' OR 'wellbeing' OR 'well-being' OR 'mental disorder' OR 'mental diagnosis' OR 'mental illness' OR 'mental problem' OR 'mental condition' OR 'psychological' OR 'anxiety' OR 'stress' OR 'distress' OR 'depression' OR 'depressive' OR 'burnout' OR 'burned out' OR 'compassion fatigue' OR 'grief' OR 'moral distress' OR 'moral injury' OR 'depersonalisation' OR 'depersonalization' OR 'emotional turmoil' OR 'cynicism' OR 'cynical' OR 'fatigue' OR 'insomnia' OR 'exhaustion' OR 'suicide' OR 'suicidal' (second search box)
PubMed (advanced search)	'veterinary nurse' OR 'veterinary nursing' OR 'veterinary professional' OR 'veterinary practice' (first search box) AND 'mental health' OR 'wellbeing' OR 'well-being' OR 'mental disorder' OR 'mental diagnosis' OR 'mental illness' OR 'mental problem' OR 'mental condition' OR 'psychological' (second search box)
Google Scholar	'veterinary nurse' OR 'veterinary nursing' AND 'mental health' OR 'wellbeing' OR 'well-being' OR 'mental disorder' OR 'mental diagnosis' OR 'mental illness' OR 'mental problem' OR 'mental condition' OR 'psychological'

tested, it was not a contemporaneous comparison and the scores were not sufficiently low to be indicative of depression.¹² In another investigation, which examined the effects of sustained bullying in the veterinary profession,¹³ 390 VNs reported experiencing at least one form of bad behaviour or bullying, almost one-third of whom related it to poor mental health in free-text responses. In a recent Mind Matters and VN Futures survey,¹⁴ completed by around 650 SVNs, recently qualified VNs and clinical coaches, 96% agreed that bullying and incivility were problems in the profession, 81% agreed that their work was stressful and 75% thought that the demands of work and studying affected their wellbeing.

However, to date, there are no comprehensive reviews of empirical studies assessing the mental health of VNs and SVNs, and therefore no clear evidence synthesis. This review addresses this gap by focusing on two research questions: (1) how prevalent are mental health problems in VNs and SVNs? and (2) what types of mental health problems do VNs and SVNs experience? It conceptualises 'mental health problems' in a broad sense, including but not limited to clinically diagnosed mental health disorders or illnesses.

METHODS

A scoping review was conducted in line with the framework proposed by Arksey and O'Malley¹⁵ and expanded by Levac et al.¹⁶ Scoping reviews are optimal for mapping existing evidence and identifying knowledge gaps, especially for topics that are emerging and underexplored, in contrast to systematic reviews that focus on more specific questions and critical appraisal.^{15,17} Where appropriate, the review was reported in line with the Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) checklist.¹⁸

Information sources and search strategy

Four electronic databases (MEDLINE, PsycINFO, PubMed and Web of Science) were used, in addition to Google Scholar, which has been found to add value to literature searches.^{19,20} Searches were restricted to journal articles, but no publication date limit was applied. The search terms are provided in Table 1. The number of these was reduced for PubMed and Google Scholar, as initial searches generated 2,657,174 publications from PubMed and about 9750 from Google Scholar. The terms 'veterinary technician', 'veterinary technologist' and 'vet tech' were not included. Although these roles in the United States and Canada are similar to those of the VN in the UK, Ireland and Australasia, the definitions and terminologies vary between states and countries. For clarity and consistency, this review focused only on VNs. The final search produced 229 publications from MEDLINE, 23 from PsycINFO, 110 from PubMed, 296 from Web of Science and about 1460 from Google Scholar. Manual searching of reference lists was conducted on all publications subjected to full-text screening.

Inclusion criteria

Inclusion criteria (Figure 1) were developed iteratively during screening and selection, in line with Levac et al.'s¹⁶ recommendations.

Screening and selection procedure

Titles of the 658 publications from the four databases and the first 950 publications from Google Scholar were screened in March 2023 by the first author (Naomi King). The last 400 of these 950 publications were clearly irrelevant, so the remaining Google Scholar publications were excluded without screening. Of the 1608 title-screened publications, 1414 were




FIGURE 1 Inclusion criteria and sub-criteria. SVN, student veterinary nurse; VN, veterinary nurse




FIGURE 2 Flow diagram of the screening and selection procedure

removed because they were duplicates or did not meet the inclusion criteria. A further 162 of the remaining 194 publications were removed after abstract screening because they did not meet the inclusion criteria. Thirty-two publications were then subjected to full-text screening, and another 21 publications were removed because they did not meet inclusion criteria (i) to (iii). A second search of all databases and Google Scholar was conducted in September 2023 by the first author, and seven new publications were screened in the same way. Figure 2 outlines the full screening and selection procedure. The final sample comprised 13 publications.

RESULTS

Overview of publications reviewed

Of the 13 studies reviewed, the majority ($n = 8$) were published between 2020 and 2023, and all but one were published between 2016 and 2023 (Table S1). Seven studies recruited only VNs and/or SVNs, while six included other veterinary professional roles. Studies were located in Australasia ($n = 8$), the UK ($n = 4$) and Portugal ($n = 1$). Ten collected data relevant to this review from validated questionnaires, one of which also included an open-text question. Of

the remaining studies, one used non-validated questions, one used semi-structured interviews and one gathered open-text responses. The majority of studies had relatively small sample sizes; six analysed relevant data from fewer than 100 VNs and/or SVNs, four from fewer than 170, two from fewer than 290 and one from 992. No study directly aimed to investigate whether VNs and/or SVNs experienced mental health problems in general. Instead, they focused on specific, pre-defined mental health problems and/or explored potential causes, contexts, coping strategies or outcomes.

The findings are summarised below in five categories: mental health and wellbeing, burnout, stress, compassion fatigue and moral distress. In our synthesis, we have not included issues such as job dissatisfaction, disillusionment or poor physical health; these issues were examined by some of the studies reviewed, but while they are associated with poor mental health, they are not mental health problems in themselves. We also excluded positive aspects of mental health, such as compassion satisfaction, as low levels do not necessarily denote poor mental health and people can experience compassion satisfaction simultaneously with compassion fatigue.²¹ The relevant data are shown in Tables S2–S6.

Mental health and wellbeing

Four studies examined the broad concepts of mental health and wellbeing, three of which included only VNs/SVNs (Table S2). Bedford and Anscombe-Skirrow²² focused primarily on disillusionment resulting from workplace bullying, but they also reported how the open-text responses of 273 VNs and SVNs showed that bullying led to ‘impairment of self-confidence’, ‘hindrance to working ability through decreased concentration and enjoyment’ and ‘colleague and task avoidance’. ‘Mental health’ and ‘reduced self-worth’ were mentioned by around 54% and 69% of participants, respectively, but it was unclear whether these were researcher-created categorisations or terms used by participants. No indications of question phrasing, analysis methods, evidence (e.g., participant quotes) or interpretations were provided. Deacon and Brough²³ examined the psychological impact of exposure to patient death and client bereavement using interview data from 26 VNs and former VNs who self-identified as having experienced occupational stress. Participants reported a range of distressing emotions, as well as high levels of ‘psychological strain’ and symptoms consistent with burnout and post-traumatic stress. However, there were some nuances; for some, compassionate euthanasia, for example, led not only to adverse emotions but also to a strong sense of job satisfaction and increased personal resilience. Van Soest and Fritschi²⁴ focused predominantly on physiological health hazards encountered by 147 VNs (145 of whom were female) from Australia and Tasmania but also assessed ‘job-related affective well-

being’ using the anxiety-contentment and depression-enthusiasm axes of Warr’s²⁵ scale. The mean scores were similar (not significance tested) to those of Warr’s 847 female workers in comparably skilled jobs in the UK, suggesting that the VNs did not experience poorer mental health. However, these two populations were 14 years apart and in different countries.

The study by Mair et al.²⁶ examined the ‘mental wellbeing’ of 451 equine veterinary staff, including 20 equine VNs/SVNs, in the UK during the COVID-19 pandemic using the WEMWBS. The VNs/SVNs had a lower mean score, suggesting poorer wellbeing, than the veterinary surgeons in the same study and the equine veterinary staff in a pre-pandemic 2019 survey,²⁷ but these differences in means were not significance tested. The authors noted that while a low score could be interpreted as indicative of possible depression, it should be interpreted extremely cautiously due to the very low number of VNs and SVNs surveyed.

Overall, only Bedford and Anscombe-Skirrow’s²² study suggested that poor mental health was common in VNs/SVNs, but this was based on open-text responses about the effects of bullying, which could not reliably identify poor mental health or capture problems unrelated to bullying. In contrast, Van Soest and Fritschi’s²⁴ study suggested that female VNs did not experience lower wellbeing than other female workers, but the comparison with a non-contemporaneous population was unreliable. Deacon and Brough²³ and Mair et al.²⁶ presented results from very small, specific participant samples, so their findings, while interesting, cannot be generalised.

Burnout

Eight studies examined the specific concept of ‘burnout’ (Table S3). The World Health Organization defines this as an occupational phenomenon resulting from chronic workplace stress, characterised by three dimensions: feelings of energy depletion or exhaustion; increased mental distance from one’s job, or feelings of negativism or cynicism related to one’s job; and reduced professional efficacy.²⁸ However, there are many other conceptualisations, and a systematic review of research on ‘occupational burnout’ found 88 unique definitions across 248 studies.²⁹ The term is also often used in common parlance to describe commonplace experiences such as tiredness and loss of creativity, which is inconsistent with formal definitions.³⁰ Figure 3 summarises conceptualisations of burnout used in the reviewed studies.

Five studies, two focusing solely on VNs/SVNs, used the Professional Quality of Life Scale (ProQOL^{31,32}). Beetham et al.³³ reported a mean burnout score of 28.94, indicating moderate risk, in 166 VNs surveyed during the COVID-19 pandemic. Having expected a higher score, the authors suggested that the ProQOL may have failed to capture the true levels of stress in a pandemic or that the VNs simply had good coping




FIGURE 3 Summary of how the measuring scales used by the reviewed studies conceptualise burnout

mechanisms and resilience. Despite almost identical results (mean score of 28.84 in 992 VNs), Smith³⁴ concluded that as 92.8% of VNs were at moderate/high risk of burnout, working as a VN puts workers at high risk of suffering from it. This highlights the fact that ProQOL results can be interpreted in different ways.

Three studies used the ProQOL in mixed veterinary professional groups. Foote³⁵ presented data from 370 veterinary professionals, including 169 VNs, and

found similar mean burnout scores (not significance tested) for VNs, veterinary surgeons, patient care assistants and receptionists. These findings were similar to those of Beetham et al.³³ and Smith,³⁴ and were interpreted as showing ‘moderate levels’ of burnout. Rohlf et al.³⁶ found no significant difference between mean burnout scores for 93 VNs and 43 veterinarians. Scotney et al.³⁷ reported a mean burnout score of 24.8 for all 229 participants, and although a smaller

percentage of the 67 VNs (14.9%) than the 69 veterinarians (34.8%) fell in the high-risk burnout category, variation across groups was not significant. The authors gave two self-contradictory interpretations of their findings, observing in the abstract and results that 'low burnout was reported by 78% of participants', having combined the low and moderate scores, but stating in the discussion that VNs were an 'at-risk group for burnout', having combined the moderate and high scores.

Of the three studies using alternative scales, one focused solely on VNs. Deacon and Brough³⁸ reported that 53% of the 144 VNs who had completed the work-related subscale of the Copenhagen Burnout Inventory³⁹ had a high degree of burnout, referring to Kristensen et al.'s³⁹ criteria, although these criteria are unclear and variously interpreted elsewhere.^{40,41} The mean score in Deacon and Brough's participants (49.48) was higher (not significance tested) than the mean score (33.0) for human healthcare professionals, including hospital doctors, nurses, midwives and social workers.³⁹ Deacon and Brough³⁸ concluded that high levels of burnout were prevalent in VNs.

Ashton-James and McNeillage⁴² surveyed 249 veterinary professionals, including 77 VNs, from a single specialist Australian veterinary hospital during the COVID-19 pandemic. A subset of 239 respondents completed the Maslach Burnout Inventory – General Survey.⁴³ The clinical staff, who accounted for 67.1% of participants and included VNs, were significantly more emotionally exhausted than non-clinical staff, but there were no significant differences on the cynicism or professional efficacy subscales. For the VNs, only their emotional exhaustion mean score exceeded the authors' threshold for high levels. Varela and Correia⁴⁴ analysed data from 229 veterinarians and 96 VNs who had completed a Portuguese adaptation of the Oldenburg Burnout Inventory,⁴⁵ for which higher scores on the 1–5 response scale indicated a higher risk of burnout. The mean scores were identical between VNs and veterinarians for the exhaustion subscale and similar for the disengagement subscale (not significance tested).

Overall, only Deacon and Brough's³⁸ study suggested that VNs experienced high levels of burnout, although it must be noted that their participants were employed in one Australian state, and the comparative norms were from a study conducted 12 years earlier in Denmark. In studies using the ProQOL, all mean scores for VNs fell on the lower side of the moderate-risk category, but these scores were variously interpreted. Stamm³² does not define the ProQOL's use of 'moderate' but does affirm that 'moderate to low' burnout is optimal, suggesting that only high scores are a cause for concern.

Stress

Five studies examined 'stress' (Table S4). Two investigated general forms of stress, focusing solely on VNs.

Harvey and Cameron⁴⁶ stated that just over two-thirds of their 288 VNs had responded to what appeared to be a single, bespoke item about self-defined stress, and while 49% reported feeling stressed 'always', 'most of the time' or 'half of the time', 46% reported feeling stressed just 'some of the time' and 5% 'never'. The authors interpreted this as showing a 'high incidence of stress', despite acknowledging that participants who did not experience stress may have ignored the question and that an objective means of determining stress levels was required. Van Soest and Fritsch²⁴ reported that, in response to an open-text question about any occupational health issues not raised in the questionnaire, 'many' of their 147 VNs mentioned 'mental stress' but provided no further detail. In both studies, the term 'stress' was open to subjective interpretation by participants.

Three studies used the ProQOL to investigate secondary traumatic stress, described by Stamm³² as the effects of 'work-related, secondary exposure to people who have experienced extremely or traumatically stressful events'. Symptoms may include fear, sleep difficulties, intrusive images and an inability to separate one's private life from one's life as a helper. Smith³⁴ reported a mean secondary traumatic stress score of 25.52, on the low side of the moderate category, in 992 VNs, but stated that 68.1% were at moderate/high risk of secondary traumatic stress, indicating a problematic level. Rohlf et al.³⁶ found no significant difference between mean secondary traumatic stress scores for VNs and veterinarians, which were both in the moderate-risk category. Scotney et al.³⁷ reported a mean secondary traumatic stress score of 24.6 for all participants, with no significant differences between occupational groups.

Overall, only Harvey and Cameron's⁴⁶ study provided some evidence of stress in VNs, but this was based on a single question in which stress was not defined and which almost one-third of participants chose not to answer. Similarly, Van Soest and Fritsch²⁴ did not define 'mental stress', and their reporting of results was ambiguous. The studies analysing secondary traumatic stress found, as with burnout, that all of the VNs' mean scores fell at the lower end of the moderate-risk category.

Compassion fatigue

Three studies investigated 'compassion fatigue' (Table S5), characterised by Stamm³² as 'the negative aspects of providing care to those who have experienced extreme or traumatic stressors'. Two of these studies^{33,35} appeared to have used an older version of the ProQOL, which conceptualised compassion fatigue and secondary traumatic stress as synonymous and interchangeable³¹ (Figure 3). This differs from the current (fifth) version of the ProQOL used by the three studies discussed in the above subsection,^{34,36,37} which instructs researchers to report separate subscores for burnout and secondary traumatic stress,

as opposed to a combined compassion fatigue score.

Beetham et al.³³ reported a mean compassion fatigue score of 26.74 in 166 VNs, which was at the lower end of the moderate-risk category. Foote³⁵ did not report a mean score for their 169 VNs but indicated that it was between the other groups' mean scores of 24.67 and 28.00 (not significance tested), therefore mirroring Beetham et al.'s³³ findings. Harvey and Cameron⁴⁶ used a single, bespoke item about self-defined compassion fatigue to which approximately two-thirds of their 288 VNs responded, and while 33.5% reported experiencing it 'always', 'most of the time' or 'half of the time', 48.5% reported 'some of the time' and 18% reported 'never'. This was interpreted by the authors as demonstrating a 'high incidence' of compassion fatigue; however, they again acknowledged that participants who did not experience compassion fatigue may have ignored the question and that a more objective measure was required, as the term could be misunderstood or variously interpreted. Overall, there was no clear indication from any study that VNs experienced concerning levels of compassion fatigue.

Moral distress

Two studies examined 'moral distress' (Table S6), described as distress occurring when a person is 'unable to carry out what they believe to be the right course of action because of real or perceived constraints on that action',⁴⁷ and considered to be prevalent among healthcare providers.⁴⁸ Deacon and Brough's²³ VN interviewees 'often' struggled with moral distress due to conflict between needing to respect clients' wishes and wanting to act in patients' best interests, and were 'fraught with emotional anguish' when euthanasia duties conflicted with personal morals. These findings were not intended to be generalised to a wider VN/SVN population. Using the Measure of Moral Distress for Healthcare Professionals,⁴⁸ Foote³⁵ found that VNs appeared to experience moderate levels of moral distress, while veterinary surgeons, patient care assistants and veterinary receptionists experienced low levels. This is consistent with human healthcare research suggesting that nurses' moral distress is intensified by their lower position in the staff hierarchy and their lack of power in decision making.⁴⁹ However, the mean moral distress score for the VNs in Foote's³⁵ study nevertheless fell on the lower side of the moderate category, and apparent differences across groups were not significance tested.

DISCUSSION

Our first question was, 'How prevalent are mental health problems in VNs and SVNs?' While evidence from five of the 13 reviewed studies suggested that

some VNs and SVNs experienced some form of poor mental health, these findings lacked generalisability or transferability for reasons such as small participant numbers and recruitment of participants from specific groups (e.g., equine staff, those who self-identified as having occupational stress), locations (e.g., one Australian state) and contexts (e.g., the COVID-19 pandemic). These limitations, along with the shortcomings of convenience sampling and the likelihood of self-selection bias, were acknowledged in some articles.

No study set out to estimate the prevalence of poor mental health in general, so findings relevant to this review were largely based on single validated scales, individual non-validated items or open-text questions, which provided either very specific results (e.g., pertaining to one particular mental health problem) or ambiguous results (e.g., the proportion of participants mentioning 'mental health'). The only qualitative study²³ purposively recruited participants who self-identified as being affected by occupational stress and focused on the effects of exposure to patient death and client bereavement. Participants in all studies may have experienced mental health problems that fell outside the scope of the research and were therefore overlooked.

There was considerable inconsistency and ambiguity across studies in interpretation of findings (e.g., varying interpretations of similar ProQOL results), and even some self-contradictory interpretations. Where open-text questions or unvalidated items were used to explore issues such as mental health,²² mental stress,²⁴ and stress and compassion fatigue,⁴⁴ it was unclear whether or not these terms were introduced and defined by the researchers; if not, such terms are open to various interpretations. Stress is a particularly ambiguous concept, as although intense or chronic forms can negatively affect people's mental health, minor, short-term forms are experienced by everyone, and can have positive effects such as improving motivation.⁵⁰

Even validated scales may not fully capture participants' experiences; this was noted by Beetham et al.,³³ in relation to the use of the ProQOL during the COVID-19 pandemic. However, others have found the ProQOL to be questionable in regular circumstances. In a systematic meta-analysis of 27 international, peer-reviewed studies examining the ProQOL's psychometric structure, Hotchkiss and Wong⁵¹ reported several factorial and internal structure issues and called for the development of 'more parsimonious, reliable and valid measures'. The ProQOL manual itself advises users that their scores may not accurately reflect their professional quality of life and that, unless persistent, high-risk burnout scores may simply indicate that they are having a bad day or need some time off. For those who fall in the high-risk secondary traumatic stress category, Stamm³² states, 'While higher scores do not mean that you do have a problem, they are an indication that you may want to examine how you feel about your work and your work environment'.

Several studies did not test the significance of apparent differences when comparing VN mean values of scales with those of other populations, meaning that any conclusions drawn may be incorrect.

Our second question was, 'What types of mental health problems do VNs and/or SVN experience?' While several studies concluded that VNs had a high risk of experiencing burnout, stress or secondary traumatic stress, compassion fatigue and/or moral distress, only one study by Deacon and Brough³⁸ provided relatively clear evidence. None of the studies focusing on specific problems included SVN, and none examined anxiety or depression, which are considered the most common mental disorders globally.^{52,53}

Multiple, incompatible definitions of mental health problems presented another difficulty, and there were often inconsistencies within studies. For example, Scotney et al.³⁷ characterised burnout in the introduction and discussion as incorporating the three components of emotional exhaustion, depersonalisation and low personal accomplishment, consistent with the original Maslach Burnout Inventory. However, in their study, they used the ProQOL, which conceptualises burnout as an element of compassion fatigue with no subcomponents of its own. Beetham et al.³³ and Foote³⁵ cited the ProQOL version 5 but presented data for compassion satisfaction, burnout and compassion fatigue separately, in line with an older version.³¹ Beetham et al.³³ also referred to secondary traumatic stress as an umbrella term for compassion fatigue and burnout, a conceptualisation that did not match any versions of the ProQOL or the publication they cited,⁵⁴ but later suggested that the term was synonymous with compassion fatigue.³¹ Several studies presented oversimplified descriptions of the problem(s) they set out to measure, implying that there was general acceptance of a single definition rather than numerous, heterogeneous views.

Our review excluded grey literature, such as reports, theses, conference presentations, working papers and media articles, which can provide valuable insights and reduce publication bias.^{55,56} However, they also vary considerably in quality and rigour, can be time and resource consuming to find and evaluate, and are difficult to compare to academic journal articles due to diverse lengths and formats.^{55,56} A preliminary search of grey literature prior to this review found very little empirical research on the mental health of VNs and SVN, with the exception of the reports mentioned in the introduction.^{10,13,14} We also excluded studies that did not provide appropriate descriptive statistics, or present relevant data separately for VNs and/or SVN if the sample included other participants. While these articles may have provided interesting findings, they could not contribute to answering our research questions. Additionally, potential risk factors, including demographics, individual characteristics (e.g., self-efficacy, emotional intelligence), coping strategies and sources of support, were not reviewed. Before examining these issues, baseline estimates of the prevalence of poor mental health in VNs and SVN

are required. Future reviews could expand our boundaries to include United States and Canadian veterinary technicians and veterinary technologists.

CONCLUSIONS

As observed by Davidson,^{57,58} veterinary nursing is a profession often overlooked by clients, colleagues and employers. Our review suggests that it has also been overlooked by researchers, highlighting considerable gaps in the evidence supporting our understanding of the mental health of VNs and SVN. Future research should include the assessment of this in relation to other occupational and national populations, which would help to establish whether the VN profession is different in any way. Longitudinal studies would provide a more consistent overview and counterbalance temporary issues. Recognition of the crucial role that VNs play in interprofessional teams, and the need to protect their mental health in order to maintain effective performance, job satisfaction, career commitment and optimal clinical outcomes, should be promoted.

AUTHOR CONTRIBUTIONS

Conceptualisation and funding acquisition: all authors. *Investigation and writing—original draft:* Naomi King and Jacqueline M. Cardwell. *Supervision:* Jacqueline M. Cardwell. *Writing—review and editing:* all authors.

ACKNOWLEDGEMENTS

This study was funded by the IDEXX Foundation.

CONFLICT OF INTEREST STATEMENT

The authors declare they have no conflicts of interest.

DATA AVAILABILITY STATEMENT


Data sharing is not applicable as no original data were generated.

ETHICS STATEMENT


No ethical approval was required for this study.

ORCID

Naomi King  <https://orcid.org/0000-0002-9887-3031>

Elisa G. Lewis  <https://orcid.org/0000-0002-5759-9734>

Tierney Kinnison  <https://orcid.org/0000-0001-6845-2140>

Claudia Civali  <https://orcid.org/0000-0002-6745-2074>

Stephen Anthony May  <https://orcid.org/0000-0003-1813-7745>

Jacqueline M. Cardwell  <https://orcid.org/0000-0002-9065-3253>

REFERENCES

- Pohl R, Botscharow J, Böckelmann I, Thielmann B. Stress and strain among veterinarians: a scoping review. *Ir Vet J.* 2022;75(1):15.

2. Moir F, Van den Brink A. Current insights in veterinarians' psychological wellbeing. *N Z Vet J.* 2020;68(1):3–12.
3. Cardwell JM, Lewis EG, Smith KC, Holt ER, Baillie S, Allister R, et al. A cross-sectional study of mental health in UK veterinary undergraduates. *Vet Rec.* 2013;173(11):266.
4. Lewis EG, Cardwell JM. A comparative study of mental health and wellbeing among UK students on professional degree programmes. *J Furth High Educ.* 2019;43(9):1226–38.
5. Hafen M, Reisbig AMJ, White MB, Rush BR. Predictors of depression and anxiety in first-year veterinary students: a preliminary report. *J Vet Med Educ.* 2006;33(3):432–40.
6. Karaffa KM, Hancock TS. Mental health experiences and service use among veterinary medical students. *J Vet Med Educ.* 2019;46(4):449–58.
7. Kinnison T, May SA. Evidence-based healthcare: the importance of effective interprofessional working for high quality veterinary services, a UK example. *Vet Evid.* 2016;1(4).
8. Bonnema L. A registered veterinary nurse's (RVN's) secret struggle ...: a short communication on mental health in the profession. *Vet Nurs J.* 2017;32(3):63–65.
9. Hunt G. Understanding and overcoming the effects of compassion fatigue within the veterinary profession. *Vet Nurs J.* 2017;32(5):141–43.
10. Robinson D, Edwards M, Akehurst G, Cockett J, Graham KA, Martin A. The 2019 survey of the veterinary nurse profession: a report for the Royal College of Veterinary Surgeons. Brighton, UK: Institute for Employment Studies; 2019. Accessed 20 Mar 2024. Available from: <https://www.rcvs.org.uk/news-and-views/publications/the-2019-survey-of-the-veterinary-nursing-profession/>
11. NHS Digital. Health survey for England, 2016. 2017. Accessed 20 Mar 2024. Available from: <https://digital.nhs.uk/data-and-information/publications/statistical/health-survey-for-england/health-survey-for-england-2016>
12. Warwick Medical School. Collect, score, analyse and interpret WEMWBS. 2023. Accessed 20 Mar 2024. Available from: <https://warwick.ac.uk/fac/sci/med/research/platform/wemwbs/using/howto/>
13. Everitt S. Behaviour in veterinary practice. 2017. Accessed 20 Mar 2024. Available from: <https://www.vetsurgeon.org/behaviour.pdf>
14. Mind Matters, VN Futures. Report of the student veterinary nursing wellbeing discussion forum. 2021. Accessed 20 Mar 2024. Available from: <https://vetmindmatters.org/wp-content/uploads/2022/01/SVN-wellbeing-discussion-forum-2021-report.pdf>
15. Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol.* 2005;8(1):19–32.
16. Levac D, Colquhoun H, O'Brien KK. Scoping studies: advancing the methodology. *Implement Sci.* 2010;5(1):69.
17. Munn Z, Peters MDJ, Stern C, Tufanaru C, McArthur A, Aromataris E. Systematic review or scoping review? Guidance for authors when choosing between a systematic or scoping review approach. *BMC Med Res Methodol.* 2018;18(1):143.
18. Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. *Ann Intern Med.* 2018;169(7):467–73.
19. Bramer WM, Rethlefsen ML, Kleijnen J, Franco OH. Optimal database combinations for literature searches in systematic reviews: a prospective exploratory study. *Syst Rev.* 2017;6(1):245.
20. Gusenbauer M, Haddaway NR. Which academic search systems are suitable for systematic reviews or meta-analyses? Evaluating retrieval qualities of Google Scholar, PubMed, and 26 other resources. *Res Synth Methods.* 2020;11(2):181–217.
21. Dehlin M, Lundh LG. Compassion fatigue and compassion satisfaction among psychologists: can supervision and a reflective stance be of help? *J Pers Oriented Res.* 2018;4(2):95–107.
22. Bedford S, Anscombe-Skirrow E. Bullying and disillusionment in veterinary nursing. *Vet Nurs J.* 2018;33(9):250–56.
23. Deacon RE, Brough P. Companion animal death and client bereavement: a qualitative investigation of veterinary nurses' caregiving experiences. *Death Stud.* 2021;45(10):805–16.
24. Van Soest E, Fritschi L. Occupational health risks in veterinary nursing: an exploratory study. *Aust Vet J.* 2004;82(6):346–50.
25. Warr P. The measurement of well-being and other aspects of mental health. *J Occup Psychol.* 1990;63(3):193–210.
26. Mair TS, Mountford DR, Radley R, Lockett E, Parkin TD. Mental wellbeing of equine veterinary surgeons, veterinary nurses and veterinary students during the COVID-19 pandemic. *Equine Vet Educ.* 2021;33(1):15–23.
27. Robinson D, Edwards M, Mason B, Cockett J, Graham KA, Martin A. The 2019 survey of the veterinary profession: a report for the Royal College of Veterinary Surgeons. Brighton, UK: Institute for Employment Studies; 2019. Accessed 20 Mar 2024. Available from: <https://www.rcvs.org.uk/news-and-views/publications/the-2019-survey-of-the-veterinary-profession/>
28. World Health Organization. Burn-out an 'occupational phenomenon': international classification of diseases. 2019. Accessed 20 Mar 2024. Available from: <https://www.who.int/news/item/28-05-2019-burn-out-an-occupational-phenomenon-international-classification-of-diseases>
29. Guseva Canu I, Marca SC, Dell'Oro F, Balázs Á, Bergamaschi E, Besse C, et al. Harmonized definition of occupational burnout: a systematic review, semantic analysis, and Delphi consensus in 29 countries. *Scand J Work Environ Health.* 2021;47(2):95–107.
30. Morgan K. Why we may be measuring burnout all wrong. 2021. Accessed 20 Mar 2024. Available from: <https://www.bbc.com/worklife/article/20210426-why-we-may-be-measuring-burnout-all-wrong>
31. Stamm BH. The ProQOL manual. Derwood: Sidran Press; 2005.
32. Stamm BH. The concise ProQOL manual. New York: Eastwoods LLC; 2010.
33. Beetham L, Cameron K, Harvey L. Compassion fatigue, compassion satisfaction and burnout during a pandemic. *Vet Nurs.* 2021;12(9):436–43.
34. Smith N. A questionnaire based study to assess compassion fatigue in UK practising veterinary nurses. *Vet Nurs.* 2016;7(7):418–25.
35. Foote A. Burnout, compassion fatigue and moral distress in veterinary professionals. *Vet Nurs.* 2023;14(2):90–99.
36. Rohlf VI, Scotney R, Monaghan H, Bennett P. Predictors of professional quality of life in veterinary professionals. *J Vet Med Educ.* 2022;49(3):372–81.
37. Scotney RL, McLaughlin D, Keates HL. An investigation of the prevalence of compassion fatigue, compassion satisfaction and burnout in those working in animal-related occupations using the Professional Quality of Life (ProQoL) scale. *Vet Nurs.* 2019;10(5):276–84.
38. Deacon RE, Brough P. Veterinary nurses' psychological wellbeing: the impact of patient suffering and death. *Aust J Psychol.* 2017;69(2):77–85.
39. Kristensen TS, Borritz M, Villadsen E, Christensen KB. The Copenhagen Burnout Inventory: a new tool for the assessment of burnout. *Work Stress.* 2005;19(3):192–207.
40. Creedy DK, Sidebotham M, Gamble J, Pallant J, Fenwick J. Prevalence of burnout, depression, anxiety and stress in Australian midwives: a cross-sectional survey. *BMC Pregnancy Childbirth.* 2017;17(1):13.
41. Thrush CR, Gathright MM, Atkinson T, Messias EL, Benjamin Guise J. Psychometric properties of the Copenhagen Burnout Inventory in an academic healthcare institution sample in the U.S. *Eval Health Prof.* 2021;44(4):400–405.
42. Ashton-James CE, McNeillage AG. A mixed methods investigation of stress and wellbeing factors contributing to burnout and job satisfaction in a specialist small animal hospital. *Front Vet Sci.* 2022;9:942778.
43. Schaufeli WB, Leiter MP, Maslach C, Jackson SE. Maslach Burnout Inventory—General Survey (MBI-GS). In: Maslach C, Jackson SE, Leiter MP, editors. *Maslach Burnout Inventory manual*. 3rd ed. Palo Alto: Consulting Psychologists Press; 1996. p. 19–26.
44. Varela M, Correia I. Empathy and burnout in veterinarians and veterinary nurses: identifying burnout protectors. *Anthrozoos.* 2023;36(1):15–34.

45. Bakker AB, Demerouti E, Verbeke W. Using the job demands-resources model to predict burnout and performance. *Hum Resour Manage.* 2004;43(1):83–104.
46. Harvey LC, Cameron KE. Stress and compassion fatigue in veterinary nurses in New Zealand. *Vet Nurs.* 2020;11(1):42–46.
47. Helmers A, Palmer KD, Greenberg RA. Moral distress: developing strategies from experience. *Nurs Ethics.* 2020;27(4):1147–56.
48. Epstein EG, Whitehead PB, Prompahakul C, Thacker LR., Hamric AB. Enhancing understanding of moral distress: the Measure of Moral Distress for Health Care Professionals. *AJOB Empir Bioeth.* 2019;10(2):113–24.
49. Burston AS, Tuckett AG. Moral distress in nursing. *Nurs Ethics.* 2013;20(3):312–24.
50. National Health Service (NHS). Dealing with stress. 2023. Accessed 20 Mar 2024. Available from: <https://www.nhs.uk/every-mind-matters/mental-health-issues/stress/>
51. Hotchkiss JT, Wong MYC. Factorial structure of the ProQOL—systematic meta-analysis and integration of 27 international factor analysis studies. *Trends Psychol.* 2022:1–32.
52. Institute of Health Metrics and Evaluation, Global Health Data Exchange. GBD results. 2019. Accessed 20 Mar 2024. Available from: <https://vizhub.healthdata.org/gbd-results/>
53. World Health Organization (WHO). Mental disorders. 2023. Accessed 20 Mar 2024. Available from: <https://www.who.int/news-room/fact-sheets/detail/mental-disorders/bib>>
54. Huggard PK, Huggard EJ. When the caring gets tough: compassion fatigue and veterinary care. *VetScript.* 2008;5:14–16.
55. Adams J, Hillier-Brown FC, Moore HJ, Lake AA, Araujo-Soares V, White M, et al. Searching and synthesising 'grey literature' and 'grey information' in public health: critical reflections on three case studies. *Syst Rev.* 2016;5(1):164.
56. Paez A. Gray literature: an important resource in systematic reviews. *J Evid Based Med.* 2017;10(3):233–40.
57. Davidson J. Is veterinary nursing a visible profession? Part one. *Vet Nurs.* 2017;8(7):403–7.
58. Davidson J. Is veterinary nursing a visible profession? Part two. *Vet Nurs.* 2017;8(9):516–18.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: King N, Lewis EG, Kinnison T, Langridge A, Civai C, May SA, et al. Mental health of veterinary nurses and student veterinary nurses: A scoping review. *Vet Rec.* 2024;e4091. <https://doi.org/10.1002/vetr.4091>