



# City Research Online

## City St George's, University of London

**Citation:** Castro-Alvaredo, O. A., Negro, S. & Sailis, F. (2024). Completing the bootstrap program for TT deformed massive integrable quantum field theories. *Journal of Physics A: Mathematical and Theoretical*, 57(26), 265401. doi: 10.1088/1751-8121/ad5395

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/33380/>

**Link to published version:** <https://doi.org/10.1088/1751-8121/ad5395>

**Copyright and Reuse:** Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

# Completing the Bootstrap Program for $T\bar{T}$ -Deformed Massive Integrable Quantum Field Theories

Olalla A. Castro-Alvaredo<sup>♥</sup>, Stefano Negro<sup>♣</sup> and Fabio Sails<sup>◇</sup>

<sup>♥</sup> <sup>◇</sup> Department of Mathematics, City, University of London, 10 Northampton Square EC1V 0HB, UK

<sup>♣</sup> Department of Mathematics, University of York, York YO10 5DD, UK

## Abstract

In recent years a considerable amount of attention has been devoted to the investigation of 2D quantum field theories perturbed by certain types of irrelevant operators. These are the composite field  $T\bar{T}$  – constructed out of the components of the stress-energy tensor – and its generalisations – built from higher-spin conserved currents. The effect of such perturbations on the infrared and ultraviolet properties of the theory has been extensively investigated. In the context of integrable quantum field theories, a fruitful perspective is that of factorised scattering theory. In fact, the above perturbations were shown to preserve integrability. The resulting deformed scattering matrices – extensively analysed with the thermodynamic Bethe ansatz – provide the first step in the development of a bootstrap program. In this paper we present a systematic approach to computing matrix elements of operators in generalised  $T\bar{T}$ -perturbed models, based on employing the standard form factor program. We show that for theories with diagonal scattering and certain types of fields the deformed form factors, factorise into the product of the undeformed ones and of a perturbation- and theory-dependent term. From these solutions, correlation functions can be obtained and their asymptotic properties studied.

**Keywords:**  $T\bar{T}$ -Perturbed Theories, Integrable Quantum Field Theory, Form Factor Program

<sup>♥</sup> o.castro-alvaredo@city.ac.uk

<sup>♣</sup> stefano.negro@york.ac.uk

<sup>◇</sup> fabio.sails@city.ac.uk

July 22, 2024

# 1 Introduction

Generalised  $T\bar{T}$  deformations of 2D quantum field theory (QFT) [1–4] admit a variety of interesting interpretations. From coupling the original QFT to two-dimensional topological gravity [5] or to random geometry [6] to a state-dependent change of coordinates [7], among others. For these reasons,  $T\bar{T}$  perturbations and their generalisations have been intensely studied in 2D quantum field theory [1, 2, 7–10], in the framework of the ODE/IM correspondence [11] (see [12] for a review), employing thermodynamic Bethe ansatz (TBA) [3, 4, 13–18], perturbed CFT [6, 19–25], string theory [26–29], holography [30–37], quantum gravity [5, 38–42], out-of-equilibrium CFT [43–46], in long-range spin chains [47–50], and in the context of the generalised hydrodynamics approach [51–53]. A generalisation of this family of deformations has also been proposed for quantum-mechanical systems [54–56] and higher-dimensional field theories [57–59].

In this paper, we approach the study of generalised  $T\bar{T}$  deformations from the viewpoint of integrable quantum field theory (IQFT) [60, 61], integrability techniques and the bootstrap program. Here we define the latter as consisting of the following steps: 1) computation of the exact scattering matrix, 2) computation of matrix elements (form factors) of local fields, 3) computation of correlation functions, and 4) consistency checks of those form factors and correlation functions, which include the TBA approach and Zamolodchikov’s  $c$ -theorem [62]. To date point 1) and a large part of point 4) (i.e. the TBA treatment) are very well developed, whereas points 2) and 3) and most of 4) are very much outstanding. In particular, although much work on the TBA analysis of the theories has been done, it has not yet been possible to compare TBA results with information obtained from correlation functions. In this paper, we show how these outstanding steps may be carried out for a large family of IQFTs, fields, and perturbations.

It is well known that in IQFT perturbations of  $T\bar{T}$ -type lead to a modification of the exact two-body scattering matrix by a multiplicative or CDD [63] factor. The deformed  $S$ -matrix is

$$S_{\alpha}(\theta) = \Phi_{\alpha}(\theta)S_{\mathbf{0}}(\theta), \quad (1)$$

where  $\alpha = (\alpha_s)_{s \in \mathcal{S} \subset \mathbb{N}}$  is a, possibly infinite, vector,  $S_{\mathbf{0}}(\theta)$  is some undeformed, consistent, factorised  $S$ -matrix in a theory with a single particle, and

$$\Phi_{\alpha}(\theta) = \exp \left[ -i \sum_{s \in \mathcal{S}} \alpha_s \sinh(s\theta) \right]. \quad (2)$$

A generalisation to more complicated spectra is possible along the same lines as presented here. For simplicity, we have taken the fundamental mass scale  $m = 1$ , but it is useful to notice that the combination  $\alpha_s m^{2s}$  is dimensionless.  $\mathcal{S}$  is a set of spin values, typically those of local conserved charges. Since  $\Phi_{\alpha}(\theta)$  is a CDD factor, the theory described by this new  $S$ -matrix is still integrable. The computation of the exact matrix elements (form factors) of local operators corresponding to the family of  $S$ -matrices (1) has not been attempted systematically. As we shall see, the unusual UV physics of  $T\bar{T}$ -perturbed theories, will also emerge clearly when pursuing the form factor program. In this sense, form factors provide a new useful tool to understand this physics. For example, we will find parameter regions, where a distinct asymptotics of correlators can be read off from a form factor expansion and consistently interpreted in the context of the physics of  $T\bar{T}$  perturbations. Pursuing this program will not only shed light on the properties of the correlators of  $T\bar{T}$ -perturbed IQFTs but also reveal the mathematical structure of a new

family of solutions to the form factor equations. Form factors and correlation functions of a theory with similar features were first studied in [64].

The main results of our work are: (1) We find closed solutions for the form factors of local and semi-local fields in generalised  $T\bar{T}$ -perturbed IQFTs and, in some parameter regions, convergent two-point function expansions. (2) When all  $\alpha_i < 0$  two-point functions can be expanded in terms of form factors of the corresponding fields and the expansion is convergent both for long and short distances. For long distances (IR) the theory flows to a trivial (massive) fixed point, just as in the unperturbed case, whereas for short distance (UV) two-point functions still scale as power-laws and the  $c$ -theorem gives a finite result which is a function of the couplings  $\alpha$ . (3) If  $\alpha_s > 0$ , where  $s$  is the largest spin involved in the sum (1), the two-point function is divergent in the UV limit, and also in the IR limit for momenta above a certain threshold which is universally characterised by Lambert's  $W$ -function.

Our paper is organised as follows: In Section 2 we review the form factor equations and minimal form factors in  $T\bar{T}$ -perturbed theories. In Section 3 we present our main result, namely a derivation of the factorisation property of form factors, leading to close formulae for the form factors of  $T\bar{T}$ -perturbed theories with diagonal scattering. In Section 4 we discuss the asymptotics of correlation functions in different parameter regimes. In Section 5 we discuss how Zamolodchikov's  $c$ -theorem can still be evaluated for  $T\bar{T}$ -perturbed theories, giving rise to a function that flows with the RG parameter  $|\alpha|$ . We conclude and discuss some open problems and subtleties of our derivation in Section 6.

## 2 Form Factor Bootstrap

Let

$$F_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \alpha) = \langle 0 | \mathcal{O}(0) | \theta_1, \dots, \theta_n \rangle, \quad (3)$$

be a matrix element of a field  $\mathcal{O}$  between the ground state  $|0\rangle$  and an  $n$ -particle in-state  $|\theta_1, \dots, \theta_n\rangle$  characterised by rapidities  $\{\theta_i\}_n$ . Since we are considering only theories with a single particle we do not need to consider additional quantum numbers in this case. The form factor equations for a diagonal theory without bound states are [65–67]

$$F_n^{\mathcal{O}}(\theta_1, \dots, \theta_i, \theta_{i+1}, \dots, \theta_n; \alpha) = S_{\alpha}(\theta_i - \theta_{i+1}) F_n^{\mathcal{O}}(\theta_1, \dots, \theta_{i+1}, \theta_i, \dots, \theta_n; \alpha), \quad (4)$$

$$F_n^{\mathcal{O}}(\theta_1 + 2\pi i, \theta_2, \dots, \theta_n; \alpha) = \gamma_{\mathcal{O}} F_n^{\mathcal{O}}(\theta_2, \dots, \theta_n, \theta_1; \alpha), \quad (5)$$

and

$$\lim_{\bar{\theta} \rightarrow \theta} (\bar{\theta} - \theta) F_{n+2}^{\mathcal{O}}(\bar{\theta} + i\pi, \theta, \theta_1, \dots, \theta_n; \alpha) = i(1 - \gamma_{\mathcal{O}} \prod_{j=1}^n S_{\alpha}(\theta - \theta_j)) F_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \alpha), \quad (6)$$

with the two first equations constraining the monodromy of the form factors and the last equation, the kinematic residue equation, specifying their pole structure. In (6) we find the parameter  $\gamma_{\mathcal{O}}$  named the factor of local commutativity [68, 69] which can be a non-trivial phase in theories possessing an internal symmetry (such as the Ising model which we have studied in more detail in [70]). The equations (4)-(6) are typically solved by the following ansatz

$$F_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \alpha) = H_n^{\alpha} Q_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \alpha) \prod_{i < j}^n \frac{F_{\min}(\theta_{ij}; \alpha)}{e^{\theta_i} + e^{\theta_j}}. \quad (7)$$

where  $\theta_{ij} := \theta_i - \theta_j$  and  $F_{\min}(\theta; \boldsymbol{\alpha})$  is the minimal form factor, that is, a minimal solution (i.e. without poles in the physical strip) of the form factor equations for two particles,

$$F_{\min}(\theta; \boldsymbol{\alpha}) = S_{\boldsymbol{\alpha}}(\theta)F_{\min}(-\theta; \boldsymbol{\alpha}) = F_{\min}(2\pi i - \theta; \boldsymbol{\alpha}). \quad (8)$$

Note that for spinless fields, form factors are functions of rapidity differences only, hence the minimal form factor depends on a single rapidity variable. For the same reason, one-particle form factors when non-vanishing, must be rapidity-independent. By construction, the product over minimal form factors in (7) provides a solution to equations (4) and (5). That means that the functions  $Q_n^{\mathcal{O}}(\{\theta_i\}_n; \boldsymbol{\alpha})$  must be symmetric and  $2\pi i$ -periodic in all rapidities. In other words, they should be functions of the exponentials  $\{e^{\theta_i}\}_n$ . The parameters  $H_n^{\boldsymbol{\alpha}}$  are normalisation constants and the factors  $e^{\theta_i} + e^{\theta_j}$  in the denominator ensure the correct kinematic pole structure.

Based on the ansatz (7) it is clear that the first step towards solving the form factor equations is the computation of the minimal form factor. For unperturbed IQFTs with diagonal scattering, as considered here, there is a systematic approach for solving the problem, which typically leads to an integral representation of the minimal form factor. It is easy to show that, given a solution for the undeformed theory,  $F_{\min}(\theta; \mathbf{0})$ , a consistent solution for the deformed model can be written as

$$F_{\min}(\theta; \boldsymbol{\alpha}) := \varphi(\theta; \boldsymbol{\alpha})C(\theta; \boldsymbol{\beta})F_{\min}(\theta; \mathbf{0}), \quad (9)$$

with

$$\varphi(\theta; \boldsymbol{\alpha}) := \exp \left[ \frac{\theta - i\pi}{2\pi} \sum_{s \in \mathcal{S}} \alpha_s \sinh(s\theta) \right]. \quad (10)$$

The formula (10) for the minimal form factor of  $T\bar{T}$ -deformed massive integrable theories with diagonal scattering was, to our knowledge, first derived by I. M. Szécsényi in an unpublished work, while its massless version appeared in an investigation in Nambu-Goto theory [3], which later turned out to be the  $T\bar{T}$  deformation of a free, massless, 2-dimensional scalar field theory.

The function  $C(\theta; \boldsymbol{\beta})$  is a minimal form factor ‘CDD’ factor, that is a function such that any solution can be multiplied by it, while still solving all required equations. It takes the form

$$C(\theta; \boldsymbol{\beta}) := \exp \left( \sum_{s \in \mathcal{S}'} \beta_s \cosh(s\theta) \right), \quad (11)$$

where  $\mathcal{S}'$  is a subset of the integers. Note that the sum over  $s$  of  $\alpha_s \sinh(s\theta)$  is a sum over the one-particle eigenvalues of local conserved quantities of spin  $s$ . However, those one-particle eigenvalues are more generically linear combinations of the form  $\alpha_s \sinh(s\theta) + \beta_s \cosh(s\theta)$ . Thus the fact that we can always add a sum over  $s$  of  $\beta_s \cosh(s\theta)$  can be seen as a reflection of this ambiguity in our choice of conserved quantities. It has been recently shown that the structure of the minimal form factor (9) is in fact very natural and also found in standard IQFTs such as the sinh-Gordon model [71].

At this stage the usual procedure to obtain higher particle form factors would be to substitute the ansatz (7) which solves (4) and (5) into equation (6) and find recursive equations for the functions  $Q_n^{\mathcal{O}}(\{\theta_i\}_n; \boldsymbol{\alpha})$  and the constants  $H_n^{\boldsymbol{\alpha}}$ . We have done this for the Ising field theory and obtained closed solutions for all local fields in [70]. However, also in [70] we observed that the solutions had a factorised structure, as the form factors of the undeformed theory, times a new function of the parameters  $\boldsymbol{\alpha}$  (in [70] we set all  $\boldsymbol{\beta} = 0$ ). In the next section, we show that this factorisation holds beyond the example of the Ising field theory.

### 3 Form Factor Factorisation

We can report that, for any IQFT with diagonal scattering matrix and a single-particle spectrum, all solutions have the general structure

$$F_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \boldsymbol{\alpha}) = F_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \mathbf{0}) \Upsilon_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \boldsymbol{\alpha}) , \quad (12)$$

with  $\Upsilon_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \boldsymbol{\alpha})$  a non-trivial function of  $\boldsymbol{\alpha}$  and  $\Upsilon_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \mathbf{0}) = 1$ . Here, as we did in [70] for the Ising field theory, we will restrict ourselves to the case  $\boldsymbol{\beta} = \mathbf{0}$ .

The function  $\Upsilon_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \boldsymbol{\alpha})$  can be computed by plugging (12) into (7). This gives rise to the following equations

$$H_n^{\boldsymbol{\alpha}} = H_n^{\mathbf{0}} \mathcal{H}_n^{\boldsymbol{\alpha}} , \quad F_{\min}(\theta; \boldsymbol{\alpha}) = F_{\min}(\theta; \mathbf{0}) \varphi(\theta; \boldsymbol{\alpha}) , \quad (13)$$

$$Q_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \boldsymbol{\alpha}) = Q_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \mathbf{0}) \Theta_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \boldsymbol{\alpha}) , \quad (14)$$

or equivalently

$$\Upsilon_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \boldsymbol{\alpha}) = \mathcal{H}_n^{\boldsymbol{\alpha}} \Theta_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \boldsymbol{\alpha}) \prod_{i < j}^n \varphi(\theta_i - \theta_j; \boldsymbol{\alpha}) . \quad (15)$$

The quantities  $H_n^{\mathbf{0}}$ ,  $Q_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \mathbf{0})$  and  $F_{\min}(\theta; \mathbf{0})$  are supposed to satisfy the axioms for the undeformed theory, which means

$$\mathcal{H}_n^{\mathbf{0}} = \Theta_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \mathbf{0}) = \varphi(\theta; \mathbf{0}) = 1 . \quad (16)$$

Imposing the form factor equations for the deformed theory we recover the equations<sup>1</sup>

$$\varphi(\theta; \boldsymbol{\alpha}) = \Phi_{\boldsymbol{\alpha}}(\theta) \varphi(-\theta; \boldsymbol{\alpha}) = \varphi(2\pi i - \theta; \boldsymbol{\alpha}) , \quad (17)$$

which have already been solved to (9), and

$$\lim_{\bar{\theta} \rightarrow \theta} \Upsilon_{n+2}^{\mathcal{O}}(\bar{\theta} + \pi i, \theta, \theta_1, \dots, \theta_n; \boldsymbol{\alpha}) = \frac{1 - \gamma_{\mathcal{O}} \prod_{j=1}^n S_{\boldsymbol{\alpha}}(\theta - \theta_j)}{1 - \gamma_{\mathcal{O}} \prod_{j=1}^n S_{\mathbf{0}}(\theta - \theta_j)} \Upsilon_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \boldsymbol{\alpha}) . \quad (18)$$

This equation in turn splits into

$$\mathcal{H}_{n+2}^{\boldsymbol{\alpha}} = \mathcal{H}_n^{\boldsymbol{\alpha}} , \quad (19)$$

$$\Theta_{n+2}^{\mathcal{O}}(\theta + i\pi, \theta, \theta_1, \dots, \theta_n; \boldsymbol{\alpha}) = \mathbb{S}_n(\theta - \theta_1, \dots, \theta - \theta_n; \gamma_{\mathcal{O}}) \Theta_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \boldsymbol{\alpha}) , \quad (20)$$

where we introduced the quantity

$$\mathbb{S}_n(\theta_1, \dots, \theta_n; \gamma_{\mathcal{O}}) = \frac{\prod_{j=1}^n S_{\boldsymbol{\alpha}}(\theta_j)^{-1/2} - \gamma_{\mathcal{O}} \prod_{j=1}^n S_{\boldsymbol{\alpha}}(\theta_j)^{1/2}}{\prod_{j=1}^n S_{\mathbf{0}}(\theta_j)^{-1/2} - \gamma_{\mathcal{O}} \prod_{j=1}^n S_{\mathbf{0}}(\theta_j)^{1/2}} . \quad (21)$$

Notice that as a consequence of the unitarity  $S_{\boldsymbol{\alpha}}(-\theta) = S_{\boldsymbol{\alpha}}(\theta)^{-1}$  and crossing symmetry  $S_{\boldsymbol{\alpha}}(i\pi - \theta) = S_{\boldsymbol{\alpha}}(\theta)$  of the  $S$ -matrix, the function (21) satisfies the following properties

$$\mathbb{S}_n(\theta_1 \pm i\pi, \dots, \theta_n \pm i\pi; \gamma_{\mathcal{O}}) = \mathbb{S}_n(\theta_1, \dots, \theta_n; \gamma_{\mathcal{O}}^{-1}) , \quad (22)$$

<sup>1</sup>Note that if  $\gamma_{\mathcal{O}} = 1$  and  $S(\theta) = 1$ , the equation for  $\Upsilon$  would involve division by zero. Extra care is needed in those cases.

$$\mathbb{S}_{n+2}(\beta, \beta \pm i\pi, \theta_1, \dots, \theta_n; \gamma_{\mathcal{O}}) = \mathbb{S}_n(\theta_1, \dots, \theta_n; \gamma_{\mathcal{O}}), \quad (23)$$

which will be useful momentarily.

The recursion (19) for the constant is trivially solved

$$\mathcal{H}_n^\alpha = 1, \quad (24)$$

while the equation (20) for  $\Theta$  requires some more effort, especially for a general undeformed  $S$ -matrix. Let us give here the solution

$$\Theta_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \alpha) = \sqrt{\prod_{i=1}^n \mathbb{S}_n(\theta_i - \theta_1, \dots, \theta_i - \theta_n; \gamma_{\mathcal{O}}^{(-1)^i})}, \quad (25)$$

and check that it indeed solves equation (20). The left-hand-side is given by the product of the following three expressions

$$\sqrt{\mathbb{S}_{n+2}(0, i\pi, \theta + i\pi - \theta_1, \dots, \theta + i\pi - \theta_n; \gamma_{\mathcal{O}}^{-1})} = \sqrt{\mathbb{S}_n(\theta - \theta_1, \dots, \theta - \theta_n; \gamma_{\mathcal{O}})}, \quad (26)$$

$$\sqrt{\mathbb{S}_{n+2}(-i\pi, 0, \theta - \theta_1, \dots, \theta - \theta_n; \gamma_{\mathcal{O}})} = \sqrt{\mathbb{S}_n(\theta - \theta_1, \dots, \theta - \theta_n; \gamma_{\mathcal{O}})}, \quad (27)$$

and

$$\sqrt{\prod_{i=1}^n \mathbb{S}_{n+2}(\theta_i - \theta - i\pi, \theta_i - \theta, \theta_i - \theta_1, \dots, \theta_i - \theta_n; \gamma_{\mathcal{O}}^{(-1)^i})} = \Theta_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \alpha), \quad (28)$$

where we used the properties (22) and (23). Taking the product of these three terms indeed yields the right-hand-side of (20) as long as  $\gamma_{\mathcal{O}} = \gamma_{\mathcal{O}}^{-1}$  or  $\gamma_{\mathcal{O}} = \pm 1$ . More explicitly, we can write the final expression

$$\begin{aligned} \Theta_n^{\mathcal{O}}(\{\theta_i\}_n; \alpha) &= \prod_{i=1}^n \sqrt{\frac{\prod_{j=1}^n S_{\alpha}(\theta_{ij})^{1/2} - \gamma_{\mathcal{O}} \prod_{j=1}^n S_{\alpha}(\theta_{ij})^{-1/2}}{\prod_{j=1}^n S_0(\theta_{ij})^{1/2} - \gamma_{\mathcal{O}} \prod_{j=1}^n S_0(\theta_{ij})^{-1/2}}} \\ &= \prod_{i=1}^n \sqrt{\frac{\sin\left(\frac{1}{2} \sum_{j=1}^n [\delta(\theta_{ij}) - i \log \Phi_{\alpha}(\theta_{ij})] - \frac{\omega_{\mathcal{O}}}{2}\right)}{\sin\left(\frac{1}{2} \sum_{j=1}^n \delta(\theta_{ij}) - \frac{\omega_{\mathcal{O}}}{2}\right)}}, \end{aligned} \quad (29)$$

where  $S_0(\theta) = S_0(-\theta)^{-1} = e^{i\delta(\theta)}$  and  $\gamma_{\mathcal{O}} = e^{i\omega_{\mathcal{O}}}$ . We also assumed that the  $S$ -matrix is fermionic in the sense that  $S_0(0) = S_0(i\pi) = -1$ , so  $\delta(0) = \delta(i\pi) = \pi$ . As noted above, the solution (29) also requires  $\gamma_{\mathcal{O}} = \pm 1$ . Thus, strictly speaking, it is valid for local and semi-local fields <sup>2</sup>.

<sup>2</sup>Note that care must be taken for free theories to avoid a zero denominator. This can be done consistently. An example was given in [70].

### 3.1 Example

The factorised structure (12) emerges naturally for free theories. For instance, for the order and disorder fields  $(\mu, \sigma)$  in the Ising field theory with  $\gamma_\mu = -1$  and  $\gamma_\sigma = 1$  we have that

$$\Theta_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \boldsymbol{\alpha}) = \sqrt{\prod_{j=1}^n \cos\left(\frac{i}{2} \sum_{i=1}^n \log(\Phi_{\boldsymbol{\alpha}}(\theta_{ij}))\right)}, \quad (30)$$

with  $n$  even for  $\mu$  and odd for  $\sigma$ . Note that we have  $\Theta_0^{\mathcal{O}}(\boldsymbol{\alpha}) = \Theta_1^{\mathcal{O}}(\boldsymbol{\alpha}) = 1$ . The unperturbed form factors in this case are well-known to be proportional to the simple product  $\prod_{i < j} \tanh \frac{\theta_{ij}}{2}$  for both fields, with  $\mu$  admitting only even particle numbers and  $\sigma$  only odd ones. Note that for free theories,  $\mathbb{Z}_2$ -symmetry is preserved by the generalised T $\bar{\text{T}}$  perturbations. In particular, this means that for the Ising field theory, the trace of the stress-energy tensor, usually denoted by  $\Theta$  will have non-vanishing even particle form factors, even for particle numbers higher than two. Indeed, these form factors will be also proportional to a factor  $\Theta_{2n}^{\Theta}(\{\theta_i\}_{2n}; \boldsymbol{\alpha})$  which is identical to (30) with cosine replaced by sine.

## 4 Correlation Functions and Asymptotics

One of the most interesting applications of our solutions is to the computation of ground-state two-point functions in the deformed theory. These can be spanned in terms of form factors. For the purposes of this paper, we will just consider the leading contributions to the long and short-distance expansions of the logarithm of a typical (normalised) two-point function  $\langle \mathcal{O}(0) \mathcal{O}^\dagger(r) \rangle / \langle \mathcal{O} \rangle^2$  and will restrict ourselves to fields that only have even-particle form factors. Also for simplicity, we will consider only the case when  $\alpha_1 := \alpha \neq 0$  and  $\alpha_i = 0$  for  $i > 1$ . In this case, the two-particle contribution to the logarithm of the two-point function is

$$q_2^{\mathcal{O}}(r; \alpha) = \int_{-\infty}^{\infty} \frac{dx}{(2\pi)^2} |\Theta_2^{\mathcal{O}}(x; \alpha)|^2 |\hat{F}_2^{\mathcal{O}}(x; 0)|^2 e^{\frac{\alpha x}{\pi} \sinh x} K_0(2r \cosh \frac{x}{2}), \quad (31)$$

where<sup>3</sup>

$$\Theta_2^{\mathcal{O}}(\theta; \alpha) = \begin{cases} \left| \frac{\cos \frac{1}{2}(\delta(\theta) - \alpha \sinh \theta)}{\cos \frac{\delta(\theta)}{2}} \right| & \text{for } \gamma_{\mathcal{O}} = 1 \\ \left| \frac{\sin \frac{1}{2}(\delta(\theta) - \alpha \sinh \theta)}{\sin \frac{\delta(\theta)}{2}} \right| & \text{for } \gamma_{\mathcal{O}} = -1 \end{cases}, \quad (32)$$

and  $\hat{F}_2^{\mathcal{O}}(x; 0)$  is the two-particle form factor normalised by the vacuum expectation value  $\langle \mathcal{O} \rangle$ . Although this is only the two-particle contribution, it allows us to make some general observations that can then be extended to the full form factor expansion. This is discussed in more detail in [70].

First, for  $\alpha > 0$  we see that the exponential term is strongly divergent. This divergence can not be compensated for by the underformed form factor which will generally decay exponentially.

---

<sup>3</sup>Note that for the Ising model  $\cos \frac{\delta(\theta)}{2} = 0$  in (32) but this can be compensated for by appropriate normalisation of the form factors.

For short distances we can interpret this behaviour as indicative of an unusual UV theory. Indeed, it is known [52] that for  $\alpha > 0$  particles acquire a typical (positive) length which makes the UV theory ill-defined. For long distances however, there is an interplay between the length scale, the momentum and the parameter  $\alpha$ . The integral is divergent above a certain critical value of the rapidity  $x$ , a cut-off resulting from comparing the dominant exponential decay of the Bessel function and the exponential coming from the minimal form factor  $e^{-2r \cosh \frac{\Lambda}{2}} = e^{\frac{\alpha\Lambda}{\pi} \sinh \Lambda}$ . For large  $\Lambda$ , this equation is solved by the principal branch of Lambert's  $W$ -function

$$\Lambda = 2W_0\left(\frac{\pi r}{\alpha}\right). \quad (33)$$

Thus, for large distances and sufficiently small momentum, the series is still convergent, indicating that the finite length of particles is not visible when probed from a large distance at low momenta/energy. However, if the momentum is high, even with  $r$  large the size of particles is again probed and convergence is again spoiled.

For  $\alpha < 0$  on the other hand there is rapid convergence both for short and long distances. One possible interpretation of this behaviour is that for  $\alpha < 0$ , particles acquire a (negative) length. This may be understood as follows: the description in terms of positive/negative lengths comes from the relationship with the hard rod model [72]. The hard rod model is a classical model describing the interaction amongst hard rods of length  $\ell$ . The sign of  $\ell$  determines the scattering shift associated with the collision of two rods, as well as the thermodynamic properties of the system. In [72] it was shown that the non-relativistic limit of the TBA/GHD equations is described by the hard rod model. This viewpoint leads to another interpretation of the  $\text{T}\bar{\text{T}}$ -deformation, namely that its non-relativistic version corresponds to a change in particle (rod) length. Since particle (rod) length is only a model parameter related to the scattering phase, it can be chosen to be either positive or negative, hence our interpretation of  $\alpha > 0$  as positive length and  $\alpha < 0$  as negative length.

In the IR due to the presence of a mass scale, the theory flows to a trivial IR point, just as in the unperturbed case. In the UV, the form factor expansion needs to be resummed in order to obtain the exact leading behaviour. However, since  $K_0(2r \cosh \frac{\theta}{2}) \approx -\log r$  for  $r \ll 1$  and a similar behaviour is found for every contribution in the form factor series, the  $n$ -particle contribution will scale as  $-z_n^{\mathcal{O}}(\alpha) \log r$  with coefficient  $z_n^{\mathcal{O}}(\alpha)$  given in terms of a (multiple) integral in the rapidities. For instance, for two particles, the contribution scales as

$$q_2^{\mathcal{O}}(r \ll 1; \alpha < 0) = -z_2^{\mathcal{O}}(\alpha) \log r, \quad (34)$$

with

$$z_2^{\mathcal{O}}(\alpha) = \int_{-\infty}^{\infty} \frac{d\theta}{(2\pi)^2} |\Theta_2^{\mathcal{O}}(\theta; \alpha)|^2 |\hat{F}_2^{\mathcal{O}}(\theta; 0)|^2 e^{\frac{\alpha\theta}{\pi} \sinh \theta}. \quad (35)$$

The rapidly decaying  $\alpha$ -dependent exponential has the effect of producing coefficients  $z_n^{\mathcal{O}}(\alpha)$  such that  $z_n^{\mathcal{O}}(\alpha) \leq z_n^{\mathcal{O}}(0)$  for  $\alpha < 0$ , whenever  $z_n^{\mathcal{O}}(0)$  is well-defined<sup>4</sup>. In other words, the UV behaviour is characterised by a power law just as happens in CFT, but this power is  $\alpha$ -dependent and tends to zero as  $\alpha \rightarrow -\infty$ .

---

<sup>4</sup>It is worth noting that this formula only makes sense for  $\alpha = 0$  if the unperturbed form factor decays exponentially for large  $\theta$ . This is not the case for the fields  $\mu, \Theta$  in the Ising field theory, for instance.

## 5 $c$ -Theorem and Numerics

Another avenue that can be explored is Zamolodchikov's  $c$ -theorem, which generally relates the connected part of the two-point function of the trace of the stress-energy tensor<sup>5</sup> with the change in the central charge between the IR and UV fixed points

$$c(\alpha) := c^{UV} - c^{IR} = \frac{3}{2} \int_0^\infty dr r^3 \langle \Theta(0)\Theta(r) \rangle_c. \quad (36)$$

For  $\alpha = 0$ , this equation gives the value of  $c^{UV}$ , since  $c^{IR} = 0$ , in agreement with the fact that a massive theory flows in the IR to a trivial gapped fixed point. A  $\overline{\text{T}\overline{\text{T}}}$  perturbation alters the UV behaviour of the theory. For  $\alpha < 0$  we still expect to have  $c^{IR} = 0$  and so equation (36) will give some finite value which can be interpreted as the deformation of  $c^{UV}$  under the  $\overline{\text{T}\overline{\text{T}}}$  perturbation. This quantity will now depend on  $\alpha$  and behave as shown in Fig. 1. The




Figure 1: The functions  $c_2(\alpha)$  for the Ising model.

trace of the stress-energy tensor is a local field hence its two-particle form factor does not have a kinematic pole. As we have shown in [70], for the Ising model and a  $\overline{\text{T}\overline{\text{T}}}$  perturbation, it takes the form

$$F_2^\Theta(\theta; \alpha) := 2\pi \left| \frac{\sin\left(\frac{\alpha}{2} \sinh \theta\right)}{\frac{\alpha}{2} \sinh \theta} \right| \frac{F_{\mathbf{min}}(\theta; \alpha)}{F_{\mathbf{min}}(i\pi; \alpha)}. \quad (37)$$

where the oscillatory factor is of the same type found earlier. In the two-particle approximation, after integration in  $r$  and in one rapidity variable, the integral (36) gives

$$c_2(\alpha) = \frac{3}{8} \int_{-\infty}^{+\infty} dx \left[ \frac{\sin\left(\frac{\alpha}{2} \sinh x\right)}{\alpha \cosh^3 \frac{x}{2}} \right]^2 e^{\frac{\alpha}{\pi} x \sinh x}. \quad (38)$$

The resulting function is evaluated numerically in Fig. 1 where we see that it flows from the unperturbed value  $1/2$  to zero for  $-\alpha$  large. The oscillatory dependence is entirely washed out by the rapidly decaying exponential.

<sup>5</sup>Note that the field  $\Theta$  – the trace of the stress-energy tensor – should not be confused with the function  $\Theta_n^\Theta(\theta; \alpha)$  introduced earlier.

## 6 Conclusions, Discussion and Outlook

In this paper we have proposed a form factor program for IQFTs perturbed by  $T\bar{T}$  and descendants thereof. We have shown that the form factor equations can be solved exactly for a large class of theories and fields, and that these solutions span correlation functions in the usual way. We find that, compared to the unperturbed theory, all form factor solutions are “dressed” by oscillatory functions of the rapidities and perturbation parameters.

Our derivation is subject a number of constraints and choices that we would like to now discuss in some detail. As we have seen, we requires the factors  $\gamma_{\mathcal{O}} = \pm 1$  and have chosen  $\beta = 0$ . In addition, in order for (21) to hold, we have assumed all spins  $s$  to be odd. This assumption is rather natural since the local conserved charges of most IQFTs are indeed odd. In that case, the following condition holds,

$$\varphi(\theta; \alpha)\varphi(\theta + i\pi; \alpha) = \Phi_{\alpha}^{1/2}(\theta).$$

This constraint can however be easily lifted by modifying the definition of the function (21).

An additional implicit assumption, built into the ansatz (7) is that all form factors have kinematic poles. However, local fields, such as the trace of the stress-energy tensor, do not have a kinematic pole for their two-particle form factor but have kinematic poles for higher particle numbers. We may then ask what changes in this derivation for local fields. In fact, for interacting theories, the derivation presented here goes through with  $\gamma_{\mathcal{O}} = 1$  and for particle numbers  $n > 2$ , since the two-particle and higher particle solutions have slightly different structures. In fact, in order to make the solution  $\Theta_n^{\mathcal{O}}(\theta_1, \dots, \theta_n; \alpha)$  found here consistent with a two-particle form factor without kinematic pole and with the kinematic residue equations we will typically need to multiply our solution (29) by an additional function  $g_n(\theta_1, \dots, \theta_n; \alpha)$  with the property that it solves all form factor equations trivially, that is

$$g_{n+2}(\theta + i\pi, \theta, \theta_1, \dots, \theta_n; \alpha) = g_n(\theta_1, \dots, \theta_n; \alpha), \quad (39)$$

while being symmetric and  $2\pi i$ -periodic in all variables. This function will be specific to each theory. For the Ising field theory we have discussed the construction of this function in [70].

An unusual feature of our form factor solutions is that the function (29) which acts as a kind of “dressing” of the unperturbed form factors, involves a square root. We think that this may be related to the fact that a  $T\bar{T}$  perturbation can be interpreted in the TBA sense as a state-dependent change of the inverse temperature (or generalised temperatures for more general perturbations). It seems therefore natural that a relationship between finite temperature form factors and  $T\bar{T}$ -deformed form factors should exist. Interestingly, form factors at finite temperature also involve square roots, as can be seen for instance for the Ising model in [73, 74]. Closely related are finite volume form factors, which are also typically dressed by square root factors [75, 76]. Both for finite temperature and for finite volume form factors, the square root factors are functions of TBA quantities. It would be very interesting to investigate this connection further. This viewpoint also gives a new interpretation to the  $c$ -function presented in Fig.1, which has a very similar form as the TBA scaling function, if we replace inverse temperature by  $-\alpha$ .

An alternative setting in which the non-analyticity of form factors may be observed is in the choice of the minimal form factor, a subtlety that is not discussed in this paper, where we chose to set  $\beta = 0$  in (11). For now, it remains an open question whether there is a “natural” choice

for the parameters  $\beta$  in the context of  $T\bar{T}$ -perturbed theories. However, what we have learned from [71] is that a minimal form factor structure characterised by (10) and (11) is an integral part of the “standard” minimal form factor construction that has been employed for IQFTs such as the sinh-Gordon model, since the work of Karowski and Weiss [65]. The caveat however is that for a conventional IQFT the parameters  $\beta$  are entirely fixed, largely by the requirement of analyticity. In other words, most choices of (11) will also lead to non-analyticity of the minimal form factor. Thus, analyticity issues seem ubiquitous in the context of constructing form factors for  $T\bar{T}$ -perturbed theories and the question remains as to whether these issues admit a physical interpretation and/or should be fixed by a careful choice of the minimal form factor.

The asymptotics of correlation functions is consistent with the picture that point-like degrees of freedom (particles) in unperturbed theories become objects of finite (positive, for  $\alpha > 0$ , negative for  $\alpha < 0$ ) length after the perturbation. In this light, it is clear the UV theory is not a standard QFT whereas in the presence of a mass scale, the IR theory is as usual trivial. Correlation functions obtained from form factors reflect these features by diverging at short distances with  $\alpha > 0$  and converging very rapidly for  $\alpha < 0$ , while displaying power-law scaling with  $\alpha$ -dependent exponents.

Together with [70, 71, 77], this work provides a new viewpoint on generalised  $T\bar{T}$ -perturbed theories as well as new computational tools. Many different problems can now be addressed, from the study of correlation functions in interacting theories, to that of entanglement measures. There remain also many interesting open questions, e.g. what is the interpretation of the function  $c(\alpha)$  and of the powers  $z^{\mathcal{O}}(\alpha)$  characterising the short-distance scaling of correlators? We hope to return to these problems in the near future.

**Acknowledgments:** The authors thank John Donahue, Benjamin Doyon, Fedor Smirnov, Roberto Tateo and Alexander Zamolodchikov for useful discussions.. Olalla A. Castro-Alvaredo thanks EPSRC for financial support under Small Grant EP/W007045/1. The work of Stefano Negro is partially supported by the NSF grant PHY-2210349 and by the Simons Collaboration on Confinement and QCD Strings. Fabio Sailis is grateful for his PhD Studentship which is funded by City, University of London. This project was partly inspired by a meeting at the Kavli Institute for Theoretical Physics (Santa Barbara) in September 2022. Olalla A. Castro-Alvaredo and Stefano Negro thank the Institute for financial support from the National Science Foundation under Grant No. NSF PHY-1748958, and hospitality during the conference “Talking Integrability: Spins, Fields and Strings”, August 29-September 1 (2022) and the related extended program on “Integrability in String, Field and Condensed Matter Theory”, August 22-October 14 (2022).

## References

- [1] F. A. Smirnov and A. B. Zamolodchikov, *On space of integrable quantum field theories*, *Nucl. Phys. B* **915** (2017) 363–383 [[1608.05499](#)].
- [2] A. Cavaglià, S. Negro, I. M. Szécsényi and R. Tateo,  *$T\bar{T}$ -deformed 2D Quantum Field Theories*, *JHEP* **10** (2016) 112 [[1608.05534](#)].
- [3] S. Dubovsky, R. Flauger and V. Gorbenko, *Solving the Simplest Theory of Quantum Gravity*, *JHEP* **09** (2012) 133 [[1205.6805](#)].

- [4] M. Caselle, D. Fioravanti, F. Gliozzi and R. Tateo, *Quantisation of the effective string with TBA*, *JHEP* **07** (2013) 071 [[1305.1278](#)].
- [5] S. Dubovsky, V. Gorbenko and M. Mirbabayi, *Asymptotic fragility, near AdS<sub>2</sub> holography and  $T\bar{T}$* , *JHEP* **09** (2017) 136 [[1706.06604](#)].
- [6] J. Cardy, *The  $T\bar{T}$  deformation of quantum field theory as random geometry*, *JHEP* **10** (2018) 186 [[1801.06895](#)].
- [7] R. Conti, S. Negro and R. Tateo, *The  $T\bar{T}$  perturbation and its geometric interpretation*, *JHEP* **02** (2019) 085 [[1809.09593](#)].
- [8] R. Conti, L. Iannella, S. Negro and R. Tateo, *Generalised Born-Infeld models, Lax operators and the  $T\bar{T}$  perturbation*, *JHEP* **11** (2018) 007 [[1806.11515](#)].
- [9] R. Conti, S. Negro and R. Tateo, *Conserved currents and  $T\bar{T}_s$  irrelevant deformations of 2D integrable field theories*, *JHEP* **11** (2019) 120 [[1904.09141](#)].
- [10] S. Dubovsky, S. Negro and M. Porrati, *Topological Gauging and Double Current Deformations*, [2302.01654](#).
- [11] F. Aramini, N. Brizio, S. Negro and R. Tateo, *Deforming the ODE/IM correspondence with  $T\bar{T}$* , *JHEP* **03** (2023) 084 [[2212.13957](#)].
- [12] P. Dorey, C. Dunning, S. Negro and R. Tateo, *Geometric aspects of the ODE/IM correspondence*, *J. Phys. A* **53** (2020), no. 22 223001 [[1911.13290](#)].
- [13] G. Hernández-Chifflet, S. Negro and A. Sfondrini, *Flow Equations for Generalized  $T\bar{T}$  Deformations*, *Phys. Rev. Lett.* **124** (2020), no. 20 200601 [[1911.12233](#)].
- [14] G. Camilo, T. Fleury, M. Lencsés, S. Negro and A. Zamolodchikov, *On factorizable S-matrices, generalized  $T\bar{T}$ , and the Hagedorn transition*, *JHEP* **10** (2021) 062 [[2106.11999](#)].
- [15] L. Córdova, S. Negro and F. I. Schaposnik Massolo, *Thermodynamic Bethe Ansatz past turning points: the (elliptic) sinh-Gordon model*, *JHEP* **01** (2022) 035 [[2110.14666](#)].
- [16] A. LeClair, *Thermodynamics of perturbations of some single particle field theories*, *J. Phys. A* **55** (2022), no. 18 185401 [[2105.08184](#)].
- [17] A. LeClair, *deformation of the Ising model and its ultraviolet completion*, *J. Stat. Mech.* **2111** (2021) 113104 [[2107.02230](#)].
- [18] C. Ahn and A. LeClair, *On the classification of UV completions of integrable  $T\bar{T}$  deformations of CFT*, *JHEP* **08** (2022) 179 [[2205.10905](#)].
- [19] M. Guica, *An integrable Lorentz-breaking deformation of two-dimensional CFTs*, *SciPost Phys.* **5** (2018), no. 5 048 [[1710.08415](#)].
- [20] J. Cardy,  *$T\bar{T}$  deformation of correlation functions*, *JHEP* **12** (2019) 160 [[1907.03394](#)].

- [21] O. Aharony and T. Vaknin, *The  $TT^*$  deformation at large central charge*, *JHEP* **05** (2018) 166 [[1803.00100](#)].
- [22] O. Aharony, S. Datta, A. Giveon, Y. Jiang and D. Kutasov, *Modular invariance and uniqueness of  $T\bar{T}$  deformed CFT*, *JHEP* **01** (2019) 086 [[1808.02492](#)].
- [23] M. Guica and R. Monten, *Infinite pseudo-conformal symmetries of classical  $T\bar{T}$ ,  $J\bar{T}$  and  $JT_a$  - deformed CFTs*, *SciPost Phys.* **11** (2021) 078 [[2011.05445](#)].
- [24] M. Guica,  *$J\bar{T}$ -deformed CFTs as non-local CFTs*, [2110.07614](#).
- [25] M. Guica, R. Monten and I. Tsiaras, *Classical and quantum symmetries of  $T\bar{T}$ -deformed CFTs*, [2212.14014](#).
- [26] M. Baggio and A. Sfondrini, *Strings on NS-NS Backgrounds as Integrable Deformations*, *Phys. Rev. D* **98** (2018), no. 2 021902 [[1804.01998](#)].
- [27] A. Dei and A. Sfondrini, *Integrable  $S$  matrix, mirror TBA and spectrum for the stringy  $AdS_3 \times S^3 \times S^3 \times S^1$  WZW model*, *JHEP* **02** (2019) 072 [[1812.08195](#)].
- [28] S. Chakraborty, A. Giveon and D. Kutasov,  *$T\bar{T}$ ,  $J\bar{T}$ ,  $T\bar{J}$  and String Theory*, *J. Phys. A* **52** (2019), no. 38 384003 [[1905.00051](#)].
- [29] N. Callebaut, J. Kruthoff and H. Verlinde,  *$T\bar{T}$  deformed CFT as a non-critical string*, *JHEP* **04** (2020) 084 [[1910.13578](#)].
- [30] L. McGough, M. Mezei and H. Verlinde, *Moving the CFT into the bulk with  $T\bar{T}$* , *JHEP* **04** (2018) 010 [[1611.03470](#)].
- [31] A. Giveon, N. Itzhaki and D. Kutasov,  *$T\bar{T}$  and LST*, *JHEP* **07** (2017) 122 [[1701.05576](#)].
- [32] V. Gorbenko, E. Silverstein and G. Torroba,  *$dS/dS$  and  $T\bar{T}$* , *JHEP* **03** (2019) 085 [[1811.07965](#)].
- [33] P. Kraus, J. Liu and D. Marolf, *Cutoff  $AdS_3$  versus the  $T\bar{T}$  deformation*, *JHEP* **07** (2018) 027 [[1801.02714](#)].
- [34] T. Hartman, J. Kruthoff, E. Shaghoulian and A. Tajdini, *Holography at finite cutoff with a  $T^2$  deformation*, *JHEP* **03** (2019) 004 [[1807.11401](#)].
- [35] M. Guica and R. Monten,  *$T\bar{T}$  and the mirage of a bulk cutoff*, *SciPost Phys.* **10** (2021), no. 2 024 [[1906.11251](#)].
- [36] Y. Jiang, *Expectation value of  $T\bar{T}$  operator in curved spacetimes*, *JHEP* **02** (2020) 094 [[1903.07561](#)].
- [37] G. Jafari, A. Naseh and H. Zolfi, *Path Integral Optimization for  $T\bar{T}$  Deformation*, *Phys. Rev. D* **101** (2020), no. 2 026007 [[1909.02357](#)].
- [38] S. Dubovsky, V. Gorbenko and G. Hernández-Chifflet,  *$T\bar{T}$  partition function from topological gravity*, *JHEP* **09** (2018) 158 [[1805.07386](#)].

- [39] A. J. Tolley,  $T\bar{T}$  deformations, massive gravity and non-critical strings, *JHEP* **06** (2020) 050 [[1911.06142](#)].
- [40] L. V. Iliesiu, J. Kruthoff, G. J. Turiaci and H. Verlinde,  $JT$  gravity at finite cutoff, *SciPost Phys.* **9** (2020) 023 [[2004.07242](#)].
- [41] S. Okumura and K. Yoshida,  $T\bar{T}$ -deformation and Liouville gravity, *Nucl. Phys. B* **957** (2020) 115083 [[2003.14148](#)].
- [42] S. Ebert, C. Ferko, H.-Y. Sun and Z. Sun,  $T\bar{T}$  in  $JT$  Gravity and  $BF$  Gauge Theory, *SciPost Phys.* **13** (2022), no. 4 096 [[arXiv:2205.07817](#)].
- [43] D. X. Horváth, *Hydrodynamics of massless integrable RG flows and a non-equilibrium c-theorem*, *JHEP* **10** (2019) 020 [[1905.08590](#)].
- [44] M. Medenjak, G. Policastro and T. Yoshimura,  $T\bar{T}$ -Deformed Conformal Field Theories out of Equilibrium, *Phys. Rev. Lett.* **126** (2021), no. 12 121601 [[2011.05827](#)].
- [45] M. Medenjak, G. Policastro and T. Yoshimura, *Thermal transport in  $T\bar{T}$ -deformed conformal field theories: From integrability to holography*, *Phys. Rev. D* **103** (2021), no. 6 066012 [[2010.15813](#)].
- [46] R. Travaglini, M. Mazzoni and O. A. Castro-Alvaredo, *Generalised Hydrodynamics of  $T\bar{T}$ -Deformed Integrable Quantum Field Theories*, [2405.06564](#).
- [47] T. Bargheer, N. Beisert and F. Loebbert, *Boosting Nearest-Neighbour to Long-Range Integrable Spin Chains*, *J. Stat. Mech.* **0811** (2008) L11001 [[0807.5081](#)].
- [48] T. Bargheer, N. Beisert and F. Loebbert, *Long-Range Deformations for Integrable Spin Chains*, *J. Phys. A* **42** (2009) 285205 [[0902.0956](#)].
- [49] B. Pozsgay, Y. Jiang and G. Takács,  $T\bar{T}$ -deformation and long range spin chains, *JHEP* **03** (2020) 092 [[1911.11118](#)].
- [50] E. Marchetto, A. Sfondrini and Z. Yang,  $T\bar{T}$  Deformations and Integrable Spin Chains, *Phys. Rev. Lett.* **124** (2020), no. 10 100601 [[1911.12315](#)].
- [51] B. Doyon, J. Durnin and T. Yoshimura, *The Space of Integrable Systems from Generalised  $T\bar{T}$ -Deformations*, *SciPost Phys.* **13** (2022), no. 3 072 [[2105.03326](#)].
- [52] J. Cardy and B. Doyon,  $T\bar{T}$  deformations and the width of fundamental particles, *JHEP* **04** (2022) 136 [[2010.15733](#)].
- [53] B. Doyon, F. Hübner and T. Yoshimura, *Generalised  $T\bar{T}$ -deformations of classical free particles*, [2312.14855](#).
- [54] D. J. Gross, J. Kruthoff, A. Rolph and E. Shaghoulian,  $T\bar{T}$  in  $AdS_2$  and Quantum Mechanics, *Phys. Rev. D* **101** (2020), no. 2 026011 [[1907.04873](#)].
- [55] D. J. Gross, J. Kruthoff, A. Rolph and E. Shaghoulian, *Hamiltonian deformations in quantum mechanics,  $T\bar{T}$ , and the SYK model*, *Phys. Rev. D* **102** (2020), no. 4 046019 [[1912.06132](#)].

- [56] G. Coppa, F. Giordano, S. Negro and R. Tateo, “The Generalised Born Oscillator and the Berry-Keating Hamiltonian.” To appear.
- [57] G. Bonelli, N. Doroud and M. Zhu,  *$T\bar{T}$ -deformations in closed form*, *JHEP* **06** (2018) 149 [[1804.10967](#)].
- [58] M. Taylor,  *$TT$ - deformations in general dimensions*, [1805.10287](#).
- [59] R. Conti, J. Romano and R. Tateo, *Metric approach to a  $T\bar{T}$ -like deformation in arbitrary dimensions*, *JHEP* **09** (2022) 085 [[2206.03415](#)].
- [60] A. B. Zamolodchikov, *Integrable field theory from conformal field theory*, *Adv. Stud. Pure Math.* **19** (1989) 641–674.
- [61] S. Negro, *Integrable structures in quantum field theory*, *J. Phys. A* **49** (2016), no. 32 323006 [[1606.02952](#)].
- [62] A. B. Zamolodchikov, *Irreversibility of the flux of the renormalization group in a 2-d field theory*, *JETP Lett.* **43** (1986) 730–732.
- [63] L. Castillejo, R. H. Dalitz and F. J. Dyson, *Low’s scattering equation for the charged and neutral scalar theories*, *Phys. Rev.* **101** (1956) 453–458.
- [64] G. Mussardo and P. Simon, *Bosonic-type s-matrix, vacuum instability and CDD ambiguities*, *Nucl. Phys.* **B578** (2000) 527–551.
- [65] M. Karowski and P. Weisz, *Exact s matrices and form-factors in (1+1)-dimensional field theoretic models with soliton behavior*, *Nucl. Phys.* **B139** (1978) 455–476.
- [66] F. Smirnov, *Form factors in completely integrable models of quantum field theory*, *Adv. Series in Math. Phys.* **14** (1992) World Scientific, Singapore.
- [67] G. Mussardo, *Statistical Field Theory: An Introduction to Exactly Solved Model in Statistical Physics*, *Oxford University Press* (2010).
- [68] V. P. Yurov and A. B. Zamolodchikov, *Correlation functions of integrable 2-d models of relativistic field theory. ising model*, *Int. J. Mod. Phys.* **A6** (1991) 3419–3440.
- [69] O. Babelon and D. Bernard, *From form-factors to correlation functions: The Ising model*, *Phys. Lett. B* **288** (1992) 113–120 [[hep-th/9206003](#)].
- [70] O. A. Castro-Alvaredo, S. Negro and F. Sailis, *Form factors and correlation functions of  $T\bar{T}$ -deformed integrable quantum field theories*, *JHEP* **09** (2023) 048 [[2306.01640](#)].
- [71] O. A. Castro-Alvaredo, S. Negro and I. M. Szécsényi, *On the representation of minimal form factors in integrable quantum field theory*, *Nucl. Phys. B* **1000** (2024) 116459 [[2311.16955](#)].
- [72] B. Doyon, H. Spohn and T. Yoshimura, *A geometric viewpoint on generalized hydrodynamics*, *Nucl. Phys. B* **926** (2018) 570–583 [[1704.04409](#)].

- [73] B. Doyon, *Finite-temperature form-factors in the free Majorana theory*, *J. Stat. Mech.* **0511** (2005) P11006 [[hep-th/0506105](#)].
- [74] B. Doyon, *Finite-temperature form-factors: A Review*, *SIGMA* **3** (2007) 011 [[hep-th/0611066](#)].
- [75] B. Pozsgay and G. Takacs, *Form factors in finite volume. II. Disconnected terms and finite temperature correlators*, *Nucl. Phys. B* **788** (2008) 209–251 [[0706.3605](#)].
- [76] B. Pozsgay and G. Takacs, *Form-factors in finite volume I: Form-factor bootstrap and truncated conformal space*, *Nucl. Phys. B* **788** (2008) 167–208 [[0706.1445](#)].
- [77] O. A. Castro-Alvaredo, S. Negro and F. Sails, *Entanglement entropy from form factors in  $\overline{T\bar{T}}$ -deformed integrable quantum field theories*, *JHEP* **11** (2023) 129 [[2306.11064](#)].