



City Research Online

City St George's, University of London

Citation: Chavez-Ugalde, I. Y., de Vocht, F., Jago, R., Adams, J., Ong, K. K., Forouhi, N. G., Colombet, Z., Ricardo, L. I. C., van Sluijs, E. & Toumpakari, Z. (2024). Ultra-processed food consumption in UK adolescents: distribution, trends, and sociodemographic correlates using the National Diet and Nutrition Survey 2008/09 to 2018/19. *European Journal of Nutrition*, 63(7), pp. 2709-2723. doi: 10.1007/s00394-024-03458-z

This is the published version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/33562/>

Link to published version: <https://doi.org/10.1007/s00394-024-03458-z>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).



Ultra-processed food consumption in UK adolescents: distribution, trends, and sociodemographic correlates using the National Diet and Nutrition Survey 2008/09 to 2018/19

Irazu Yanaina Chavez-Ugalde^{1,2,3} · Frank de Vocht^{2,3,4} · Russell Jago^{2,3,4,5} · Jean Adams^{1,3} · Ken K. Ong¹ · Nita G. Forouhi¹ · Zoé Colombet⁶ · Luiza I. C. Ricardo¹ · Esther van Sluijs^{1,3} · Zoi Toumpakari⁵

Received: 23 November 2023 / Accepted: 17 June 2024
© The Author(s) 2024

Abstract

Purpose We quantified levels of ultra-processed food (UPF) consumption and investigated consumption patterns in a representative sample of UK adolescents.

Methods We used data from 4-day food diaries from adolescents in the UK National Diet and Nutrition Survey (NDNS) (2008/09–2018/19). UPF were identified using the NOVA classification. We estimated the percentage of Total Energy Intake (%TEI) and the absolute weight (grams). Linear regression models quantified differences in UPF consumption across survey years and its association with participant's individual characteristics. This was an analysis of the repeated cross-sectional data from the UK NDNS Rolling Programme waves 1–11 (2008/09–2018/19). A total of 2991 adolescents (11–18y) with complete information on dietary intake were included.

Results Mean UPF consumption was 861 (SD 442) g/d and this accounted for 65.9% (SD 13.4%) of TEI. Between 2008 and 2019, mean UPF consumption decreased from 996 to 776 g/d [−211 (95%CI −302; −120)] and from 67.7% to 62.8% of TEI [−4.8% (95%CI −8.1; −1.5)]. Higher %TEI was consumed by adolescents with lower socioeconomic status; white ethnicity and living in England North. A higher weight of UPF consumption (g/d) was associated with being male, white, age 18y, having parents with routine or manual occupation, living in England North, and living with obesity.

Conclusion Average energy intake from UPF has decreased over a decade in UK adolescents. We observed a social and regional patterning of UPF consumption, with higher consumption among adolescents from lower socioeconomic backgrounds, from a white ethnicity and living in England North. Our findings suggest inequalities associated with UPF intake and factors that might lie beyond individual choice.

Keywords Ultra-processed food · Adolescents · National diet and nutrition survey · Food consumption

Abbreviations

CI	Confidence interval	MVPA	Moderate-to-vigorous physical activity
HIC	High income country	NDNS	National diet and nutrition survey
LMIC	Low-middle-income country	NS-SEC	National statistics socioeconomic class
		RPAQ	Recent physical activity questionnaire
		TEI	Total energy intake

Esther van Sluijs and Zoi Toumpakari joint last authors.

✉ Irazu Yanaina Chavez-Ugalde
yanaina.chavez-ugalde@mrc-epid.cam.ac.uk

¹ MRC Epidemiology Unit, Institute of Metabolic Science, University of Cambridge School of Clinical Medicine, Cambridge Biomedical Campus, Cambridge CB2 0QQ, UK

² Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

³ National Institute for Health and Care Research (NIHR) School for Public Health Research (SPHR), Newcastle, UK

⁴ NIHR Applied Research Collaboration West (NIHR ARC West), University Hospitals Bristol and Weston NHS Foundation Trust, Bristol, UK

⁵ Centre for Exercise, Nutrition and Health Sciences, School for Policy Studies, University of Bristol, Bristol, UK

⁶ Department of Public Health and Policy, University of Liverpool, Liverpool, UK

UK	United Kingdom
UPF	Ultra-processed food(s)
USA	United States of America
zBMI	Standardised body mass index
ZC	Zoé Colombet

Background

The consumption of ultra-processed foods (UPF) has been proposed as one of the key drivers of the global rise in chronic diet-related diseases [1]. UPF are often manufactured from cheap industrial substances extracted or derived from foods, such as fats and oils, free sugars, and amino acids, which are then mixed with cosmetic additives, like colours, stabilisers, humectants, emulsifiers which are not used in domestic kitchens. Diets high in UPFs are linked to indicators of poor dietary quality, including elevated levels of added sugars, saturated fat, and sodium, as well as increased energy density and decreased fibre, protein, and micronutrient content [2, 3]. Such unhealthy dietary patterns have been implicated in the rise of chronic diseases through inflammatory pathways [4]. There is a growing body of evidence linking the consumption of UPF with poor dietary quality, and diet related diseases such as obesity in children, adolescents and adults [7], and chronic non-communicable diseases, such as cardiovascular disease and cancer [8] and all-cause mortality in adults [9]. A recent systematic umbrella review of published meta-analyses found that increased consumption of UPFs is correlated with increased risks of negative health outcomes, particularly relating to cardiometabolic, common mental disorders, and mortality [16]. Moreover, the detrimental health effects associated with UPFs may not be attributed to their nutrient profile and calorie density but also to physical alterations in the food matrix and chemical properties resulting from industrial processing methods, ingredients (such as additives and non-sugar sweeteners), and by-products [5]. Some examples of UPF are soft drinks, breakfast cereals, reconstituted meat products, packaged breads, ready-to-eat foods. UPF are durable, convenient, ready-to-eat, hyper-palatable, have attractive packaging and are strongly marketed to children and adolescents [6]. These results underscore the need to create and assess the efficacy of population-wide and public health interventions aimed at minimizing the intake of UPFs to enhance human health.

Adolescence is a key transitional stage when major changes in practices that influence health occur [10]. Adolescents' search for novelty and openness to change also makes them a vulnerable group for commercial marketing [6]. Adolescents' food patterns and practices are strongly driven by their food environments and eating contexts, their autonomy, peer influence and social norms [11]. These food

patterns can additionally be influenced by commercially targeted activities, such as marketing and advertising, product placement, and pricing strategies such as food promotions and discounts [12].

Evidence across different countries suggests that adolescents are the highest consumers on average of UPF compared to other age groups [13]. In Canada it is estimated that UPF contribute approximately 55% of the total caloric intake in children and adolescents (2–18y) versus 45% in adults [14]. In the USA, children (6–11y) (69.0%) and adolescents (12–19y) (67.7%) consumed a significantly higher percentage of energy from UPF than those aged 2–5 years (61.1%) [15].

Although consumption of UPF is becoming more prevalent worldwide [1, 17], there are notable cross-country and socioeconomic status (SES) differences [18]. In nationally representative samples from high-income countries UPF contribute more than 50% of energy intake [19], and up to 30% in middle-income countries [20]. Even though consumption of UPF is much higher in high-income compared to upper-middle income countries, among these latter countries adolescents are still the highest UPF consumption age group. Evidence from high- and upper-middle income countries signals a distinction in social patterning of UPF consumption according to the nutritional transition stage of each country [1].

Globally, the availability and sales of UPF have increased over time (2006–2019) [1], and evidence from the USA (2010–2018) and Korea (1999–2018) suggests that consumption of UPF among adolescents has also increased over time [15, 21]. However, evidence from a representative sample of UK population (> 1.5 years) did not find evidence for an increase in the energy share of UPF between 2008 and 2019 [22], yet changes in weight consumption (g/d) from UPF were not explored. Examining the changes in consumption of UPF over time can aid us in understanding the behavioural trends within a shifting food landscape. Considering that adolescents are the highest consumers of UPF, and considering evidence from other countries indicating an increase in consumption in this age group over time, we explored the 11 data waves in NDNS to determine if consumption patterns of energy (TEI%) and weight (g/d) followed similar trends.

Previous studies on UPF consumption have mostly expressed UPF as percentage of energy intake. However, the French NutriNet-Santé study [23] proposed expressing UPF by weight (grams) given that this can capture non-nutritional factors relating to processing of food and foods and drinks that contribute little to energy intake (e.g., additives, non-sugar sweeteners, neo-formed contaminants and endocrine-disrupting chemicals from packaging materials [24]). The associations observed between UPF and health might be influenced by various mechanisms. For example,

research shows that weight measures (g/d) of UPF consumption have a stronger association with type 2 diabetes and cardiovascular disease. Conversely, energy intake from UPF was found to be more strongly associated with body fat accumulation [25]. Furthermore, research shows that UPF remains strongly associated with obesity and health-related outcomes after adjusting for quality and dietary pattern [23].

Currently there is limited evidence on UPF consumption among adolescents. Quantifying levels of UPF consumption and exploring consumption patterns is essential to enable policies that prevent unhealthy dietary patterns in adolescent years and prevent diet-related diseases later in life. To address this research gap, this study aimed to quantify the levels UPF consumption and investigate consumption patterns in a representative sample of UK adolescents. To achieve this, we carried out the following objectives: we calculated UPF consumption and its contribution to relative energy intake (% kcal/day) and absolute food weight intake (g/day); we described UPF consumption across the 11 NDNS survey waves (2008–2019); and investigated the association between sociodemographic characteristics and UPF consumption.

Methods

We conducted an analysis using repeated cross-sectional data from 11- to 18-year-old participants (adolescents) in the UK National Diet and Nutrition Survey Rolling Programme (NDNS) waves 1–11 (2008/09–2018/19). Data were downloaded from the UK Data Service [26]. This study is reported according to the Strengthening the Reporting of Observational studies in Epidemiology – Nutritional Epidemiology (STROBE-nut). Parental consent was obtained for participants aged 11 to 15 and direct consent was obtained for participants aged 16 to 18 years. Additional ethical approval for this secondary analysis of anonymised data was not required.

Study design and population

The NDNS is an annual rolling programme (RP) of cross-sectional surveys conducted in the UK (England, Wales, Scotland, and Northern Ireland) assessing the food consumption, nutrient intake and nutritional status of the general UK population aged 1.5 years and above living in private households. Additionally, data collected in England is presented by regions: England North, Central/Midlands, England South (including London). NDNS survey year starts in April and data collection runs from April until March the following year. Population, sampling and recruitment are described in detail elsewhere [27]. Briefly, NDNS's continuous rolling programme seeks to collect data from

a representative sample of the UK population with 1000 participants every year (500 adults, 500 children and adolescents) with provision of an additional sample to achieve country-level representativeness.

Sampling follows a multistage probability design to generate a new random sample of private households in the UK every year. Each year a random sample from small geographical areas [(i.e., primary sampling units (PSUs))] is selected. Within these PSUs, private addresses are randomly selected from the Postcode Address File and the selected households receive a visit by the study team. Among the participating household one child and one adult are randomly selected to take part. Participants within households are then asked to complete a food diary across 4 days to record all foods and beverages consumed inside and outside of the house. Those who record at least 3 days are then invited for further physical measurements.

Inclusion criteria

Individuals were included in the analysis if they took part in NDNS waves 1 to 11, were aged between ≥ 11 and ≤ 18 years at data collection and completed at least three out of four food diary days.

The current study combines NDNS data from waves 1 to 11 (2008/09 to 2018/19). Overall, at an individual level, 53% of those selected to take part, including adults and children, completed at least three food diary days. From the sample of included individuals, 3,270 were adolescents (11- to 18-year-olds).

Dietary data collection & processing

Dietary assessment in NDNS waves 1–11 was designed to provide full description, detail and quantification of all food and drink consumed during the dietary assessment period, with the ability to capture habitual consumption when conducted over a number of days. Seasonal variations in food and drink consumption are addressed by the continuous fieldwork design.

Trained interviewers collected sociodemographic information through interviews and administered the food diaries to adolescent participants. Adolescents were instructed (in the case of 11- and 12-year-olds, their parent or carer) by the trained interviewer to record location, time and quantity of all the food and drink they consumed inside and outside of the home over four consecutive days. Recording of the four-day food diary would start on selected days to ensure even representation of all days of the week across the whole sample. After the food diaries were completed, the interviewer returned to collect the diary, reviewed the data with the participant and added missing details to improve completeness. Participants received monetary gift vouchers following

completion of at least three of the four dietary recordings. Household measures and nutritional information from labels were used to estimate portion sizes consumed. Food diaries were entered in the Dietary assessment system DINO (Diet In Nutrients Out) that uses food composition data from the NDNS Nutrient Databank [28, 29]. Additional details of the coding of food intake data and calculation of energy intakes and food composition in NDNS data are described in detail elsewhere [28–31].

Across the 11 years of data collection, NDNS participants reported on 60 main food groups, 154 subsidiary food groups and 4,944 food items. Classification of foods and beverages according to their level of processing was conducted using the NOVA food classification system which considers the nature, extent and purpose of industrial food manufacturing processing [32]. The NOVA classification includes four categories: (1) unprocessed or minimally processed foods, (2) processed culinary ingredients, (3) processed foods and (4) UPF. Figure 1 shows these four classifications and examples of each food group. More details on the NOVA classification can be found elsewhere [4, 32].

All foods and drinks in the NDNS Nutrient Databank were coded and grouped independently by two researchers (YCU and ZC) based on main food group first (e.g., ‘Pasta rice and other cereals’), then coded according to subsidiary food groups (e.g., pasta, manufactured products and ready meals) and then by individual food item (e.g., pasta, spaghetti, canned in tomato sauce) based on their recipe information and the ingredient list if the classification of a whole main or subgroup was not possible (e.g., composite dishes). This allowed us to classify each underlying or main constituent ingredient into the corresponding NOVA group as previously suggested on how to disaggregate composite food codes in NDNS data [30, 31]. Food supplements (i.e., vitamins and minerals) were not coded. Previous studies that have used either the NOVA classification system and/or NDNS data served as a useful coding framework to classify dietary data in this study [7, 34, 35].

This study was specifically interested in the consumption of UPF. Thus, we classified group 1, 2 and 3 as non-UPF, and group 4 as UPF and focused on the UPF

Fig. 1 The 4 groups in the NOVA food classification system [based on information from [33]]

1. Unprocessed/ minimally processed foods

Foods which are fresh or processed without adding any substances to the original food.

E.g., Beans, rice, frozen meat, pasteurized or power milk, plain yogurt, fresh, dried or frozen vegetables, grains, flour or pasta.

2. Processed culinary ingredients

Substances obtained directly from group 1 or products found in nature and that are used to prepare or cook group 1 foods.

E.g., Sugar, salt, vegetable oils, butter and vinegar.

3. Processed foods

Foods that have added substances from group 2 that have been added to group 1.

E.g., Vegetables in brine, cheese, breads made only with flour, water and salt, fruits in syrup, canned fish.

4. Ultra-processed foods

Industrial formulations that result from a sequence of physical and chemical processes applied to foods and their constituents or products synthesised in a laboratory based on organic materials.

E.g., Soft drinks, sweetened fruit drinks, sweet or savoury packaged snacks, packaged breads and buns, cookies, confectionery, ice cream, sweetened yogurts, dairy beverages, breakfast cereals, ready-to-eat / - heat meals, sausages or other reconstituted meats, and prepared frozen or shelf-stable dishes.

category (NOVA 4) to describe UPF consumption among UK adolescents.

NOVA classification level of agreement

The level of agreement between both coders was 97% after independently classifying all food and drink items using the full NOVA classification system (NOVA 1, 2, 3 and 4). Most of this disagreement (152 of 4,784 food and drink items, 3.1%) was between NOVA categories 1, 2 and 3. Only 10 of 4,784 (0.2%) food items caused a disagreement in UPF classification (NOVA 4). These 10 UPF were mainly composite dishes (i.e., beef curry takeaway, cottage pie with instant mash potato, black pudding in batter takeaway, fish pie with crust flaky pastry, vegetable pie with crust, *vol au vents* made with mushroom sauce and pastry), sauces (i.e., chilli pickle sweet, golden syrup), and ice-cream (i.e., ice-cream with double cream and purchased sorbet). These 10 food items categorised as NOVA 4 (UPF) by the first researcher (YCU) based on the individual food name and searches for lists of ingredients in branded products, whilst the second researcher (ZC) classified them based on the food name without searching for list of ingredients in branded items. Once the lists of ingredients were reviewed and discussed both researchers reached a 100% agreement.

Variables of interest

The outcomes of interest were the relative energy intake from UPF (NOVA 4) (% kcal/d) and the absolute weight of UPF consumed (g/d). These were calculated based on the relative energy and total absolute weight of foods and drinks classified as UPF in the NDNS food individual level dietary dataset reported by each participant. Individual-level sociodemographic characteristics included sex at birth, age, parent's occupation social class, ethnic group and UK region. Other individual characteristics included weight categories derived from standardised body mass index (zBMI), survey year, and moderate-to-vigorous-physical activity (MVPA).

BMI was calculated from nurse measured height and weight. BMI values were standardised for sex and age and categorised [normal weight (≤ 1 standard deviation (sd)), overweight (> 1 sd) and obese (> 2 sd)] based on the 1990 British Growth Reference (UK90).

MVPA data was self-reported using the Recent Physical Activity Questionnaire (RPAQ) [36], which assessed type, amount of physical activity and screen time, and is validated for use in older adolescents and adults. Data was therefore collected only in 16–18-year-olds, and among these, only 56% provided MVPA data (18% of the entire sample, $n = 575$ out of $n = 3270$). Daily time spent in MVPA was summed in each individual (min/day). As MVPA was not normally

distributed we created quartiles (< 21 min/day; 21–52 min/day; 52–124 min/day; > 124 min/day).

Other sociodemographic variables were categorised as follows: sex (male and female), age in completed whole years (8 categories from 11 to 18 years), parent's occupation social class (higher managerial, administrative, and professional occupations; intermediate occupations; routine and manual occupations; we used the three level National Statistics Socioeconomic classification (NS-SEC) [37]), ethnic group (white, non-white), and region [England North, England Central/Midlands, England South (including London), Scotland, Wales, and Northern Ireland].

Statistical analysis

Study weights were provided by NDNS, and datasets were re-scaled and adjusted for the adolescent subsample based on NDNS study weights guide documentation [38]. These weights were applied using the *svy* prefix in Stata for all analysis to account for non-response and sampling error and to provide estimates representative of the UK adolescent population.

For the description of the sample, we reported weighted percentages (with 95% CIs) of the distribution of individual sociodemographic characteristics alongside the distribution and means of relative UPF energy consumption (%kcal/day) and absolute weight of UPF consumption (g/d) in the overall sample and for each sociodemographic characteristic. We displayed the mean of all available days of food diary for each individual.

We pooled all survey years and used multivariable linear regression models to test if each of our two outcome variables – dietary contribution of UPF (%kcal/d and g/d) – differed across each of the variables of interest by doing individual stepwise models (i.e., sociodemographic categories and individual characteristics separately) adjusting for age, sex and survey year as covariates. Each category within a variable was compared to the reference group (i.e., sex (male), age (11 years) parent's occupation social class (higher managerial, administrative, and professional occupations), ethnic group (white), and region (England North)). We also used linear regression analysis to evaluate if mean UPF consumption (%kcal/d and g/d) changed across NDNS survey years by comparing each survey year (2 to 11) vs year 1 (2008/09) and adjusting for age and sex. Given the clustered regional sampling design of NDNS, we used survey analysis procedures to incorporate sample weights in these models.

Missing data

Variables with complete data ($n = 3,270$) were sex, age, survey year, and region. Variables with missing data were

ethnicity (0.1%), parents' occupation (4.5%), BMI z-score (4.3%), and MVPA (82.4%). We used a complete case analysis for all analyses ($n=2,991$ for all variables, except for MVPA $n=534$, for which a separate analysis was done using this subsample).

Additional analysis

We ran an analysis adjusting for total energy intake in addition to the primary analysis to explore whether individual characteristics were associated with relative energy (%kcal/d) and absolute weight (g/d) independently from total energy intake.

All data analysis was conducted in Stata Statistical Software: Release 17.0 (StataCorp LP., College Station, TX, USA).

Results

Table 1 presents descriptive data of the 2991 participants with complete data for the variables of interest except MVPA ($n=534$). Of all adolescents, 51.3% were females, 42.9% had parents with a higher managerial, administrative and professional occupation, 66% had normal weight, 83.3% were from a white ethnic group, 43.7% lived in England South (including London), and of participants with physical activity data (16 to 18 year olds) 26.7% reported more than 124 min/day of MVPA.

Overall, adolescents reported a mean total energy intake per day of 1741 kcal/day (95%CI 1717.6; 1764.4) and 65.9% (95%CI 65.2; 66.5) of these calories come from UPF. In terms of food weight, adolescents consumed a mean of 2004 (95%CI 1967; 2041) g/day, of which 861 g/day (95%CI 840; 882) was UPF (43% of total food weight consumed). Table 2 shows the percentage of energy from UPF and grams per day of UPF consumed by adolescents by individual characteristics.

Associations of UPF consumption with time and sociodemographic variables

UPF consumption across NDNS survey years

Figures 2 and 3 (and Supplementary Table S1) show mean UPF consumption and CI's across NDNS survey years (2009/09 – 2018/19) (adjusted for age and sex). There was evidence for a difference in relative energy (%kcal/day) and absolute weight (g/day) consumption from UPF across survey years ($p < 0.001$).

In Fig. 2 we observed that comparing survey year 2 (2009/10) through 7 (2014/15) versus survey year 1 (i.e., reference year) there was no evidence of a difference in the

absolute UPF as percentage of total energy intake (%TEI). However, %TEI from UPF was significantly lower from survey year 8 (2015/16) through 10 (2017/19) (vs year 1 – 2008/09) with the largest reduction in survey year 8 (-5.8%kcal/day).

In Fig. 3 we observed that the highest absolute weight from UPF consumed (g/d) was seen in year 1 (2008/09) (993.7 g/d). Similar to relative energy intake (%kcal/d), we observed a steady decline with some random variation in UPF weight across survey years.

Individual characteristics and %TEI from UPF

After adjusting for age, sex and survey year, results show (Fig. 4 and Supplementary Table S1) that parents' occupation, ethnic group and UK region were associated with percentage of energy consumption from UPF. A higher %TEI from UPF was consumed by adolescents whose parents had routine and manual occupations or intermediate occupations compared to adolescents with parents who had higher managerial occupations [intermediate: 2.0% (95%CI 0.4; 3.5); routine and manual parental occupation: 4.6% (95%CI 3.2; 6.1)]. Adolescents from a non-white ethnicity (vs white) reported lower %TEI from UPF [-8.0% (95%CI -9.8; -6.1)] as well as those living in England South (including London) (vs England North) [-3.2% (95%CI -4.9; -1.5)].

Individual characteristics and absolute weight from UPF

Figure 5 (and Supplementary Table S1) shows that higher weight of UPF consumption (g/d) was associated with age between 17 and 18 years (vs 11-year-olds), with the highest consumption observed in 18-year-olds [115.1 g/day, (95%CI 26.6; 203.5)]. Adolescents whose parents had a routine and manual occupation (vs higher managerial) [78.9 g/day, (95%CI 34.6; 123.3)], and adolescents living with obesity (vs normal weight) [90.3 g/day (95%CI 39.0; 141.5)] reported a higher UPF weight consumption. In contrast, lower weight of UPF consumption (g/d) was associated with female sex (vs male) [-169.2 g/day, (95%CI -206.8; -131.7)], non-white ethnicity (vs white) [-247.2, (95%CI -292.8; -201.6)], and living in South England and Northern Ireland (vs. North England) [-99.2, (95%CI -150.6; -47.8); -76.8, (95%CI -135.0; -18.6)].

Additional analysis

After additionally adjusting for total energy intake, all associations with %TEI from UPF persisted and in addition 18-year-olds (vs. 11-year-olds) had a lower %TEI (-3.0%TEI, 95%CI -5.7, -0.3). For UPF weight

Table 1 Descriptive characteristics of a weighted sample in the NDNS adolescents' sample waves 1–11 (2008/09–2018/19) (n = 2991)

Individual characteristics	%N (weighted) ^a	95%CI
<i>Sex</i>		
Male	51.3	(49.0, 53.6)
Female	48.7	(46.4, 51.0)
<i>Age</i>		
11	12.3	(10.9, 13.9)
12	12.8	(11.3, 14.5)
13	12.8	(11.3, 14.5)
14	12.6	(11.1, 14.2)
15	11.4	(10.1, 12.8)
16	14.6	(13.0, 16.4)
17	13.7	(12.0, 15.5)
18	9.9	(8.6, 11.3)
<i>Parent's occupation social class^b</i>		
Higher managerial, administrative, and professional	42.9	(40.6, 45.3)
Intermediate	22.8	(20.9, 24.8)
Routine and manual	34.3	(32.1, 36.6)
<i>Survey year (survey wave for data collection)</i>		
(1) 2008/09	8.0	(6.5, 9.9)
(2) 2009/10	10.1	(8.2, 12.4)
(3) 2010/11	9.7	(7.7, 12.1)
(4) 2011/12	8.4	(6.8, 10.4)
(5) 2012/13	8.6	(6.8, 10.9)
(6) 2013/14	9.2	(7.4, 11.4)
(7) 2014/15	9.4	(7.5, 11.7)
(8) 2015/16	8.6	(6.8, 10.9)
(9) 2016/17	9.7	(7.7, 12.2)
(10) 2017/18	9.0	(7.1, 11.3)
(11) 2018/19	9.2	(7.3, 11.5)
<i>Weight category^c</i>		
Normal weight	66.0	(63.6, 68.3)
Overweight	13.7	(12.1, 15.4)
Obese	20.3	(18.4, 22.4)
<i>Ethnic group</i>		
White	83.3	(81.2, 85.1)
Non-white	16.8	(14.9, 18.8)
<i>Region</i>		
England North	22.9	(21.2, 24.7)
England Central/Midlands	17.7	(16.1, 19.3)
England South (including London)	43.7	(41.8, 45.7)
Scotland	7.7	(6.4, 9.3)
Wales	4.7	(4.2, 5.3)
Northern Ireland	3.2	(2.9, 3.5)
<i>MVPA^d</i>		
< 21 min/day	26.9	(21.9, 32.6)
21–52 min/day	22.4	(18.0, 27.5)
52–124 min/day	24.0	(19.3, 29.4)
> 124 min/day	26.7	(21.1, 33.2)

BMI z-score standardised body mass index, *MVPA* Moderate-to-vigorous physical activity, *NDNS* National Diet and Nutrition Survey, *NS-SEC* National Statistics Socioeconomic Class, *UPF* ultra-processed food, *NOVA4* classification group

^aPercentages and means are weighed based on non-selection and non-response survey weights provided by NDNS year 2008–2019

^bParents occupation is based on the three-class NS-SEC. A small number of households were excluded from this classification where the household has never worked or was classified under other

Table 1 (continued)^cBMI z-score was created by standardising BMI for sex and age based on the 1990 British Growth Reference (UK90)^dMVPA: There was self-reported data available only for 16–18-year-olds, n=534

consumption, after adjusting for total energy intake associations were attenuated for age, survey years 4, 5, 6 and 10, and for MVPA > 124 min/day (Supplementary Table S2).

Discussion

In this repeated cross-sectional study of a nationally representative sample of UK adolescents, we found that mean UPF among 11- to 18-year-olds was 861 g/day and accounted for 65.9% of their TEI. After adjusting for age and sex, mean consumption of UPF (both %TEI and weight) declined between 2008/09/ and 2018/19. Percentage of energy from UPF was lower by 5 percentage points and weight consumption from UPF was lower by 211.2 g per day when comparing the first vs last NDNS survey waves (wave 1: 2008/09 vs wave 11:2018/19). After adjusting for age, sex and survey year, adolescents with lower socioeconomic status (having parents in an intermediate or routine and manual occupations), from a white ethnicity, and living in England North had higher %TEI from UPF. Additionally, being male, being between 17 and 18 years, having a lower socioeconomic status (having parents in a routine and manual occupation), living with obesity, from a white ethnicity, and living in England North were associated with a higher weight of UPF consumption (g/d). This increased consumption in weight of UPFs but not calories from UPFs could be explained by health behaviour changes and increased health awareness as adolescents age, and by adolescents living with obesity consuming a larger weight of UPFs that are low in fat, sugar and/or calories (e.g., zero calorie soft drinks) (22). In additional analysis, adjusting for total energy intake, patterns of association with UPF weight became more consistent to those with %TEI from UPF.

These findings are consistent with previous analyses of the same NDNS data in 11 to 18-year-olds using survey years 2008/16 and 2008/17 (68% contribution of UPF to total energy intake [35, 39] and with analyses of data from adolescents living in other high-income countries (55% in Canada [14], and 68% in USA [15]). Although UPF are becoming more prevalent worldwide [1, 17, 18], there are cross-country differences that should be acknowledged. A recent multi-country study assessing adolescents' UPF consumption in upper-middle and high-income countries found that consumption across these countries ranged between 19 to 36% in upper-middle-income countries (i.e., Argentina, Brazil, Colombia, and Mexico), and from 34 to 68% in high-income countries (i.e., Australia, Chile, USA

and UK) with UK adolescents being the highest consumers of UPF [40]. The high consumption of UPF among HIC such as the UK may be partly explained by a combination of social, cultural, economic, and marketing factors [18]. Urbanisation in the UK, as in other HIC, can increase access to a greater diversity of and cheaper foods, including UPF, and increased exposure to commercial marketing with a wide offer of ready-to-eat products [41]. These offer convenient solutions to longer working hours, changes in family structure and contribute to shifting from home food preparations to more ready-to-consume foods [42]. An analysis of household availability of UPF in nineteen European countries showed that in the UK 50.4% of total purchased dietary energy comes from UPF, in contrast with 10.2% in Portugal, 13.4% in Italy and 46.2% in Germany [43].

Our findings suggest a decline of UPF consumption (%TEI and weight) between survey years 1 and 11 in the NDNS (2008/09 – 2018/19). Time trend analysis in NDNS (years 1 to 9) show a decline in total energy intake in this age group, especially between survey years 7 and 8 (2014/15). Additionally, the general downward trend in energy consumption from UPF is consistent with other study findings; a study using a controlled interrupted time series analysis between 2014 and 2019 found a 10% reduction in free sugars purchased per household per week from SSBs between 2014 and 2019 [44], however they did not find a reduction in volume (mL) purchased from SSBs. Another study using data from adults and children in NDNS found a reduction in energy share by 1.4% from sweetened beverages between 2008/09 to 2018/19, also without a reduction in volume (mL) consumed [22]. We hypothesise that this reduction could be partly explained by an increased public awareness and health concerns associated with sugar consumption, government-led campaigns and SSB reformulation to reduce sugar content. However, in this study we did not analyse UPF subgroup consumption and cannot attribute this drop in consumption exclusively to SSBs or free sugars. The reduction in UPF weight consumption in this study might be attributable to other UPF subgroups, or alternatively to a change in NDNS main food group classification or subsidiary group classification in the NDNS Nutrient Databank.

Interestingly, in our sample UPF contributed proportionately more to TEI (65%) than to food weight (43%), which reflects the overall higher energy density of UPF. Energy density is associated with weight gain, type 2 diabetes, and obesity [45]. Some associations were apparent

Table 2 Description of UPF consumption (%kcal and grams per day) of a weighted sample in the NDNS adolescents' sample waves 1–11 (2008/09–2018/19)

% energy from UPF per day ^a	95%CI	Grams of consumption of UPF per day ^a	95%CI	
<i>Sex</i>				
Male	66.0	(65.0, 66.9)	941.5	(911.9, 971.2)
Female	65.8	(64.8, 66.7)	776.3	(748.6, 804.0)
<i>Age</i>				
11	65.6	(63.9, 67.3)	797.0	(741.8, 852.3)
12	66.1	(64.3, 67.8)	824.9	(770.1, 879.7)
13	68.0	(66.1, 69.8)	842.9	(794.5, 891.4)
14	67.2	(65.5, 68.8)	860.1	(803.3, 916.9)
15	66.0	(64.4, 67.7)	862.9	(807.0, 918.8)
16	66.0	(64.4, 67.5)	878.3	(821.3, 935.2)
17	64.4	(62.1, 66.6)	908.8	(852.2, 965.4)
18	63.4	(61.3, 65.5)	918.7	(849.0, 988.3)
<i>Parents' occupation social class^b</i>				
Higher managerial, administrative, and professional	63.8	(62.8, 64.8)	826.3	(796.3, 856.2)
Intermediate	65.9	(64.6, 67.3)	838.4	(796.9, 879.9)
Routine and manual	68.4	(67.3, 69.6)	919.6	(883.4, 955.7)
<i>Survey year (survey wave for data collection)</i>				
(1) 2008/09	67.7	(65.6, 69.8)	995.5	(928.1, 1,063.0)
(2) 2009–2010	68.1	(66.2, 69.9)	948.9	(889.4, 1008.5)
(3) 2010–2011	67.9	(66.4, 69.4)	915.8	(845.4, 986.2)
(4) 2011–2012	67.5	(65.5, 69.5)	880.8	(814.9, 946.7)
(5) 2012–2013	67.0	(65.0, 69.1)	907.0	(846.5, 967.6)
(6) 2013–2014	67.0	(64.8, 69.2)	876.7	(816.1, 937.3)
(7) 2014–2015	67.3	(65.4, 69.1)	913.2	(826.8, 999.5)
(8) 2015–2016	62.0	(59.8, 64.2)	746.0	(680.8, 811.3)
(9) 2016–2017	62.8	(60.2, 65.4)	700.7	(640.7, 760.6)
(10) 2017–2018	62.8	(60.1, 65.4)	820.7	(745.6, 895.9)
(11) 2018–2019	64.6	(62.1, 67.0)	775.9	(710.7, 841.0)
<i>Weight category^c</i>				
Normal weight	65.8	(64.9, 66.6)	841.5	(816.3, 866.8)
Overweight	66.3	(64.7, 67.9)	861.4	(810.8, 912.1)
Obese	65.9	(64.5, 67.4)	924.1	(873.0, 975.2)
<i>Ethnic group</i>				
White	67.3	(66.6, 67.9)	905.8	(883.2, 928.5)
Non-white	59.0	(57.3, 60.8)	638.4	(596.4, 680.3)
<i>Region</i>				
England North	67.4	(66.1, 68.8)	910.8	(866.6, 954.9)
England Central/Midlands	66.8	(65.3, 68.2)	923.7	(866.2, 981.3)
England South (including London)	64.1	(62.9, 65.2)	803.9	(771.2, 836.6)
Scotland	67.8	(65.7, 69.9)	891.6	(830.0, 953.2)
Wales	67.1	(65.5, 68.8)	882.9	(822.3, 943.5)
Northern Ireland	67.9	(66.6, 69.1)	833.9	(792.5, 875.3)
<i>MVPA^d</i>				
<21 min/day	65.1	(61.4, 68.8)	865.1	(776.1, 954.1)
21–52 min/day	65.5	(62.7, 68.3)	921.4	(820.8, 1022.0)
52–124 min/day	65.3	(62.1, 68.5)	845.8	(725.5, 966.2)
> 124 min/day	66.5	(63.7, 69.3)	1052.2	(962.7, 1141.7)

BMI z-score: standardised body mass index; CI: Confidence Interval; MVPA: Moderate-to-vigorous physical activity; NDNS: National Diet and Nutrition Survey; NS-SEC: National Statistics Socioeconomic Class; UPF: ultra-processed food, NOVA4 classification group

^aResults are weighed based on non-selection and non-response survey weights provided by NDNS year 2008–2019

^bParents occupation is based on the three-class NS-SEC. A small number of households were excluded from this classification where the household has never worked or was classified under other

Table 2 (continued)

^cBMI z-score was created by standardising BMI for sex and age based on the 1990 British Growth Reference (UK90)

^dMVPA: There was data available only for 16–18-year-olds, n = 534




Fig. 2 Mean relative energy consumption (%kcal/day) and CI's from UPF across NDNS survey years (adjusted for age and sex)




Fig. 3 Mean grams of UPF consumption (plus 95% confidence interval) across NDNS survey years (adjusted for age and sex)

with food weight but not with %TEI from UPF, for example with sex (higher in males vs. females) and age (higher in older adolescents). These differences largely reflect the higher TEI consumed by males and older adolescents. However, after accounting for TEI we still observed an independent significant effect. We need to understand mechanisms of harm, if any, to help guide finding a correct metric of exposure.

Similar to other findings, we observed that being from a white ethnicity is associated with a significantly higher consumption of UPF [15, 39]. However, the relationship between ethnicity and UPF is complex and multifactorial, and the observed higher UPF consumption among white ethnicity adolescents could be due to other factors associated

with ethnicity, for example, cultural and other economic factors (e.g., home ownership [46]).

Our findings add to the body of knowledge that in HIC, lower socioeconomic groups consume higher levels of UPF [1]. This may be partly due to the greater affordability (i.e., price per calorie) of food with less nutritional value across country incomes and regions [47] alongside targeted marketing to specific population subgroups [48]. Additionally, reduced at-home facilities for food preparation and lack of time to cook at home may lead to an increased consumption of more processed and convenience foods (e.g., pre-prepared meals) [49]. Whilst in HIC most UPF are relatively cheaper than less processed foods, the opposite patterns are seen in LMIC. As an example, soft drinks are relatively inexpensive in HIC, whilst they are relatively expensive in LMIC [1, 18].

There is currently no universally agreed “safe” levels of dietary share from UPF. Therefore, future research to understand the mechanisms of harms to health may substantially improve nutritional quality of adolescent diets and contribute to the prevention of diet related NCDs.

Strengths and limitations

To the best of our knowledge, this is the first study to characterise and present data associating to both %TEI and weight of UPF consumption (g/d) in a representative sample of UK adolescents.

This study has several additional strengths. Due to the consistent methods of data collection over time, the data across waves could be mixed resulting in a relatively large sample size. This study used individual-level dietary data, had 3 or 4 food diary days for each individual and had relatively high numbers of individuals among each group, which is likely to give a more accurate assessment of total dietary intake versus other methods [50, 51]. For example, food records allow researchers to have high levels of detail on dietary intake, they were completed in real-time, which avoids reliance on recall, a common limitation of food frequency questionnaires and 24-h recalls. Additionally, food records, when compared to direct observation and doubly labelled water, perform much better than other self-reported methods and capture about 80% of energy intake [50, 52]. Weighting of the sample was applied to reduce non-response and sampling bias, therefore, the study results can be generalisable to the UK adolescent population. Additionally, this study included a weight measure of UPF to capture non-nutritional factors relating to processing of food (i.e.,

% Kcal/day vs. Reference




Fig. 4 Linear regression of the adjusted associations between participants’ characteristics (vs reference category) and consumption of UPF defined as relative energy (%kcal/day) (adjusted for age, sex and survey year)

additives, non-sugar sweeteners, neo-formed contaminants), and foods and drinks that do not contribute to energy intake (e.g., artificially sweetened beverages). Our results show that there were more individual characteristics associated with weight of UPF consumption (g/d) than UPF contribution to TEI. The inclusion of this measure could further enhance our knowledge about the risks involved by diets high in UPF beyond their contribution to TEI, but this should be systematically tested in studies of UPF and health outcomes.

To assess the variability and potential misclassification of UPF two researchers blindly classified all food and drink items in the food files within NDNS. We reached a 97% agreement in classification across all NOVA groups, and 99.8% agreement for classification of UPF. The variability of our classification of UPF showed that the energy contribution ranged from 65.9% under the current more conservative approach (YCU) to 70% when we applied the more liberal approach. Other studies that have assessed the variability

Grams/day vs. Reference




Fig. 5 Linear regression of the adjusted associations between participants' characteristics (vs reference category) and consumption of UPF defined as absolute weight (g/day) (adjusted for age, sex and survey year)

and potential misclassification of UPF showed that <10% of individual foods and beverages reported in NHANES in the US (24-h recall) were at risk of misclassification [54]. This up to 6 percentage points in variation range provides confidence in the current approach used to classify foods within NDNS according to the degree of processing using the NOVA classification system using food diaries.

Some limitations should be considered. We could not include a measure of household income due to the way

this variable was collected in waves 9 to 11 limiting our knowledge of the impact of household income on UPF consumption among adolescents. However, other proxy for SES (i.e., parent's social occupation social class) showed higher energy (%kcal/d) UPF consumption among adolescents with parents with intermediate and routine and manual occupations, and higher weight (g/d) UPF consumption only in adolescents with parents with routine and manual occupations. Due to the observational nature

of our findings and the proxy measures used for SES more research is needed to unpick socio-economic patterning of consumption among adolescents.

Classification of food items according to the NOVA system was time-consuming because NDNS food diaries were not designed to capture UPF. Some of the main groups and subsidiary food groups were classified easily. However, composite food dishes had to be classified on an individual basis. Based on previous studies that have used either the NOVA classification system and/or NDNS data served as a practical coding framework [7, 8, 34, 35]. Since the food records utilised were not designed to classify or evaluate foods according to their industrial processing, some items may have been misclassified.

MVPA was self-reported using an instrument validated for use only in older adolescents and adults (53). Therefore, the only available data was for 16–18-year-olds, considerably reducing our sample size for this variable. We therefore used the limited data available on MVPA as an exploratory analysis to identify potential associations between UPF consumption and levels of physical activity. Although we did find that the most physically active adolescents consumed a higher weight of UPF compared to the least physically active adolescents, due to the small sample size this association needs to be interpreted with caution. It should be explored further with a larger sample size and with data from younger adolescents (< 16 years).

Conclusion

This study showed that UK adolescents aged 11 to 18 years, living in England North, from the lowest SES group and with white ethnicity have the highest energy and weight intakes of UPF. Additionally, a higher weight consumption from UPF was observed in adolescents who were male, aged 17 to 18 years, and living with obesity. We found that the average energy intake and food weight from UPF has decreased in UK adolescents between year 1 and 11 of NDNS survey waves. However, it remains among the highest levels across high-income countries (e.g., Canada 55.0% and USA 67.7% of TEI). Estimating “safe” levels of dietary share from UPF and understanding the mechanisms of harms to health may substantially improve nutritional quality of adolescent diets and contribute to the prevention of diet-related NCDs.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00394-024-03458-z>.

Acknowledgements The data supporting the results of this study were provided by the National Diet and Nutrition Survey database. We thank all investigators and participants in the study for their contribution.

Author contributions YCU data collection, processing, design, data analysis, writing – first draft; FdV, ZT, RJ – conceptualisation, data processing; EvS, JA, KO, NF, ZT, FdV, RJ – editing, reviewing and supervision; ZC – coding and data processing; LICR – data visualisation. All authors have read and agreed to the final manuscript version.

Funding This study was part of YCU PhD studentship funded by the National Institute for Health and Care Research (NIHR) School for Public Health Research (SPHR) (Grant Reference Number PD-SPH-2015), supervised by Frank de Vocht, Russell Jago, Zoi Toumpakari and Martin White. The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care. The NIHR School for Public Health Research is a partnership between the Universities of Sheffield; Bristol; Cambridge; Imperial; and University College London; The London School for Hygiene and Tropical Medicine (LSHTM); LiLaC – a collaboration between the Universities of Liverpool and Lancaster; and Fuse—The Centre for Translational Research in Public Health a collaboration between Newcastle, Durham, Northumbria, Sunderland and Tees-side Universities. YCU is now a postdoctoral research associate at the Medical Research Council (MRC) to the MRC Epidemiology Unit, University of Cambridge [grant number MC_UU_00006/5] and this study received funding for publication by the MRC Epidemiology Unit. EVS acknowledge support from the MRC Epidemiology Unit (MC_UU_00006/5). NGF acknowledges support from the MRC Epidemiology Unit (MC_UU_00006/3), the National Institute of Health and Care Research (NIHR) Cambridge Biomedical Research Centre (NIHR203312), and she is an NIHR Senior Investigator. The views expressed are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care. JA is supported by the MRC Epidemiology Unit, University of Cambridge [Medical Research Council grant number MC/UU/00006/7] mrc.ukri.org. KO is supported by the Medical Research Council (MC_UU_00006/2). FdV and RJ are partially funded by the National Institute for Health Research Applied Research Collaboration West (NIHR ARC West). The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Data availability The data used in this study are derived from the National Diet and Nutrition Survey (NDNS), conducted by Public Health England (PHE) and the Food Standards Agency (FSA) in the United Kingdom. Access to NDNS data is managed by the UK Data Service (UKDS) under license from PHE and FSA. Researchers interested in accessing NDNS data should consult the UK Data Service website (<https://www.ukdataservice.ac.uk/>) for information on data access procedures and conditions.

Declarations

Competing interests Yanaina Chavez-Ugalde (YCU) – Has no competing interests to declare that are relevant to the content of this article. Frank De Vocht (FdV) – Has no competing interests to declare that are relevant to the content of this article. Russell Jago (RJ) – Has no competing interests to declare that are relevant to the content of this article. Jean Adams (JA) – Has no competing interests to declare that are relevant to the content of this article. Ken K. Ong (KO) – Has no competing interests to declare that are relevant to the content of this article. Nita Forouhi (NF) – Has no competing interests to declare that are relevant to the content of this article. Zoé Colombet (ZC) – Has no competing interests to declare that are relevant to the content of this article. Luíza I.C. Ricardo (LICR) – Has no competing interests to declare that are relevant to the content of this article. Esther Van Sluijs (EvS) – Has no competing interests to declare that are relevant to the content of this article. Zoi Toumpakari (ZT) – Has no competing interests to declare that are relevant to the content of this article.

Ethical standards This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving research study participants for collection of NDNS data was provided by the Oxfordshire A Research Ethics Committee. All participants provided written informed consent to take part in NDNS. Additional ethical approval for this secondary analysis of anonymised data was not required.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Baker P, Machado P, Santos T, Sievert K, Backholer K, Hadjikakou M et al (2020) Ultra-processed foods and the nutrition transition: Global, regional and national trends, food systems transformations and political economy drivers. *Obesity Rev* 21(12):e13126
- Martini D, Godos J, Bonaccio M, Vitaglione P, Grosso G (2021) Ultra-processed foods and nutritional dietary profile: a meta-analysis of nationally representative samples. *Nutrients* 13(10):3390
- Machado P, Cediél G, Woods J, Baker P, Dickie S, Gomes FS et al (2022) Evaluating intake levels of nutrients linked to non-communicable diseases in Australia using the novel combination of food processing and nutrient profiling metrics of the PAHO Nutrient Profile Model. *Eur J Nutr* 61(4):1801–1812
- Monteiro CA, Jaime PC, Cannon G, Levy RB, Louzada MLC, Rauber F et al (2019) Ultra-processed foods: what they are and how to identify them. *Public Health Nutr* 22(5):936–941
- Dicken SJ, Batterham RL (2021) The role of diet quality in mediating the association between ultra-processed food intake, obesity and health-related outcomes: a review of prospective cohort studies. *Nutrients* 14(1):23
- Norman J, Kelly B, Boyland E, McMahon A-T (2016) The impact of marketing and advertising on food behaviours: evaluating the evidence for a causal relationship. *Curr Nutr Rep* 5(3):139–149
- Chang K, Khandpur N, Neri D, Touvier M, Huybrechts I, Millett C et al (2021) Association between childhood consumption of ultraprocessed food and adiposity trajectories in the avon longitudinal study of parents and children birth cohort. *JAMA Pediatr* 175(9):e211573
- Rauber F, da Costa Louzada ML, Steele EM, Millett C, Monteiro CA, Levy RB (2018) Ultra-processed food consumption and chronic non-communicable diseases-related dietary nutrient profile in the UK (2008–2014). *Nutrients* 10(5):587
- Kim H, Hu EA, Rebholz CM (2019) Ultra-processed food intake and mortality in the USA: results from the Third National Health and Nutrition Examination Survey (NHANES III, 1988–1994). *Public Health Nutr* 22(10):1777–1785
- Olstad DL, Kirkpatrick SI (2021) Planting seeds of change: reconceptualizing what people eat as eating practices and patterns. *Int J Behav Nutr Phys Act* 18(1):32
- Patton GC, Sawyer SM, Santelli JS, Ross DA, Afifi R, Allen NB et al (2016) Our future: a Lancet commission on adolescent health and wellbeing. *Lancet (London, England)* 387(10036):2423–2478
- Chavez-Ugalde Y, Jago R, Toumpakari Z, Egan M, Cummins S, White M et al (2021) Conceptualising the commercial determinants of dietary behaviors associated with obesity: a systematic review using principles from critical interpretative synthesis. *Obes Sci Pract* 7(4):473–486
- Leonie E, Machado P, Zinöcker M, Baker P, Lawrence M (2020) Ultra-processed foods and health outcomes: a narrative review. *Nutrients* 12(7):1955
- Moubarac J-C, Batal M, Louzada ML, Martinez Steele E, Monteiro CA (2017) Consumption of ultra-processed foods predicts diet quality in Canada. *Appetite* 108:512–520
- Wang L, Martínez Steele E, Du M, Pomeranz JL, O'Connor LE, Herrick KA et al (2021) Trends in consumption of ultraprocessed foods among US Youths Aged 2–19 Years, 1999–2018. *JAMA* 326(6):519–530
- Lane MM, Gamage E, Du S, Ashtree DN, McGuinness AJ, Gauci S et al (2024) Ultra-processed food exposure and adverse health outcomes: umbrella review of epidemiological meta-analyses. *BMJ*. 384:e077310
- Monteiro CA, Moubarac JC, Cannon G, Ng SW, Popkin B (2013) Ultra-processed products are becoming dominant in the global food system. *Obes Rev* 14(S2):21–28
- Moodie R, Bennett E, Kwong E, Santos TM, Pratiwi L, Williams J et al (2021) Ultra-processed profits: the political economy of countering the global spread of ultra-processed foods - a synthesis review on the market and political practices of transnational food corporations and strategic public health responses. *Int J Health Policy Manag* 10(12):968–982
- Martínez Steele E, Baraldi LG, Louzada ML, Moubarac JC, Mozaffarian D, Monteiro CA (2016) Ultra-processed foods and added sugars in the US diet: evidence from a nationally representative cross-sectional study. *BMJ Open* 6(3):e009892
- Marrón-Ponce JA, Sánchez-Pimienta TG, Louzada M, Batis C (2018) Energy contribution of NOVA food groups and sociodemographic determinants of ultra-processed food consumption in the Mexican population. *Public Health Nutr* 21(1):87–93
- Shim JS, Shim SY, Cha HJ, Kim J, Kim HC (2021) Socioeconomic characteristics and trends in the consumption of ultra-processed foods in Korea from 2010 to 2018. *Nutrients* 13(4):1120
- Madruga M, Martínez Steele E, Reynolds C, Levy RB, Rauber F (2023) Trends in food consumption according to the degree of food processing among the UK population over 11 years. *Br J Nutr* 130(3):476–483
- Julia C, Martinez L, Allès B, Touvier M, Hercberg S, Méjean C et al (2018) Contribution of ultra-processed foods in the diet of adults from the French NutriNet-Santé study. *Public Health Nutr* 21(1):27–37
- Lustig RH, Collier D, Kassotis C, Roepke TA, Kim MJ, Blanc E et al (2022) Obesity I: overview and molecular and biochemical mechanisms. *Biochem Pharmacol* 199:115012
- Schulze K (2020) Ultra-processed foods and cardiometabolic health. University of Cambridge, Cambridge
- National Diet and Nutrition Survey Years 1–8, 2008/09–2015/16. [data collection] [Internet]. UK Data Service. 2018. Available from: <https://beta.ukdataservice.ac.uk/datacatalogue/studies/study?id=6533&type=Data%20catalogue#!/details>.
- Public Health England. Appendix A. Dietary data collection and editing. In National Diet and Nutrition Survey. Results from Years 1–4 (Combined) of the Rolling Programme (2008/2009–2011/2012). London, UK: Public Health England; 2014.

28. Fitt E, Cole D, Ziauddeen N, Pell D, Stickley E, Harvey A et al (2015) DINO (Diet in Nutrients Out)-An integrated dietary assessment system. *Public Health Nutr* 18(2):234–241
29. Public Health England (2015) McCance and Widdowson's the composition of foods integrated dataset 2015. In: *Public Health England*, editor. London, UK
30. Fitt E, Mak TN, Stephen AM, Prynne C, Roberts C, Swan G et al (2010) Disaggregating composite food codes in the UK National Diet and Nutrition Survey food composition databank. *Eur J Clin Nutr* 64(3):S32–S6
31. MRC Human Nutrition Research (2017) Food Standards Agency Standard Recipes Database, 1992–2012 [data collection]. In: *UK Data Service*, editor
32. Monteiro CA, Jaime PC, Cannon G, Moubarac JC, Levy RB, Louzada MLC (2018) The UN decade of nutrition, the NOVA food classification and the trouble with ultra-processing. *Public Health Nutr* 21(1):5–17
33. Monteiro CA, Cannon G, Lawrence M, Costa Louzada ML, Pereira Machado P (2019) Ultra-processed foods, diet quality, and health using the NOVA classification system. *FAO, Rome*
34. Martines RM, Machado PP, Neri DA, Levy RB, Rauber F (2019) Association between watching TV whilst eating and children's consumption of ultraprocessed foods in United Kingdom. *Mater Child Nutr* 15(4):n/a–n/a
35. Rauber F, Martins CA, Azeredo CM, Leffa PS, Louzada MLC, Levy RB (2022) Eating context and ultraprocessed food consumption among UK adolescents. *Br J Nutr* 127(1):112–122
36. Golubic R, May AM, Benjaminsen Borch K, Overvad K, Charles M-A, Diaz MJT et al (2014) Validity of electronically administered recent physical activity questionnaire (RPAQ) in Ten European Countries. *PLoS One* 9(3):e92829
37. Rose D, Pevalin D, Or K (2005) *The National Statistics Socio-economic Classification: origins, development and use*: Palgrave Macmillan UK
38. UK Data Archive Study (2020) *Weights Guide: Combining data from Years 1–4, Years 5&6, Years 7&8 and Years 9–11*, National Diet and Nutrition Survey
39. Parnham JC, Chang K, Rauber F, Levy RB, Millett C, Laverly AA et al (2022) The ultra-processed food content of school meals and packed lunches in the United Kingdom. *Nutrients* 14(14):2961
40. Neri D, Steele EM, Khandpur N, Cediel G, Zapata ME, Rauber F et al (2022) Ultraprocessed food consumption and dietary nutrient profiles associated with obesity: a multicountry study of children and adolescents. *Obesity Reviews*. 23(S1):e13387
41. Popkin BM, Adair LS, Ng SW (2012) Global nutrition transition and the pandemic of obesity in developing countries. *Nutr Rev* 70(1):3–21
42. Smith LP, Ng SW, Popkin BM (2013) Trends in US home food preparation and consumption: analysis of national nutrition surveys and time use studies from 1965–1966 to 2007–2008. *Nutr J* 12:45
43. Monteiro CA, Moubarac JC, Levy RB, Canella DS, Louzada M, Cannon G (2018) Household availability of ultra-processed foods and obesity in nineteen European countries. *Public Health Nutr* 21(1):18–26
44. Pell D, Mytton O, Penney TL, Briggs A, Cummins S, Penn-Jones C et al (2021) Changes in soft drinks purchased by British households associated with the UK soft drinks industry levy: controlled interrupted time series analysis. *BMJ* 372:n254
45. Wang J, Luben R, Khaw K-T, Bingham S, Wareham NJ, Forouhi NG (2008) Dietary energy density predicts the risk of incident Type 2 Diabetes: The European Prospective Investigation of Cancer (EPIC)-Norfolk Study. *Diabetes Care* 31(11):2120–2125
46. Poti JM, Mendez MA, Ng SW, Popkin BM (2015) Is the degree of food processing and convenience linked with the nutritional quality of foods purchased by US households? *Am J Clin Nutr* 101(6):1251–1262
47. Headey DD, Alderman HH (2019) The relative caloric prices of healthy and unhealthy foods differ systematically across income levels and Continents. *J Nutr* 149(11):2020–2033
48. Kelly B, Vandevijvere S, Freeman B, Jenkin G (2015) New media but same old tricks: food marketing to children in the digital age. *Curr Obes Rep* 4(1):37–45
49. Adams J, White M (2015) Characterisation of UK diets according to degree of food processing and associations with socio-demographics and obesity: cross-sectional analysis of UK National Diet and Nutrition Survey (2008–12). *Int J Behav Nutr Phys Act* 12(1):160
50. Black AE, Prentice AM, Goldberg GR, Jebb SA, Bingham SA, Livingstone MB et al (1993) Measurements of total energy expenditure provide insights into the validity of dietary measurements of energy intake. *J Am Diet Assoc* 93(5):572–579
51. Bingham SA (1991) Limitations of the various methods for collecting dietary intake data. *Ann Nutr Metab* 35(3):117–127
52. Park Y, Dodd KW, Kipnis V, Thompson FE, Potischman N, Schoeller DA et al (2018) Comparison of self-reported dietary intakes from the Automated Self-Administered 24-h recall, 4-d food records, and food-frequency questionnaires against recovery biomarkers. *Am J Clin Nutr* 107(1):80–93
53. Mindell J, Stamatakis E, Brage S (2009) Measuring Energy Expenditure in NDNS 2008–2012. [Google Scholar](#)
54. Steele EM, O'Connor LE, Juul F, Khandpur N, Galastri Baraldi L, Monteiro CA et al (2023) Identifying and estimating ultraprocessed food intake in the US NHANES according to the nova classification system of food processing. *J Nutr* 153(1):225–241