



# City Research Online

## City St George's, University of London

**Citation:** Naderi, K., Jameel, A., Low, S., Wagh, V., Bhogal, M., Ritchie, A., Robbie, S., Hammond, C., Mohamed, M., Stanojcic, N., et al (2024). 'Off the shelf' toric intraocular lenses to allow better access in public healthcare: a randomised control study. *Eye*, 38(13), pp. 2651-2652. doi: 10.1038/s41433-024-03068-3

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/33701/>

**Link to published version:** <https://doi.org/10.1038/s41433-024-03068-3>

**Copyright and Reuse:** Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

## **‘Off the Shelf’ Toric Intraocular Lenses to allow better access in Public Healthcare: A Randomised Control Study**

Khayam Naderi <sup>1,2</sup> FRCOphth, Ashmal Jameel <sup>1,2</sup> FRCOphth, Sancy Low <sup>1</sup> FRCOphth PhD, Vijay Wagh<sup>1</sup> FRCOphth MD, Mani Bhogal <sup>1</sup> FRCOphth PhD, Ailsa Ritchie<sup>1</sup> FRCOphth, Scott Robbie <sup>1</sup> FRCOphth PhD, Chris Hammond <sup>1,2</sup> FRCOphth MD, Moin Mohamed <sup>1</sup> FRCOphth PhD, Nick Stanojcic <sup>1,2</sup> FRCOphth MD, Elodie Azan <sup>1</sup> FRCOphth PhD, Lily Lai <sup>1</sup> FRCOphth, Chris Hull <sup>3</sup> PhD, David O’Brart <sup>1,2</sup> FRCOphth MD.

1. Department of Ophthalmology, St. Thomas’ Hospital, Lambeth Palace Road, London, SE1 7EH
2. King’s College, London, WC2R 2LS
3. Department of Optometry and Visual Science, School of Health and Psychological Sciences, City, University of London, Northampton Square, London EC1V 0HB.

Corresponding author. Professor David O’Brart, Department of Ophthalmology, St. Thomas’ Hospital, Lambeth Palace Road, London, SE1 7EH. Email [david.obrart@gstt.nhs.uk](mailto:david.obrart@gstt.nhs.uk)

**RUNNING HEAD:** Two-cylinder power ‘Off the Shelf’ Toric Intraocular Lenses

**ARTICLE TYPE:** Brief Communication

**Key words:** Toric intraocular lenses, Cataract Surgery, National Health Service

**Financial Disclosure of all authors:** Professor O’Brart holds non-commercial research grants with Rayner Ltd. and J&J Inc. The authors declare no conflicts of interest.

**Disclosure:** This study was supported by a non-commercial grant from Rayner Ltd.

Despite benefits (1), toric intraocular lenses (TIOs) comprise only 0.4% of IOLs implanted in the NHS (2). Reasons include typically higher IOL prices, delivery logistics/costs as on-site storage of TIOs is impractical due to large numbers of TIO spherical and cylindrical combinations, and increased consultation/operative and administration times (3). By using TIOs with only two-cylinder powers, on-site TIO banks might be feasible due to lower numbers of lens combinations and TIO unit price possibly reduced by bulk manufacture (3).

This was an ethics approved (19/WA/0272), single-centre, prospective, single-masked, randomised case-control study, which adhered to the tenets of the Declaration of Helsinki and UK Data Protection Act (2018). Inclusion criteria were age over 18 with full capacity and symptomatic cataract(s), without other significant ocular pathologies and regular corneal astigmatism (1.50-5.00D).

Forty-seven patients were randomised to a fully tailored' (FT) and 44 to an 'off the shelf' group (OTS). The refractive target was emmetropia, with the aspheric RayOne single-piece, hydrophilic acrylic TIO (Rayner, Worthing, West Sussex, UK) being implanted. FT patients received a TIO with full correction of corneal astigmatism. OTS patients received either a 2.00D or 4.00D cylinder TIO, with residual calculated astigmatism corrected by opposite clear corneal incisions (OCCI), the degree of astigmatism correction being estimated at 0.25D and 0.50D with 2.4-millimetre (mm) and 2.75mm incision sizes respectively (4-5). Patients were reviewed at 4 weeks (4W) and 6 months (6M). Primary outcome measures were monocular uncorrected distance visual acuity (UDVA), best-corrected distance visual acuity (BDVA), and residual subjective refractive cylinder (D). Secondary outcomes were patient-reported outcome measures (PROMs), TIO rotational stability, and adverse events. Statistical analysis was performed using GraphPad Prism (GraphPad Software, San Diego, California USA). Alpins vector analysis was performed using the online VekTrAK software (<http://www.assort.com>). Data-set normality was assessed using the Shapiro-Wilk test. Parametric data were analysed with student unpaired t-test with statistical significance set at  $p < 0.05$ .

There were no differences in baseline demographics (table 1). At 4W, mean UDVA (+/- standard deviation) was 0.14 (0.11) in the OTS and 0.15 (0.14) in the FT group ( $p=0.65$ ); mean BDVA was 0.02 (0.1) in OTS and 0.00 (0.085) in FT ( $p=0.27$ ); mean RC was 0.73D (0.39) in OTS, and 0.85D (0.49) in FT ( $p=0.23$ ). At 6M, mean UDVA was 0.16 (0.15) in OTS and 0.10 (0.13) in FT ( $p=0.075$ ); mean BDVA was 0.01 (0.10) in OTS and -0.01 (0.09) in FT ( $p=0.34$ ); mean RC was 0.89D (0.50) in OTS and 0.84D (0.42) in FT ( $p=0.59$ ). There were no differences in PROMs between the groups (table 2). No cases required further surgery to reposition the TIO.

In this study our OTS approach was not significantly inferior to a FT method. Using TIOs with only 2.0DC and 4.0DC cylinder powers could facilitate increased access to TIOs in the NHS/public health sector and the developing world by allowing 'on-site' TIO banks, reducing costs in administrative time, lens transport/delivery, inventory wastage and giving manufacturers the opportunity to produce 2.00D and 4.00DC TIOs in high volume, allowing possible reductions in production costs.

1. Kessel L, Andresen J, Tendal B, Erngaard D, Flesner P, Hjortdal J. Toric Intraocular Lenses in the Correction of Astigmatism During Cataract Surgery: A Systematic Review and Meta-analysis. *Ophthalmology*. 2016 Feb;123(2):275-286.
2. The Royal College of Ophthalmologists. *National Ophthalmology Database Audit: Feasibility study of Post-cataract Posterior Capsule Opacification*, 2021. Available from: <https://www.nodaudit.org.uk/u/docs/20/rijbxkcubs/RCOphth%20NOD%20PCO%20Report%202021.pdf> [Accessed 20<sup>th</sup> January 2023].
3. Stanojic N, Roberts H, Wagh V, Zuberbuhler B, O'Brart D. A randomised, prospective study of 'off-the-shelf' use of toric intraocular lenses for cataract patients with pre-existing corneal astigmatism in the NHS. *Eye (Lond)*. 2020 Oct;34(10):1809-1819.8u7
4. Kavşut, Bilal & Özcan, Sait & Tunçbilek, Zehra & Solmaz, Nilgün & Önder, Feyza. (2018). Cataract Surgery Outcomes: Comparison of 2.4 mm and 2.8 mm Clear Corneal Incisions. *Van Medical Journal*. 25. 349-354.
5. Ernest P, Hill W, Potvin R. Minimizing surgically induced astigmatism at the time of cataract surgery using a square posterior limbal incision. *J Ophthalmol*. 2011;2011:243170.

Parameter	Off The Shelf (n=47) Mean (SD)	Fully Tailored (n=44) Mean (SD)	p-value (95% Confidence Interval)
Age (Years)	69.13 (11.86)	69.45 (11.97)	0.90 (-4.64 to 5.29)
Gender (Male:Female)	25:22	12:32	0.018
Pre-operative UDVA (logMAR)	0.92 (0.52)	0.83 (0.43)	0.37 (-0.29 to 0.11)
Pre-operative BDVA (logMAR)	0.55 (0.45)	0.46 (0.37)	0.32 (-0.26 to 0.86)
Axial Length (mm)	24.21 (1.47)	24.64 (1.55)	0.18 (-2.00 to 1.06)
Pre-operative Biometric K1-K2 (D)	2.31 (0.62)	2.33 (0.64)	0.89 (-0.24 to 0.28)
Pre-operative Pentacam K1-K2 (D)	2.25 (0.67)	2.23 (0.73)	0.92 (-0.31 to 0.28)
Endothelial Cell Count	2498 (263.7)	2515 (280.9)	0.76 (-96.25 to 130.6)
IOP (mmHg)	15.43 (3.11)	14.43 (3.47)	0.15 (-3.27 to 0.38)
CATPROM-5 Rasch-calibrated Score	0.809 (2.57)	0.301 (2.63)	0.35 (-1.59 to 0.57)
EQ-5D-3L Value Score	0.888 (0.15)	0.808 (0.24)	0.06 (-0.16 to 0.0033)
EQ-5D-3L VAS	83.77 (18.18)	79.52 (18.77)	0.28 (-11.94 to 3.45)

Table 1: Baseline Demographics of the two study groups.

SD=Standard deviation; UDVA=Uncorrected distance visual acuity; BDVA= Best-corrected distance visual acuity; K=Keratometry; D=Dioptres; IOP=Intraocular Pressure (measured using icare); VAS=Visual Analogue Scale.

Fisher's exact test to compare proportions of each gender between the two groups,

Unpaired t-test used to compare means between two groups. Mean (+/-Standard deviation)

	4 weeks Follow-up (mean, SD)			6 months Follow-up (mean, SD)		
	OTS (n=47)	FT (n=44)	p-value (95% CI)	OTS (n=43)	FT (n=41)	p-value (95% CI)
UDVA (logMAR)	0.14 (0.11)	0.15 (0.14)	0.65 (-0.41 to 0.066)	0.16 (0.15)	0.10 (0.13)	0.075 (-0.12 to 0.0057)
BDVA (logMAR)	0.023 (0.098)	0.0011 (0.085)	0.27 (-0.060 to 0.017)	0.01 (0.10)	-0.01 (0.09)	0.34 (-0.063 to 0.022)
Residual Subjective Refractive Cylinder (D)	0.73 (0.39)	0.85 (0.49)	0.23 (-0.073 to 0.30)	0.89 (0.50)	0.84 (0.42)	0.59 (-0.25 to 0.15)
SE (D)	-0.45 (0.52)	-0.60 (0.47)	0.17 (-0.35 to 0.062)	-0.30 (0.58)	-0.37 (0.39)	0.53 (-0.28 to 0.15)
Pentacam K1-K2 (D)	1.93 (0.74)	2.06 (0.70)	0.40 (-0.17 to 0.43)	1.98 (0.62)	2.12 (0.75)	0.35 (-0.16 to 0.44)
Endothelial Cell Count	2191 (416)	2002 (580)	0.08 (-399.7 to 20.49)	2259 (399.4)	2163 (372.1)	0.26 (-265.3 to 72.50)
IOP (mmHg)	13.04 (3.83)	12.73 (3.82)	0.70 (-1.91 to 1.28)	11.81 (2.68)	12.05 (3.18)	0.72 (-1.04 to 1.51)
CATPROM-5 Score	-4.30 (2.63)	-4.23 (2.81)	0.90 (-1.07 to 1.01)	-6.15 (3.03)	-5.19 (2.91)	0.14 (-0.33 to 2.25)
EQ-5D-3L Value Score	0.93 (0.12)	0.91 (0.13)	0.49 (-0.071 to 0.034)	0.88 (0.18)	0.84 (0.20)	0.31 (-0.12 to 0.041)
EQ-5D-3L VAS	88.23 (11.15)	85.74 (12.94)	0.33 (-7.54 to 2.56)	87.12 (15.38)	87.73 (11.49)	0.84 (-5.36 to 6.57)
TIA (D)	2.31 (0.62)	2.33 (0.64)	0.89 (-0.24 to 0.28)	2.28 (0.59)	2.29 (0.63)	0.28 (-0.25 to 0.28)
SIA (D)	2.46 (0.82)	2.54 (1.05)	0.68 (-0.31 to 0.47)	2.35 (0.98)	2.46 (0.93)	0.60 (-0.31 to 0.52)
DV (D)	0.73 (0.39)	0.84 (0.37)	0.27 (-0.079 to 0.28)	0.89 (0.50)	0.84 (0.52)	0.59 (-0.25 to 0.15)
AOE (Degrees)	2.02 (7.59)	2.84 (9.52)	0.65 (-2.76 to 4.40)	4.65 (9.96)	2.90 (9.28)	0.41 (-5.93 to 2.44)
MOE (D)	0.15 (0.57)	0.22 (0.67)	0.62 (-0.19 to 0.32)	0.073 (0.73)	0.17 (0.59)	0.52 (-0.20 to 0.39)
CI	1.07 (0.27)	1.08 (0.31)	0.89 (-0.11 to 0.13)	1.03 (0.34)	1.07 (0.29)	0.58 (-0.099 to 0.18)
IOS	0.34 (0.21)	0.38 (0.24)	0.46 (-0.059 to 0.13)	0.51 (0.23)	0.38 (0.20)	0.53 (-0.12 to 0.063)
COA	1.00 (0.31)	1.01 (0.56)	0.58 (-0.13 to 0.24)	1.20 (1.06)	1.05 (0.20)	0.41 (-0.51 to 0.21)

Table 2: Outcomes at 4-weeks and 6-months follow-up between the two groups.

SD=Standard deviation; CI=Confidence Interval; OTS=Off The Shelf; FT=Fully Tailored; UDVA=Uncorrected distance visual acuity; BDVA= Best-corrected distance visual acuity; K=Keratometry; D=Dioptres; IOP=Intraocular Pressure (measured using icare); VAS=Visual Analogue Scale; TIA=Target Induced Astigmatism; SIA=Surgically Induced Astigmatism; DV=Difference Vector; AOE=Angle of Error; MOE=Magnitude of Error; CI=Correction Index; IOS=Index of Success; COA=Coefficient of Adjustment.

Unpaired t-test used to compare means between two groups. Mean (+/-Standard Deviation)