



City Research Online

City, University of London Institutional Repository

Citation: Bowman, C., De Visscher, M., Farrell, N., Hazi, A. & Norton, E. (2025). Oriented Temperley-Lieb algebras and Combinatorial Kazhdan-Lusztig theory. *Canadian Journal of Mathematics*, pp. 1-43. doi: 10.4153/S0008414X24001032

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/34006/>

Link to published version: <https://doi.org/10.4153/S0008414X24001032>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk

ORIENTED TEMPERLEY–LIEB ALGEBRAS AND COMBINATORIAL KAZHDAN–LUSZTIG THEORY

CHRIS BOWMAN, MAUD DE VISSCHER, NIAMH FARRELL, AMIT HAZI, AND EMILY NORTON

ABSTRACT. We define oriented Temperley–Lieb algebras for Hermitian symmetric spaces. This allows us to explain the existence of closed combinatorial formulae for the Kazhdan–Lusztig polynomials for these spaces.

1. INTRODUCTION

To each parabolic Coxeter system, (W, P) , we have an associated family of “anti-spherical Kazhdan–Lusztig polynomials”, $n_{\lambda, \nu}(q)$, indexed by pairs of cosets for $P \leq W$. These polynomials are some of the most important combinatorial objects in Lie theory and representation theory and they can be computed (at least in theory) via a recursive, non-positive formula. Deodhar proposed a (non-recursive!) combinatorial approach to studying these polynomials in [Deo90].

Libedinsky–Williamson categorified the anti-spherical Kazhdan–Lusztig polynomials by interpreting them as composition factor multiplicities of simple modules within standard modules for the *anti-spherical Hecke category*, $\mathcal{H}_{(W, P)}$ [LW]. In more detail: we first fix a reduced word $\underline{\mu}$ for each $\mu \in {}^P W$, for $\lambda \in {}^P W$ the standard $\mathcal{H}_{(W, P)}$ -module $\Delta(\lambda)$ has *light leaves basis* enumerated by $\cup_{\mu \in {}^P W} \text{Path}(\lambda, \underline{\mu})$ the set of all paths (or “Bruhat strolls”) in the coset graph for (W, P) which terminate at λ such that the steps in these paths are “coloured by” $\underline{\mu}$. This basis is graded according to the degree statistic for the underlying paths, we record this in the matrix

$$\Delta^{(W, P)} := (\Delta_{\lambda, \underline{\mu}}(q))_{\lambda, \mu \in {}^P W} \quad \Delta_{\lambda, \underline{\mu}}(q) = \sum_{S \in \text{Path}(\lambda, \underline{\mu})} q^{\deg(S)}$$

which is a (square) lower uni-triangular matrix. This matrix can be factorised *uniquely* as a product of lower uni-triangular matrices

$$N^{(W, P)} := (n_{\lambda, \nu}(q))_{\lambda, \nu \in {}^P W} \quad B^{(W, P)} := (b_{\nu, \underline{\mu}}(q))_{\nu, \mu \in {}^P W}$$

such that $n_{\lambda, \nu}(q) \in q\mathbb{Z}[q]$ for $\lambda \neq \nu$ and $b_{\nu, \underline{\mu}}(q) \in \mathbb{Z}[q + q^{-1}]$. The polynomial $n_{\lambda, \nu}(q)$ is the anti-spherical Kazhdan–Lusztig polynomial for $\lambda \leq \nu \in {}^P W$. Over the complex field, the polynomial $n_{\lambda, \nu}(q)$ counts the graded composition factor multiplicity $[\Delta(\lambda) : L(\nu)]$ and the polynomials $b_{\nu, \underline{\mu}}(q)$ describe the graded character of the simple module $L(\nu)$ [EW14, LW]. This provides an *innately positive* interpretation of the coefficients of the polynomial $n_{\lambda, \nu}(q)$ and thus proves the famous Kazhdan–Lusztig positivity conjecture [EW14] and its anti-spherical counterpart [LW].

Libedinsky–Williamson proposed that this extra $\mathcal{H}_{(W, P)}$ -structure should provide new insight toward Deodhar’s goal of a counting formula for the Kazhdan–Lusztig polynomials. They ask in [LW21, Problem 1.2] whether it is possible to construct an explicit basis of “canonical light leaves” for a \mathbb{Z} -module $\mathbb{N}_{\lambda, \underline{\mu}}$ whose graded rank is equal to $n_{\lambda, \nu}(q)$. Each canonical light leaf basis element of degree k would then be a generator of some composition factor of $\Delta(\lambda)$ isomorphic to $L(\nu)\langle k \rangle$. We solve this problem in the case of Hermitian symmetric pairs (see Figure 1) by introducing an oriented Temperley–Lieb algebra of type (W, P) for all Hermitian symmetric pairs (W, P) .

Theorem A. *Let (W, P) be a Hermitian symmetric pair. For all $\lambda, \nu \in {}^P W$, the space $\mathbb{N}_{\lambda, \underline{\mu}}$ has basis indexed by the set of “standard” basis elements in the anti-spherical module for the oriented Temperley–Lieb algebra of type (W, P) . These elements can be described in a closed*

combinatorial (non-iterative) fashion. Moreover, this construction is entirely independent of the choice of a reduced word $\underline{\nu}$.




FIGURE 1. Enumeration of nodes in the parabolic Dynkin diagram of the Hermitian symmetric pairs. Namely, types $(A_n, A_{k-1} \times A_{n-k})$, (C_n, A_{n-1}) and (B_n, B_{n-1}) , (D_{n+1}, A_n) and (D_{n+1}, D_n) and (E_6, D_5) and (E_7, E_6) respectively. The single node not belonging to the parabolic is highlighted in pink in each case.

The use of Temperley–Lieb style combinatorics for calculating Kazhdan–Lusztig polynomials goes back to work of Brundan and Stroppel in type $(A_n, A_k \times A_{n-k-1})$ [BS11a, BS10, BS11b, BS12] and Cox and De Visscher in type (D_n, A_{n-1}) [CD11]. In this paper, we generalise these ideas to all Hermitian symmetric pairs and lift the combinatorics to a higher structural level; we do this by interpreting these polynomials as graded-dimensions of “anti-spherical modules” for oriented Temperley–Lieb algebras of type (W, P) . The definition of these algebras and their anti-spherical modules is simple and uniformly given in terms of the underlying root system, see Definitions 3.3 and 3.9. We also relate these newly defined oriented Temperley–Lieb algebras of type (W, P) to the generalised Temperley–Lieb algebras of type W introduced by Fan and Graham. In Section 4, we give a proof of Theorem A for types (D_n, D_{n-1}) , (B_n, B_{n-1}) and the exceptional types (E_6, D_5) and (E_7, E_6) . From Section 5 onwards we focus solely on the remaining types, namely $(W, P) = (A_n, A_k \times A_{n-k-1})$, (D_n, A_{n-1}) and (C_n, A_{n-1}) . In Section 6 we prove that the oriented Temperley–Lieb algebras admit a diagrammatic visualisation, and use this in Section 7 to understand the graded structure of the algebra by way of *closed combinatorial formulas*. In Section 8 we apply these ideas to the anti-spherical module and hence prove Theorem A for the remaining types.

Further rewards of our approach will be harvested in the companion paper [BDHN]. In [BDHN], we establish isomorphisms interrelating Hecke categories and use these isomorphisms in order to construct the basic algebras of these Hecke categories and prove that they are standard Koszul (this uses the results of this paper in order to deduce the required graded vector space dimension counts) and to prove that the p -Kazhdan–Lusztig polynomials are entirely independent of the prime $p \geq 0$.

Acknowledgements. *The first and fourth authors are grateful for funding from EPSRC grant EP/V00090X/1 and the Royal Commission for the Exhibition of 1851, respectively. The authors are thankful to Jon Brundan and Catharina Stroppel for interesting and informative conversations and to the MFO workshop “Character Theory and Categorification” for providing an excellent environment for collaboration.*

We are also very thankful to the anonymous referee for their incredibly detailed comments, which improved the exposition of the paper a great deal.

2. KAZHDAN-LUSZTIG POLYNOMIALS AND DEODHAR'S DEFECT

Let (W, S_W) be a Coxeter system: W is the group generated by the finite set S_W subject to the relations $(st)^{m_{st}} = 1$ for $s, t \in S_W$, $m_{st} \in \mathbb{N} \cup \{\infty\}$ satisfying $m_{st} = m_{ts}$, and $m_{st} = 1$ if and only if $s = t$. Let $\ell : W \rightarrow \mathbb{N}$ be the corresponding length function. Consider $S_P \subseteq S_W$ a subset and (P, S_P) its corresponding Coxeter system. We say that P is the parabolic subgroup corresponding to $S_P \subseteq S_W$. Let ${}^P W \subseteq W$ denote a set of minimal length coset representatives in $P \backslash W$. For $\underline{w} = s_{i_1} s_{i_2} \cdots s_{i_\ell}$ an expression in the generators $s_{i_j} \in S_W$ for $0 \leq j \leq \ell$, we define a subexpression of \underline{w} to be an expression of the form $\underline{w}^{\underline{k}} := s_{i_1}^{k_1} s_{i_2}^{k_2} \cdots s_{i_\ell}^{k_\ell}$ where $\underline{k} = (k_1, k_2, \dots, k_\ell) \in \{0, 1\}^\ell$. We let \leq denote the (strong) Bruhat order on ${}^P W$: namely $y \leq w$ if for some reduced expression \underline{w} for w , there exists a reduced expression \underline{y} for y such that \underline{y} is a subexpression of \underline{w} .

We define a directed graph $\mathcal{G}_{(W,P)}$ with vertex set ${}^P W$ and edges defined as follows. For $\lambda, \mu \in {}^P W$ we have an edge $\lambda \rightarrow \mu$ if $\mu = \lambda s_i > \lambda$ for some $s_i \in S_W$. (Note that this is the Hasse diagram of the poset $({}^P W, \leq_r)$ where \leq_r denotes the (weak) right Bruhat order). Examples are given in Figures 2 and 3.

The identity element $1 \in W$ is the minimal coset representative of the identity coset P , and for convenience, we will denote it by \emptyset instead (the empty word in the generators).




FIGURE 2. The graph $\mathcal{G}_{(W,P)}$ for $(W, P) = (A_3, A_1 \times A_1)$ and $(A_4, A_1 \times A_2)$ respectively. (We haven't drawn the direction on the edges but all arrows are pointing upwards).

We now define $\widehat{\mathcal{G}}_{(W,P)}$ to be the directed graph having the same set of vertices as $\mathcal{G}_{(W,P)}$ but replacing each edge in $\mathcal{G}_{(W,P)}$ between λ and λs_i by four directed edges

$$\lambda \xrightarrow{s_i} \lambda, \quad \lambda s_i \xrightarrow{s_i} \lambda s_i, \quad \lambda \xrightarrow{s_i} \lambda s_i, \quad \lambda s_i \xrightarrow{s_i} \lambda$$

and, in order to keep the notation to a minimum, we will simply label the edge by the subscript of the reflection (not the reflection itself). We assign a degree to each edge in $\widehat{\mathcal{G}}_{(W,P)}$ by setting

$$\deg(\lambda \xrightarrow{i} \lambda s_i) = \deg(\lambda s_i \xrightarrow{i} \lambda) = 0 \quad \deg(\lambda \xrightarrow{i} \lambda) = \begin{cases} 1 & \text{if } \lambda s_i > \lambda \\ -1 & \text{if } \lambda s_i < \lambda \end{cases}$$

Given a path (or ‘‘Bruhat stroll’’) on $\widehat{\mathcal{G}}_{(W,P)}$

$$T : \lambda_1 \xrightarrow{i_1} \lambda_2 \xrightarrow{i_2} \lambda_3 \xrightarrow{i_3} \cdots \xrightarrow{i_{k-1}} \lambda_k,$$

we say that the degree $\deg(\mathbb{T})$ is the sum of the degrees of each edge in \mathbb{T} . (The degree is also sometimes known as the “Deodhar defect”.) We also define the weight of \mathbb{T} , denoted by $\omega(\mathbb{T})$ to be the expression

$$\omega(\mathbb{T}) := s_{i_1} s_{i_2} s_{i_3} \dots s_{i_{k-1}}.$$

We write $\text{Path}_{(W,P)}$ for the set of all paths on $\widehat{\mathcal{G}}_{(W,P)}$. For $\lambda, \nu \in {}^P W$, we let $\text{Path}(\lambda \rightarrow \nu)$ denote the set of all paths in $\text{Path}_{(W,P)}$ beginning at λ and ending at ν . When $\lambda = \emptyset$ we set $\text{Path}(\nu) := \text{Path}(\emptyset \rightarrow \nu)$. Let \underline{w} be an expression in the generators S_W . We set $\text{Path}(\lambda \rightarrow \nu, \underline{w})$ to be the set of paths $\mathbb{T} \in \text{Path}(\lambda \rightarrow \nu)$ with $\omega(\mathbb{T}) = \underline{w}$. When $\lambda = \emptyset$ we set $\text{Path}(\nu, \underline{w}) := \text{Path}(\emptyset \rightarrow \nu, \underline{w})$.

Throughout the paper, we fix one reduced expression $\underline{\mu}$ for each $\mu \in {}^P W$. The set of paths $\text{Path}(\lambda, \underline{\mu})$ for $\lambda, \mu \in {}^P W$ will play a crucial role. Examples of such paths are given in Figure 4. We will see in particular that, for Hermitian symmetric pairs, the set $\text{Path}(\lambda, \underline{\mu})$ consists of either 0 or 1 elements.




FIGURE 3. The graph $\mathcal{G}_{(W,P)}$ for types $(W,P) = (D_5, D_4)$ and (B_4, B_3) . The general case (D_{n+1}, D_n) and (B_n, B_{n-1}) is no more difficult (see [BDHN, Section 1]) – merely extend the top and bottom vertical chains of the graph.




FIGURE 4. On the left we depict the unique path in $\text{Path}(\alpha, \underline{\alpha})$ corresponding with a choice of reduced word $\underline{\alpha}$ and on the right we depict the unique element of $\text{Path}(\beta, \underline{\alpha})$ for $\alpha = s_2 s_3 s_4 s_1 s_2 s_3$ and $\beta = s_2 s_1$. These are paths on $\widehat{\mathcal{G}}_{(A_4, A_1 \times A_2)}$ (also known as “Bruhat strolls”) but we depict only the edges in $\mathcal{G}_{(A_4, A_1 \times A_2)}$ (for readability).

The following path theoretic definition of Kazhdan–Lusztig polynomials was for a long time talked about implicitly in the literature, see for example [Deo90] (in particular Proposition 3.5 and Section 4, and also Section 5 for the parabolic setting). It is explicitly proven to be equivalent to the classical definition of these polynomials in [Soe97, Proposition 3.3].

Definition 2.1. Let $\lambda, \mu \in {}^P W$. We set $b_{\lambda, \underline{\lambda}}(q) = 1 = n_{\lambda, \lambda}(q)$. For $\lambda \neq \mu$, we recursively define the polynomials

$$b_{\lambda, \underline{\mu}}(q) \in \mathbb{Z}[q + q^{-1}] \quad n_{\lambda, \underline{\mu}}(q) \in q\mathbb{Z}[q]$$

by induction on the Bruhat order \leq as follows

$$b_{\lambda, \underline{\mu}}(q) + n_{\lambda, \underline{\mu}}(q) = \sum_{S \in \text{Path}(\lambda, \underline{\mu})} q^{\deg(S)} - \sum_{\lambda < \nu < \mu} n_{\lambda, \nu}(q) b_{\nu, \underline{\mu}}(q). \quad (2.1)$$

The polynomials $n_{\lambda, \underline{\mu}}(q)$ are called the anti-spherical Kazhdan–Lusztig polynomials associated to $\lambda, \mu \in {}^P W$.

We can reformulate the above in terms of matrix multiplication. We define the matrix of light leaves polynomials

$$\Delta^{(W, P)} := (\Delta_{\lambda, \underline{\mu}}(q))_{\lambda, \mu \in {}^P W} \quad \Delta_{\lambda, \underline{\mu}}(q) = \sum_{S \in \text{Path}(\lambda, \underline{\mu})} q^{\deg(S)}$$

to be the (square) lower uni-triangular matrix whose entries record the degrees of paths in $\text{Path}(\lambda, \underline{\mu})$. This matrix can be factorised *uniquely* as a product $\Delta^{(W, P)} = N^{(W, P)} \times B^{(W, P)}$ of lower uni-triangular matrices

$$N^{(W, P)} := (n_{\lambda, \nu}(q))_{\lambda, \mu \in {}^P W} \quad B^{(W, P)} := (b_{\nu, \underline{\mu}}(q))_{\nu, \mu \in {}^P W}$$

such that $n_{\lambda, \nu}(q) \in q\mathbb{Z}[q]$ for $\lambda \neq \nu$ and $b_{\nu, \underline{\mu}}(q) \in \mathbb{Z}[q + q^{-1}]$.

Example 2.2. The matrix Δ in type $(A_3, A_1 \times A_1)$ is depicted below.

Δ	$s_2 s_1 s_3 s_2$	$s_2 s_1 s_3$	$s_2 s_1$	$s_2 s_3$	s_2	\emptyset
$s_2 s_1 s_3 s_2$	1	\cdot	\cdot	\cdot	\cdot	\cdot
$s_2 s_1 s_3$	q	1	\cdot	\cdot	\cdot	\cdot
$s_2 s_1$	\cdot	q	1	\cdot	\cdot	\cdot
$s_2 s_3$	\cdot	q	\cdot	1	\cdot	\cdot
s_2	q	q^2	q	q	1	\cdot
\emptyset	q^2	\cdot	\cdot	\cdot	q	1

The factorisation of this matrix is trivial, with $N = \Delta$ and $B = \text{Id}_{6 \times 6}$ the identity matrix.

Example 2.3. For (C_3, A_2) the factorisation $\Delta = N \times B$ is given below. The rows of the matrix can be taken to be ordered with respect to any total refinement of the Bruhat order (there are two such total orders), see Figure 9 for the corresponding graph $\mathcal{G}_{(W,P)}$.

$$\begin{pmatrix} 1 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ q & 1 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ 1 & q & 1 & \cdot & \cdot & \cdot & \cdot & \cdot \\ q & \cdot & q & 1 & \cdot & \cdot & \cdot & \cdot \\ q & \cdot & q & \cdot & 1 & \cdot & \cdot & \cdot \\ q^2 & q & q^2 & q & q & 1 & \cdot & \cdot \\ q & q^2 & \cdot & \cdot & 1 & q & 1 & \cdot \\ q^2 & \cdot & \cdot & \cdot & q & \cdot & q & 1 \end{pmatrix} = \begin{pmatrix} 1 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ q & 1 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & q & 1 & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & q & 1 & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & q & \cdot & 1 & \cdot & \cdot & \cdot \\ \cdot & q & q^2 & q & q & 1 & \cdot & \cdot \\ q & q^2 & \cdot & \cdot & \cdot & q & 1 & \cdot \\ q^2 & \cdot & \cdot & \cdot & \cdot & \cdot & q & 1 \end{pmatrix} \begin{pmatrix} 1 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & 1 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ 1 & \cdot & 1 & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & 1 & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & 1 & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & 1 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 1 & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 1 \end{pmatrix}$$

Example 2.4. The matrices Δ for exceptional types of Hermitian symmetric pairs are given in Section 4. Again, we have that B is the identity matrix in these cases.

The paths $\mathbf{s} \in \text{Path}(\lambda, \underline{\mu})$ enumerate a “light leaf basis” of the Hecke category. We refer to [BDHN] for an algorithmic construction of these basis elements in the language of this paper, see also [LW]. We are now able to restate Libedinsky–Williamson’s goal (from the introduction) more precisely using the language of paths. They ask if it is possible to produce (via a closed combinatorial algorithm) a set $\text{NPath}(\lambda, \underline{\nu}) \subseteq \text{Path}(\lambda, \underline{\nu})$ and a canonical basis for a space

$$\bigoplus_{\mathbf{s} \in \text{NPath}(\lambda, \underline{\nu})} \mathbb{R}\mathbf{s} \xrightarrow{\sim} \mathbb{N}_{\lambda, \underline{\nu}}$$

so that, upon taking graded dimensions, we get

$$n_{\lambda, \underline{\nu}} = \sum_{\mathbf{s} \in \text{NPath}(\lambda, \underline{\nu})} q^{\deg(\mathbf{s})}.$$

In this paper, we answer this question for (W, P) a Hermitian symmetric pair (see Figure 1 for a list of such pairs). In fact, we go further and produce closed combinatorial descriptions of canonical bases for spaces

$$\bigoplus_{\mathbf{s} \in \text{NPath}(\lambda, \underline{\nu})} \mathbb{R}\mathbf{s} \xrightarrow{\sim} \mathbb{N}_{\lambda, \underline{\nu}} \qquad \bigoplus_{\mathbf{s} \in \text{BPath}(\lambda, \underline{\nu})} \mathbb{R}\mathbf{s} \xrightarrow{\sim} \mathbb{B}_{\lambda, \underline{\nu}}$$

so that, upon taking graded dimensions, we get

$$n_{\lambda, \underline{\nu}} = \sum_{\mathbf{s} \in \text{NPath}(\lambda, \underline{\nu})} q^{\deg(\mathbf{s})} \qquad b_{\nu, \underline{\mu}} = \sum_{\mathbf{s} \in \text{BPath}(\nu, \underline{\mu})} q^{\deg(\mathbf{s})}$$

for subsets $\text{NPath}(\lambda, \underline{\nu}) \subseteq \text{Path}(\lambda, \underline{\nu})$ and $\text{BPath}(\nu, \underline{\mu}) \subseteq \text{Path}(\nu, \underline{\mu})$.

We note further that, for Hermitian symmetric pairs, the subsets $\text{NPath}(\lambda, \underline{\nu})$ and $\text{BPath}(\nu, \underline{\mu})$ are independent of the choice of reduced expressions for μ and ν . This follows from the fact that the elements of ${}^P W$ are fully commutative (see Section 3.1 below).

3. THE ORIENTED TEMPERLEY–LIEB ALGEBRAS

We will assume from now on that (W, P) is a Hermitian symmetric pair, that is it is one of the following infinite families $(A_n, A_{k-1} \times A_{n-k})$ with $1 \leq k \leq n$, (D_n, A_{n-1}) , (D_n, D_{n-1}) , (B_n, B_{n-1}) , (C_n, A_{n-1}) for $n \geq 2$ or is one of the exceptional cases (E_6, D_5) , (E_7, E_6) . The corresponding Coxeter graphs of these pairs are recorded in Figure 1.

3.1. The oriented Temperley–Lieb algebras and strong full commutativity. For w, w' two expressions in the generators $s_i \in S_W$, we say that w' is a subword of w if there are expressions u and v such that $w = uw'v$. One of the crucial property of Hermitian symmetric pairs for this paper is that the elements of ${}^P W$ are fully commutative (as defined by Stembridge [Ste96, Introduction]). We recall that an element $w \in W$ is called fully commutative if and only if any two reduced expression of w are related by applying only the commutation relations in W . Equivalently, no reduced expressions of w contains $s_i s_j s_i$ as a subword for $m_{i,j} = 3$ or $s_i s_j s_i s_j$ as a subword for $m_{i,j} = 4$. In fact, the elements of ${}^P W$ satisfy the following slightly stronger property.

Definition 3.1. *We say that an element $w \in W$ is strongly fully commutative if no reduced expression of w contains $s_i s_j s_i$ as a subword for any $s_i, s_j \in S_W$ with either $m_{i,j} = 3$ or $m_{i,j} = 4$ when α_i is a short root. We denote by W_{sfc} the set of all strongly fully commutative elements of W .*

Lemma 3.2. *Let (W, P) be a Hermitian symmetric pair. Then every element of ${}^P W$ is strongly fully commutative.*

Proof. This is well-known and can be seen for example from the explicit description of the elements of ${}^P W$ in terms of tilings given in [EHP14, Appendix: Diagrams of Hermitian types]. \square

Definition 3.3. *Let (W, P) be a Hermitian symmetric pair. The oriented Temperley–Lieb algebra of type (W, P) , $\text{TL}_{(W,P)}(q)$, is defined to be the unital associative $\mathbb{Z}[q, q^{-1}]$ -algebra generated by elements*

$$\{1_\lambda \mid \lambda \in {}^P W\} \cup \{E_i \mid s_i \in S_W\}$$

subject to the following relations. The idempotent relations,

$$\begin{aligned} 1 &= \sum_{\lambda \in {}^P W} 1_\lambda, & 1_\lambda E_i 1_\lambda &= 0 \text{ if } \lambda s_i \notin {}^P W, \\ 1_\lambda 1_\mu &= \delta_{\lambda, \mu} 1_\lambda, & 1_\lambda E_i 1_\mu &= 0 \text{ if } \mu \notin \{\lambda, \lambda s_i\} \end{aligned} \quad (3.1)$$

for all $\lambda, \mu \in {}^P W$. For all $s_i \in S_W$, any $\lambda \in {}^P W$ with $\lambda s_i \in {}^P W$ and $\mu, \nu \in \{\lambda, \lambda s_i\}$ we have

$$1_\mu E_i 1_\lambda E_i 1_\nu = q^{\ell(\lambda s_i) - \ell(\lambda)} 1_\mu E_i 1_\nu. \quad (3.2)$$

If $m_{i,j} = 2$ or 3, then

$$E_i E_j = E_j E_i, \quad E_i E_j E_i = E_i, \quad (3.3)$$

respectively. If $m_{i,j} = 4$ and α_i is a short root then we have that

$$1_\mu E_i 1_\lambda E_j 1_\lambda E_i 1_\nu = 1_\mu E_i 1_\nu, \quad (3.4)$$

for any $\lambda \in {}^P W$ with $\lambda s_i, \lambda s_j \in {}^P W$ and $\mu, \nu \in \{\lambda, \lambda s_i\}$.

Remark 3.4. *It follows from the relations (3.1) that $\text{TL}_{(W,P)}(q)$ is generated by the elements 1_λ for $\lambda \in {}^P W$ and $1_\lambda E_i 1_\mu$ for all $\lambda \in {}^P W$ with $\lambda s_i \in {}^P W$ and $\mu \in \{\lambda, \lambda s_i\}$.*

Remark 3.5. *Note that for $m_{i,j} = 2$ we have $1_\lambda E_i 1_\mu E_j 1_\nu \neq 0$ if and only if $\lambda, \lambda s_i, \lambda s_j \in {}^P W$ and either $\mu = \lambda$ and $\nu \in \{\lambda, \lambda s_j\}$, or $\mu = \lambda s_i$ and $\nu \in \{\lambda s_i, \lambda s_i s_j\}$. So we have that the first relation in (3.3) is equivalent to*

$$1_\lambda E_i 1_\lambda E_j 1_\nu = 1_\lambda E_j 1_\nu E_i 1_\nu, \quad \text{and} \quad 1_\lambda E_i 1_{\lambda s_i} E_j 1_\nu = 1_\lambda E_j 1_{\nu s_i} E_i 1_\nu$$

for $\nu \in \{\lambda, \lambda s_j\}$ and $\nu \in \{\lambda s_i, \lambda s_i s_j\}$ respectively, and every $\lambda \in {}^P W$.

Remark 3.6. Note that for $m_{i,j} = 3$, using Lemma 3.2, we have $E_i 1_\mu E_j 1_\nu E_i \neq 0$ implies $\nu = \mu$ and $\mu s_i, \mu s_j \in {}^P W$. Now the second relation in (3.3) is equivalent to

$$1_\lambda E_i E_j E_i 1_\eta = 1_\lambda E_i 1_\eta$$

for all $\lambda, \eta \in {}^P W$ with $\lambda s_i, \eta s_i \in {}^P W$. Note that for each such λ , using Lemma 3.2, we have either $\lambda s_j \in {}^P W$ or $\lambda s_i s_j \in {}^P W$ but not both. Thus the second relation in (3.3) is also equivalent to

$$1_\lambda E_i 1_\lambda E_j 1_\lambda E_i 1_\eta = 1_\lambda E_i 1_\eta$$

for all $\lambda \in {}^P W$ with $\lambda s_i, \lambda s_j \in {}^P W$ and $\eta = \{\lambda, \lambda s_i\}$, and

$$1_\lambda E_i 1_{\lambda s_i} E_j 1_{\lambda s_i} E_i 1_\eta = 1_\lambda E_i 1_\eta$$

for all $\lambda \in {}^P W$ with $\lambda s_i, \lambda s_i s_j \in {}^P W$ and $\eta = \{\lambda, \lambda s_i\}$.

Remark 3.7. Note that for $m_{i,j} = 4$ with α_i being a short root, we have

$$1_\lambda E_i E_j E_i 1_\mu = 1_\lambda E_i 1_\lambda E_j 1_\lambda E_i 1_\mu + 1_\lambda E_i 1_{\lambda s_i} E_j 1_{\lambda s_i} E_i 1_\mu = 2(1_\lambda E_i 1_\mu)$$

for $\lambda, \lambda s_i, \lambda s_j, \lambda s_i s_j \in {}^P W$ and $\mu \in \{\lambda, \lambda s_i\}$. This implies that $E_i E_j E_i = 2E_i$ and so $E_i E_j E_i E_j = 2E_i E_j$ and $E_j E_i E_j E_i = 2E_j E_i$.

3.2. Path basis and the anti-spherical module. For any path

$$\mathbb{T} : \lambda_1 \xrightarrow{i_1} \lambda_2 \xrightarrow{i_2} \lambda_3 \xrightarrow{i_3} \dots \xrightarrow{i_{k-1}} \lambda_k$$

on $\widehat{\mathcal{G}}_{(W,P)}$, we write

$$E_{\mathbb{T}} := 1_{\lambda_1} E_{i_1} 1_{\lambda_2} E_{i_2} 1_{\lambda_3} \dots 1_{\lambda_{k-1}} E_{i_{k-1}} 1_{\lambda_k}$$

to be the corresponding element in $\text{TL}_{(W,P)}(q)$.

Recall that we denote by W_{sfc} the set of all strongly fully commutative elements of W . We now fix one reduced expression \underline{w} for each $w \in W_{sfc}$.

Theorem 3.8. The algebra $\text{TL}_{(W,P)}(q)$ has a $\mathbb{Z}[q, q^{-1}]$ -basis given by the set

$$\{E_{\mathbb{T}} : \mathbb{T} \in \text{Path}_{(W,P)} \text{ with } \omega(\mathbb{T}) = \underline{w} \text{ for some } w \in W_{sfc}\}.$$

Proof. It is clear from relations (3.1) that

$$1_{\lambda_1} E_{i_1} 1_{\lambda_2} E_{i_2} 1_{\lambda_3} \dots 1_{\lambda_{k-1}} E_{i_{k-1}} 1_{\lambda_k} \neq 0$$

implies that

$$\mathbb{T} : \lambda_1 \xrightarrow{i_1} \lambda_2 \xrightarrow{i_2} \lambda_3 \xrightarrow{i_3} \dots \xrightarrow{i_{k-1}} \lambda_k$$

is a path on $\widehat{\mathcal{G}}_{(W,P)}$. Thus we have that $\text{TL}_{(W,P)}(q)$ is spanned by elements of the form $E_{\mathbb{T}}$ where $\mathbb{T} \in \text{Path}_{(W,P)}$. Now it follows from relations (3.2)–(3.4) that any such $E_{\mathbb{T}} = q^x E_{\mathbb{S}}$ for some $x \in \mathbb{Z}$ and some path \mathbb{S} such that $\omega(\mathbb{S}) = \underline{w}$ for some $w \in W_{sfc}$. Set $\text{Path}_{(W,P)}^{sfc}$ to be the set of all $\mathbb{T} \in \text{Path}_{(W,P)}$ such that $\omega(\mathbb{T}) = \underline{w}$ for some $w \in W_{sfc}$. It remains to show that the set $\{E_{\mathbb{T}} : \omega(\mathbb{T}) \in \text{Path}_{(W,P)}^{sfc}\}$ is linearly independent. We do this by constructing a formal $\text{TL}_{(W,P)}(q)$ -module M with $\mathbb{Z}[q, q^{-1}]$ -basis labelled by $[\mathbb{T}]$ for $\mathbb{T} \in \text{Path}_{(W,P)}^{sfc}$. To define the action on this module we will need to ‘reduce’ any path on $\widehat{\mathcal{G}}_{(W,P)}$ using the following local operations.

◦ If $m_{ij} = 2$ then for $\nu \in \{\lambda, \lambda s_j\}$ we set

$$[\lambda \xrightarrow{i} \lambda \xrightarrow{j} \nu] \implies [\lambda \xrightarrow{j} \nu \xrightarrow{i} \lambda]$$

and for $\nu \in \{\lambda s_i, \lambda s_i s_j\}$ we set

$$[\lambda \xrightarrow{i} \lambda s_i \xrightarrow{j} \nu] \implies [\lambda \xrightarrow{j} \nu s_i \xrightarrow{i} \nu].$$

◦ If $m_{ij} = 3$ or $m_{ij} = 4$ and α_i is a short root then for $\eta \in \{\lambda, \lambda s_i\}$ we set

$$[\lambda \xrightarrow{i} \lambda \xrightarrow{j} \lambda \xrightarrow{i} \eta] \quad \Longrightarrow \quad [\lambda \xrightarrow{i} \eta]$$

and

$$[\lambda \xrightarrow{i} \lambda s_i \xrightarrow{j} \lambda s_i \xrightarrow{i} \eta] \quad \Longrightarrow \quad [\lambda \xrightarrow{i} \eta].$$

◦ For for $\mu, \nu \in \{\lambda, \lambda s_i\}$, we set

$$[\mu \xrightarrow{i} \lambda \xrightarrow{i} \nu] \quad \Longrightarrow \quad q^{\ell(\lambda s_i) - \ell(\lambda)} [\mu \xrightarrow{i} \nu].$$

(Note that these follow exactly the relations given in Definition 3.3, see also Remark 3.5, Remark 3.6 and Remark 3.7.) It is clear that starting with any path $\mathsf{T} \in \text{Path}_{(W,P)}$, applying these operations repeatedly, we obtain a uniquely defined power $q^{x(\mathsf{T})}$ and a unique path $\text{rex}(\mathsf{T}) \in \text{Path}_{(W,P)}^{sfc}$. Now, for any $\mathsf{T} \in \text{Path}_{(W,P)}^{sfc}$ with

$$\mathsf{T} : \lambda_1 \xrightarrow{i_1} \lambda_2 \xrightarrow{i_2} \lambda_3 \xrightarrow{i_3} \dots \xrightarrow{i_{k-1}} \lambda_k = \mu$$

we set

$$[\mathsf{T}]1_\nu = \delta_{\mu\nu}[\mathsf{T}] \quad \text{and} \quad [\mathsf{T}]1_\mu E_i 1_\nu = q^{x(\mathsf{T}')} [\text{rex}(\mathsf{T}')]]$$

for $\nu \in \{\mu, \mu s_i\}$ and $\mu, \mu s_i \in {}^P W$, where

$$\mathsf{T}' = \lambda_1 \xrightarrow{i_1} \lambda_2 \xrightarrow{i_2} \lambda_3 \xrightarrow{i_3} \dots \xrightarrow{i_{k-1}} \lambda_k = \mu \xrightarrow{i} \nu.$$

As the relations used to reduce the path correspond precisely to the relations defining the oriented Temperley–Lieb algebra, it is clear that this turns M into a $\text{TL}_{(W,P)}(q)$ -module.

We are now ready to prove that the set $\{E_{\mathsf{T}} : \mathsf{T} \in \text{Path}_{(W,P)}^{sfc}\}$ is linearly independent. Assume that

$$\sum_{\mathsf{T}} a_{\mathsf{T}} E_{\mathsf{T}} = 0 \quad \text{for some } a_{\mathsf{T}} \in \mathbb{Z}[q, q^{-1}].$$

We need to show that $a_{\mathsf{T}} = 0$ for all T . Fix $\lambda \in {}^P W$ and consider the trivial path $[\lambda] \in M$. Then we have

$$[\lambda] \left(\sum_{\mathsf{T}} a_{\mathsf{T}} E_{\mathsf{T}} \right) = \sum_{\substack{\nu \in {}^P W \\ \mathsf{T} \in \text{Path}(\lambda \rightarrow \nu)}} a_{\mathsf{T}} [\mathsf{T}] = 0.$$

As $\{[\mathsf{T}] : \mathsf{T} \in \text{Path}(\lambda \rightarrow \nu), \nu \in {}^P W\}$ is linearly independent in M , we deduce that $a_{\mathsf{T}} = 0$ for all $\mathsf{T} \in \text{Path}(\lambda \rightarrow \nu)$, all $\nu \in {}^P W$. But this holds for all $\lambda \in {}^P W$ so we are done. \square

Definition 3.9. We define the anti-spherical $\text{TL}_{(W,P)}(q)$ -module to be the right module $1_{\emptyset} \text{TL}_{(W,P)}(q)$.

Corollary 3.10. The anti-spherical module $1_{\emptyset} \text{TL}_{(W,P)}(q)$ has a $\mathbb{Z}[q, q^{-1}]$ -basis given by

$$\{E_{\mathsf{T}} \mid \mathsf{T} \in \text{Path}(\lambda, \underline{\mu}) : \lambda, \mu \in {}^P W\}.$$

Proof. By Theorem 3.8 we have that the anti-spherical module has a basis given by all E_{T} where T is a path on $\widehat{\mathcal{G}}_{(W,P)}$ starting at \emptyset with $\omega(\mathsf{T}) = \underline{w}$ some strongly fully commutative element $w \in W$. Note that \underline{w} must start with the unique $s \notin S_P$, as T starts at \emptyset . Moreover, as w is fully commutative, any other reduced expression \underline{w}' for w is obtained from \underline{w} by applying only the commutation relations and so \underline{w}' is also the weight of a path on $\widehat{\mathcal{G}}_{(W,P)}$ starting at \emptyset . In particular, \underline{w}' also starts with the unique $s \notin S_P$. This implies that $w = \mu \in {}^P W$. \square

3.3. Grading. We can view $\mathrm{TL}_{(W,P)}(q)$ as a \mathbb{Z} -algebra in the usual way, by considering q and q^{-1} as additional central generators. The next proposition shows that, as such, $\mathrm{TL}_{(W,P)}(q)$ is a \mathbb{Z} -graded algebra.

Proposition 3.11. *Set $\deg(\mathbf{1}_\lambda) = 0$ for all $\lambda \in {}^P W$,*

$$\deg(\mathbf{1}_\lambda E_i \mathbf{1}_{\lambda s_i}) = 0, \quad \deg(\mathbf{1}_\lambda E_i \mathbf{1}_\lambda) = \begin{cases} 1 & \text{if } \lambda s_i > \lambda \\ -1 & \text{if } \lambda s_i < \lambda \end{cases}$$

for all $\lambda \in {}^P W$, $s_i \in S_W$ with $\lambda s_i \in {}^P W$, $\deg(q) = 1$ and $\deg(q^{-1}) = -1$.

This defines a \mathbb{Z} -grading on the \mathbb{Z} -algebra $\mathrm{TL}_{(W,P)}(q)$. In particular, we have $\deg(E_\top) = \deg(\top)$ for all $\top \in \mathrm{Path}_{(W,P)}$.

Proof. We need to check that the defining relations (3.1)–(3.4) are (equivalent to) homogeneous relations. For relations (3.1), there is nothing to prove. We now consider relation (3.2). We need to show that for any $\mu, \nu \in \{\lambda, \lambda s_i\}$ we have

$$\deg(\mathbf{1}_\mu E_i \mathbf{1}_\lambda E_i \mathbf{1}_\nu) = (\ell(\lambda s_i) - \ell(\lambda)) + \deg(\mathbf{1}_\mu E_i \mathbf{1}_\nu).$$

If $\mu = \nu = \lambda$ we have

$$\deg(\mathbf{1}_\lambda E_i \mathbf{1}_\lambda E_i \mathbf{1}_\lambda) = 2(\ell(\lambda s_i) - \ell(\lambda)), \quad \text{and} \quad \deg(\mathbf{1}_\lambda E_i \mathbf{1}_\lambda) = \ell(\lambda s_i) - \ell(\lambda)$$

as required. If $\mu = \nu = \lambda s_i$ we have

$$\deg(\mathbf{1}_{\lambda s_i} E_i \mathbf{1}_\lambda E_i \mathbf{1}_{\lambda s_i}) = 0, \quad \text{and} \quad \deg(\mathbf{1}_{\lambda s_i} E_i \mathbf{1}_{\lambda s_i}) = \ell(\lambda) - \ell(\lambda s_i)$$

as required. Finally if $\mu \neq \nu$ then we have

$$\deg(\mathbf{1}_\mu E_i \mathbf{1}_\lambda E_i \mathbf{1}_\nu) = \ell(\lambda s_i) - \ell(\lambda), \quad \text{and} \quad \deg(\mathbf{1}_\mu E_i \mathbf{1}_\nu) = 0$$

as required. Next consider the leftmost relation in (3.3). By Remark 3.5, we need to show that the relations

$$\begin{aligned} \mathbf{1}_\lambda E_i \mathbf{1}_\lambda E_j \mathbf{1}_\nu &= \mathbf{1}_\lambda E_j \mathbf{1}_\nu E_i \mathbf{1}_\lambda & \text{for } \nu \in \{\lambda, \lambda s_j\}, \quad \text{and} \\ \mathbf{1}_\lambda E_i \mathbf{1}_{\lambda s_i} E_j \mathbf{1}_\nu &= \mathbf{1}_\lambda E_j \mathbf{1}_{\nu s_i} E_i \mathbf{1}_\nu & \text{for } \nu \in \{\lambda s_i, \lambda s_i s_j\} \end{aligned}$$

are homogeneous. The former is trivial for $\nu = \lambda$ and follows from the fact that $\lambda s_j s_i > \lambda s_j$ if and only if $\lambda s_i > \lambda$ for $\nu = \lambda s_j$. The latter is trivial for $\nu = \lambda s_i s_j$ and follows from the fact that $\lambda s_i s_j > \lambda s_i$ if and only if $\lambda s_j > \lambda$ for $\nu = \lambda s_i$.

We now consider the rightmost relation in (3.3). By Remark 3.6, this relation is equivalent to

$$\mathbf{1}_\lambda E_i \mathbf{1}_\lambda E_j \mathbf{1}_\lambda E_i \mathbf{1}_\eta = \mathbf{1}_\lambda E_i \mathbf{1}_\eta$$

for all $\lambda \in {}^P W$ with $\lambda s_i, \lambda s_j \in {}^P W$ and $\eta = \{\lambda, \lambda s_i\}$, and

$$\mathbf{1}_\lambda E_i \mathbf{1}_{\lambda s_i} E_j \mathbf{1}_{\lambda s_i} E_i \mathbf{1}_\eta = \mathbf{1}_\lambda E_i \mathbf{1}_\eta$$

for all $\lambda \in {}^P W$ with $\lambda s_i, \lambda s_i s_j \in {}^P W$ and $\eta = \{\lambda, \lambda s_i\}$. The former is homogeneous as $\lambda s_i > \lambda$ if and only if $\lambda s_j < \lambda$ and so

$$\deg(\mathbf{1}_\lambda E_i \mathbf{1}_\lambda E_j \mathbf{1}_\lambda) = 0.$$

To see that the latter is also homogeneous, observe that $\lambda s_i s_j > \lambda s_i$ if and only if $\lambda s_i > \lambda$ and so

$$\deg(\mathbf{1}_{\lambda s_i} E_j \mathbf{1}_{\lambda s_i}) = \deg(\mathbf{1}_\lambda E_i \mathbf{1}_\lambda) \quad \text{and} \quad \deg(\mathbf{1}_{\lambda s_i} E_j \mathbf{1}_{\lambda s_i} E_i \mathbf{1}_{\lambda s_i}) = 0.$$

Finally, consider relation (3.4). For $\mu = \lambda$ and $\nu \in \{\lambda, \lambda s_i\}$ we have to show that

$$\mathbf{1}_\lambda E_i \mathbf{1}_\lambda E_j \mathbf{1}_\lambda E_i \mathbf{1}_\nu = \mathbf{1}_\lambda E_i \mathbf{1}_\nu$$

is homogeneous, and for $\mu = \lambda s_i$ and $\nu \in \{\lambda, \lambda s_i\}$ we need to show that

$$\mathbf{1}_{\lambda s_i} E_i \mathbf{1}_\lambda E_j \mathbf{1}_\lambda E_i \mathbf{1}_\nu = \mathbf{1}_{\lambda s_i} E_i \mathbf{1}_\nu$$

is homogeneous. These look exactly the same as the relations for $m_{i,j} = 3$ above (replacing λs_i with λ for the second equation) and the same arguments apply here. \square

We immediately obtain the following result.

Corollary 3.12. (1) *The anti-spherical module $1_{\emptyset} \text{TL}_{(W,P)}(q)$ is a graded $\text{TL}_{(W,P)}(q)$ -module with homogeneous basis $\{E_{\mathbb{T}} : \mathbb{T} \in \text{Path}(\lambda, \underline{\mu}), \lambda, \mu \in {}^P W\}$ satisfying*

$$\deg E_{\mathbb{T}} = \deg \mathbb{T}.$$

(2) *The light leaves matrix can be computed follows. For any $\lambda, \mu \in {}^P W$ we have*

$$\Delta_{\lambda, \mu} = \begin{cases} \sum_{\mathbb{T} \in \text{Path}(\lambda, \underline{\mu})} q^{\deg(E_{\mathbb{T}})} & \text{if } \text{Path}(\lambda, \underline{\mu}) \neq \emptyset \\ 0 & \text{otherwise.} \end{cases}$$

Thus the anti-spherical module for the oriented Temperley–Lieb algebra gives us a model to study the light leaves matrix and its factorisation for all Hermitian symmetric pairs. This will be done in details in each type (W, P) in the next few sections but first we take a short detour to relate our oriented Temperley–Lieb algebras to the generalised Temperley–Lieb algebras associated to W .

3.4. Relationship with Fan–Graham’s Temperley–Lieb algebras. We now relate our (W, P) -Temperley–Lieb algebra to the generalised Temperley–Lieb algebra associated to W introduced by Fan (in the simply-laced type) and Graham (in the non-simply laced type).

Definition 3.13. *The generalised Temperley–Lieb algebra $\text{TL}_W(q)$ is defined as the $\mathbb{Z}[q, q^{-1}]$ -algebra generated by*

$$\{U_i \mid s_i \in S_W\}$$

subject to the following relations: For all $s_i \in S_W$ we have

$$U_i^2 = (q + q^{-1})U_i. \quad (3.5)$$

Furthermore, we have that

$$U_i U_j = U_j U_i, \quad U_i U_j U_i = U_i, \quad U_i U_j U_i U_j = 2U_i U_j \quad (3.6)$$

for $m_{i,j} = 2, 3$ or 4 , respectively.

Proposition 3.14. *There is a $\mathbb{Z}[q, q^{-1}]$ -algebra homomorphism from $\text{TL}_W(q)$ to $\text{TL}_{(W,P)}(q)$ defined by $U_i \mapsto E_i$ for all $s_i \in S_W$.*

Proof. We need to check that the E_i ’s satisfy the relations (3.5) and (3.6). The leftmost two relations of (3.6) are given by (3.3). The rightmost relation in (3.6) follows from Remark 3.7. It remains to show (3.5). We have

$$E_i = \sum_{\lambda} 1_{\lambda} E_i (1_{\lambda} + 1_{\lambda s_i})$$

where the sum is taken over all $\lambda \in {}^P W$ such that $\lambda s_i \in {}^P W$. Now we have

$$E_i^2 = \sum_{\lambda} 1_{\lambda} E_i (1_{\lambda} + 1_{\lambda s_i}) E_i (1_{\lambda} + 1_{\lambda s_i}) = (q + q^{-1}) \sum_{\lambda} 1_{\lambda} E_i (1_{\lambda} + 1_{\lambda s_i}) = (q + q^{-1}) E_i$$

as required. Here the first equality follows by (3.1) and the second by (3.2), the third is trivial. \square

Remark 3.15. *Note that this homomorphism is not injective in general. To see this, take for example W of type A_3 and P of type A_2 , then ${}^P W = \{1, s_3, s_3 s_2, s_3 s_2 s_1\}$. Then we claim that $E_1 E_3 = 0$. To see this, note that $E_1 1_{\lambda} \neq 0$ implies that $\lambda = s_3 s_2$ or $s_3 s_2 s_1$ but $1_{\lambda} E_3 \neq 0$ implies $\lambda = 1$ or s_3 . However, $U_1 U_3 \neq 0$ in $\text{TL}_W(q)$.*

4. LIGHT LEAVES MATRIX FACTORISATION FOR THE TRIVIAL AND EXCEPTIONAL TYPES

Corollary 3.12 provides a way of studying the light leaves matrix using the oriented Temperley–Lieb algebra and its anti-spherical module for all Hermitian symmetric pairs (W, P) . In Sections 5 to 8, we will construct a diagrammatic version of the oriented Temperley–Lieb algebras in types $(A_n, A_k \times A_{n-k-1})$, (D_n, A_{n-1}) and (C_n, A_{n-1}) which will provide closed combinatorial formulas for the light leaves matrix and its factorisation. This could also be done in types (B_n, B_{n-1}) and (D_n, D_{n-1}) but the extra effort is unwarranted as the light leaves matrices and their factorisation can be easily described without them. This will be done in this section, together with the exceptional types (E_6, D_5) and (E_7, E_6) , which are best tackled with a computer (although (E_6, E_5) is manageable by hand as well). This provides a proof of Theorem A in these trivial and exceptional types.

4.1. Exceptional types. We first consider the exceptional Hermitian symmetric pairs. One can calculate the Δ matrix for type (E_6, D_5) easily by hand. For type (E_7, E_6) this is a much larger calculation, but can be readily done using the Coxeter 3 package in SAGE which wraps Folklo Ducloux’s original work in C^{++} . The matrices Δ are recorded in Figures 5 and 6. In both cases, note that all off-diagonal entries belong to $q\mathbb{N}_0[q]$ and so $\Delta = N$ and $B = \text{Id}$. This gives a proof of Theorem A in these cases by setting every basis element $E_\tau \in 1_\emptyset \text{TL}_{(W,P)}(q)$ to be standard.

4.2. Type (D_n, D_{n-1}) . In this case, there is precisely one element in ${}^P W$ of each length $0 \leq \ell \leq 2(n-1)$, $\ell \neq n-1$ and precisely two elements of length $n-1$. Ordering the rows and columns of the matrix Δ by decreasing length we have

$$\Delta = \begin{bmatrix} 1 & \cdot & \cdots & \cdot & \cdot & \cdot & \cdot & \cdot & \cdots & \cdot & \cdot \\ q & 1 & \cdots & \cdot & \cdot & \cdot & \cdot & \cdot & \cdots & \cdot & \cdot \\ \vdots & & \ddots & & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \cdot & \cdot & \cdots & 1 & \cdot & \cdot & \cdot & \cdot & \cdots & \cdot & \cdot \\ \cdot & \cdot & \cdots & q & 1 & \cdot & \cdot & \cdot & \cdots & \cdot & \cdot \\ \hline \cdot & \cdot & \cdots & \cdot & q & 1 & \cdot & \cdot & \cdots & \cdot & \cdot \\ \cdot & \cdot & \cdots & \cdot & q & \cdot & 1 & \cdot & \cdots & \cdot & \cdot \\ \hline \cdot & \cdot & \cdots & q & q^2 & q & q & 1 & \cdot & \cdots & \cdot \\ \cdot & \cdot & \cdots & q^2 & \cdot & \cdot & \cdot & q & 1 & \cdots & \cdot \\ \vdots & & \ddots & & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ q & q^2 & \cdots & \cdot & \cdot & \cdot & \cdot & \cdot & \cdots & 1 & \cdot \\ q^2 & \cdot & \cdots & \cdot & \cdot & \cdot & \cdot & \cdot & \cdots & q & 1 \end{bmatrix}$$

that is, the top left and bottom right $(n-1) \times (n-1)$ -matrices have non-zero entries on the diagonal and sub-diagonal only; the bottom left $(n-1) \times (n-1)$ -matrix has non-zero entries on the anti-diagonal and sup-anti-diagonal only. In this case the matrix factorisation is trivial, with $B = \text{Id}$ and $N = \Delta$. This gives a proof of Theorem A in type (D_n, D_{n-1}) by setting every basis elements in the anti-spherical module for $\text{TL}_{(D_n, D_{n-1})}(q)$ to be standard.

4.3. Type (B_n, B_{n-1}) . In this type there is precisely one element $\lambda \in {}^P W$ of each length. Ordering the rows and columns of the matrix Δ by decreasing length it is easy to see that

$$\Delta = \begin{bmatrix} 1 & \cdot & \cdots & \cdot & \cdot & \cdot & \cdot & \cdot & \cdots & \cdot & \cdot \\ q & 1 & \cdots & \cdot & \cdot & \cdot & \cdot & \cdot & \cdots & \cdot & \cdot \\ \vdots & & \ddots & & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \cdot & \cdot & \cdots & 1 & \cdot & \cdot & \cdot & \cdot & \cdots & \cdot & \cdot \\ \cdot & \cdot & \cdots & q & 1 & \cdot & \cdot & \cdot & \cdots & \cdot & \cdot \\ \hline \cdot & \cdot & \cdots & 1 & q & 1 & \cdot & \cdots & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdots & q & \cdot & q & 1 & \cdots & \cdot & \cdot & \cdot \\ \vdots & & \ddots & & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 1 & q & \cdots & \cdot & \cdot & \cdot & \cdot & \cdots & 1 & \cdot & \cdot \\ q & \cdot & \cdots & \cdot & \cdot & \cdot & \cdot & \cdots & q & 1 & \cdot \end{bmatrix}$$




FIGURE 5. The light leaves matrix $\Delta_{\lambda,\mu}$ of type (E_6, D_5) . For purposes of space, we record q^i simply as i , and we record each zero polynomials as a dot. For example, the matrix is uni-triangular with diagonal entries $q^0 = 1$. The rows and columns are ordered by a total refinement of the Bruhat order in which we prefer to add the reflection with largest possible subscript. More specifically, the order is as follows $\emptyset, s_1, s_1s_2, s_1s_2s_3, s_1s_2s_3s_6, s_1s_2s_3s_4, s_1s_2s_3s_6s_4, s_1s_2s_3s_4s_5, s_1s_2s_3s_6s_4s_5, s_1s_2s_3s_6s_4s_3, s_1s_2s_3s_6s_4s_5s_3, s_1s_2s_3s_6s_4s_5s_3s_4, s_1s_2s_3s_6s_4s_3s_2, s_1s_2s_3s_6s_4s_5s_3s_2, s_1s_2s_3s_6s_4s_5s_3s_4s_2, s_1s_2s_3s_6s_4s_5s_3s_4s_2s_3, s_1s_2s_3s_6s_4s_5s_3s_4s_2s_3s_6, s_1s_2s_3s_6s_4s_3s_2s_1, s_1s_2s_3s_6s_4s_5s_3s_2s_1, s_1s_2s_3s_6s_4s_5s_3s_4s_2s_1, s_1s_2s_3s_6s_4s_5s_3s_4s_2s_3s_1, s_1s_2s_3s_6s_4s_5s_3s_4s_2s_3s_6s_1, s_1s_2s_3s_6s_4s_5s_3s_4s_2s_3s_1s_2, s_1s_2s_3s_6s_4s_5s_3s_4s_2s_3s_6s_1s_2, s_1s_2s_3s_6s_4s_5s_3s_4s_2s_3s_6s_1s_2s_3, s_1s_2s_3s_6s_4s_5s_3s_4s_2s_3s_6s_1s_2s_3s_4, s_1s_2s_3s_6s_4s_5s_3s_4s_2s_3s_6s_1s_2s_3s_4s_5$.

that is, the top left and bottom right $n \times n$ -matrices have non-zero entries on the diagonal and sub-diagonal only; the bottom left $n \times n$ -matrix has non-zero entries on the anti-diagonal and sub-anti-diagonal only. We then immediately deduce that the matrix factorisation is as follows

$$\left[\begin{array}{c|ccc} 1 & \dots & & \\ q & 1 & \dots & \\ \vdots & & \ddots & \\ & & & 1 \\ & & & q & 1 \end{array} \right] \left[\begin{array}{c|ccc} 1 & \dots & & \\ & 1 & \dots & \\ \vdots & & \ddots & \\ & & & 1 \\ & & & 1 \end{array} \right] \left[\begin{array}{c|ccc} 1 & \dots & & \\ & 1 & \dots & \\ \vdots & & \ddots & \\ & & & 1 \\ & & & 1 \end{array} \right] \left[\begin{array}{c|ccc} & & & 1 \\ & & & q & 1 \\ & & & & \\ \vdots & & \ddots & & \\ & & & & 1 \\ & & & & q & 1 \end{array} \right] \left[\begin{array}{c|ccc} 1 & \dots & & \\ & 1 & \dots & \\ \vdots & & \ddots & \\ & & & 1 \\ & & & 1 \end{array} \right] \left[\begin{array}{c|ccc} & & & 1 \\ & & & q & 1 \\ & & & & \\ \vdots & & \ddots & & \\ & & & & 1 \\ & & & & q & 1 \end{array} \right] \left[\begin{array}{c|ccc} 1 & \dots & & \\ & 1 & \dots & \\ \vdots & & \ddots & \\ & & & 1 \\ & & & 1 \end{array} \right] \left[\begin{array}{c|ccc} & & & 1 \\ & & & q & 1 \\ & & & & \\ \vdots & & \ddots & & \\ & & & & 1 \\ & & & & q & 1 \end{array} \right] \left[\begin{array}{c|ccc} 1 & \dots & & \\ & 1 & \dots & \\ \vdots & & \ddots & \\ & & & 1 \\ & & & 1 \end{array} \right]$$

where we note that the matrix on the left is the same as the matrix of Kazhdan-Lusztig polynomials of type (A_{2n-1}, A_{2n-2}) , this will be explained in our companion paper [BDHN]. This gives a proof of Theorem A in type (B_n, B_{n-1}) by setting every basis element E_τ corresponding to non-zero entries in the matrix N (on the left) to be standard.




FIGURE 6. The light leaves matrix Δ of type (E_7, E_6) . For purposes of space, we record q^i simply as i , and we record each zero polynomial as a dot. For example, the matrix is untriangular with diagonal entries $q^0 = 1$. The rows and columns are ordered by a total refinement of the Bruhat order in which add the reflection with largest possible subscript.

From now, until the end of Section 8, we focus solely on the remaining cases, namely $(W, P) = (A_n, A_{k-1} \times A_{n-k}), (C_n, A_{n-1})$ and (D_n, A_{n-1}) .

5. BRUHAT GRAPHS IN CLASSICAL TYPE

We now introduce an elementary way of visualising the graphs $\mathcal{G}_{(W,P)}$ for classical Hermitian symmetric pairs $(W, P) = (A_n, A_{k-1} \times A_{n-k}), (C_n, A_{n-1})$ and (D_n, A_{n-1}) . We will use these in the next section to define a diagrammatic visualisation of the oriented Temperley–Lieb algebra in these types.

We start by recalling the description of the Coxeter groups of type A_n, C_n and D_n as groups of (signed) permutations. The Coxeter group of type A_n is the symmetric group consisting of all permutations of $\{1, 2, \dots, n + 1\}$. The simple reflections s_i , for $1 \leq i \leq n$ are given by

$$s_i = (i, i + 1).$$

The Coxeter group of type C_n is the signed permutation group, namely the group of all permutations w of $\{\pm 2, \pm 3, \dots, \pm(n+1)\}$ such that $w(-i) = -w(i)$ for all $2 \leq i \leq n+1$. The simple reflections s_i for $i = 1', 2, \dots, n$ are given by

$$s_{1'} = (2, -2) \quad \text{and} \quad s_i = (i, i+1)(-i, -(i+1)) \text{ for } 2 \leq i \leq n.$$

The Coxeter group of type D_n is the even signed permutation group, that is the subgroup of the group of all signed permutations on $\{\pm 1, \pm 2, \dots, \pm n\}$ consisting of all elements flipping an even number of signs. The simple reflections s_i for $i = 1'', 1, 2, \dots, n-1$ are given by

$$s_{1''} = (1, -2)(-1, 2) \quad \text{and} \quad s_i = (i, i+1)(-i, -(i+1)) \text{ for } 1 \leq i \leq n-1.$$

Remark 5.1. *The choice of labelling in type C_n might seem slightly unnatural at this point but (apart from giving a uniform definition of the generators s_i for $2 \leq i \leq n-1$ in all types) it is required for compatibility with the diagrammatic Temperley-Lieb algebra of type C given in Section 6.*

Given this description of the Coxeter groups as (signed) permutation groups, we have the following natural visualisation of the cosets ${}^P W$. We will represent elements of ${}^P W$ as horizontal lines with $n(+1)$ points in positions $1, 2, \dots, n, (n+1)$ labelled with the symbols $\{\wedge, \vee, \circ\}$. The generators of W will act on these as follows: For $i \geq 1$, s_i swaps the labels in positions i and $i+1$, the generator $s_{1'}$ flips (through the horizontal axis) the label in second position, and $s_{1''}$ swaps and flips (through the horizontal line) the labels in first and second positions. Now, we start by representing the identity coset, which we denote by \emptyset (for the empty word in the generators), as follows:

- If $(W, P) = (A_n, A_{k-1} \times A_{n-k})$, draw \emptyset as the horizontal line containing $n+1$ points with the first k points labelled by \wedge and the last $n-k+1$ points labelled by \vee .
- If $(W, P) = (C_n, A_{n-1})$ draw \emptyset as the horizontal line with $n+1$ points with the first point labelled by \circ and the last n points labelled by \vee .
- If $(W, P) = (D_n, A_{n-1})$ draw \emptyset as the horizontal line with n points all labelled by \vee .

The elements of ${}^P W$ are then obtained as the elements of the orbit of \emptyset under the action of W .






FIGURE 7. We depict the identity coset \emptyset along the bottom of the diagram, the coset λ along the top of the diagram, and its corresponding reduced expression $s_5 s_4 s_3 s_2 s_1 s_6 s_5 s_4 s_3 s_7 s_6 s_8 s_7$.

Moreover, we can construct the graph $\mathcal{G}_{(W,P)}$ starting from \emptyset as follows.

- Start by drawing \emptyset at the bottom.
- If applying a simple reflection does result in a new coset, then record this in the next level up in the diagram. We record the colour of the reflection as an edge relating the two points in the graph.

We repeat the above until the process terminates. This is best illustrated via the examples in Figures 8 and 9.

FIGURE 8. The graph $\mathcal{G}_{(W,P)}$ for $(W, P) = (A_4, A_1 \times A_2)$.FIGURE 9. The graphs $\mathcal{G}_{(W,P)}$ for $(W, P) = (D_4, A_3)$ and (C_3, A_2) .

The Bruhat order on ${}^P W$ can also easily be visualised in this setting, namely $\lambda < \mu$ if either λ contains strictly fewer \wedge arrows than μ or if λ has the same number of \wedge arrows as μ and μ is obtained from λ by moving \wedge arrows to the right.

Remark 5.2. *It is worth taking a moment to consider which diagrams can appear elements of ${}^P W$. It is clear that there are $\binom{n+1}{k}$ cosets for $(A_n, A_{k-1} \times A_{n-k})$ and that these are given by all possible diagrams with k \wedge -arrows and $n - k + 1$ \vee -arrows. There are 2^n cosets for (C_n, A_{n-1}) and these are given by freely choosing the \wedge versus \vee decorations on the vertices $\{2, \dots, n+1\}$ (the decoration on vertex 1 is always a \circ). There are 2^{n-1} cosets for (D_n, A_{n-1}) and these are given by freely choosing the \wedge versus \vee decorations on the vertices $\{1, 2, \dots, n\}$ subject to the condition that the total number of \wedge -arrows is even.*

For the remainder of the paper we will freely identify cosets with their diagrams without further mention.

Remark 5.3. *In type (D_n, A_{n-1}) , the description we gave of the cosets (and graph $\mathcal{G}_{(W,P)}$) uses the parabolic subgroup of type A_{n-1} as the subgroup generated by $\{s_1, s_2, \dots, s_{n-1}\}$. We could have used instead the parabolic subgroup of type A_{n-1} generated by $\{s_{1'}, s_2, \dots, s_{n-1}\}$. This gives an alternative construction of $\mathcal{G}_{(W,P)}$ in type (D_n, A_{n-1}) , starting by defining the coset \emptyset as the horizontal line with n points where the first one is labelled by \wedge and all others are labelled by \vee . So the cosets now must have an odd number of \wedge -arrows.*

6. DIAGRAMMATIC ORIENTED TEMPERLEY-LIEB ALGEBRAS IN CLASSICAL TYPE

We now introduce a visualisation of oriented Temperley-Lieb algebras for (W, P) of classical type, namely $(A_n, A_k \times A_{n-k-1})$, (D_n, A_{n-1}) and (C_n, A_{n-1}) .

6.1. Green’s diagrammatic Temperley–Lieb algebras. We first recall Green’s diagrammatic realisation of the generalised Temperley-Lieb algebras of type W (see [Gre98]).

Definition 6.1. (1) *An n -tangle is a rectangular frame with n vertices on the northern and southern boundaries which are paired-off by n non-crossing strands and a finite number of non-crossing closed loops. Strands and loops can be decorated by a finite number of beads if they are left exposed (i.e. can be deformed to touch the left boundary of the frame). We refer to a strand connecting a northern and southern vertex as a **propagating strand**. We refer to any strand connecting two northern vertices (or two southern vertices) to each other as an **arc**. Two n -tangles are equal if there exists an isotopy of the plane fixing the boundaries of the frame carrying one n -tangle to the other. We denote the set of all such n -tangles by \mathbb{DT}_n .*

- (2) *We call an n -tangle $d \in \mathbb{DT}_n$ **undecorated** if it has no beads on its strands or loops.*
- (3) *For R any commutative ring, we have that $R\mathbb{DT}_n$ has the structure of an R -algebra where the multiplication is given by the vertical concatenation of n -tangles. Specifically, for $d, d' \in \mathbb{DT}_n$, we define the product dd' simply by placing d' above d .*




FIGURE 10. Examples of undecorated tangles.




FIGURE 11. Examples of tangles. In the first and third diagrams we decorate every left-exposed strand.

We define e_i for $i \geq 1$ to be the undecorated n -tangle with a single pair of arcs connecting the i th and $(i + 1)$ th northern (respectively southern) vertices, and with $(n - 2)$ vertical strands. We set $e_{1'} = e_{1''}$ to be the n -tangle which has a single pair of arcs connecting the 1st and 2nd northern (respectively southern) vertices, both of which carry a single bead, and with $(n - 2)$ vertical undecorated strands. We use the distinct subscripts $1'$ and $1''$ to remind the reader of

the corresponding Coxeter labels in Figure 1 for what follows. Examples of these elements are depicted in Figure 12.




FIGURE 12. The n -tangle $e_{1'} = e_{1''}$, e_1 , e_2 , and e_3 for $n = 5$.

We now recall the diagrammatic visualisation of the generalised Temperley–Lieb algebras $\text{TL}_W(q)$ given in Definition 3.13. From now on, we take $R = \mathbb{Z}[q, q^{-1}]$.

Theorem 6.2 (Kauffman [Kau87]). *The Temperley–Lieb algebra $\text{TL}_{A_n}(q)$ of type A_n , is isomorphic to the subquotient of the algebra $R\text{DT}_{n+1}$ generated by e_1, \dots, e_n subject to the relation*

$$\boxed{\bigcirc} = q + q^{-1}$$

It has a basis given by the set of all undecorated $(n+1)$ -tangles with no loops, which we denote by $\text{DT}[A_n]$.

Theorem 6.3 (Green [Gre98]). *The generalised Temperley–Lieb algebra $\text{TL}_{C_n}(q)$ of type C_n is isomorphic to the subquotient of the algebra $R\text{DT}_{n+1}$ generated by $e_{1'}, e_2, \dots, e_n$ subject to the relations*

$$\boxed{\begin{array}{c} \bullet \\ | \\ \bullet \end{array}} = \boxed{\bullet} \quad \boxed{\bigcirc} = \boxed{\bigcirc} = q + q^{-1}$$

It has a basis given by the set of all $(n+1)$ -tangles with no loops and at most one decoration on each strand and satisfying one of the following (mutually exclusive) conditions.

- (1) The leftmost northern vertex is connected to the leftmost southern vertex by an undecorated strand.
- (2) The leftmost northern vertex is connected to the leftmost southern vertex by a decorated strand and there is at least one northern and one southern arc.
- (3) The strands emerging from the leftmost northern and southern vertices are distinct and both decorated.

We denote the set of all such $(n+1)$ -tangles by $\text{DT}[C_n]$.

Theorem 6.4 (Green [Gre98]). *The generalised Temperley–Lieb algebra $\text{TL}_{D_n}(q)$ of type D_n is isomorphic to the subquotient of the algebra $R\text{DT}_n$ generated by $e_{1''}, e_1, e_2, \dots, e_{n-1}$ subject to relations*

$$\boxed{\begin{array}{c} \bullet \\ | \\ \bullet \end{array}} = \boxed{|} \quad \boxed{\bigcirc} = q + q^{-1} \quad \boxed{\begin{array}{c} \bullet \\ \bigcirc \\ \bullet \end{array}} = \boxed{\begin{array}{c} \bullet \\ \bigcirc \\ \bullet \end{array}}$$

It has a basis given by the set of all n -tangles with at most one decoration on each strand or loop, and which satisfy one of the following (mutually exclusive) conditions.

- (1) It contains one decorated loop and no other loops or decorations, and there is at least one northern and one southern arc.
- (2) It contains no loops and the number of decorations is even.

We denote the set of all such n -tangles by $\text{DT}[D_n]$.

6.2. Diagrammatic oriented Temperley–Lieb algebras. Recall from Section 5 that we can represent each coset representative $\lambda \in {}^P W$ by its coset diagram, which we also denote by λ . For $W \in \{A_n, C_n, D_n\}$, we let $\mathbb{DT}[W]$ be the set of tangles defined in Theorems 6.2 to 6.4. Now starting with any tangle in $\mathbb{DT}[W]$ we will form so-called oriented tangles by considering elements of the form $\lambda d \mu$ for $\lambda, \mu \in {}^P W$ by placing λ , respectively μ on the southern, respectively northern, boundary of d . Each strand of d in $\lambda d \mu$ now connects two symbols from the set $\{\wedge, \vee, \circ\}$ from the boundaries. A strand connecting two symbols from the set $\{\wedge, \vee\}$ is said to be oriented if one arrow points into the strand and the other arrow point out of the strand. We say that a strand connecting two symbols from the set $\{\wedge, \vee\}$ is flip-oriented if either both arrows point into the strand or both arrows point out of the strand. A strand connecting \circ with one symbol from the set $\{\wedge, \vee\}$ is said to be both oriented and flip-oriented.

Definition 6.5. For $d \in \mathbb{DT}[W]$ and $\lambda, \mu \in {}^P W$ we say that $\lambda d \mu$ is an oriented tangle of type (W, P) if the following conditions holds.

- Every undecorated strand is oriented.

Moreover, if $(W, P) = (D_n, A_{n-1})$ then

- every decorated strand is flip-oriented, and
- there are no loops.

If $(W, P) = (C_n, A_{n-1})$ then

- there are no decorations on the strand connecting \circ to \circ .

We denote by $\mathbb{ODT}[W, P]$ the set of all oriented tangles of type (W, P) and refer to these as oriented Temperley–Lieb diagrams of type (W, P) .

Examples of oriented tangles are given in Figures 13 and 14.




FIGURE 13. Oriented tangles of type $(A_8, A_4 \times A_3)$ obtained from the diagrams of Figure 10.

Remark 6.6. Note that, by definition, we have that $\lambda e_i \mu \in \mathbb{ODT}[W, P]$ if and only if $\lambda \xrightarrow{i} \mu$ is an edge in the graph $\widehat{\mathcal{G}}_{(W, P)}$.

The following proposition follows directly from Remark 5.2.

Proposition 6.7. Let $d \in \mathbb{DT}[W]$. Then there exists $\lambda, \mu \in {}^P W$ such that $\lambda d \mu \in \mathbb{ODT}[W, P]$ if and only if

- $(W, P) = (A_n, A_k \times A_{n-k-1})$ and d has at most $\min\{k+1, n-k\}$ northern/southern arcs.
- $(W, P) = (C_n, A_{n-1})$ and d contains no decoration on a strand connecting \circ to \circ (i.e. d satisfies condition (1) or (3) from Theorem 6.3).
- $(W, P) = (D_n, A_{n-1})$ and d contains no loops (i.e. d satisfies condition (2) from Theorem 6.4).


We denote the set of all tangles described above by $\mathbb{DT}[W, P]$.

Definition 6.8. The diagrammatic oriented Temperley–Lieb algebra of type (W, P) , denoted by $\text{TL}_{(W, P)}^{\wedge, \vee}(q)$, is the $\mathbb{Z}[q, q^{-1}]$ -algebra with basis $\mathbb{ODT}[W, P]$ and multiplication defined as follows. For $\lambda d \mu, \lambda' d' \mu' \in \mathbb{ODT}[W, P]$ we have

$$(\lambda d \mu)(\lambda' d' \mu') = 0 \quad \text{if } \mu \neq \lambda' \quad (6.1)$$

and vertical concatenation, denoted by $\lambda d \mu d' \mu'$, if $\mu = \lambda'$, subject to the following relations.

- (1) **Closed loop relation.** Remove any closed loop and replace it by q if its rightmost vertex is labelled by \vee , and by q^{-1} if its rightmost vertex is labelled by \wedge .



$$\begin{array}{c} \text{Loop with } x \\ \text{Loop with } x \text{ and left loop} \\ \text{Loop with } x \text{ and right loop} \end{array} = \begin{cases} q & \text{if } x = \vee \\ q^{-1} & \text{if } x = \wedge \end{cases}$$


Once all closed loops have been removed, delete all remaining symbols coming from $\mu = \lambda'$ and apply the following relations.

- (2) **Double bead relation.** We have that




in types C_n and D_n respectively.

- (3) **Non-simply laced bead relation.** In type C_n , we have that



Remark 6.9. Note that the multiplication is associative. Indeed, consider the product of three oriented tangles forming a closed loop as pictured below. Then we have that the red zigzag-strand is not left-exposed and so must be oriented. This implies that the symbols at x and y are either both \wedge or both \vee and hence we will get the same result whichever way we apply the two multiplications.



Therefore $\text{TL}_{(W,P)}^{\wedge\vee}(q)$ is a unital associative algebra with identity $\sum_{\lambda \in PW} \lambda 1 \lambda$ where 1 is the tangle in $\mathbb{DT}(W)$ containing only undecorated propagating strands (that is, the identity element in $\text{TL}_W(q)$).




FIGURE 14. Oriented tangles of type (C_9, A_8) , (D_{10}, A_9) , and (C_8, A_7) respectively obtained by orienting the diagrams from Figure 11.

Definition 6.10. We define the parity specialisation map

$$A_{n-1}C_n \rightarrow A_nD_{n+1} \quad : \lambda \mapsto \bar{\lambda}$$

by replacing the \circ in λ by either an \wedge or \vee arrow in such a way that the resulting total number of \wedge arrows is even.

We note that the parity specialisation map is a bijection between our diagrammatic coset representatives, by definition.

In Section 8, we will see that whilst the matrices of light leaves polynomials of types (D_{n+1}, A_n) and (C_n, A_{n-1}) are genuinely distinct, the underlying Kazhdan–Lusztig polynomials are the same (see also [Boe88, ES16a]). The key to understanding this phenomenon will be the following:

Proposition 6.11. *There is a surjective algebra homomorphism*

$$\varphi : \mathrm{TL}_{(C_n, A_{n-1})}^{\wedge \vee}(q) \rightarrow \mathrm{TL}_{(D_{n+1}, A_n)}^{\wedge \vee}(q)$$

by setting $\varphi(\lambda d \mu)$ to be the oriented tangle obtained from $\bar{\lambda} d \bar{\mu}$ by removing the beads from all decorated oriented strands.

Proof. Clearly the map is surjective. We now verify that φ preserves the relations. The closed loops relation is trivially preserved as it only involves the rightmost decoration on the loop, which is unchanged by the parity specialisation map. Equation (6.1) is also trivially preserved. We now verify the bead relations of Definition 6.8. We slightly abuse notation by setting d to be the tangle consisting of a single decorated propagating strand and by setting u to be the tangle consisting of a single undecorated propagating strand. We check that

$$\varphi(\lambda d \mu d \nu) = \varphi \left(\begin{array}{c} \lambda \\ \bullet \\ \mu \\ \bullet \\ \nu \end{array} \right) = \varphi \left(\begin{array}{c} \lambda \\ \bullet \\ \mu \end{array} \right) \varphi \left(\begin{array}{c} \mu \\ \bullet \\ \nu \end{array} \right) = \varphi(\lambda d \mu) \varphi(\mu d \nu)$$

by breaking this up into three cases depending on $\lambda, \mu, \nu \in \{\circ, \wedge, \vee\}$. Note that $\bar{\lambda}, \bar{\mu}, \bar{\nu}$ depend on the entirety of the coset diagram, not just the label of the strand that we are considering (as they are calculated by the overall parity). The three cases are as follows:

(i) If $\bar{\lambda} = \bar{\mu} = \bar{\nu}$ then $\bar{\lambda} d \bar{\mu}$, $\bar{\mu} d \bar{\nu}$, and $\bar{\lambda} d \bar{\nu}$ are oriented strands and so

$$\varphi(\lambda d \mu d \nu) = \varphi(\lambda d \nu) = \bar{\lambda} u \bar{\nu} = (\bar{\lambda} u \bar{\mu})(\bar{\mu} u \bar{\nu}) = \varphi(\lambda d \mu) \varphi(\mu d \nu).$$

(ii) If $\bar{\lambda} = \bar{\mu} \neq \bar{\nu}$ then $\bar{\lambda} d \bar{\mu}$ is oriented and $\bar{\lambda} d \bar{\nu}$, $\bar{\mu} d \bar{\nu}$ are unoriented strands and so

$$\varphi(\lambda d \mu d \nu) = \varphi(\lambda d \nu) = \bar{\lambda} d \bar{\nu} = (\bar{\lambda} u \bar{\mu})(\bar{\mu} d \bar{\nu}) = \varphi(\lambda d \mu) \varphi(\mu d \nu).$$

(iii) If $\bar{\lambda} \neq \bar{\mu} \neq \bar{\nu}$ then $\bar{\lambda} d \bar{\nu}$ is oriented and $\bar{\lambda} d \bar{\mu}$, $\bar{\mu} d \bar{\nu}$ are unoriented strands and so

$$\varphi(\lambda d \mu d \nu) = \varphi(\lambda d \nu) = \bar{\lambda} u \bar{\nu} = (\bar{\lambda} d \bar{\mu})(\bar{\mu} d \bar{\nu}) = \varphi(\lambda d \mu) \varphi(\mu d \nu).$$

The result follows. \square

We now state the main result of this section, which establishes that the abstract and diagrammatically defined algebras are, in fact, isomorphic. Recall, from Remark 6.6 that for any edge $\lambda \xrightarrow{i} \mu$ in the graph $\widehat{\mathcal{G}}_{(W,P)}$ we have an element $\lambda e_i \mu \in \mathrm{TL}_{(W,P)}^{\wedge \vee}(q)$. More generally, for any path $T = \lambda_1 \xrightarrow{i_1} \lambda_2 \xrightarrow{i_2} \dots \lambda_{i_{k-1}} \xrightarrow{i_{k-1}} \lambda_k$ on the graph $\widehat{\mathcal{G}}_{(W,P)}$ we can form the product

$$e_T := \lambda_1 e_{i_1} \lambda_2 e_{i_2} \dots \lambda_{i_{k-1}} e_{i_{k-1}} \lambda_k \in \mathrm{TL}_{(W,P)}^{\wedge \vee}(q).$$

Theorem 6.12. *Let $(W, P) = (A_n, A_k \times A_{n-k-1}), (C_n, A_{n-1})$ or (D_n, A_{n-1}) . There is an isomorphism of $\mathbb{Z}[q, q^{-1}]$ -algebras*

$$\vartheta : \mathrm{TL}_{(W,P)}(q) \longrightarrow \mathrm{TL}_{(W,P)}^{\wedge \vee}(q)$$

defined by $\vartheta(1_\lambda) = \lambda 1_\lambda$, and $\vartheta(1_\lambda E_i 1_\mu) = \lambda e_i \mu$. In particular, for any $T \in \mathrm{Path}_{(W,P)}$ we have $\vartheta(E_T) = e_T$.

Remark 6.13. *Note that we have $\vartheta(E_i) = \vartheta(\sum_\lambda (1_\lambda E_i 1_\lambda + \lambda E_i (1_{\lambda s_i}))) = \sum_\lambda (\lambda e_i \lambda + \lambda e_i (\lambda s_i))$ where the sum is over all $\lambda \in {}^P W$ with $\lambda s_i \in {}^P W$.*

We will prove this theorem in the rest of this section.

Proposition 6.14. *The map ϑ is a $\mathbb{Z}[q, q^{-1}]$ -algebra homomorphism.*

Proof. We need to check that the relations (3.1)–(3.4) are preserved under ϑ . The two leftmost relations in (3.1) are clear by definition of the multiplication in $\text{TL}_{(W,P)}^{\wedge\vee}(q)$. The two rightmost ones follow from the description of the graph $\mathcal{G}_{(W,P)}$ given in Section 5. Relation (3.2) is satisfied by the closed loop relation noting that $\vee\vee < \wedge\wedge$ and $\wedge\vee < \vee\wedge$. The leftmost relation in (3.3) holds as the corresponding tangles are isotopic for $\{i, j\} \neq \{1, 1''\}$. Note that when $\{i, j\} = \{1, 1''\}$ in type (D_n, A_{n-1}) , we have that $E_1 1_\lambda E_{1''} = E_{1''} 1_\lambda E_1 = 0$ for all $\lambda \in {}^P W$, so $E_1 E_{1''} = E_{1''} E_1 = 0$ and there is nothing to check. The rightmost relation in (3.3) is also satisfied using the fact that the corresponding tangles are isotopic and the double bead relation when $\{i, j\} = \{1'', 2\}$ in type (D_n, A_{n-1}) . Relation (3.4) only applies when $i = 2$ and $j = 1'$ in type (C_n, A_{n-1}) . Now, it is easy to see that it holds in the diagrammatic algebra using isotopy, the double bead relations in type C_n and the non-simply laced bead relation. \square

To show that ϑ is an isomorphism, we will need to show that every oriented Temperley–Lieb diagram can be written as a product of generators. We start with the non-oriented diagrams.

6.3. Closed and iterative constructions of Temperley–Lieb diagrams. For $W = A_{n-1}, C_{n-1}$ or D_n , we consider a tiling of a vertical strip of the plane with square tiles labelled by the simple reflections $s \in S_W$ as illustrated in Figure 15.




FIGURE 15. The Temperley–Lieb algebra tiling pictures for types A_9 , D_{10} , and C_9 respectively.

Now, for $(W, P) = (A_{n-1}, A_{k-1} \times A_{n-k-1})$, (C_{n-1}, A_{n-2}) or (D_n, A_{n-1}) and $d \in \mathbb{DT}[W, P]$ we have that d has n vertices on the northern edge, labelled by $1, \dots, n$ (from left to right) and n vertices on the southern edge, labelled by $1', \dots, n'$ (from left to right). We write $i' \geq j'$ whenever $i \geq j$. Each such diagram d will correspond to a finite region on our tiling with boundary given by the path

$$\pi(d) = (\pi(1), \pi(2), \dots, \pi(n), \pi(n'), \pi((n-1)'), \dots, \pi(2'), \pi(1'))$$

defined as follows: Start at the leftmost corner of a tile labelled by 1, or $1'$ if $W = C_{n-1}$, then for $1 \leq k \leq n$ we have

$$\pi(k) = \begin{cases} \text{NE} & \text{if } k \text{ is connected to a vertex } l > k \text{ or } l' \geq k' \text{ by an undecorated strand} \\ \text{SE} & \text{otherwise.} \end{cases}$$

$$\pi(k') = \begin{cases} \text{NW} & \text{if } k' \text{ is connected to a vertex } l > k \text{ or } l' > k' \text{ by an undecorated strand} \\ \text{SW} & \text{otherwise.} \end{cases}$$

Examples are given in Figure 16 and 17. We have drawn vertical lines through all tiles not included in $R(d)$. We see that the vertical line starting at vertex i , respectively i' meets the path π at $\pi(i)$, respectively $\pi(i')$ for each $1 \leq i \leq n$.

Given a subset $X \subseteq \{1, \dots, n, n', \dots, 1'\}$ we set

$$N_\pi(X) = \{k \in X \mid \pi(k) = \text{NE}\} \sqcup \{k' \in X \mid \pi(k') = \text{NW}\}$$

$$S_\pi(X) = \{k \in X \mid \pi(k) = \text{SE}\} \sqcup \{k' \in X \mid \pi(k') = \text{SW}\}.$$

First note that the path $\pi(d)$ takes n steps to the East and n steps to the West. So the path starts and finishes on the left boundary of the tiling. Moreover, the number of steps to the North is precisely the number of undecorated strands in d . So if d is an undecorated n -tangle, then the path starts and ends at the same point; this is because

$$|N_\pi\{1, \dots, n, n', \dots, 1'\}| = n \quad |S_\pi\{1, \dots, n, n', \dots, 1'\}| = 2n - n = n.$$

But if d has at least one bead, then the path will end strictly below where it started, this is because in this case

$$|N_\pi\{1, \dots, n, n', \dots, 1'\}| < n \quad |S_\pi\{1, \dots, n, n', \dots, 1'\}| > 2n - n = n.$$

More generally, the second half of the path (going West) is always weakly to the South of the first half of the path (going East). To see this, observe that for $1 \leq k \leq n$, the difference in height in the path after k steps and after $2n - k$ steps is equal to

$$|S_\pi\{k+1, \dots, n, n', \dots, (k+1)'\}| - |N_\pi\{k+1, \dots, n, n', \dots, (k+1)'\}| \geq 0$$

where the inequality follows by definition as every l (or l') in $N_\pi\{k+1, \dots, n, n', \dots, (k+1)'\}$ is connected to a vertex in $S_\pi\{k+1, \dots, n, n', \dots, (k+1)'\}$. Thus $\pi(d)$ defines a region $R(d)$ in the tiling which contains a finite set of tiles t_1, \dots, t_r (which are labelled by simple reflections in S_W).




FIGURE 16. The tilings of the diagrams from Figure 13. In each case the path begins at the western-most point, which is denoted with a circle; the path then follows the orientation depicted on the diagram. An example of the associated e_w for the rightmost diagram is $e_w = e_7 e_4 e_6 e_8 e_1 e_3 e_5 e_7 e_2 e_4$.

We now explain how the region $R(d)$ defines a reduced word of a strongly fully commutative element of W . Enumerate the tiles in $R(d)$, t_{i_1}, \dots, t_{i_r} , in such a way that for each j , the tiles in $R(d)$ to the SW and SE of t_{i_j} appear before t_{i_j} . Taking the corresponding ordered product of simple reflections gives a word $w = \underline{w}(d)$ in the elements of S_W . There is, of course, more than one way of enumerating the tiles in $R(d)$ in this fashion and these give (all the) different reduced expressions for the same element of W (as they differ only by commutation relations).




FIGURE 17. The tiling of the two leftmost diagrams from Figure 14 and an additional one in type (C_9, A_8) . The path begins at the northerly western-most point; the path then follows the orientation depicted on the diagram until terminating at the southerly western-most point (both of which are denoted with circles). Some of the strands in type C carry two dots which can be simplified to one dot.

Conversely, any expression $\underline{w} = s_{i_1} s_{i_2} \dots s_{i_k}$ with $s_{i_j} \in S_W$ defines a region $R(\underline{w})$ in the tiling by picking any horizontal line in the tiling and stacking the boxes in order, starting from s_{i_1} , then s_{i_2} , \dots , and finally s_{i_k} . (This is an alternative description of the ‘heaps’ introduced by Stembridge [Ste96]).

- Proposition 6.15.** (1) Let \underline{w} is a reduced expression for a strongly fully commutative element of W such that $\underline{w} = \omega(\mathbb{T})$ for some $T \in \text{Path}_{(W,P)}$. Then $R(\underline{w})$ does not have a boundary containing an inadmissible section of the form depicted in Figure 18.
- (2) For any $d \in \text{DT}[W, P]$, the region $R(d)$ does not have a boundary containing an inadmissible section of the form depicted in Figure 18. In particular, $\underline{w}(d)$ is a reduced expression for a strongly fully commutative element of W .
- (3) For any $d \in \text{DT}[W, P]$, if $\underline{w}(d) = \underline{w} = s_{i_1} s_{i_2} \dots s_{i_k}$ then we have

$$d = \mathbf{e}_{\underline{w}} := \mathbf{e}_{i_1} \mathbf{e}_{i_2} \dots \mathbf{e}_{i_k}.$$

Moreover, if \underline{w}' is another reduced expression for the same strongly fully commutative element of W then $\mathbf{e}_{\underline{w}'} = \mathbf{e}_{\underline{w}}$.

- Proof.* (1) The conditions on i, j follow from Definition 3.1 and the definition of a reduced expression (see Section 2). For the conditions involving 1 and 1'', we note that any path on $\widehat{\mathcal{G}}_{(D_n, A_{n-1})}$ containing an s_1 -step and an $s_{1''}$ -step must contain an s_2 -step in between.
- (2) Note that the path $\pi(d)$ takes all steps to the East first, followed by all steps to the West, this implies that boundary of $R(d)$ does not contain any inadmissible sections and hence $\underline{w}(d)$ is a reduced expression for a fully commutative element of W .
- (3) Now recall that for each simple reflection $s_i \in S_W$, we have a generator $\mathbf{e}_i \in \text{TL}_W(q)$, and so any reduced expression $\underline{w} = s_{i_1} s_{i_2} \dots s_{i_k}$ for a strongly fully commutative element of W define a product of the generators, $\mathbf{e}_{\underline{w}} = \mathbf{e}_{i_1} \mathbf{e}_{i_2} \dots \mathbf{e}_{i_k} \in \text{TL}_W(q)$. These are pictured in Figures 16 and 17 for $\underline{w} = \underline{w}(d)$. Again, it is clear that for a different choice of reduced word \underline{w}' we have $\mathbf{e}_{\underline{w}'} = \mathbf{e}_{\underline{w}}$ (as these differ only by commutation relations). We claim that $\mathbf{e}_{\underline{w}} = d$. To see this, we first partition the region $R(d)$ by splitting it along the horizontal




FIGURE 18. We emphasise the inadmissible section of the border-region by drawing it thickly. In type A_{n-1} we require that there is no such region for any $i, j \in \{1, 2, \dots, n-1\}$. In type D_n we require that there is no such region for any $i \in \{2, \dots, n-1\}$ or $j \in \{2, \dots, n-1\}$. In type C_{n-1} we require that there is no such region for any $i \in \{1', 2, \dots, n-1\}$ or $j \in \{2, \dots, n-1\}$. We note that the latter two regions (involving a 1 and a $1''$ tile) occur only in type D_{n-1} .

lines through the vertices of the tiles. This gives a partition of $R(d)$ into horizontal strips. If a strip intersects the left boundary in more than one point, merge it with the strip above or below so that the new wider strip now contains precisely two vertices of tiles on the left boundary. These strips are shown in Figures 16 and 17 (by alternating between grey and white shading). It is clear that the strands in $e_{\underline{w}}$ are in one-to-one correspondence with the strips of $R(d)$. Each strip contains precisely two edges of the path $\pi(d)$ and joins the corresponding vertices in d ; one thus recovers the (decorated) n -tangle d .

□

Proposition 6.16. *Let \underline{w} be a reduced expression for a strongly fully commutative element of W . Suppose that we have $\lambda, \mu \in {}^P W$ with $\lambda e_{\underline{w}} \mu \in \text{ODT}[W, P]$. Then there is a unique path T on $\widehat{\mathcal{G}}_{(W, P)}$ starting at λ and ending at μ with $w(\mathsf{T}) = \underline{w}$. Moreover, we have $\lambda e_{\underline{w}} \mu = e_{\mathsf{T}}$.*

Proof. Having fixed the orientation at the top and bottom of the product of generators $e_{\underline{w}}$, there are only two situations in which we have a choice of orientation for strand segments in this product. The first one is when the diagram contains a closed loop. These are formed by a tile configuration of the form depicted on the left of Figure 19. The second one is when $(W, P) = (C_{n-1}, A_{n-2})$ and we have a tile configuration of the form depicted on the right of Figure 19. By Proposition 6.15, neither of these can happen. Therefore the top and bottom orientations uniquely determine the orientation of every strand segment in the diagram, proving the result. □




FIGURE 19. On the left is the tile configuration for a closed loop. On the right the tile configuration of a propagating strand connecting the first northern and southern vertices in type (C_{n-1}, A_{n-2}) . The orientation of the closed loop is not determined by the orientation at the northern and southern edges of the diagram (obviously). The orientation of strand segment *between* the two decorations is also not determined by the orientation at the northern and southern edges of the diagram. The colouring of tiles should be compared with that of Figure 18.

Corollary 6.17. *The map ϑ is a $\mathbb{Z}[q, q^{-1}]$ -module isomorphism.*

Proof. Using Propositions 6.15 and 6.16, we have that every oriented Temperley–Lieb diagram $\lambda d\mu = \lambda e_{\underline{w}}\mu = e_{\mathbb{T}} = \vartheta(E_{\mathbb{T}})$ and so the map ϑ is surjective. To show that it is injective, note that the basis elements $E_{\mathbb{T}}$ for $\mathrm{TL}_{(W,P)}(q)$ given in Theorem 3.8 are mapped to distinct basis elements in $\mathrm{TL}_{(W,P)}^{\wedge\vee}(q)$. \square

This completes the proof of Theorem 6.12. From now on, we identify $\mathrm{TL}_{(W,P)}(q) = \mathrm{TL}_{(W,P)}^{\wedge\vee}(q)$ and use the diagrammatic notation for its elements.

As noted earlier, if \underline{w} and \underline{w}' are two reduced expressions for the same strongly fully commutative element w , then we have $e_{\underline{w}} = e_{\underline{w}'}$. So we will denote this element by e_w . In particular, for each $\mu \in {}^P W$ we have the corresponding element $e_{\mu} \in \mathrm{TL}_W(q)$. (We will give a closed combinatorial description of this element in Section 8.) Now, restricting our attention to the anti-spherical module and using Corollary 3.10 gives the following.

Corollary 6.18. *The anti-spherical module $\mathcal{O}\mathrm{TL}_{(W,P)}(q)$ has basis given by*

$$\{e_{\mathbb{T}} = \mathcal{O}e_{\mu}\lambda \mid \mathbb{T} \in \mathrm{Path}(\lambda, \underline{\mu}), \lambda, \mu \in {}^P W\}$$

In particular we have

$$|\mathrm{Path}(\lambda, \underline{\mu})| \leq 1 \quad \text{for all } \lambda, \mu \in {}^P W.$$

Thus, the diagrammatic oriented Temperley–Lieb algebra setting (and its anti-spherical module in particular) provides a diagrammatic model for studying the matrix $\Delta_{\lambda,\mu}$. More precisely, we have shown that

$$\Delta_{\lambda,\mu} = \begin{cases} q^{\deg(\mathcal{O}e_{\mu}\lambda)} & \text{if } \mathcal{O}e_{\mu}\lambda \in \mathcal{O}\mathrm{DT}[W, P] \\ 0 & \text{otherwise.} \end{cases}$$

In the next section, we investigate the degree of oriented Temperley–Lieb diagrams.

7. A CLOSED COMBINATORIAL INTERPRETATION OF THE GRADING

The isomorphism ϑ given in Theorem 6.12 gives a grading on the diagrammatic oriented Temperley–Lieb algebra. Explicitly, we have that the degree of any oriented Temperley–Lieb diagram $\lambda d\mu = \lambda e_w\mu = e_{\mathbb{T}}$ is equal to $\deg(\mathbb{T})$. This is computed as the sum of the degree of each step in the path \mathbb{T} . We now provide a closed combinatorial description of this degree in terms of the diagram $\lambda d\mu$ itself.

Theorem 7.1. *We define the degree of an oriented Temperley–Lieb diagram by assigning a degree to each of its strands and then summing over all strands. We define the degree of a northern arc whose rightmost vertex is labelled by \vee to be +1, and the degree of a southern arc whose rightmost vertex is labelled by \wedge to be –1. All other strands are defined to have degree 0 (in particular, all propagating strands have degree 0). Diagrammatically, we record the degree of an oriented Temperley–Lieb diagram as follows*

$$\# \left\{ \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} \right\} - \# \left\{ \begin{array}{c} \curvearrowleft \\ \curvearrowright \end{array} \right\}$$

where the dots are to emphasise that the lefthand-side of the arc does not contribute to the degree. Then for any oriented Temperley–Lieb diagram $\lambda d\mu = e_{\mathbb{T}}$ we have $\deg(\lambda d\mu) = \deg(\mathbb{T})$.

Proof. We first check that the degree of the generators $\lambda e_i\mu$ (for $\mu = \lambda$ or λs_i) are correct. Note that we have $\lambda < \lambda s_i \in {}^P W$ precisely when $\lambda = \dots \wedge \vee \dots$ and $\lambda s_i = \dots \vee \wedge \dots$ for $i \in \{1, \dots, n-1\}$, or $\lambda = \vee \vee \dots$ and $\lambda s_{1'} = \wedge \wedge \dots$, or $\lambda = \circ \vee \dots$ and $\lambda s_{1'} = \circ \wedge \dots$. So the degrees of the generators are given by

$$\lambda e_i\lambda = \begin{cases} 1 + 0 = 1 & \text{if } \lambda < \lambda s_i \\ 0 - 1 = -1 & \text{if } \lambda > \lambda s_i \end{cases} \quad \lambda e_i(\lambda s_i) = \begin{cases} 0 + 0 = 0 & \text{if } \lambda < \lambda s_i \\ 1 - 1 = 0 & \text{if } \lambda > \lambda s_i \end{cases}.$$

These are illustrated in Figure 20 for $i \in \{1, \dots, n-1\}$. Comparing these with Proposition 3.11 proves the result for the generators.




FIGURE 20. The four generators $\lambda_{e_i\mu}$ for $i \in \{1, \dots, n-1\}$ of degree $1 = 1 + 0$, $-1 = 0 - 1$, $0 = 0 + 0$, and $0 = 1 - 1$ respectively. We record the degrees of the strand segments within the tile (with $+$ = 1, $-$ = -1 and o = 0).

Now as explained in Subsection 6.3, we can write any oriented Temperley-Lieb diagram as a product of generators e_τ with $\omega(T)$ a reduced word for a strongly fully commutative element of W using a region in the tiling for W . Moreover, we have seen how each strand of the diagram corresponds to a horizontal strip in the tiling. We can then obtain the degree of strand in the product by adding the degree of each small arc coming from the generators. We now run through the various types of strands and check that the result holds in each case.

First observe that any northern or southern arc passes through an odd number of tiles and any propagating strand passes through an even number of tiles. The undecorated northern arcs are illustrated in Figure 21.




FIGURE 21. The undecorated northern arcs.

In Figures 21 to 25, we highlight the tiles through which the strand “wiggles” and the degree of the strand within a given tile (which we have already calculated in terms of the generators. Note that, reading from left to right, the strand oscillates between being either the “top” or “bottom” of a given generator tile. Assume the arc is undecorated, northern, and clockwise-oriented. The degree contribution as the arc passes through these tiles is given by

$$+1 - 1 + 1 - 1 + 1 - 1 \dots + 1 = 1$$

(notice that the strand is locally either a clockwise northern arc or an anticlockwise southern arc at each step). If an undecorated northern arc is anti-clockwise oriented then it has degree

$$0 + \dots + 0 = 0.$$

Thus the degrees match up. The degrees of undecorated southern arcs can be computed similarly. The degree of undecorated propagating strands can easily be seen to be zero as illustrated in Figure 24.




FIGURE 22. Degree 1 decorated northern strands. The pink highlights an odd number of boxes, the blue highlights an even number. In this way, note that the blue section of the strand never contributes to the degree, regardless of the orientation.




FIGURE 23. Degree zero decorated northern strands.

The decorated northern arcs of degree 1 in type (C_{n-1}, A_{n-2}) are illustrated in Figures 22 to 25. Swapping the orientation of the rightmost vertex from \vee to \wedge , we see that the pink strip now doesn't contribute to the degree either, and so these northern arcs have degree zero. The decorated southern arcs can be dealt with in a similar way.

It is easy to see that any (decorated) propagating strand also has degree zero as it goes through an even number of tiles.




FIGURE 24. The undecorated propagating strands, all of which are of degree 0.




FIGURE 25. The decorated propagating strands, all of which are of degree 0.

Finally, the strands in a digram of type (D_n, A_{n-1}) are the same as those in type (C_{n-1}, A_{n-2}) except that they contain at most one decoration and we only need to consider arcs which are flip-oriented. So the result holds for these as well. \square

Corollary 7.2. *The anti-spherical module for $\mathrm{TL}_{(W,P)}(q)$ is non-negatively graded, that is*

$$\deg(\emptyset e_\mu \lambda) \geq 0 \quad \text{for all } \lambda, \mu \in {}^P W.$$

Thus, the non-zero entries in the light leaves matrix $\Delta_{\lambda,\mu}$ are non-negative powers of q .

Proof. Any northern arc or propagating strand is non-negatively graded, thus it suffices to consider the southern arcs (which can be negatively graded). Now recall the coset diagram for \emptyset in types $(W, P) = (A_n, A_k \times A_{n-k-1}), (C_n, A_{n-1}), (D_n, A_{n-1})$ from Section 5. We see that any southern arc must have rightmost vertex labelled by \vee . So the southern arcs all have degree zero. This proves the result. \square

8. FACTORISATION OF THE LIGHT LEAVES MATRIX

We now explain how to construct the element $e_\mu \in \mathrm{TL}_W(q)$ for $\mu \in {}^P W$ via an efficient closed combinatorial algorithm, which has its origins in [BS12, ES16b, Mar15, CD11]. This allows us to enumerate the elements $\emptyset e_\mu \lambda$ and hence the paths $\mathrm{NPath}(\lambda, \underline{\mu}) \subseteq \mathrm{Path}(\lambda, \underline{\mu})$ and $\mathrm{BPath}(\nu, \underline{\mu}) \subseteq \mathrm{Path}(\nu, \underline{\mu})$ corresponding to decomposition numbers and bases of simple modules

for the Hecke category respectively (in other words, solving Libedinsky–Williamson’s question for Hermitian symmetric pairs).

We have seen that every strongly fully commutative element w in W with $\underline{w} = \omega(\mathbb{T})$ for some $\mathbb{T} \in \text{Path}_{(W,P)}$ corresponds to a region $R(w)$ in the tiling of W . That region is determined by the path $\pi(w)$ walking along its boundary, starting with its northern boundary going East and coming back along its southern boundary going West. When $w = \mu \in {}^P W$, the region $R(\mu)$ has a particularly simple shape.

Lemma 8.1. *We have that $w \in {}^P W$ if and only if the region $R(w)$ has the following property. In type $(A_{n-1}, A_{k-1} \times A_{n-k-1})$, the last n steps of the path $\pi(w)$ are given by $((SW)^{n-k}, (NW)^k)$. In types (C_{n-1}, A_{n-2}) and (D_n, A_{n-1}) , the last n steps of the path $\pi(w)$ are given by $((SW)^n)$.*

Proof. Note that the minimal length coset representatives $\mu \in {}^P W$ are characterised by the fact that every reduced expression for μ starts with $s \notin P$. From that characterisation, it is clear that if $\pi(w)$ has the stated form then $w \in {}^P W$. Examples of $\pi(\mu)$ for $\mu \in {}^P W$ are given in the first two pictures in Figure 16 and Figure 17. Now suppose that $\pi(w)$ does not have the stated form then $\pi(w)$ would have two consecutive steps $\pi((n-i)') = SW$ and $\pi((n-i-1)') = NW$ (with $i \neq k$ in type $(A_{n-1}, A_{k-1} \times A_{n-k-1})$). But this would imply that there is a reduced expression for w starting with some $s \in P$ and so $w \notin {}^P W$. This is illustrated in the rightmost picture of Figure 16 and Figure 17 where we indeed observe that there is a reduced word for w starting with $s \in P$ in each case. \square

Therefore for $\mu \in {}^P W$, the region $R(\mu)$ is completely determined by its northern boundary. In fact, we have the following natural correspondence between the northern boundary of $R(\mu)$ and the coset diagram of μ .

Lemma 8.2. *Let $\mu \in {}^P W$. The i -th vertex of the coset diagram of μ is labelled by \wedge , respectively \vee , if and only if the i -th step in $\pi(\mu)$ is given by SE , respectively NE . In type (C_{n-1}, A_{n-2}) the first vertex is always labelled by \circ and the first step is always SE .*

Proof. We proceed by induction on $\ell(\mu)$. For $\ell(\mu) = 0$ we have $\mu = \emptyset$ and the result is clear from the description of the coset diagram for \emptyset given in Section 5. Now assume that the result holds for $\lambda \in {}^P W$ and let $\mu = \lambda s_i > \lambda$. If $i = 1, \dots, n-1$ then the coset diagrams for λ and λs_i only differ in position i and $i+1$ and we have $\lambda = \dots \wedge \vee \dots$ and $\lambda s_i = \dots \vee \wedge \dots$. The corresponding paths $\pi(\lambda)$ and $\pi(\lambda s_i)$ are depicted in Figure 26. We see that the i -th and $(i+1)$ -th steps in λ , respectively λs_i , are given by SE, NE , respectively NE, SE and so the result follows by induction. For $i = 1'$, the coset diagrams of λ and $\lambda s_{1'}$ only differ in the second position and we have $\lambda = \circ \vee \dots$ and $\lambda s_{1'} = \circ \wedge \dots$. The corresponding paths $\pi(\lambda)$ and $\pi(\lambda s_{1'})$ are depicted in Figure 26. We see that the second step in λ is given by NE while the second step in $\lambda s_{1'}$ is given by SE , and so the result follows by induction. Finally, if $i = 1''$ then the coset diagrams for λ and $\lambda s_{1''}$ only differ in the first two positions and we have $\lambda = \vee \vee \dots$ and $\lambda s_{1''} = \wedge \wedge \dots$. The corresponding paths $\pi(\lambda)$ and $\pi(\lambda s_{1''})$ are depicted in Figure 26. We see that the first two steps in λ , respectively $\lambda s_{1''}$, are given by NE, NE , respectively SE, SE , and so the result follows by induction. \square

Using this correspondence, we easily obtain the following closed combinatorial algorithm to construct \mathbf{e}_μ .

Proposition 8.3. *For each $\mu \in {}^P W$, the diagram $\mathbf{e}_\mu \in \text{TL}_W(q)$ can be constructed as follows. Place the coset diagram μ on the northern boundary and \emptyset on the southern boundary. Then*

- (1) *Repeatedly connect neighbouring northern vertices (in the sense that they are next to each other or only have vertices already connected by an arc between them) labelled by \vee and \wedge by a northern anti-clockwise arc.*

We are left with (in type (C_{n-1}, A_{n-2}) one vertex labelled by \circ followed by) some vertices labelled by \wedge followed by some vertices labelled by \vee .




FIGURE 26. The effect of applying a reflection s_i for $1 \leq i \leq n-1$; the reflection $s_i = s_{1'}$ (in type C); and the reflection $s_i = s_{1''}$ (in type D) respectively. In each case we depict the pair of weights λ and λ_{s_i} and the corresponding northern edges of the paths $\pi(\lambda)$ and $\pi(\lambda_{s_i})$. We highlight in blue and pink the difference between λ and λ_{s_i} (both on the level of tiles and coset diagrams).

- (2) (a) In type $(A_{n-1}, A_{k-1} \times A_{n-k-1})$, draw undecorated propagating strands on all remaining vertices.
- (b) In type (D_n, A_{n-1}) , starting from the left, connect neighbouring vertices labelled with \wedge 's with decorated northern arcs. Then draw a decorated propagating strand from the remaining vertex labelled by \wedge (if it exists), and undecorated propagating strands on all remaining vertices labelled by \vee .
- (c) In type (C_{n-1}, A_{n-2}) , if $\mu = \emptyset$, then draw an undecorated propagating strand from each northern vertex. Otherwise, view the first label \circ of μ as a \wedge , then follow exactly the same procedure as in type (D_n, A_{n-1}) .

Now there is a unique way of completing the diagram e_μ such that $\emptyset e_\mu \mu \in \text{ODT}[W, P]$.

Example 8.4. A couple of illustrative large examples of the construction in Proposition 8.3 are given in Figures 27 and 28. The complete set of all e_μ for $\mu \in {}^P W$ for (W, P) of type (D_4, A_3) and (C_3, A_2) are given in Figure 29. In all examples, we will only picture the top of the diagram as this completely determines e_μ .




FIGURE 27. Two examples of the construction of e_μ , the former is of type $(A_8, A_4 \times A_3)$ and the latter is of type (C_8, A_7) .




FIGURE 28. An example of the construction of e_μ in type (D_{19}, A_{18}) .

Proposition 8.3 gives us an efficient way of finding the degree of $\emptyset e_\mu \lambda$ as the number of northern arcs whose rightmost vertex is labelled by \vee . As this does not depend on the bottom

of the diagram (as all southern arcs have degree zero), we will only depict the top of $\varnothing e_\mu \lambda$. Examples are given in Figures 30 and 31.




FIGURE 29. The construction of e_μ for all $\mu \in {}^P W$ in types (D_4, A_3) and (C_3, A_2) .




FIGURE 30. The elements $\varnothing e_\mu \lambda$ from the first column of Figure 31 in order.













						
	1
	q	1
	.	q	1	.	.	.
	.	q	.	1	.	.
	q	q^2	q	q	1	.
	q^2	.	.	.	q	1

FIGURE 31. The $q^{\deg(\varnothing e_\mu \lambda)}$ for $\lambda, \mu \in {}^P W$ for $(W, P) = (A_3, A_1 \times A_1)$.

We have already seen in Corollary 7.2 that the entries in the light leaves matrix $\Delta_{\lambda\mu}$ are either zero or non-negative powers of q . In the simply-laced cases, we can say more.

Theorem 8.5. *Assume $(W, P) = (A_{n-1}, A_{n-2})$ or (D_n, A_{n-1}) . Then for any $\lambda, \mu \in {}^P W$ with $\varnothing e_\mu \lambda \in \text{ODT}[W, P]$ we have $\deg(\varnothing e_\mu \lambda) = 0$ if and only if $\lambda = \mu$. In particular, we have $\Delta_{\lambda\mu} = 1$ if and only if $\lambda = \mu$ and so the matrix of light leaves has a trivial factorisation*

$$B = \text{Id} \quad \text{and} \quad \Delta = N.$$

Proof. Fix $\mu \in {}^P W$ and consider all $\lambda \in {}^P W$ with $\varnothing e_\mu \lambda \in \text{ODT}[W, P]$ and $\deg(\varnothing e_\mu \lambda) = 0$. Note that this forces all undecorated northern arcs to be anti-clockwise, all undecorated propagating strands to be oriented, all northern decorated arcs to be labelled by two \wedge 's and all decorated propagating strands to be flip-oriented. This gives only one choice for λ , namely $\lambda = \mu$. \square

Thus, setting every basis element $\varnothing e_\mu \lambda$ for the anti-spherical module to be standard proves Theorem A in type $(W, P) = (A_{n-1}, A_{n-2})$ and (D_n, A_{n-1}) . We now state and proof Theorem A in the only remaining type of Hermitian symmetric pair, namely (C_n, A_{n-1}) .

Theorem 8.6. *Let $\lambda, \mu \in {}^{A_{n-1}} C_n$. We say that $\varnothing e_\mu \lambda \in \text{ODT}[C_n, A_{n-1}]$ is standard if every decorated strand is flip-oriented.*

We define the matrices N and B in type $(W, P) = (C_n, A_{n-1})$ as follows:

$$N_{\lambda, \mu} = \begin{cases} q^{\deg(\varnothing e_\mu \lambda)} & \text{if } \varnothing e_\mu \lambda \in \text{ODT}[C_n, A_{n-1}] \text{ is standard} \\ 0 & \text{otherwise} \end{cases} \quad (8.1)$$

$$B_{\lambda, \mu} = \begin{cases} 1 & \text{if } \varnothing e_\mu \lambda \in \text{ODT}[C_n, A_{n-1}] \text{ and } \deg(\varnothing e_\mu \lambda) = 0 \\ 0 & \text{otherwise.} \end{cases} \quad (8.2)$$

Then the matrix of light leaves in type (C_n, A_{n-1}) factorises as

$$\Delta = NB.$$

Proof. Fix $\mu \in {}^{A_{n-1}} C_n$ and consider

$$C_\mu = \{\lambda \in {}^{A_{n-1}} C_n : \varnothing e_\mu \lambda \in \text{ODT}[C_n, A_{n-1}]\}.$$

Define an equivalence relation on C_μ by setting $\lambda \sim_\mu \eta$ for $\lambda, \eta \in C_\mu$ if $\varnothing e_\mu \lambda$ and $\varnothing e_\mu \eta$ differ only in one northern arc where the differing arcs are given by

$$\left\{ \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} \right\}, \left\{ \begin{array}{c} \curvearrowright \\ \curvearrowright \end{array} \right\} \text{ or } \left\{ \begin{array}{c} \curvearrowright \\ \bullet \end{array} \right\}, \left\{ \begin{array}{c} \curvearrowleft \\ \bullet \end{array} \right\} \text{ or } \left\{ \begin{array}{c} \curvearrowright \\ \bullet \end{array} \right\}, \left\{ \begin{array}{c} \curvearrowleft \\ \bullet \end{array} \right\} \text{ or } \left\{ \begin{array}{c} \circ \\ \bullet \end{array} \right\}, \left\{ \begin{array}{c} \circ \\ \bullet \end{array} \right\}. \quad (8.3)$$

We extend \sim_μ by transitivity. Note that, under this relation, each equivalence class contains a unique element ν satisfying $\deg(\varnothing e_\mu \nu) = 0$ (obtained by orienting all northern arcs with rightmost vertex labelled by \wedge).

Now, let $\lambda, \mu \in {}^{A_{n-1}}C_n$. We claim that

$$\Delta_{\lambda, \mu} = \sum_{\eta \in {}^{A_{n-1}}C_n} N_{\lambda, \eta} B_{\eta, \mu} = \begin{cases} N_{\lambda, \nu} B_{\nu, \mu} & \text{if } \varnothing e_\mu \lambda \in \mathbb{ODT}[C_n, A_{n-1}] \\ 0 & \text{otherwise} \end{cases}$$

where ν is the unique element satisfying $\lambda \sim_\mu \nu$ and $\deg(\varnothing e_\mu \nu) = 0$. Suppose that $N_{\lambda, \eta} \neq 0$ and $B_{\eta, \mu} \neq 0$ for some η . This implies that $\varnothing e_\mu \eta \in \mathbb{ODT}[C_n, A_{n-1}]$ with $\deg(\varnothing e_\mu \eta) = 0$, that is all northern arcs in $\varnothing e_\mu \eta$ have rightmost vertex labelled by \wedge . Thus we have that e_η has the same arcs as e_μ but with beads corresponding to oriented decorated northern arcs not connected to the symbol \circ in $\varnothing e_\mu \eta$ removed. Now, $N_{\lambda, \eta} \neq 0$ means that $\varnothing e_\eta \lambda \in \mathbb{ODT}[C_n, A_{n-1}]$ is standard. This implies that we also have $\varnothing e_\mu \lambda \in \mathbb{ODT}[C_n, A_{n-1}]$. Now we claim we have $\lambda \sim_\mu \eta$. The fact that the northern arcs are related as in (8.3) follows by definition. That the orientation of the propagating lines of $\varnothing e_\mu \lambda$ and $\varnothing e_\mu \eta$ coincide follows from the fact that the orientation on the southern boundary is given by \varnothing in both cases and if e_μ contains a decorated propagating line then its northern label is given by the parity condition. Now as $\eta \sim_\mu \lambda$ and $\deg(\varnothing e_\mu \eta) = 0$ implies that $\eta = \nu$. Thus we have shown that

$$\sum_{\eta \in {}^{A_{n-1}}C_n} N_{\lambda, \eta} B_{\eta, \mu} = N_{\lambda, \nu} B_{\nu, \mu}.$$

Finally note that, as $\nu \sim_\mu \lambda$, a decorated arc in e_μ is oriented in $\varnothing e_\mu \lambda$ if and only if it is oriented in $\varnothing e_\mu \nu$. Now, as noted above, e_ν is obtained from e_μ by removing all beads on these oriented decorated arcs but the position of the arcs are the same and so $\deg(\varnothing e_\mu \lambda) = \deg(\varnothing e_\nu \lambda)$. Thus we get

$$\Delta_{\lambda, \mu} = q^{\deg(\varnothing e_\mu \lambda)} = q^{\deg(\varnothing e_\nu \lambda)} = N_{\lambda, \nu} B_{\nu, \mu}.$$

as required. \square

Remark 8.7. Note that the unique element ν described in the proof above is precisely the element satisfying $\varphi(\varnothing e_\mu \lambda) = \overline{\varnothing e_\nu \lambda}$ where $\varphi : \text{TL}_{(C_n, A_{n-1})}(q) \rightarrow \text{TL}_{(D_{n+1}, A_n)}(q)$ is the homomorphism defined in Proposition 6.11.

Example 8.8. There are 6 non-standard basis elements $\varnothing e_\mu \lambda$ in the anti-spherical module of type (C_3, A_2) (this can be deduced from Example 2.3 and Theorem 8.6), these are depicted in Figure 32.




FIGURE 32. The six non-standard diagrams of type (C_3, A_2) . The first and third of these diagrams have degree zero.

Corollary 8.9. Under the parity specialisation map ${}^{A_{n-1}}C_n \rightarrow {}^{A_n}D_{n+1} : \lambda \mapsto \bar{\lambda}$ given in Definition 6.10 we have that

$$N_{\lambda, \mu}^{(C_n, A_{n-1})} = N_{\bar{\lambda}, \bar{\mu}}^{(D_{n+1}, A_n)}$$

Proof. We need to show that $\varnothing e_\mu \lambda \in \text{ODT}[C_n, A_{n-1}]$ is standard if and only if $\overline{\varnothing} e_{\bar{\mu}} \bar{\lambda} \in \text{ODT}[D_{n+1}, A_n]$. First note that if $\varnothing e_\mu \lambda \in \text{ODT}[C_n, A_{n-1}]$ is not standard then $\overline{\varnothing} e_{\bar{\mu}} \bar{\lambda} \notin \text{ODT}[D_{n+1}, A_n]$. Now assume that $\varnothing e_\mu \lambda \in \text{ODT}[C_n, A_{n-1}]$ is standard. The result is trivial for $\mu = \varnothing$ so we assume that $\mu \neq \varnothing$. This implies that the strand coming out of the first northern vertex in e_μ , call it S , is decorated. By definition, $e_{\bar{\mu}}$ is equal to e_μ if S is flip-oriented in $\overline{\varnothing} e_{\bar{\mu}} \bar{\lambda}$, and $e_{\bar{\mu}}$ is obtained from e_μ by removing the bead on S if it is oriented. Now, except possibly for S the orientations of the propagating strands in $\overline{\varnothing} e_{\bar{\mu}} \bar{\lambda}$ and in $\overline{\varnothing} e_\mu \lambda$ coincide and each northern arc is oriented, respectively flip-oriented, in the former if and only if it is in the latter. As $\bar{\lambda}$ and $\bar{\mu}$ both have an even number of \wedge arrow, this implies that S is oriented, respectively flip-oriented, in the former, if and only if it is in the latter. Hence we have that $\overline{\varnothing} e_{\bar{\mu}} \bar{\lambda} \in \text{ODT}[D_{n+1}, A_n]$. \square

APPENDIX A. SINGULAR KAZHDAN–LUSZTIG THEORY

In this section, for any $\tau \in S_W$, we calculate the singular Kazhdan–Lusztig polynomials “lying on a τ -hyperplane” for simply-laced Hermitian symmetric pairs (W, P) . This is a technical result that we will need in [BDHN] and we therefore adopt the following notation from that paper. We set $\mathcal{P}_{(W,P)} = {}^P W$ and we let

$$\mathcal{P}_{(W,P)}^\tau := \{\mu \in {}^P W \mid \mu\tau < \mu\} \subset \mathcal{P}_{(W,P)}.$$

We believe the following proposition was first proven by Enright–Shelton, where any singular category \mathcal{O} for a Hermitian symmetric pair is proven to be Morita equivalent to a regular category \mathcal{O} for a Hermitian symmetric pair of smaller rank. It is explicitly stated as [BDHN, Proposition 5.4] in our companion paper, however we note that the proof offered there is simply given by inspection and follows from the combinatorial description of the cosets (this combinatorics can be found explicitly recorded in many other places too, even if the bijection itself cannot, see for instance [EHP14, Appendix]). Thus while the following is a reference to a paper whose results depend on the current work, this is not a circular argument.

Proposition A.1. *Let (W, P) be a simply laced Hermitian symmetric pair and let $\tau \in S_W$. There is an order preserving bijection*

$$\psi_\tau : \mathcal{P}_{(W,P)}^\tau \rightarrow \mathcal{P}_{(W,P)^\tau}$$

where $(W, P)^\tau = (W^\tau, P^\tau)$ is defined by

- $(A_n, A_k \times A_{n-k-1})^\tau = (A_{n-2}, A_{k-1} \times A_{n-k-2})$;
- $(D_n, A_{n-1})^\tau = (D_{n-2}, A_{n-3})$;
- $(D_n, D_{n-1})^\tau = (A_1, A_0)$;
- $(E_6, D_5)^\tau = (A_5, A_4)$;
- $(E_7, E_6)^\tau = (D_6, D_5)$.

For $\tau = s_i$ in types $(W, P) = (A_n, A_k \times A_{n-k})$ and (D_n, A_{n-1}) , this map is given by deleting the pair of symbols in the i th and $(i+1)$ th positions in the coset diagram (for $\tau = s_{1''}$, delete the pair of symbols in position 1 and 2).

The explicit element $\psi_\tau(\lambda)$ in type (D_n, D_{n-1}) and exceptional types can be described in terms of tilings (see [BDHN, Proposition 5.4]) or deduced directly from the Bruhat graphs without much effort.

Theorem A.2. *Let (W, P) be a simply-laced Hermitian symmetric pair, $\tau \in S_W$ and $\lambda, \mu \in \mathcal{P}_{(W,P)}^\tau$. Then we have a degree-preserving bijection*

$$\text{Path}(\lambda, \mu) \rightarrow \text{Path}(\psi_\tau(\lambda), \psi_\tau(\mu)).$$

Proof. For types $(D_n, D_{n-1})^\tau = (A_1, A_0)$, $(E_6, D_5)^\tau = (A_5, A_4)$, and $(E_7, E_6)^\tau = (D_6, D_5)$ the result can be checked directly by examining the light leaves matrices given in Section 4. In types $(A_n, A_k \times A_{n-k-1})$ with $\tau = s_i \in S_W$ arbitrary and in type (D_n, A_{n-1}) and for any $\tau = s_i \neq s_{1''}$, note that $\lambda, \mu \in \mathcal{P}_{(W,P)}^\tau$ if and only if the i -th and $(i+1)$ -th vertex in their coset




FIGURE 33. The element λ from Figure 7 (with the vertices we will remove under the map ψ_τ highlighted in blue) and its image $\psi_\tau(\lambda)$. The tri-colouring of the rightmost node of the Coxeter graph on the righthand-side is explained in [BDHN] but is not important here.




FIGURE 34. The element μ from Figures 7 and 33 with its associated diagram e_μ and their image under ψ_τ .

diagram are labelled by $\vee \wedge$. This implies that the diagrams $e_T = \emptyset e_\mu \lambda$ for $T \in \text{Path}(\lambda, \underline{\mu})$ are precisely those which have an anti-clockwise oriented northern arc connecting i and $i+1$. In type (D_n, A_{n-1}) with $\tau = s_{1''}$, note that $\lambda, \mu \in \mathcal{P}_{(W,P)}^\tau$ if and only if the first and second vertex of their coset diagram are labelled by $\wedge \wedge$. This implies that the diagrams $e_T = \emptyset e_\mu \lambda$ for $T \in \text{Path}(\lambda, \underline{\mu})$ are precisely those which have flip-oriented decorated northern arc connecting the first and second vertex, both labelled by \wedge . For the purposes of this proof, we will only consider the top half of the diagram $\emptyset e_\mu \lambda$, for ease of exposition (the bottom half plays no significant role, see Proposition 8.3).

In all cases, the northern arc identified above joins two adjacent vertices and has degree 0. Now, if $(W, P) = (A_n, A_k \times A_{n-k-1})$, then removing this oriented arc produces an oriented Temperley-Lieb diagram of type $(A_{n-2}, A_{k-1} \times A_{n-k-2})$ of the same degree. If $(W, P) = (D_n, A_{n-1})$ and $\tau = s_{1''}$ then removing this flip-oriented decorated northern arc produces an oriented Temperley-Lieb diagram of type (D_{n-2}, A_{n-3}) of the same degree. Finally, if $(W, P) = (D_n, A_{n-1})$ and $\tau = s_i$ for $i \in \{1, \dots, n-1\}$ then removing this oriented (undecorated) northern arc produces an oriented Temperley-Lieb diagram of the same degree for the isomorphic copy of $TL_{(D_{n-2}, A_{n-3})}(q)$ where each coset diagram has an odd number of \wedge arrows (as discussed in Remark 5.3).

This map is clearly a bijection, as starting from any oriented Temperley-Lieb diagram for the anti-spherical module in rank $n-2$, we can insert the degree zero arc in the position corresponding to τ to obtain an oriented Temperley-Lieb diagram for the anti-spherical module in rank n and these maps are clearly inverse to each other. □

REFERENCES

- [BDHN] C. Bowman, M. De Visscher, A. Hazi, and E. Norton, *The anti-spherical Hecke categories for Hermitian symmetric pairs*, [arXiv:2208.02584](https://arxiv.org/abs/2208.02584).
- [Boe88] B. Boe, *Kazhdan-Lusztig polynomials for Hermitian symmetric spaces*, *Trans. Amer. Math. Soc.* **309** (1988), no. 1, 279–294.
- [BS10] J. Brundan and C. Stroppel, *Highest weight categories arising from Khovanov’s diagram algebra. II. Koszulity*, *Transform. Groups* **15** (2010), no. 1, 1–45.
- [BS11a] ———, *Highest weight categories arising from Khovanov’s diagram algebra I: cellularity*, *Mosc. Math. J.* **11** (2011), no. 4, 685–722, 821–822.
- [BS11b] ———, *Highest weight categories arising from Khovanov’s diagram algebra III: category \mathcal{O}* , *Represent. Theory* **15** (2011), 170–243.
- [BS12] ———, *Gradings on walled Brauer algebras and Khovanov’s arc algebra*, *Adv. Math.* **231** (2012), no. 2, 709–773.
- [CD11] A. Cox and M. De Visscher, *Diagrammatic Kazhdan-Lusztig theory for the (walled) Brauer algebra*, *J. Algebra* **340** (2011), 151–181.

- [Deo90] V. Deodhar, *A combinatorial setting for questions in Kazhdan-Lusztig theory*, *Geom. Dedicata* **36** (1990), no. 1, 95–119. MR 1065215
- [EHP14] T. Enright, M. Hunziker, and A. Prueett, *Diagrams of Hermitian type, highest weight modules, and syzygies of determinantal varieties*, *Symmetry: representation theory and its applications*, *Progr. Math.*, vol. 257, Birkhäuser/Springer, New York, 2014, pp. 121–184.
- [ES16a] M. Ehrig and C. Stroppel, *Diagrammatic description for the categories of perverse sheaves on isotropic Grassmannians*, *Selecta Math. (N.S.)* **22** (2016), no. 3, 1455–1536. MR 3518556
- [ES16b] M. Ehrig and C. Stroppel, *Koszul gradings on Brauer algebras*, *Int. Math. Res. Not. IMRN* (2016), no. 13, 3970–4011.
- [EW14] B. Elias and G. Williamson, *The Hodge theory of Soergel bimodules*, *Ann. of Math. (2)* **180** (2014), no. 3, 1089–1136.
- [Gre98] R. M. Green, *Generalized Temperley-Lieb algebras and decorated tangles*, *J. Knot Theory Ramifications* **7** (1998), no. 2, 155–171.
- [Kau87] L. H. Kauffman, *State models and the Jones polynomial*, *Topology* **26** (1987), 395–407.
- [LW] N. Libedinsky and G. Williamson, *The anti-spherical category*, [arXiv:1702.00459](https://arxiv.org/abs/1702.00459).
- [LW21] ———, *Kazhdan-Lusztig polynomials and subexpressions*, *J. Algebra* **568** (2021), 181–192. MR 4166587
- [Mar15] P. P. Martin, *The decomposition matrices of the Brauer algebra over the complex field*, *Trans. Amer. Math. Soc.* **367** (2015), no. 3, 1797–1825.
- [Soe97] W. Soergel, *Kazhdan-Lusztig polynomials and a combinatoric for tilting modules*, *Represent. Theory* **1** (1997), 83–114 (electronic).
- [Ste96] J. Stembridge, *On the fully commutative elements of Coxeter groups*, *J. Algebraic Combin.* **5** (1996), no. 4, 353–385.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, HESLINGTON, YORK, UK
Email address: `Chris.Bowman-Scargill@york.ac.uk`

DEPARTMENT OF MATHEMATICS, CITY, UNIVERSITY OF LONDON, LONDON, UK
Email address: `Maud.DeVisscher@city.ac.uk`

LEIBNIZ UNIVERSITÄT HANNOVER, WELFENGARTEN 1, D-30167 HANNOVER, GERMANY
Email address: `nifarrel@tcd.ie`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, HESLINGTON, YORK, UK
Email address: `Amit.Hazi@york.ac.uk`

SCHOOL OF MATHEMATICS, STATISTICS AND ACTUARIAL SCIENCE, UNIVERSITY OF KENT, CANTERBURY, UK
Email address: `E.Norton@kent.ac.uk`