



## City Research Online

### City, University of London Institutional Repository

---

**Citation:** Simmonds, M., Llewellyn, A., Walker, R., Fulbright, H., Walton, M., Hodgson, R., Bojke, L., Stewart, L., Dias, S., Rush, T., et al (2024). Anti-VEGF drugs compared with laser photocoagulation for the treatment of diabetic retinopathy: a systematic review and meta-analysis. *Health Technology Assessment*, pp. 1-71. doi: 10.3310/pcgv5709

This is the published version of the paper.

This version of the publication may differ from the final published version.

---

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/34327/>

**Link to published version:** <https://doi.org/10.3310/pcgv5709>

**Copyright:** City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

**Reuse:** Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

---

City Research Online:

<http://openaccess.city.ac.uk/>

[publications@city.ac.uk](mailto:publications@city.ac.uk)

---



## Research Article

# Anti-VEGF drugs compared with laser photocoagulation for the treatment of diabetic retinopathy: a systematic review and meta-analysis

Mark Simmonds<sup>1\*</sup>, Alexis Llewellyn<sup>1</sup>, Ruth Walker<sup>1</sup>, Helen Fulbright<sup>1</sup>, Matthew Walton<sup>1</sup>, Rob Hodgson<sup>1</sup>, Laura Bojke<sup>2</sup>, Lesley Stewart<sup>1</sup>, Sofia Dias<sup>1</sup>, Thomas Rush<sup>3</sup>, John G Lawrenson<sup>4</sup>, Tunde Peto<sup>5</sup> and David Steel<sup>6</sup>

<sup>1</sup>Centre for Reviews and Dissemination, University of York, York, UK

<sup>2</sup>Centre for Health Economics, University of York, York, UK

<sup>3</sup>Patient Representative, Belfast, UK

<sup>4</sup>Department of Optometry and Visual Sciences, City University of London, London, UK

<sup>5</sup>Centre for Public Health, Queen's University Belfast, Belfast, UK

<sup>6</sup>Biosciences Institute, Newcastle University, Newcastle upon Tyne, UK

\*Corresponding author [mark.simmonds@york.ac.uk](mailto:mark.simmonds@york.ac.uk)

Published December 2024

DOI: 10.3310/PCGV5709

## Abstract

**Background:** Diabetic retinopathy is a major cause of sight loss in people with diabetes. The most severe form, proliferative diabetic retinopathy, carries a high risk of vision loss, vitreous haemorrhage, macular oedema and other harms. Panretinal photocoagulation is the primary treatment for proliferative diabetic retinopathy. Anti-vascular endothelial growth factor drugs are used to treat various eye conditions and may be beneficial for people with diabetic retinopathy.

**Objective:** To investigate the efficacy and safety of anti-vascular endothelial growth factor therapy for the treatment of diabetic retinopathy when compared to panretinal photocoagulation.

**Methods:** A systematic review and network meta-analysis of all published randomised controlled trials comparing anti-vascular endothelial growth factor (alone or in combination with panretinal photocoagulation) to panretinal photocoagulation in people with diabetic retinopathy. The database searches were updated in May 2023. Trials where the primary focus was treatment of macular oedema or vitreous haemorrhage were excluded.

**Results:** A total of 14 trials were included: 3 of aflibercept, 5 of bevacizumab and 6 of ranibizumab. Two trials were of patients with non-proliferative diabetic retinopathy; all others were in proliferative diabetic retinopathy. Overall, anti-vascular endothelial growth factor was slightly better than panretinal photocoagulation at preventing vision loss, measured as best corrected visual acuity, at up to 2 years follow-up [mean difference in the logarithm of the minimum angle of resolution  $-0.089$  (or 3.6 Early Treatment Diabetic Retinopathy Study letters), 95% confidence interval  $-0.180$  to  $-0.019$ ]. There was no clear evidence of any difference between the anti-vascular endothelial growth factors, but the potential for bias complicated the comparison. One trial found no benefit of anti-vascular endothelial growth factor over panretinal photocoagulation after 5 years. Anti-vascular endothelial growth factor was superior to panretinal photocoagulation at preventing macular oedema (relative risk 0.29, 95% confidence interval 0.18 to 0.49) and vitreous haemorrhage (relative risk 0.77, 95% confidence interval 0.61 to 0.99). There was no clear evidence that the effectiveness of anti-vascular endothelial growth factor varied over time.

**Conclusions:** Anti-vascular endothelial growth factor injections reduce vision loss when compared to panretinal photocoagulation, but the benefit is small and unlikely to be clinically meaningful. Anti-vascular endothelial growth factor may have greater benefits for preventing complications such as macular oedema. Observational studies extending follow-up beyond the 1-year duration of most trials are needed to investigate the longer-term effects of repeated anti-vascular endothelial growth factor injections.

**Funding:** This article presents independent research funded by the National Institute for Health and Care Research (NIHR) Health Technology Assessment programme as award number NIHR132948. A plain language summary of this research article is available on the NIHR Journals Library Website <https://doi.org/10.3310/PCGV5709>.

## Background

Diabetes is a major cause of poor health that affects over 4 million people in the UK. Older people, men, people of South Asian ethnicity and more deprived populations are at higher risk.<sup>1</sup> Diabetic retinopathy is a 'chronic progressive, potentially sight-threatening disease of the retinal microvasculature'<sup>2,3</sup> that is a major complication of diabetes and a common cause of sight loss. Diabetic retinopathy impairs the sight of more than 1700 people in the UK each year.<sup>4</sup> The most severe form, proliferative diabetic retinopathy (PDR), places the patients at a high risk of vitreous haemorrhage, retinal detachment, neovascular glaucoma and vision loss.<sup>5,6</sup>

Panretinal (laser) photocoagulation (PRP) is the primary treatment for PDR, where a laser is applied to vascular abnormalities to prevent proliferation of new blood vessels or encourage regression in those with established new vessels. PRP is delivered over the entire periphery of the retina, by placing 1200–1600 laser burns per session, usually over two or three treatment sessions. It is known to be effective and long-lasting<sup>7</sup> but can have side effects including peripheral visual field loss, impaired night time and colour vision, and blurred vision. There is a small risk of central scotomata if laser burns are inadvertently placed at or near the foveal centre or if the laser scar extends centrally.<sup>8</sup>

Anti-vascular endothelial growth factor (anti-VEGF) drugs have been proposed as alternative to PRP. In the UK, the National Institute for Health and Care Excellence (NICE) has approved ranibizumab and aflibercept for the treatment of diabetic macular oedema (DMO),<sup>9,10</sup> and they are the standard treatment for wet age-related macular degeneration. However, whether they are beneficial for the treatment of diabetic retinopathy remains to be established. There are concerns that effects may not be long-lasting, and patients may have worse outcomes than those who had laser photocoagulation without repeated re-treatment and long-term follow-up.<sup>11,12</sup> They have rare but potentially serious adverse effects including: ocular hypertension, retinal detachment, endophthalmitis and other intraocular inflammation, and cataracts.<sup>13</sup>

International Council of Ophthalmology guidelines on diabetic eye care<sup>14</sup> support laser photocoagulation and 'appropriate use of anti-VEGF drugs' for the management of diabetic retinopathy. When this project commenced,

there was no current NICE guidance for the use of anti-VEGF drugs in people with diabetic retinopathy but without macular oedema. NICE guidance is under development,<sup>15</sup> and this review and meta-analysis was conducted to help inform it.

Given the uncertainty around whether anti-VEGF should be used to treat diabetic retinopathy, and the need for clear guidance on this topic, a systematic assessment of the relevant evidence and appropriate synthesis were needed. In order to synthesise data from mixed comparator studies, a network meta-analysis (NMA) approach was required to assess the value, effectiveness and rank of all relevant anti-VEGF interventions.

This paper presents a systematic review and NMA of all published randomised controlled trials (RCTs) of the three main anti-VEGFs used to treat diabetic retinopathy: aflibercept, bevacizumab and ranibizumab. While all three drugs act similarly to inhibit VEGF and slow the growth of blood vessels in the eye, they are different at molecular and receptor level, and so may differ in both efficacy and safety. This is why it is important to compare the three anti-VEGFs in a NMA.

The project was funded by the National Institute for Health and Care Research (Project number NIHR132948). The main project included a systematic review and meta-analysis incorporating individual patient data (IPD) from high-quality trials. Other components of the project included a wider assessment of anti-VEGF studies, including non-randomised studies, and an economic analysis of the cost-effectiveness of using anti-VEGF to treat diabetic retinopathy. The review was registered on PROSPERO (CRD42021272642) and the full protocol is available online from the NIHR (<https://fundingawards.nihr.ac.uk/award/NIHR132948>).

## Methods

The aim of this project was to systematically review all RCTs where anti-VEGFs were used to treat diabetic retinopathy. The review was conducted following the Centre for Reviews and Dissemination guidance on undertaking systematic reviews<sup>16</sup> and reported according to the principles of the overarching Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement.<sup>17</sup>

### Inclusion criteria

All RCTs that recruited people with diabetic retinopathy (proliferative and non-proliferative); patients with a principal indication for treatment of DMO or vitreous haemorrhage were excluded. The technologies of interest were any anti-VEGF therapy, anti-VEGF combined with PRP, PRP alone and sham injection.

A full list of outcomes of interest were reported in the review protocol. This paper focuses particularly on best corrected visual acuity (BCVA), as this was the only outcome reported in all trials. The appendices to this paper report evidence on all protocol-specified outcomes reported in the trials.

### Review methods

An Information Specialist (HF) designed a preliminary search strategy in Ovid MEDLINE which consisted of terms for the condition (diabetic retinopathy), that were combined with terms for the intervention (anti-VEGF, angiogenesis inhibitors, or specific drugs used for the treatment of diabetic retinopathy). A RCT study filter was applied. No date or language limits were applied. The final MEDLINE strategy was adapted for use in all resources searched. All search strategies are presented in full in [Appendix 1](#).

The searches were performed on 27 August 2021 and were updated on 13 July 2022 and again on 26 May 2023. The following databases were searched: Ovid MEDLINE(R) ALL, EMBASE (Ovid), Science Citation Index Expanded (Web of Science), Conference Proceedings Citation Index Science (Web of Science), Cochrane Central Register of Controlled Trials [CENTRAL (Wiley)], Cochrane Database of Systematic Reviews (Wiley), Database of Abstracts of Reviews of Effects [DARE [Centre for Reviews and Dissemination (CRD)]], PROSPERO (CRD) and Epistemonikos. The following trial registries were searched: World Health Organization (WHO) International Clinical Trials Registry Platform (ICTRP), ClinicalTrials.gov and the EU Clinical Trials Registry. Search results were imported into EndNote 20 (Clarivate Analytics, Philadelphia, PA, USA) and deduplicated.

Two researchers (RW, AL) independently screened all titles and abstracts retrieved for consideration of the full text. The reviewers then screened full texts of potentially eligible studies to determine inclusion. Disagreements were resolved through discussion or with a third reviewer (MS).

A data extraction form was developed and piloted. Data on interventions used, patient characteristics, outcomes

reported, and all outcome data were extracted for all included publications. Data extraction was completed by one reviewer and checked by a second (RW, AL). Risk of bias in all included trials was assessed using the Cochrane Risk of Bias 2 tool, focusing on the BCVA outcome, given limited reporting of other outcomes.<sup>18</sup>

### Statistical analysis

Effect estimates were pooled across trials using standard DerSimonian–Laird random-effect pairwise meta-analyses, according to the duration of follow-up. Heterogeneity was assessed in terms of  $I^2$ <sup>19</sup> and by inspecting the between-study heterogeneity standard deviations (SDs;  $\tau$ ), relative to the treatment effect size.

Network meta-analyses were performed using standard Bayesian methods of NMA in R (version 4.3.1, The R Foundation for Statistical Computing, Vienna, Austria) using the R package *multinma* (version 0.5.1).<sup>11,20</sup> This extends the standard NMA modelling approach to investigate the potential impact of patient factors (e.g. type of retinopathy) and timing of assessments on the effectiveness of anti-VEGF therapy.<sup>20</sup> Network consistency was checked by comparing the model fit and between-study heterogeneity from the NMA models to an unrelated mean effects model (similar to a model performing direct meta-analysis for each treatment comparison, but with a shared heterogeneity parameter).<sup>21</sup>

Visual acuity (BCVA) in diabetic retinopathy is commonly measured using the logarithm of the minimum angle of resolution (log-MAR) and Early Treatment Diabetic Retinopathy Study (ETDRS) scales. As both are widely used, NMAs were performed for both scales. Published data were transformed from one scale to the other, as required. This paper presents results on the log-MAR scale, with ETDRS results reported in the appendices.

The potential impact of unpublished or ongoing trials on the NMAs was investigated using threshold analysis. Threshold analysis investigates where in a NMA results might not be robust to changes in the observed evidence.<sup>22</sup>

All R code and data used for this paper are available on GitHub ([github.com/marksimmondsyork/AVID](https://github.com/marksimmondsyork/AVID)).

### Patient and public involvement

Patient and clinical representatives were involved in all stages of this project as part of our advisory group including: the funding application, protocol development, discussing the review and its findings, and writing this

paper. Further patient and stakeholder involvement was engaged through the NICE committee currently developing guidance on diabetic retinopathy management.

### Equality, diversity and inclusion

As this was a review project of existing trial data, we could not account for equality issues in this field beyond what was reported in included publications or data. We note that reporting on potential equality areas such as ethnicity or socioeconomics was limited.

## Results

### General results

Key findings for BCVA, DMO, vitrectomy, vitreous haemorrhage and adverse events are presented here. A full presentation of all analyses performed for all outcomes is provided in the appendices.

[Figure 1](#) shows the PRISMA flow chart for this review. Studies excluded from the review are listed in [Appendix 1](#). Overall, 14 RCTs were included in the meta-analyses. The searches also identified 21 other RCTs, which were unsuitable for meta-analyses. These included trials reported only as conference abstracts, not in English, published before 2010 (and therefore judged to be out-of-date), that used types of anti-VEGF not in widespread use, or did not include a PRP arm. Those trials therefore could not be reasonably included in the NMAs. These are summarised in [Appendix 1](#).

The included RCTs are summarised in [Table 1](#). Trials varied substantially in sample size from only 40 eyes up to just over 400 persons. There were six trials of ranibizumab, five of bevacizumab and three trials of aflibercept. Five trials used anti-VEGF as the intervention, while nine used anti-VEGF combined with PRP. Twelve trials were of patients with proliferative retinopathy. Two trials recruited patients with non-proliferative retinopathy; both evaluated aflibercept.<sup>23,24</sup> Trials of aflibercept and ranibizumab were conducted in Europe, North America or Brazil. All trials of bevacizumab were conducted in the Middle East or South Asia. BCVA was the only outcome reported consistently in all trials.

### Risk of bias

For the risk-of-bias assessment of the included trials, see [Table 2](#) and [Appendix 1](#). Overall, four trials were classed at low risk of bias, three moderate and seven at high risk of bias. Risk of bias across individual domains was predominately of 'some concerns', primarily due to poor reporting, although larger trials tended to be better reported. Concerns were

most common for the outcome measurement domain, due to the lack of masking of participants and outcome assessors. Other concerns included limited description of randomisation and allocation concealment processes, and missing patients and outcome data. The direction of bias was generally unpredictable. Overall, all the trials of bevacizumab were judged to be at high risk of bias. Only the larger trials of ranibizumab and aflibercept were at low risk of bias.

### Impact on vision (best corrected visual acuity)

[Figure 2](#) summarises all the data on BCVA for anti-VEGF compared to PRP, as reported across all trials. Results are shown as difference in ETDRS letters between anti-VEGF and control arms. This plot highlights significant variation in the design of the included studies, which precludes combining them all in a standard meta-analysis and demonstrates the need for NMA and meta-regression. First, some trials compare anti-VEGF to PRP directly, while others combine anti-VEGF with PRP, therefore motivating the need for NMA. Second, the time at which BCVA is measured varied enormously across trials, from 1 month to five years. Shorter trials were generally smaller in size, more likely to use bevacizumab and possibly showed larger effect sizes.


### Network meta-analyses of best corrected visual acuity in proliferative retinopathy

Given the variations in timing at which BCVA results were reported, two NMAs were performed:

1. Analysis up to and including 1 year of follow-up, using the longest follow-up in each trial
2. Analysis only of trials with 1 or 2 years' follow-up

Note that trials reporting at exactly 1 year were included in both analyses. Given the clinical differences between proliferative and non-proliferative disease, the two trials of non-proliferative disease were not included in the NMA. The network diagrams for both analyses are shown in [Figure 3](#). The size of the circles indicates the number of participants, and the width of the lines and the number of trials. Note that all the trials of bevacizumab combined with PRP had follow-up durations of < 1 year, so are not included in the analyses at 1–2 years. In both networks, there is only one trial of aflibercept and one of bevacizumab (without PRP).

[Figure 4](#) shows the results of all treatment comparisons from the NMA for data up to 1 year, and [Figure 5](#) for data from 1 to 2 years. Full results of these NMAs are given in [Appendix 2](#). In both figures, the point estimates



**FIGURE 1** Preferred Reporting Items for Systematic Reviews and Meta-Analyses flow diagram. DMO, diabetic macular edema; VH; vitreous haemorrhage; MA, meta-analysis.

are shown by the dots, with the horizontal lines being 95% credible intervals (CrIs). Negative relative effects (to the left of the vertical line) indicate favouring the first-named intervention.

For the primary comparisons with PRP at up to 1 year, all trials favoured anti-VEGF over PRP and improved vision (reduced log-MAR scores). Changes in log-MAR scores

when compared to PRP ranged from  $-0.078$  (or 3.8 ETDRS letters) for ranibizumab with PRP to  $-0.198$  (or 6.8 ETDRS letters) for bevacizumab. Results for aflibercept and bevacizumab (without PRP) were inconclusive because there was only one trial of each. Indirect comparisons between anti-VEGFs found no conclusive evidence that any one anti-VEGF was superior to the others. Heterogeneity across the network appeared to be modest,

TABLE 1 Summary of the RCTs included in the meta-analyses

Trial	Year	Anti-VEGF	Comparator	Location	Sample size	Follow-up	Population	Main outcome(s)
CLARITY <sup>23</sup>	2017	Aflibercept	PRP	UK	232 persons	1 year	PDR	BCVA, diabetic retinopathy severity, subsequent treatment, complications
DRCRN Protocol W <sup>24</sup>	2021	Aflibercept	Sham injection	USA/Canada	328 persons	2 years	Severe non-proliferative diabetic retinopathy (some DMO)	Time to proliferative diabetic retinopathy or DMO
PANORAMA <sup>25</sup>	2018	Aflibercept (every 16 weeks vs. 8 weeks)	Sham injection	International	402 persons	1 and 2 years	non-proliferative diabetic retinopathy	DR severity, subsequent treatment, complications
Marashi <sup>26</sup>	2017	Bevacizumab	PRP	Jordan/Syria	30 persons	1 year	PDR	BCVA, DR severity
Ahmad <sup>27</sup>	2012	Bevacizumab + PRP	PRP	Pakistan	54 eyes	3 months	PDR	BCVA
Ali <sup>28</sup>	2018	Bevacizumab + PRP	PRP	Pakistan	60 eyes	1 month	PDR	BCVA
Rebecca <sup>29</sup>	2021	Bevacizumab + PRP	PRP	Pakistan	76 eyes	6 months	PDR	BCVA
Roohipoor <sup>30</sup>	2016	Bevacizumab + PRP	PRP	Iran	64 eyes	10 months	PDR	BCVA
DRCRN Protocol S <sup>31</sup>	2018	Ranibizumab	PRP	USA	305 persons	2 and 5 years	PDR	DR severity, functional impact on vision, subsequent treatment, complications
Ferraz <sup>32</sup>	2015	Ranibizumab + PRP	PRP	Brazil	60 eyes	6 months	PDR	BCVA
PRIDE <sup>33</sup>	2019	Ranibizumab + PRP	PRP	Germany	106 persons	1 year	PDR	BCVA, DR severity, subsequent treatment
PROTEUS <sup>34</sup>	2018	Ranibizumab + PRP	PRP	Europe	87 persons	1 year	PDR	BCVA, subsequent treatment, complications
Sao Paulo B <sup>35</sup>	2011	Ranibizumab + PRP	PRP	Brazil	40 persons	1 year	PDR	BCVA, pain
Sao Paulo A <sup>36</sup>	2018	Ranibizumab + PRP (ETRDS)	Ranibizumab + PRP (PASCAL)	Brazil	40 eyes	1 year	PDR	BCVA

TABLE 2 Cochrane risk-of-bias assessment of outcome BCVA in the included RCTs

Trial	Risk-of-bias domain					Overall
	Randomisation	Deviation from intended intervention	Missing outcome data	Outcome measurement	Selective reporting	
Ahmad	!	!	+	-	!	High
Ali <sup>28</sup>	!	!	!	-	!	High
CLARITY <sup>23</sup>	+	+	+	!	+	Low
Ferraz <sup>32</sup>	!	!	+	+	!	Moderate
Marashi <sup>26</sup>	-	!	!	-	+	High
PANORAMA <sup>25</sup>	+	+	!	+	+	Low



TABLE 2 Cochrane risk-of-bias assessment of outcome BCVA in the included RCTs (continued)

Trial	Risk-of-bias domain					Overall
	Randomisation	Deviation from intended intervention	Missing outcome data	Outcome measurement	Selective reporting	
PRIDE <sup>33</sup>	!	+	!	-	+	Moderate
PROTEUS <sup>34</sup>	!	+	!	-	+	Moderate
Protocol S <sup>31</sup>	+	+	+	!	+	Low
Protocol W <sup>24</sup>	+	+	+	!	+	Low
Rebecca <sup>29</sup>	+	!	!	-	!	High
RECOVERY	!	+	+	-	+	Moderate
Roohipoor <sup>30</sup>	+	!	-	-	!	High
Sao Paulo A <sup>36</sup>	!	!	!	-	!	High
Sao Paulo B <sup>35</sup>	!	!	!	-	!	High
	+	Low risk				
	!	Some concerns				
	-	High risk				




FIGURE 2 All BCVA data (ETDRS letters) from all trials of anti-VEGF.

with an estimated heterogeneity standard error ( $\tau$ ) of 0.04 (95% CrI 0 to 0.14). Results for trials with a follow-up duration of 1–2 years (see Figure 5) were similar to those at up to 1 year, suggesting no obvious trend in treatment effects at up to 2 years.

Given the similarity in magnitude of effect for the various anti-VEGF agents compared to PRP, it is not surprising that the indirect comparisons between agents show no conclusive evidence of difference between any of them. There appears to be no difference between using

ranibizumab alone versus ranibizumab combined with PRP, particularly at 2 years.

Treatment rankings are shown in [Appendix 2 \(Figures 23 and 26\)](#). Given the similarity in effect sizes across the different types of anti-VEGF, it is difficult to draw conclusions from the ranking diagrams beyond the fact that PRP alone is likely to be the least effective treatment. The limited data on bevacizumab mean its ranking is very uncertain.

### Impact of follow-up time and vision at randomisation

To further examine the impact of follow-up time on the effectiveness of anti-VEGFs, we fitted a range of NMA models including time as a covariate. This meant that all trials could be combined in a single NMA, and whether the effectiveness of anti-VEGFs varied with time could be assessed. Models were also fitted including BCVA at randomisation, to account for possible variation in the effectiveness of the anti-VEGFs with initial vision (see [Appendix 2](#)).




FIGURE 3 Network diagrams at (a) up to 1 year and (b) 1–2 years.






FIGURE 4 Comparison of interventions from NMA of BCVA up to 1 year.



**FIGURE 5** Comparison of interventions from NMA of BCVA from 1 to 2 years.

Overall, results were very similar to the NMAs at up to 1 year and 1–2 years. [Figure 6](#) shows the effect estimates for anti-VEGFs compared to PRP alone from a model with

a linear association between anti-VEGF effect and both follow-up time and BCVA at randomisation. Estimates are presented for 1 year of follow-up and the mean BCVA



**FIGURE 6** Network meta-analysis of log-MAR with adjustment for follow-up time and BCVA at baseline.

at baseline across all trials (log-MAR 0.3). The pattern of effect sizes is very similar to that seen in [Figures 4 and 5](#), but CIs are wider, suggesting that adjusting for follow-up time and baseline BCVA leads to greater uncertainty.

There was no clear evidence that the relative effectiveness of anti-VEGFs varied with time or with vision at randomisation. However, it should be noted that almost all the data were for follow-up times of 2 years or less. Only one trial followed up patients for 5 years, and that found no evidence of difference between anti-VEGF (ranibizumab) and PRP after 5 years.<sup>31</sup>

### Further network meta-analyses

To further compare the anti-VEGFs to each other, simplified NMAs were performed by combining treatment arms. Two NMAs were performed:

1. Comparing anti-VEGF (of any type), anti-VEGF (any type) combined with PRP and PRP alone
2. Comparing aflibercept, ranibizumab (with or without PRP), bevacizumab (with or without PRP) and PRP alone

In both cases, NMAs included adjustment for follow-up time and BCVA at randomisation. Full results for these NMAs are presented in [Appendix 2](#). In summary, there was good evidence that, when all types of anti-VEGF were combined, anti-VEGF in general improved BCVA when compared to PRP (mean difference  $-0.089$ , 95% CrI  $-0.180$  to  $-0.019$ ), as did anti-VEGF combined with PRP compared to PRP alone (mean difference  $-0.108$ , 95% CrI  $-0.192$  to  $-0.048$ ).

When comparing the three anti-VEGFs (with or without concomitant PRP), there was no clear evidence of any difference in effectiveness between the three types of anti-VEGF; for example, there was no difference between aflibercept and ranibizumab (mean difference  $-0.003$ , 95% CI  $-0.166$  to  $0.163$ ).

### Threshold analysis

Threshold analyses of the NMAs of BCVA are reported in [Appendix 2](#). These found that the evidence for anti-VEGF

being superior to PRP was robust, but there was some uncertainty in the overall ranking of the various anti-VEGF treatments. This was probably because the evidence across the different anti-VEGFs showed very similar effectiveness.

### Other outcomes


Results on outcomes other than BCVA were inconsistently reported, with most being reported in no more than three trials. Complete results for these outcomes are presented in [Appendix 3](#). The limited data meant that NMAs were not feasible for these outcomes. A meta-analysis was performed for outcomes reported in two or more trials by assuming that the impact of anti-VEGFs is the same for all types of anti-VEGF, for anti-VEGF alone or in combination with PRP, and at all times up to 2 years. While these are strong assumptions, they may be reasonable given the results observed for BCVA, and the apparent lack of heterogeneity in the data.

Forest plots of neovascularisation of the disc (NVD) and neovascularisation elsewhere (NVE) are shown in [Appendix 3](#). These suggest that neovascularisation was reduced while using anti-VEGF. The results of meta-analyses for other non-vision outcomes are shown in [Figure 7](#). Although data were limited, the results suggest that anti-VEGF treatment substantially reduces the rate of macular oedema (DMO), the need for vitrectomy and reduces the rate of vitreous haemorrhage. No data on progression of diabetic retinopathy were reported.


### Adverse events

As with non-BCVA outcomes, adverse events were not widely reported, with little consistency across trials as to which adverse events were reported. A meta-analysis was performed for adverse event types reported in two or more trials by assuming that the impact of anti-VEGFs is the same for all types of anti-VEGF, for anti-VEGF alone or in combination with PRP, and at all times up to 2 years.

The meta-analysis results are shown in [Figure 8](#). Due to the small numbers of events, and limited numbers of trials reported each adverse event, most results are inconclusive. Anti-VEGF appeared to reduce the incidence of retinal detachment. It appeared to increase the



**FIGURE 7** Meta-analysis of non-vision outcomes. RR, relative risk.



**FIGURE 8** Meta-analyses of adverse event outcomes.

rate of ocular pain, but it was unclear whether this was procedure-related or post-intervention pain. Full results are presented in [Appendix 3](#).

### Non-proliferative retinopathy

Two trials compared aflibercept to sham injection in patients with non-proliferative retinopathy with a follow-up of 2 years (PANORAMA and Protocol W). Meta-analysis of their BCVA results found no clear evidence of any benefit of aflibercept over sham injection [mean difference (log-MAR)  $-0.02$ , 95% CI  $-0.05$  to  $0.01$ ]. Progression to macular oedema was the only other outcome reported by both trials, with strong evidence to suggest that aflibercept reduces the risk of macular oedema [relative risk (RR)  $0.283$ , 95% CI  $0.18$  to  $0.44$ ]. Protocol W reported that aflibercept reduced the rate of vitrectomy compared to sham injection (RR  $0.38$ , 95% CI  $0.24$  to  $0.60$ ). Full results are presented in [Appendix 4](#).

Protocol W found that aflibercept slowed the rate of progression to proliferative retinopathy when compared to sham injection (hazard ratio  $0.40$ , 97.5% CI  $0.28$  to  $0.57$ ). PANORAMA found that more patients on aflibercept experienced a 2 point or more improvement in Diabetic Retinopathy Severity Scale (DRSS) (RR  $4.41$ , 95% CI  $2.81$  to  $6.94$ ).

## Discussion

This systematic review included 14 trials of anti-VEGFs used to treat diabetic retinopathy. For patients with PDR, the NMA found good, but not conclusive, evidence that anti-VEGF therapy is better at maintaining vision than PRP therapy, with a benefit of around  $-0.089$  log-MAR (95% CI  $-0.179$  to  $-0.019$ ), or 3.6 ETDRS letters. This is within the region of variation that might be expected between eye tests without any intervention and is therefore unlikely

to be clinically meaningful.<sup>37</sup> There was no compelling evidence to suggest that the three anti-VEGFs (aflibercept, ranibizumab and bevacizumab) differ in effectiveness; observed differences might be due to different trial populations or potential for bias. There was no conclusive evidence that combining anti-VEGF injection with PRP therapy is more effective at improving vision than anti-VEGF alone. Anti-VEGF appears to have no impact on BCVA in people with non-proliferative disease.

A further issue is the impact of time on the effectiveness of anti-VEGF therapy. Our meta-analysis found no evidence that the effectiveness waned over the first 2 years after initialising therapy. However, the one trial with a longer follow-up (Protocol S) found no benefit of ranibizumab over PRP after 5 years.<sup>31</sup> The longer-term value of anti-VEGF therapy therefore needs further investigation, particularly regarding how anti-VEGF treatment should be repeated over long time periods. There was some evidence that the benefit of anti-VEGF over PRP may be greater in people with poorer vision at time of injection. However, it was not possible to draw any firm conclusions on this from data presented in trial publications alone.

Data on outcomes other than visual acuity were limited, and not reported consistently across trials. Given the variations in follow-up and interventions used, NMAs were not feasible, and meta-analyses had to make the strong assumption of no difference in effect between the three anti-VEGFs, and no variation over time. Given these limitations, there was some evidence that anti-VEGFs are more effective than PRP at preventing the most serious consequences of diabetic retinopathy. They reduced the incidence of macular oedema (in both PDR and NPDR patients) and vitreous haemorrhage. In patients with NPDR, there was some evidence that aflibercept slows the rate of progression to PDR and improves retinopathy severity. This suggests that

anti-VEGF may be valuable in preventing progression of diabetic retinopathy, even if its impact on vision directly is modest. Evidence on adverse events was limited due to inconsistent reporting, and small numbers of events. There was some evidence that anti-VEGF reduces the risk of retinal detachment.

Most trials were of short duration, with only one trial in PDR extending beyond 1 year. That trial found no vision benefit of anti-VEGF over PRP after 5 years, raising concerns as to the long-term efficacy of anti-VEGF therapy.

### **Patient and public perspectives**

Patient representatives noted several key areas of continued concern. Most critically was that most trials of anti-VEGF used BCVA as their primary outcome, without any consideration of how that impacted on quality of life, ability to work, drive or care for family. The lack of long-term evidence also raised concerns because there is substantial uncertainty about how PDR will be managed and treated long term.

## **Conclusion**

Anti-VEGF injection is only marginally better than PRP at maintaining vision and the benefit is unlikely to be clinically meaningful. There was no evidence of a difference in effectiveness between aflibercept, ranibizumab and bevacizumab, although data to compare these therapies were limited. There was no evidence to suggest that combining anti-VEGF with PRP improves effectiveness. Anti-VEGF may prevent, or delay, progression of macular oedema and vitreous haemorrhage. Some concern over bias in the trials remains.

The benefits of anti-VEGFs appear consistent for at least 2 years after initiation of treatment, but longer-term benefits are uncertain. There is some evidence that anti-VEGFs are less effective at maintaining visual acuity in people with less severe retinopathy, but this requires further investigation. Access to original individual-level trial data might aid in resolving this. Trials or observational studies of duration substantially longer than 1 year are needed to examine whether anti-VEGF may be beneficial in the long term, particularly with the requirement for long-term repeated anti-VEGF injections.

## **Additional information**

### **CRedit contribution statement**

**Mark Simmonds** (<https://orcid.org/0000-0002-1999-8515>): Conceptualisation (lead), Data curation (lead), Formal

analysis (lead), Funding acquisition (lead), Investigation (lead), Methodology (lead), Project administration, Writing (lead).

**Alexis Llewellyn** (<https://orcid.org/0000-0003-4569-5136>): Conceptualisation, Data curation, Formal analysis, Funding acquisition, Investigation (co-lead), Methodology, Writing.

**Ruth Walker** (<https://orcid.org/0000-0003-2765-7363>): Conceptualisation, Data curation, Formal analysis, Funding acquisition, Investigation, Writing.

**Helen Fulbright** (<https://orcid.org/0000-0002-1073-1099>): Investigation, Methodology.

**Matthew Walton** (<https://orcid.org/0000-0003-1932-3689>): Conceptualisation, Funding acquisition, Writing.

**Rob Hodgson** (<https://orcid.org/0000-0001-6962-2893>): Conceptualisation, Funding acquisition, Writing.

**Laura Bojke** (<https://orcid.org/0000-0001-7921-9109>): Conceptualisation, Funding acquisition, Writing.

**Lesley Stewart** (<https://orcid.org/0000-0003-0287-4724>): Conceptualisation, Funding acquisition, Writing.

**Sofia Dias** (<https://orcid.org/0000-0002-2172-0221>): Conceptualisation, Funding acquisition, Methodology, Writing.

**Thomas Rush**: Conceptualisation, Funding acquisition, Writing (patient and public involvement advisor).

**John G Lawrenson**: Conceptualisation, Funding acquisition, Writing.

**Tunde Peto** (<https://orcid.org/0000-0001-6265-0381>): Conceptualisation, Funding acquisition, Writing.

**David Steel** (<https://orcid.org/0000-0001-8734-3089>): Conceptualisation, Funding acquisition, Writing.

### **Acknowledgements**

We acknowledge the help and advice given by all persons involved in the NICE diabetic retinopathy guidance development process.

### **Data-sharing statement**

Data and code to reproduce the meta-analyses are available on GitHub (<https://github.com/marksimmondsyork/AVID>). For all other data requests please contact the corresponding author.

### **Ethics statement**

As this was a systematic review of existing published data, no ethics approval was required.

### Information governance statement

All data used in this paper were taken from published sources: no personal data were included.

### Disclosure of interests

**Full disclosure of interests:** Completed ICMJE forms for all authors, including all related interests, are available in the toolkit on the NIHR Journals Library report publication page at <https://doi.org/10.3310/PCGV5709>.

**Primary conflicts of interest:** Laura Bojke declares that she was on the HS&DR Researcher-Led awards panel (December 2019–December 2022). All other authors have no conflicts of interest to declare.

### Department of Health and Social Care disclaimer

This publication presents independent research commissioned by the National Institute for Health and Care Research (NIHR). The views and opinions expressed by authors in this publication are those of the authors and do not necessarily reflect those of the NHS, the NIHR, MRC, NIHR Coordinating Centre, the Health Technology Assessment programme or the Department of Health and Social Care.

This article was published based on current knowledge at the time and date of publication. NIHR is committed to being inclusive and will continually monitor best practice and guidance in relation to terminology and language to ensure that we remain relevant to our stakeholders.

### Study registration

This study is registered as PROSPERO (CRD42021272642).

### Funding

This article presents independent research funded by the National Institute for Health and Care Research (NIHR) Health Technology Assessment programme as award number NIHR132948.

This article reports on one component of the research award Anti-VEGF drugs compared with laser photocoagulation for the treatment of diabetic retinopathy: a systematic review and economic analysis. Other articles published as part of this thread are: [\[LINKS to other articles\]](#). For more information about this research please view the award page [<https://fundingawards.nihr.ac.uk/award/NIHR132948>]

### About this article

The contractual start date for this research was in August 2021. This article began editorial review in November 2023 and was accepted for publication in August 2024. The authors have been wholly responsible for all data collection, analysis

and interpretation, and for writing up their work. The Health Technology Assessment editors and publisher have tried to ensure the accuracy of the authors' article and would like to thank the reviewers for their constructive comments on the draft document. However, they do not accept liability for damages or losses arising from material published in this article.

### Copyright

Copyright © 2024 Simmonds *et al.* This work was produced by Simmonds *et al.* under the terms of a commissioning contract issued by the Secretary of State for Health and Social Care. This is an Open Access publication distributed under the terms of the Creative Commons Attribution CC BY 4.0 licence, which permits unrestricted use, distribution, reproduction and adaptation in any medium and for any purpose provided that it is properly attributed. See: <https://creativecommons.org/licenses/by/4.0/>. For attribution the title, original author(s), the publication source – NIHR Journals Library, and the DOI of the publication must be cited.

### List of abbreviations

anti-VEGF	anti-vascular endothelial growth factor
BCVA	best corrected visual acuity
CENTRAL	Cochrane Central Register of Controlled Trials
CRD	Centre for Reviews and Dissemination
DARE	Database of Abstracts of Reviews of Effects
DMO	diabetic macular oedema
DRSS	Diabetic Retinopathy Severity Scale
ETDRS	Early Treatment Diabetic Retinopathy Study
ICTRP	International Clinical Trials Registry Platform
IPD	individual patient data
log-MAR	logarithm of the minimum angle of resolution
NICE	National Institute for Health and Care Excellence
NMA	network meta-analysis
NVD	neovascularisation of the disc
NVE	neovascularisation elsewhere

PDR	proliferative diabetic retinopathy
PRISMA	Preferred Reporting Items for Systematic Reviews and Meta-Analyses
PRP	panretinal photocoagulation
RCT	randomised controlled trial
WHO	World Health Organization

## References

- Mathur R, Bhaskaran K, Edwards E, Lee H, Chaturvedi N, Smeeth L, Douglas I. Population trends in the 10-year incidence and prevalence of diabetic retinopathy in the UK: a cohort study in the Clinical Practice Research Datalink 2004–2014. *BMJ Open* 2017;**7**:e014444.
- Ghanchi F; Diabetic Retinopathy Guidelines Working Group. The Royal College of Ophthalmologists' clinical guidelines for diabetic retinopathy: a summary. *Eye (Lond)* 2013;**27**:285–7.
- Klein R, Knudtson MD, Lee KE, Gangnon R, Klein BE. The Wisconsin Epidemiologic Study of Diabetic Retinopathy: XXII the twenty-five-year progression of retinopathy in persons with type 1 diabetes. *Ophthalmology* 2008;**115**:1859–68.
- Flaxman SR, Bourne RRA, Resnikoff S, Ackland P, Braithwaite T, Cicinelli MV, *et al.*; Vision Loss Expert Group of the Global Burden of Disease Study. Global causes of blindness and distance vision impairment 1990–2020: a systematic review and meta-analysis. *Lancet Glob Health* 2017;**5**:e1221–34. [https://doi.org/10.1016/s2214-109x\(17\)30393-5](https://doi.org/10.1016/s2214-109x(17)30393-5)
- The Diabetic Retinopathy Study Research Group. Four risk factors for severe visual loss in diabetic retinopathy. The third report from the Diabetic Retinopathy Study. *Arch Ophthalmol* 1979;**97**:654–5.
- Parikh R, Shah RJ, VanHouten JP, Cherney EF. Ocular findings at initial pan retinal photocoagulation for proliferative diabetic retinopathy predict the need for future pars plana vitrectomy. *Retina* 2014;**34**:1997–2002.
- Early Treatment Diabetic Retinopathy Study Research Group. Fundus photographic risk factors for progression of diabetic retinopathy. ETDRS report number 12. *Ophthalmology* 1991;**98**:823–33.
- Royle P, Mistry H, Auguste P, Shyangdan D, Freeman K, Lois N, Waugh N. Pan-retinal photocoagulation and other forms of laser treatment and drug therapies for non-proliferative diabetic retinopathy: systematic review and economic evaluation. *Health Technol Assess* 2015;**19**:v–xxviii, 1–247.
- National Institute for Health and Care Excellence. *Aflibercept for Treating Diabetic Macular Oedema*. NICE; 2015. URL: [www.nice.org.uk/guidance/ta346/resources/aflibercept-for-treating-diabetic-macular-oedema-pdf-82602611201221](http://www.nice.org.uk/guidance/ta346/resources/aflibercept-for-treating-diabetic-macular-oedema-pdf-82602611201221) (accessed 5 July 2023).
- National Institute for Health and Care Excellence. *Ranibizumab for Treating Diabetic Macular Oedema*. NICE; 2013. URL: [www.nice.org.uk/guidance/ta274/resources/ranibizumab-for-treating-diabetic-macular-oedema-pdf-82600612458181](http://www.nice.org.uk/guidance/ta274/resources/ranibizumab-for-treating-diabetic-macular-oedema-pdf-82600612458181) (accessed 5 July 2023).
- Wubben TJ, Johnson MW, Sohn EH, Peairs JJ, Kay CN, Kim SJ, *et al.* Anti-vascular endothelial growth factor therapy for diabetic retinopathy: consequences of inadvertent treatment interruptions. *Am J Ophthalmol* 2019;**204**:13–8.
- Obeid A, Su D, Patel SN, Uhr JH, Borkar D, Gao X, *et al.* Outcomes of eyes lost to follow-up with proliferative diabetic retinopathy that received panretinal photocoagulation versus intravitreal anti-vascular endothelial growth factor. *Ophthalmology* 2019;**126**:407–13.
- Royal National Institute of Blind People (RNIB). *Anti-VEGF Treatment*. RNIB. URL: [www.rnib.org.uk/eye-health/eye-conditions/anti-vegf-treatment](http://www.rnib.org.uk/eye-health/eye-conditions/anti-vegf-treatment) (accessed 5 July 2023).
- Wong TY, Sun J, Kawasaki R, Ruamviboonsuk P, Gupta N, Lansingh VC, *et al.* Guidelines on diabetic eye care: the International Council of Ophthalmology recommendations for screening, follow-up, referral, and treatment based on resource settings. *Ophthalmology* 2018;**125**:1608–22.
- National Institute for Health and Care Excellence. *Diabetic Retinopathy (Guidance in Development)*. 2023. URL: [www.nice.org.uk/guidance/indevelopment/gid-ng10256](http://www.nice.org.uk/guidance/indevelopment/gid-ng10256) (accessed 3 October 2023).
- Centre for Reviews and Dissemination. *Systematic Reviews: CRD's Guidance for Undertaking Reviews in Health Care*. URL: [www.york.ac.uk/media/crd/Systematic\\_Reviews.pdf](http://www.york.ac.uk/media/crd/Systematic_Reviews.pdf) (accessed 17 September 2020).
- Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med* 2009;**151**:264–9, W64.
- Sterne JAC, Savovic J, Page MJ, Elbers RG, Blencowe NS, Boutron I, *et al.* Rob 2: a revised tool for assessing risk of bias in randomised trials. *BMJ* 2019;**366**:l4898.



19. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;**327**:557–60.
20. Phillippo DM, Dias S, Ades AE, Belger M, Brnabic A, Schacht A, *et al.* Multilevel network meta-regression for population-adjusted treatment comparisons. *J R Stat Soc Ser A Stat Soc* 2020;**183**:1189–210.
21. Dias S, Welton NJ, Sutton AJ, Caldwell DM, Lu G, Ades AE. Evidence synthesis for decision making 4: inconsistency in networks of evidence based on randomized controlled trials. *Med Decis Making* 2013;**33**:641–56.
22. Phillippo DM, Dias S, Welton NJ, Caldwell DM, Taske N, Ades AE. Threshold analysis as an alternative to GRADE for assessing confidence in guideline recommendations based on network meta-analyses. *Ann Intern Med* 2019;**170**:538–46.
23. Sivaprasad S, Prevost AT, Vasconcelos JC, Riddell A, Murphy C, Kelly J, *et al.*; CLARITY Study Group. Clinical efficacy of intravitreal aflibercept versus panretinal photocoagulation for best corrected visual acuity in patients with proliferative diabetic retinopathy at 52 weeks (CLARITY): a multicentre, single-blinded, randomised, controlled, phase 2b, non-inferiority trial. *Lancet* 2017;**389**:2193–203.
24. Maturi RK, Glassman AR, Josic K, Antoszyk AN, Blodi BA, Jampol LM, *et al.*; DRCR Retina Network. Effect of intravitreal anti-vascular endothelial growth factor vs sham treatment for prevention of vision-threatening complications of diabetic retinopathy: the Protocol W randomized clinical trial. *JAMA Ophthalmol* 2021;**139**:701–12.
25. Brown DM, Wyckoff CC, Boyer D, Heier JS, Clark WL, Emanuelli A, *et al.* Evaluation of intravitreal aflibercept for the treatment of severe nonproliferative diabetic retinopathy: results from the PANORAMA randomized clinical trial. *JAMA Ophthalmol* 2021;**139**:946.
26. Marashi A, Abukhalaf I, Alfaraji R, Shuman Y, Salahia A. Panretinal photocoagulation versus intravitreal bevacizumab for proliferative diabetic retinopathy treatment. *Adv Ophthalmol Vis Syst* 2017;**7**:00211. <https://doi.org/10.15406/aovs.2017.07.00211>
27. Ahmad M, Jan S. Comparison between panretinal photocoagulation and panretinal photocoagulation plus intravitreal bevacizumab in proliferative diabetic retinopathy. *J Ayub Med Coll Abbottabad* 2012;**24**:10–3.
28. Ali W, Abbasi KZ, Raza A. Panretinal photocoagulation plus intravitreal bevacizumab versus panretinal photocoagulation alone for proliferative diabetic retinopathy. *J Coll Physicians Surg Pak* 2018;**28**:923–7.
29. Rebecca MR, Shaikh FF, Jatoi SM. Comparison of efficacy of combination therapy of an intravitreal injection of bevacizumab and photocoagulation versus pan retinal photocoagulation alone in high risk proliferative diabetic retinopathy. *Pak J Med Sci* 2021;**37**:157–61.
30. Roohipoor R, Sharifian E, Ghassemi F, Riazi-Esfahani M, Karkhaneh R, Fard MA, *et al.* Choroidal thickness changes in proliferative diabetic retinopathy treated with panretinal photocoagulation versus panretinal photocoagulation with intravitreal bevacizumab. *Retina* 2016;**36**:1997–2005.
31. Gross JG, Glassman AR, Jampol LM. Panretinal photocoagulation vs intravitreal ranibizumab for proliferative diabetic retinopathy: a randomized clinical trial (vol,314 pg 2137, 2015). *JAMA* 2019;**321**:1008.
32. Ferraz DA, Vasquez LM, Preti RC, Motta A, Sophie R, Bittencourt MG, *et al.* A randomized controlled trial of panretinal photocoagulation with and without intravitreal ranibizumab in treatment-naïve eyes with non-high-risk proliferative diabetic retinopathy. *Retina* 2015;**35**:280–7.
33. Lang GE, Stahl A, Voegeler J, Quiering C, Lorenz K, Spital G, Liakopoulos S. Efficacy and safety of ranibizumab with or without panretinal laser photocoagulation versus laser photocoagulation alone in proliferative diabetic retinopathy – the PRIDE study. *Acta Ophthalmol* 2020;**98**:e530–9.
34. Figueira J, Fletcher E, Massin P, Silva R, Bandello F, Midena E, *et al.*; EVICR.net Study Group. Ranibizumab plus panretinal photocoagulation versus panretinal photocoagulation alone for high-risk proliferative diabetic retinopathy (PROTEUS study). *Ophthalmology* 2018;**125**:691–700.
35. Filho JA, Messias A, Almeida FP, Ribeiro JA, Costa RA, Scott IU, Jorge R. Panretinal photocoagulation (PRP) versus PRP plus intravitreal ranibizumab for high-risk proliferative diabetic retinopathy. *Acta Ophthalmol* 2011;**89**:e567–72.
36. Messias K, Barroso RD, Jorge R, Messias A. Retinal function in eyes with proliferative diabetic retinopathy treated with intravitreal ranibizumab and multispot laser panretinal photocoagulation. *Doc Ophthalmol* 2018;**137**:121–9.
37. Rosser DA, Cousens SN, Murdoch IE, Fitzke FW, Laidlaw DA. How sensitive to clinical change are ETDRS logMAR visual acuity measurements? *Invest Ophthalmol Vis Sci* 2003;**44**:3278–81. <https://doi.org/10.1167/iovs.02-1100>

## Appendix 1 Systematic review processes

### Database search strategies

The aim of the literature search was to identify RCTs on anti-VEGFs, angiogenesis inhibitors and other specific drugs used for the treatment of diabetic retinopathy.

An Information Specialist (HF) designed a preliminary search strategy in Ovid MEDLINE in consultation with the research team. The strategy consisted of terms for the condition (diabetic retinopathy), which were combined with terms for the intervention (anti-VEGF, angiogenesis inhibitors, or specific drugs used for the treatment of diabetic retinopathy) using the Boolean operator AND. Text word searches for terms appearing in the title and abstracts of database records were included in the strategy alongside searches of relevant subject headings. A RCT study filter was applied using the Boolean operator AND. No date or language limits were applied. The final MEDLINE strategy was adapted for use in all resources searched.

The searches were performed on 27 August 2021. The following databases were searched: Ovid MEDLINE(R) ALL, EMBASE (Ovid), Science Citation Index Expanded (Web of Science), Conference Proceedings Citation Index Science (Web of Science), Cochrane CENTRAL (Wiley), Cochrane Database of Systematic Reviews (Wiley), DARE (CRD), PROSPERO (CRD) and Epistemonikos. The following trial registries were searched: WHO ICTRP, ClinicalTrials.gov, and the EU Clinical Trials Registry.

Search results were imported into EndNote 20 and deduplicated. All search strategies are presented in full below.

The searches were updated on 13 July 2022 and again on 26 May 2023 using all the databases and strategies as used previously, except for DARE as this database is no longer updated. For each update search, the results of the databases were deduplicated against each other in a separate EndNote 20 Library before being merged with the results of the original EndNote Library and deduplicated for a second time.

### Ovid MEDLINE(R) ALL

(Includes Epub Ahead of Print, In-Process & Other Non-Indexed Citations, Ovid MEDLINE Daily and Ovid MEDLINE)

via Ovid <http://ovidsp.ovid.com/>

Date range searched: <1946–25 May 2023>

Date searched: 26 May 2023

Records retrieved: 3172

The MEDLINE strategy below includes a search filter to limit retrieval to RCTs using the Cochrane Highly Sensitive Search Strategy for identifying randomised trials in MEDLINE: sensitivity-maximising version (2008 revision); Ovid format.

Lefebvre C, Glanville J, Briscoe S, Littlewood A, Marshall C, Metzendorf MI, *et al.* Technical Supplement to *Chapter 4: Searching for and Selecting Studies*. In Higgins JPT, Thomas J, Chandler J, Cumpston MS, Li T, Page MJ, Welch VA, editors. *Cochrane Handbook for Systematic Reviews of Interventions* Version 6.2 (updated February 2021). Cochrane, 2021. Available from: [www.training.cochrane.org/handbook](http://www.training.cochrane.org/handbook).

- 1 (\*Diabetes Mellitus/ or \*Diabetes Complications/) and exp \*Retinal Diseases/ (3199)
- 2 Diabetic Retinopathy/ (29304)
- 3 ((diabet\* or DM) adj3 (retinopath\* or vitreoretinopath\* or vitreo-retinopath\* or chorioretinopath\* or chorio-retinopath\* or maculopath\*)).ti,ab,kw. (30685)
- 4 (((proliferat\* or PDR or pre-proliferat\* or preproliferat\* or non-proliferat\* or nonproliferat\* or NPDR or background) adj3 (retinopath\* or vitreoretinopath\* or vitreo-retinopath\* or chorioretinopath\* or chorio-retinopath\*)) and (diabet\* or DM)).ti,ab,kw. (7895)
- 5 (new blood vessel\* and diabet\*).ti,ab,kw. (273)
- 6 (((retin\* or subretina\* or sub-retina\* or interretina\* or inter-retina\* or vitreoretin\* or vitreo-retin\* or chorioretin\* or chorio-retin\* or choroid\* or macula\* or intraocular or intra-ocular or intravitreal or intra-vitreous) adj4 (damage\* or deteriorat\* or degenerat\* or disease\* or edema or oedema or neovascularization\*)) and diabet\*).ti,ab,kw. (13654)
- 7 ((retinal vein\* adj3 (occlu\* or obstruct\* or clos\* or stricture\* or steno\* or block\* or emboli\*)) and diabet\*).ti,ab,kw. (1473)
- 8 or/1-7 (44519)
- 9 exp Vascular Endothelial Growth Factors/ai (9366)
- 10 exp Receptors, Vascular Endothelial Growth Factor/ai (3393)
- 11 (anti adj2 VEGF\*).ti,ab,kw. (9210)
- 12 (anti-VEGF\* or antiVEGF\*).ti,ab,kw. (9455)
- 13 ((anti vascular or anti-vascular or antivascular) adj2 endothelial growth factor\*).ti,ab,kw. (5745)
- 14 (((vascular endothelial adj2 growth factor\*) or vasculotropin or VEGF\* or vascular permeability factor\* or VPF) adj2 (trap\* or inhibit\* or antagonist\*)).ti,ab,kw. (11005)
- 15 (vascular proliferation adj4 inhibit\*).ti,ab,kw. (38)
- 16 or/9-15 (28125)

- 17 Angiogenesis Inhibitors/ (28876)  
 18 exp Angiogenesis Inducing Agents/ai (118)  
 19 (angiogen\* adj2 (antagonist\* or inhibit\*)).ti,ab,kw.  
 (14831)  
 20 ((antiangiogen\* or anti angiogen\* or anti-angiogen\*)  
 adj2 (agent\* or drug\* or effect\*)).ti,ab,kw. (10949)  
 21 (angiostatic adj2 (agent\* or drug\*)).ti,ab,kw. (103)  
 22 ((neovasculari?ation or vasculari?ation) adj2 inhibit\*).  
 ti,ab,kw. (1243)  
 23 or/17-22 (45139)  
 24 Aflibercept\*.ti,ab,kw,rn. (3315)  
 25 (Eylea or Zaltrap or Ziv-Aflibercept or "AVE 0005" or  
 AVE0005 or "AVE 005" or AVE005).ti,ab,kw. (316)  
 26 Bevacizumab/ (14139)  
 27 Bevacizumab\*.ti,ab,kw,rn. (22533)  
 28 (Avastin or Mvasi or Alymsys or Aybintio or Equida-  
 cent or Onbevzi or Oyavas or Zirabev or rhuMAb-  
 VEGF or rhuMAb-VEGF or rhuMAb VEGF or "NSC  
 704865" or NSC704865).ti,ab,kw. (1675)  
 29 (IVB adj2 inject\*).ti,ab,kw. (316)  
 30 Ranibizumab/ (4684)  
 31 Ranibizumab\*.ti,ab,kw,rn. (6307)  
 32 (Lucentis or "rhuFab V2").ti,ab,kw. (456)  
 33 (IVR adj2 inject\*).ti,ab,kw. (139)  
 34 Pegaptanib\*.ti,ab,kw,rn. (671)  
 35 ("EYE 001" or EYE001 or Macugen or "NX 1838" or  
 NX1838).ti,ab,kw. (140)  
 36 or/24-35 (28353)  
 37 8 and (16 or 23 or 36) (4979)  
 38 randomized controlled trial.pt. (593242)  
 39 controlled clinical trial.pt. (95314)  
 40 randomized.ab. (604126)  
 41 placebo.ab. (238387)  
 42 drug therapy.fs. (2592996)  
 43 randomly.ab. (408822)  
 44 trial.ab. (649200)  
 45 groups.ab. (2520111)  
 46 or/38-45 (5663345)  
 47 37 and 46 (3308)  
 48 exp animals/ not humans.sh. (5123796)  
 49 47 not 48 (3190)  
 50 limit 49 to yr="2000-Current" (3182)  
 51 remove duplicates from 50 (3172)

**Key:**

/ or.sh. = indexing term (Medical Subject Heading: MeSH)

/ai = indexing term with subheading for antagonists & inhibitors

exp = exploded indexing term (MeSH)

\* or \$ = truncation

? = adds up to 1 additional character

ti,ab,kw = terms in either title, abstract or keyword fields

rn = registry number/name of substance

adj3 = terms within three words of each other (any order).

pt = publication type

fs = floating sub-heading

**EMBASE**

via Ovid <http://ovidsp.ovid.com/>

Date range searched: <1974–25 May 2023>

Date searched: 26 May 2023

Records retrieved: 2558

The EMBASE strategy below includes the Cochrane EMBASE RCT filter (Ovid format).

Glanville J, Foxlee R, Wisniewski S, Noel-Storr A, Edwards M, Dooley G. Translating the Cochrane EMBASE RCT filter from the Ovid interface to EMBASE.com: a case study. *Health Info Libr J.* 2019. doi:10.1111/hir.12269

- 1 \*diabetes mellitus/ and exp \*retina disease/ (4826)
- 2 exp diabetic retinopathy/ (53891)
- 3 ((diabet\* or DM) adj3 (retinopath\* or vitreoretino-  
 path\* or vitreo-retinopath\* or chorioretinopath\*  
 or chorio-retinopath\* or maculopath\*)).ti,ab,kw.  
 (43573)
- 4 (((proliferat\* or PDR or pre-proliferat\* or preprolif-  
 erat\* or non-proliferat\* or nonproliferat\* or NPDR  
 or background) adj3 (retinopath\* or vitreoretino-  
 path\* or vitreo-retinopath\* or chorioretinopath\* or  
 chorio-retinopath\*)) and (diabet\* or DM)).ti,ab,kw.  
 (11148)
- 5 (new blood vessel\* and diabet\*).ti,ab,kw. (391)
- 6 (((retin\* or subretina\* or sub-retina\* or interretina\*  
 or inter-retina\* or vitreoretin\* or vitreo-retin\* or  
 chorioretin\* or chorio-retin\* or choroid\* or macu-  
 la\* or intraocular or intra-ocular or intravitreal or  
 intra-vitreol) adj4 (damage\* or deteriorat\* or degn-  
 erat\* or disease\* or edema or oedema or neovascu-  
 lari?ation\*)) and diabet\*).ti,ab,kw. (20734)
- 7 ((retinal vein\* adj3 (occlu\* or obstruct\* or clos\* or  
 stricture\* or steno\* or block\* or emboli\*)) and dia-  
 bet\*).ti,ab,kw. (2199)
- 8 or/1-7 (70501)
- 9 vasculotropin inhibitor/ (7663)

- 10 (anti adj2 VEGF\*).ti,ab,kw. (15751)
- 11 (anti-VEGF\* or antiVEGF\*).ti,ab,kw. (16291)
- 12 ((anti vascular or anti-vascular or antivascular) adj2 endothelial growth factor\*).ti,ab,kw. (7400)
- 13 (((vascular endothelial adj2 growth factor\*) or vasculotropin or VEGF\* or vascular permeability factor\* or VPF) adj2 (trap\* or inhibit\* or antagonist\*).ti,ab,kw. (17346)
- 14 (vascular proliferation adj4 inhibit\*).ti,ab,kw. (50)
- 15 or/9-14 (38838)
- 16 angiogenesis inhibitor/ (20415)
- 17 (angiogen\* adj2 (antagonist\* or inhibit\*).ti,ab,kw. (20444)
- 18 ((antiangiogen\* or anti angiogen\* or anti-angiogen\*) adj2 (agent\* or drug\* or effect\*).ti,ab,kw. (15734)
- 19 (angiostatic adj2 (agent\* or drug\*).ti,ab,kw. (125)
- 20 ((neovasculari?ation or vasculari?ation) adj2 inhibit\*).ti,ab,kw. (1718)
- 21 or/16-20 (45260)
- 22 aflibercept/ (8877)
- 23 Aflibercept\*.ti,ab,kw,dy,tn. (9141)
- 24 (Eylea or Zaltrap or Ziv-Aflibercept or "AVE 0005" or AVE0005 or "AVE 005" or AVE005).ti,ab,dy,tn. (1741)
- 25 bevacizumab/ (72890)
- 26 Bevacizumab\*.ti,ab,kw,dy,tn. (75152)
- 27 (Avastin or Mvasi or Alymsys or Aybintio or Equidacent or Onbevzi or Oyavas or Zirabev or rhuMab-VEGF or rhuMab-VEGF or rhuMab VEGF or "NSC 704865" or NSC704865).ti,ab,kw,dy,tn. (11007)
- 28 (IVB adj2 inject\*).ti,ab,kw. (395)
- 29 ranibizumab/ (12442)
- 30 Ranibizumab\*.ti,ab,kw,dy,tn. (12826)
- 31 (Lucentis or "rhuFab V2").ti,ab,kw,dy,tn. (3216)
- 32 (IVR adj2 inject\*).ti,ab,kw. (197)
- 33 pegaptanib.dy,tn. (2470)
- 34 Pegaptanib\*.ti,ab,kw,dy,tn. (2544)
- 35 ("EYE 001" or EYE001 or Macugen or "NX 1838" or NX1838).ti,ab,kw,dy,tn. (1266)
- 36 or/22-35 (85594)
- 37 8 and (15 or 21 or 36) (8778)
- 38 randomized controlled trial/ (785964)
- 39 controlled clinical trial/ (469252)
- 40 Random\$.ti,ab,ot. (1968994)
- 41 randomization/ (99178)
- 42 intermethod comparison/ (297283)
- 43 placebo.ti,ab,ot. (366311)
- 44 (compare or compared or comparison).ti,ot. (604093)
- 45 ((evaluated or evaluate or evaluating or assessed or assess) and (compare or compared or comparing or comparison)).ab. (2766233)
- 46 (open adj label).ti,ab,ot. (109016)
- 47 ((double or single or doubly or singly) adj (blind or blinded or blindly)).ti,ab,ot. (274477)
- 48 double blind procedure/ (210575)
- 49 parallel group\$.ti,ab,ot. (32223)
- 50 (crossover or cross over).ti,ab,ot. (124540)
- 51 ((assign\$ or match or matched or allocation) adj5 (alternate or group or groups or intervention or interventions or patient or patients or subject or subjects or participant or participants)).ti,ab,ot. (415063)
- 52 (assigned or allocated).ti,ab,ot. (489023)
- 53 (controlled adj7 (study or design or trial)).ti,ab,ot. (450984)
- 54 (volunteer or volunteers).ti,ab,ot. (282270)
- 55 human experiment/ (650911)
- 56 trial.ti,ot. (403295)
- 57 or/38-56 (6311902)
- 58 37 and 57 (2810)
- 59 (rat or rats or mouse or mice or swine or porcine or murine or sheep or lambs or pigs or piglets or rabbit or rabbits or cat or cats or dog or dogs or cattle or bovine or monkey or monkeys or trout or marmoset\$.ti,ot. and animal experiment/ (1227092)
- 60 animal experiment/ not (human experiment/ or human/) (2577203)
- 61 59 or 60 (2645661)
- 62 58 not 61 (2689)
- 63 limit 62 to yr="2000-Current" (2686)
- 64 remove duplicates from 63 (2558)

**Key:**

/ or.sh. = indexing term (Emtree Subject Heading)

exp = exploded indexing term (Emtree)

\* or \$ = truncation

? = adds up to 1 additional character

ti,ab,kw = terms in either title or abstract fields

dy,tn = drug index terms word or drug trade name fields

adj3 = terms within three words of each other (any order).

pt = publication type

ot = original title

**Cochrane Central Register of Controlled Trials**via Wiley <http://onlinelibrary.wiley.com/>

Date range searched: Issue 5 of 12, May 2023

Date searched: 26 May 2023

Records retrieved: 1825

- #1 ([mh ^"Diabetes Mellitus"] or [mh ^"Diabetes Complications"]) and [mh "Retinal Diseases"] 250
- #2 [mh ^"Diabetic Retinopathy"] 1934
- #3 ((diabet\* or DM) NEAR/3 (retinopath\* or vitreoretinopath\* or chorioretinopath\* or maculopath\*)):ti,ab,kw 4547
- #4 (((proliferat\* or PDR or preproliferat\* or nonproliferat\* or NPDR or background) NEAR/3 (retinopath\* or vitreoretinopath\* or chorioretinopath\*)) and (diabet\* or DM)):ti,ab,kw 1326
- #5 ("new blood" NEXT vessel\* and diabet\*):ti,ab,kw 32
- #6 (((retin\* or subretina\* or interretina\* or vitreoretin\* or chorioretin\* or choroid\* or macula\* or intraocular or intravitreal) NEAR/4 (damage\* or deteriorat\* or degenerat\* or disease\* or edema or oedema or neovasculari?ation\*)) and diabet\*):ti,ab,kw 3457
- #7 ((retinal NEXT vein\* NEAR/3 (occlu\* or obstruct\* or clos\* or stricture\* or steno\* or block\* or emboli\*)) and diabet\*):ti,ab,kw 254
- #8 {OR #1-#7} 5751
- #9 [mh "Vascular Endothelial Growth Factors"/ai] 758
- #10 [mh "Receptors, Vascular Endothelial Growth Factor"/ai] 154
- #11 (anti NEAR/2 VEGF\*):ti,ab,kw 1610
- #12 (antiVEGF\*):ti,ab,kw 1523
- #13 ((anti NEXT vascular or antivascular) NEAR/2 "endothelial growth" NEXT factor\*):ti,ab,kw 699
- #14 (((("vascular endothelial" NEAR/2 growth NEXT factor\*) or vasculotropin or VEGF\* or "vascular permeability" NEXT factor\* or VPF) NEAR/2 (trap\* or inhibit\* or antagonist\*)):ti,ab,kw 2048
- #15 ("vascular proliferation" NEAR/4 inhibit\*):ti,ab,kw 1
- #16 {OR #9-#15} 3671
- #17 [mh ^"Angiogenesis Inhibitors"] 1681
- #18 [mh "Angiogenesis Inducing Agents"/ai] 0
- #19 (angiogen\* NEAR/2 (antagonist\* or inhibit\*)):ti,ab,kw 2126
- #20 ((antiangiogen\* or anti NEXT angiogen\*) NEAR/2 (agent\* or drug\* or effect\*)):ti,ab,kw 717
- #21 (angiostatic NEAR/2 (agent\* or drug\*)):ti,ab,kw 10
- #22 ((neovasculari?ation or vasculari?ation) NEAR/2 inhibit\*):ti,ab,kw 37
- #23 {OR #17-#22}2691
- #24 Aflibercept\*:ti,ab,kw 1081
- #25 (Eylea or Zaltrap or Ziv NEXT Aflibercept or "AVE 0005" or AVE0005 or "AVE 005" or AVE005):ti,ab,kw 252
- #26 [mh ^Bevacizumab] 2633
- #27 Bevacizumab\*:ti,ab,kw 7386
- #28 (Avastin or Mvasi or Alymsys or Aybintio or Equida-

cent or Onbevti or Oyavas or Zirabev or rhuMAB-VEGF or rhuMAB NEXT VEGF or "NSC 704865" or NSC704865):ti,ab,kw 941

- #29 (IVB NEAR/2 inject\*):ti,ab,kw 89
- #30 [mh ^Ranibizumab] 1049
- #31 Ranibizumab\*:ti,ab,kw 2266
- #32 (Lucentis or "rhuFab V2"):ti,ab,kw 451
- #33 (IVR NEAR/2 inject\*):ti,ab,kw 32
- #34 Pegaptanib\*:ti,ab,kw 166
- #35 ("EYE 001" or EYE001 or Macugen or "NX 1838" or NX1838):ti,ab,kw 82
- #36 {OR #24-#35}10087
- #37 #8 and (#16 or #23 or #36) 1847
- #38 (rat or rats or rodent\* or mouse or mice or "mus musculus" or "mus domesticus" or murine or murinae or bovine or sheep or ovine or "ovis aries" or porcine):ti,ab,kw 17188
- #39 #37 not #38 with Publication Year from 2000 to 2023, in Trials 1825

### Science Citation Index Expanded

via Web of Science, Clarivate Analytics <https://clarivate.com/>

Date range searched: 1900–26 May 2023

Date searched: 26 May 2023

Records retrieved: 2394

- 32 #29 NOT #30 2,394 Limited by 2000-01-01 to 2023-05-26
- 31 #29 NOT #30 2,410
- 30 TI=(animal or animals or rat or rats or rodent\* or mouse or mice or "mus musculus" or "mus domesticus" or murine or murinae or porcine or pig or pigs or piglet or piglets or sow or sows or minipig or minipigs or sheep or ovine or "ovis aries" or lamb or lambs or ewe or ewes or rabbit or rabbits or leporide or leporidae or kitten or kittens or dog or dogs or puppy or puppies or monkey or monkeys or horse or horses or foal or foals or equine or bovine or calf or calves or cattle or heifer or heifers or hamster or hamsters or chicken or chickens or livestock or alpaca\* or llama\*) 3,259,653
- 29 #27 AND #28 2,524
- 28 TS=(random\* or control\* or trial\* or "single blind" or "double blind" or "triple blind" or placebo)8,083,064
- 27 #6 AND #26 6,121
- 26 #7 OR #8 OR #9 OR #10 OR #11 OR #12 OR #13 OR #14 OR #15 OR #16 OR #17 OR #18 OR #19 OR #20 OR #21 OR #22 OR #23 OR #24 OR #25 83,065

- 25 TS=("EYE 001" or EYE001 or Macugen or "NX 1838" or NX1838) 142
- 24 TS=(Pegaptanib\*) 716
- 23 TS=(IVR NEAR/2 inject\*) 177
- 22 TS=(Lucentis or "rhuFab V2") 564
- 21 TS=(Ranibizumab\*) 9,347
- 20 TS=(IVB NEAR/2 inject\*) 307
- 19 TS=(Avastin or Mvasi or Alymsys or Aybintio or Equidacent or Onbevzi or Oyavas or Zirabev or rhuMab-VEGF or rhuMab-VEGF or "rhuMab VEGF" or "NSC 704865" or NSC704865) 3,355
- 18 TS=(Bevacizumab\*) 36,279
- 17 TS=(Eylea or Zaltrap or Ziv-Aflibercept or "AVE 0005" or AVE0005 or "AVE 005" or AVE005) 320
- 16 TS=(Aflibercept\*) 4,076
- 15 TS=((neovascularisation or neovascularization or vascularisation or vascularization) NEAR/2 inhibit\*) 1,858
- 14 TS=(angiostatic NEAR/2 (agent\* or drug\*)) 105
- 13 TS=((antiangiogen\* or "anti angiogen\*" or anti-angiogen\*) NEAR/2 (agent\* or drug\* or effect\*)) 11,802
- 12 TS=(angiogen\* NEAR/2 (antagonist\* or inhibit\*)) 19,846
- 11 TS=("vascular proliferation" NEAR/4 inhibit\*) 44
- 10 TS((((("vascular endothelial" NEAR/2 "growth factor\*") or vasculotropin or VEGF\* or "vascular permeability factor\*" or VPF) NEAR/2 (trap\* or inhibit\* or antagonist\*)) 14,540
- 9 TS=((("anti vascular" or anti-vascular or antivascular) NEAR/2 "endothelial growth factor\*") 5,018
- 8 TS=(anti-VEGF\* or antiVEGF\*) 10,111
- 7 TS=(anti NEAR/2 VEGF\*) 10,549
- 6 #1 OR #2 OR #3 OR #4 OR #5 43,073
- 5 TS=((("retinal vein\*" NEAR/3 (occlu\* or obstruct\* or clos\* or stricture\* or steno\* or block\* or emboli\*)) and diabet\*) 1,546
- 4 TS((((retin\* or subretina\* or sub-retina\* or interretina\* or inter-retina\* or vitreoretin\* or vitreo-retin\* or chorioretin\* or chorio-retin\* or choroid\* or macula\* or intraocular or intra-ocular or intravitreal or intra-vitreous) NEAR/4 (damage\* or deteriorat\* or degenerat\* or disease\* or edema or oedema or neovascularisation\*)) and diabet\*) 16,980
- 3 TS=("new blood vessel\*" and diabet\*) 288
- 2 TS((((proliferat\* or PDR or pre-proliferat\* or preproliferat\* or non-proliferat\* or nonproliferat\* or NPDR or background) NEAR/3 (retinopath\* or vitreoretinopath\* or vitreo-retinopath\* or chorioretinopath\* or chorio-retinopath\*)) and (diabet\* or DM)) 7,763
- 1 TS=((diabet\* or DM) NEAR/3 (retinopath\* or vitreoretinopath\* or vitreo-retinopath\* or chorioretinopath\* or chorio-retinopath\* or maculopath\*)) 36,053

**Key:**

TS= terms in either title, abstract, author keywords, and keywords plus fields

TI= search in title field

NEAR/3 = terms within three words of each other (any order).

\* = truncation

**Conference Proceedings Citation Index - Science**

via Web of Science, Clarivate Analytics <https://clarivate.com/>

Date range searched: 1990–26 May 2023

Date searched: 26 May 2023

Records retrieved: 86

- 32 #29 NOT #30 86 Limited by 2000-01-01 to 2023-05-26
- 31 #29 NOT #30 89
- 30 TI=(animal or animals or rat or rats or rodent\* or mouse or mice or "mus musculus" or "mus domesticus" or murine or murinae or porcine or pig or pigs or piglet or piglets or sow or sows or minipig or minipigs or sheep or ovine or "ovis aries" or lamb or lambs or ewe or ewes or rabbit or rabbits or leporide or leporidae or kitten or kittens or dog or dogs or puppy or puppies or monkey or monkeys or horse or horses or foal or foals or equine or bovine or calf or calves or cattle or heifer or heifers or hamster or hamsters or chicken or chickens or livestock or alpaca\* or llama\*) 295,290
- 29 #27 AND #28 92
- 28 TS=(random\* or control\* or trial\* or "single blind" or "double blind" or "triple blind" or placebo) 1,616,551
- 27 #6 AND #26 458
- 26 #7 OR #8 OR #9 OR #10 OR #11 OR #12 OR #13 OR #14 OR #15 OR #16 OR #17 OR #18 OR #19 OR #20 OR #21 OR #22 OR #23 OR #24 OR #25 8,998
- 25 TS=("EYE 001" or EYE001 or Macugen or "NX 1838" or NX1838) 14
- 24 TS=(Pegaptanib\*) 39
- 23 TS=(IVR NEAR/2 inject\*) 1
- 22 TS=(Lucentis or "rhuFab V2") 29
- 21 TS=(Ranibizumab\*) 564
- 20 TS=(IVB NEAR/2 inject\*) 7
- 19 TS=(Avastin or Mvasi or Alymsys or Aybintio or

- Equidacent or Onbevzi or Oyavas or Zirabev or rhuMAbVEGF or rhuMAb-VEGF or "rhuMAb VEGF" or "NSC 704865" or NSC704865) 196
- 18 TS=(Bevacizumab\*) 4,659
- 17 TS=(Eylea or Zaltrap or Ziv-Aflibercept or "AVE 0005" or AVE0005 or "AVE 005" or AVE005) 60
- 16 TS=(Aflibercept\*) 577
- 15 TS=((neovascularisation or neovascularization or vascularisation or vascularization) NEAR/2 inhibit\*) 177
- 14 TS=(angiostatic NEAR/2 (agent\* or drug\*)) 6
- 13 TS=((antiangiogen\* or "anti angiogen\*" or antiangiogen\*) NEAR/2 (agent\* or drug\* or effect\*)) 634
- 12 TS=(angiogen\* NEAR/2 (antagonist\* or inhibit\*)) 1,209
- 11 TS=("vascular proliferation" NEAR/4 inhibit\*) 6
- 10 TS((((("vascular endothelial" NEAR/2 "growth factor\*") or vasculotropin or VEGF\* or "vascular permeability factor\*" or VPF) NEAR/2 (trap\* or inhibit\* or antagonist\*)) 1,025
- 9 TS=((("anti vascular" or anti-vascular or antivascular) NEAR/2 "endothelial growth factor\*") 224
- 8 TS=(anti-VEGF\* or antiVEGF\*) 836
- 7 TS=(anti NEAR/2 VEGF\*) 869
- 6 #1 OR #2 OR #3 OR #4 OR #5 5,826
- 5 TS((((("retinal vein\*" NEAR/3 (occlu\* or obstruct\* or clos\* or stricture\* or steno\* or block\* or emboli\*)) and diabet\*) 74
- 4 TS((((("retin\*" or subretina\* or sub-retina\* or interretina\* or inter-retina\* or vitreoretin\* or vitreo-retin\* or chorioretin\* or chorio-retin\* or choroid\* or macula\* or intraocular or intra-ocular or intravitreal or intra-vitreol) NEAR/4 (damage\* or deteriorat\* or degenerat\* or disease\* or edema or oedema or neovasculari?ation\*)) and diabet\*) 2,140
- 3 TS=("new blood vessel\*" and diabet\*) 29
- 2 TS((((("proliferat\*" or PDR or pre-proliferat\* or proliferat\* or non-proliferat\* or nonproliferat\* or NPDR or background) NEAR/3 (retinopath\* or vitreoretinopath\* or vitreo-retinopath\* or chorioretinopath\* or chorio-retinopath\*)) and (diabet\* or DM)) 642
- 1 TS=((diabet\* or DM) NEAR/3 (retinopath\* or vitreoretinopath\* or vitreo-retinopath\* or chorioretinopath\* or chorio-retinopath\* or maculopath\*)) 4,723

**Key:**

TS= terms in either title, abstract, author keywords, and keywords plus fields

TI= search in title field

NEAR/3 = terms within three words of each other (any order).

\* = truncation

**Cochrane Database of Systematic Reviews**

via Wiley <http://onlinelibrary.wiley.com/>

Date range searched: Issue 5 of 12, May 2023

Date searched: 26 May 2023

Records retrieved: 14

- #1 ([mh ^"Diabetes Mellitus"] or [mh ^"Diabetes Complications"]) and [mh "Retinal Diseases"] 250
- #2 [mh ^"Diabetic Retinopathy"] 1934
- #3 ((diabet\* or DM) NEAR/3 (retinopath\* or vitreoretinopath\* or chorioretinopath\* or maculopath\*)):ti,ab,kw 4547
- #4 (((proliferat\* or PDR or preproliferat\* or nonproliferat\* or NPDR or background) NEAR/3 (retinopath\* or vitreoretinopath\* or chorioretinopath\*)) and (diabet\* or DM)):ti,ab,kw 1326
- #5 ("new blood" NEXT vessel\* and diabet\*):ti,ab,kw 32
- #6 (((retin\* or subretina\* or interretina\* or vitreoretin\* or chorioretin\* or choroid\* or macula\* or intraocular or intravitreal) NEAR/4 (damage\* or deteriorat\* or degenerat\* or disease\* or edema or oedema or neovasculari?ation\*)) and diabet\*):ti,ab,kw 3457
- #7 ((retinal NEXT vein\* NEAR/3 (occlu\* or obstruct\* or clos\* or stricture\* or steno\* or block\* or emboli\*)) and diabet\*):ti,ab,kw 254
- #8 {OR #1-#7} 5751
- #9 [mh "Vascular Endothelial Growth Factors"/ai] 758
- #10 [mh "Receptors, Vascular Endothelial Growth Factor"/ai] 154
- #11 (anti NEAR/2 VEGF\*):ti,ab,kw 1610
- #12 (antiVEGF\*):ti,ab,kw 1523
- #13 ((anti NEXT vascular or antivascular) NEAR/2 "endothelial growth" NEXT factor\*):ti,ab,kw 699
- #14 (((("vascular endothelial" NEAR/2 growth NEXT factor\*) or vasculotropin or VEGF\* or "vascular permeability" NEXT factor\* or VPF) NEAR/2 (trap\* or inhibit\* or antagonist\*)):ti,ab,kw 2048
- #15 ("vascular proliferation" NEAR/4 inhibit\*):ti,ab,kw 1
- #16 {OR #9-#15} 3671
- #17 [mh ^"Angiogenesis Inhibitors"] 1681
- #18 [mh "Angiogenesis Inducing Agents"/ai] 0
- #19 (angiogen\* NEAR/2 (antagonist\* or inhibit\*)):ti,ab,kw 2126

- #20 ((antiangiogen\* or anti NEXT angiogen\*) NEAR/2 (agent\* or drug\* or effect\*)):ti,ab,kw 717
- #21 (angiostatic NEAR/2 (agent\* or drug\*)):ti,ab,kw 10
- #22 ((neovasculari?ation or vasculari?ation) NEAR/2 inhibit\*):ti,ab,kw 37
- #23 {OR #17-#22} 2691
- #24 Aflibercept\*:ti,ab,kw 1081
- #25 (Eylea or Zaltrap or Ziv NEXT Aflibercept or "AVE 0005" or AVE0005 or "AVE 005" or AVE005):ti,ab,kw 252
- #26 [mh ^Bevacizumab] 2633
- #27 Bevacizumab\*:ti,ab,kw 7386
- #28 (Avastin or Mvasi or Alymsys or Aybintio or Equidacent or Onbevzi or Oyavas or Zirabev or rhuMAB-VEGF or rhuMAB NEXT VEGF or "NSC 704865" or NSC704865):ti,ab,kw 941
- #29 (IVB NEAR/2 inject\*):ti,ab,kw 89
- #30 [mh ^Ranibizumab] 1049
- #31 Ranibizumab\*:ti,ab,kw 2266
- #32 (Lucentis or "rhuFab V2"):ti,ab,kw 451
- #33 (IVR NEAR/2 inject\*):ti,ab,kw 32
- #34 Pegaptanib\*:ti,ab,kw 166
- #35 ("EYE 001" or EYE001 or Macugen or "NX 1838" or NX1838):ti,ab,kw 82
- #36 {OR #24-#35} 10087
- #37 #8 and (#16 or #23 or #36) 1847
- #38 (rat or rats or rodent\* or mouse or mice or "mus musculus" or "mus domesticus" or murine or murinae or bovine or sheep or ovine or "ovis aries" or porcine):ti,ab,kw 17188
- #39 #37 not #38 with Cochrane Library publication date between January 2000 and May 2023, in Cochrane Reviews 14

**Key:**

mh = exploded indexing term (MeSH)

mh ^ = unexploded indexing term (MeSH)

/ai = indexing term with subheading for antagonists & inhibitors

\* = truncation or additional characters within a word

? = adds up to 1 additional character

ti,ab,kw = terms in either title or abstract or keyword fields

near/3 = terms within three words of each other (any order)

next = terms are next to each other

**Epistemonikos**

via <https://www.epistemonikos.org/>

Date range searched: Inception – 26 May 2023

Date searched: 26 May 2023

Records retrieved: 1026

((title:(title:(diabet\* OR proliferat\* OR PDR OR pre-proliferat\* OR preproliferat\* OR non-proliferat\* OR nonproliferat\* OR NPDR OR background) AND retinopath\*)) OR abstract:(diabet\* OR proliferat\* OR PDR OR pre-proliferat\* OR preproliferat\* OR non-proliferat\* OR nonproliferat\* OR NPDR OR background) AND retinopath\*)) OR (title:(new blood vessel\* AND diabet\*)) OR abstract:(new blood vessel\* AND diabet\*)) OR abstract:(title:(diabet\* OR proliferat\* OR PDR OR pre-proliferat\* OR preproliferat\* OR non-proliferat\* OR nonproliferat\* OR NPDR OR background) AND retinopath\*)) OR abstract:(diabet\* OR proliferat\* OR PDR OR pre-proliferat\* OR preproliferat\* OR non-proliferat\* OR nonproliferat\* OR NPDR OR background) AND retinopath\*)) OR (title:(new blood vessel\* AND diabet\*)) OR abstract:(new blood vessel\* AND diabet\*)) AND (title:(anti AND VEGF\*)) OR abstract:(anti AND VEGF\*)) OR (title:(anti-VEGF\* OR antiVEGF\*)) OR abstract:(anti-VEGF\* OR antiVEGF\*)) OR (title:(("anti vascular" OR anti-vascular OR antivascular) AND "endothelial growth factor")) OR abstract:(("anti vascular" OR anti-vascular OR antivascular) AND "endothelial growth factor")) OR (title:(("vascular endothelial growth factor" OR vasculotropin OR VEGF\* OR "vascular permeability factor" OR VPF) AND (trap\* OR inhibit\* OR antagonist\*)) OR abstract:(("vascular endothelial growth factor" OR vasculotropin OR VEGF\* OR "vascular permeability factor" OR VPF) AND (trap\* OR inhibit\* OR antagonist\*)) OR (title:(angiogen\* AND (antagonist\* OR inhibit\*))) OR abstract:(angiogen\* AND (antagonist\* OR inhibit\*))) OR (title:(antiangiogen\* OR "antiangiogen" OR anti-angiogen\* OR angiostatic) AND (agent\* OR drug\* OR effect\*)) OR abstract:(antiangiogen\* OR "anti angiogen" OR anti-angiogen\* OR angiostatic) AND (agent\* OR drug\* OR effect\*)) OR (title:(Aflibercept\* OR Eylea OR Zaltrap OR Ziv-Aflibercept OR "AVE 0005" OR AVE0005 OR "AVE 005" OR AVE005 OR Bevacizumab\* OR Avastin OR Mvasi OR Alymsys OR Aybintio OR Equidacent OR Onbevzi OR Oyavas OR Zirabev OR rhuMAB-VEGF OR rhuMAB-VEGF OR "rhuMAB VEGF" OR "NSC 704865" OR NSC704865 OR Ranibizumab\* OR Lucentis OR "rhuFab V2" OR



Pegaptanib\* OR "EYE 001" OR EYE001 OR Macugen OR "NX 1838" OR NX1838)) OR abstract:(Aflibercept\* OR Eylea OR Zaltrap OR Ziv-Aflibercept OR "AVE 0005" OR AVE0005 OR "AVE 005" OR AVE005 OR Bevacizumab\* OR Avastin OR Mvasi OR Alymsys OR Aybintio OR Equidacent OR Onbevzi OR Oyavas OR Zirabev OR rhuMABVEGF OR rhuMAB-VEGF OR "rhuMAB VEGF" OR "NSC 704865" OR NSC704865 OR Ranibizumab\* OR Lucentis OR "rhuFab V2" OR Pegaptanib\* OR "EYE 001" OR EYE001 OR Macugen OR "NX 1838" OR NX1838))) OR (title:(((IVB OR IVR) AND inject\*)) OR abstract:(((IVB OR IVR) AND inject\*)))

Filter: Publication year 2000–2023

Publication type: Systematic Reviews

= 1026

#### Key:

\* = truncation

title: = searches in title field

abstract: = searches in abstract field

#### PROSPERO

via <https://www.crd.york.ac.uk/prospéro/>

Date range: Inception – 26 May 2023

Date searched: 26 May 2023

Records retrieved: 159

- #1 MeSH DESCRIPTOR Diabetic Retinopathy 107
- #2 ((diabet\* or DM) adj3 (retinopath\* or vitreoretinopath\* or vitreo-retinopath\* or chorioretinopath\* or chorio-retinopath\* or maculopath\*)) 609
- #3 (((proliferat\* or PDR or pre-proliferat\* or preproliferat\* or non-proliferat\* or nonproliferat\* or NPDR or background) adj3 (retinopath\* or vitreoretinopath\* or vitreo-retinopath\* or chorioretinopath\* or chorio-retinopath\*)) and (diabet\* or DM)) 110
- #4 (new blood vessel\* and diabet\*) 9
- #5 (((retin\* or subretina\* or sub-retina\* or interretina\* or inter-retina\* or vitreoretin\* or vitreo-retin\* or chorioretin\* or chorio-retin\* or choroid\* or macula\* or intraocular or intra-ocular or intravitreal or intravitreal) adj4 (damage\* or deteriorat\* or degenerat\* or disease\* or edema or oedema or neovascularisation\* or neovascularization\*)) AND diabet\*) 373

- #6 ((retinal vein\* adj3 (occlu\* or obstruct\* or clos\* or stricture\* or steno\* or block\* or emboli\*)) and diabet\*) 64
- #7 #1 OR #2 OR #3 OR #4 OR #5 OR #6 740
- #8 MeSH DESCRIPTOR Vascular Endothelial Growth Factors EXPLODE ALL TREES WITH QUALIFIER AI 0
- #9 MeSH DESCRIPTOR Receptors, Vascular Endothelial Growth Factor EXPLODE ALL TREES WITH QUALIFIER AI 0
- #10 (anti adj2 VEGF\*) 327
- #11 (anti-VEGF\* or antiVEGF\*) 327
- #12 ((anti vascular or anti-vascular or antivascular) adj2 endothelial growth factor\*) 153
- #13 (((vascular endothelial adj2 growth factor\*) or vasculotropin or VEGF\* or vascular permeability factor\* or VPF) adj2 (trap\* or inhibit\* or antagonist\*)) 96
- #14 (vascular proliferation adj4 inhibit\*) 0
- #15 #8 OR #9 OR #10 OR #11 OR #12 OR #13 OR #14 412
- #16 MeSH DESCRIPTOR Angiogenesis Inhibitors 40
- #17 MeSH DESCRIPTOR Angiogenesis Inducing Agents EXPLODE ALL TREES WITH QUALIFIER A IO
- #18 (angiogen\* adj2 (antagonist\* or inhibit\*)) 74
- #19 ((antiangiogen\* or anti angiogen\* or anti-angiogen\*) adj2 (agent\* or drug\* or effect\*)) 145
- #20 (angiostatic adj2 (agent\* or drug\*)) 0
- #21 ((neovascularisation\* or neovascularization\* or vascularisation\* or vascularization\*) adj2 inhibit\*) 0
- #22 #16 OR #17 OR #18 OR #19 OR #20 OR #21 224
- #23 (Aflibercept\*) 141
- #24 (Eylea or Zaltrap or Ziv-Aflibercept or AVE 0005 or AVE0005 or AVE 005 or AVE005) 22
- #25 MeSH DESCRIPTOR Bevacizumab 46
- #26 (Becavizumab\*) 445
- #27 (Avastin or Mvasi or Alymsys or Aybintio or Equidacent or Onbevzi or Oyavas or Zirabev or rhuMAB-VEGF or rhuMAB-VEGF or rhuMAB VEGF or NSC 704865 or NSC704865) 59
- #28 (IVB adj2 inject\*) 0
- #29 MeSH DESCRIPTOR Ranibizumab 7
- #30 (Ranibizumab\*) 142
- #31 (Lucentis or rhuFab V2) 23
- #32 (IVR adj2 inject\*) 0
- #33 (Pegaptanib\*) 30
- #34 (EYE 001 or EYE001 or Macugen or NX 1838 or NX1838) 5
- #35 #23 OR #24 OR #25 OR #26 OR #27 OR #28 OR #29 OR #30 OR #31 OR #32 OR #33 OR #34 500
- #36 #15 OR #22 OR #35 839
- #37 #7 AND #36 159

**Key:**

MeSH DESCRIPTOR = indexing term: Medical Subject Heading (MeSH)

QUALIFIER AI = indexing term subheading for antagonists & inhibitors

EXPLODE ALL TREES = exploded indexing term (MeSH)

\* = truncation

adj3 = terms within three words of each other (order specified).

:TI,KW = terms in either title or keyword fields

**ClinicalTrials.gov**

via <https://clinicaltrials.gov/>

Date searched: 26 May 2023

Records retrieved: 286

Two separate searches were used in Advanced Search, retrieving 286 records in total, which were imported into EndNote 20 and deduplicated.

1. **Condition or Disease:** (diabetic retinopathy)

**Other Terms:** (Aflibercept OR Eylea OR Zaltrap OR Bevacizumab OR Avastin OR Mvasi OR Alymsys OR Aybintio OR Equidacent OR Onbevzi OR Oyavas OR Zirabev OR rhuMAB VEGF OR Ranibizumab OR Lucentis OR rhuFab OR Pegaptanib OR Macugen) = **190 hits**

2. **Condition or Disease:** (diabetic retinopathy)

**Other Terms:** ((VEGF OR vascular endothelial growth factor OR vasculotropin OR vascular permeability factor or VPF) AND (anti OR trap or inhibitor or antagonist)) = **96 hits**

**European Union Clinical Trials Register**

via [www.clinicaltrialsregister.eu/ctr-search/search](http://www.clinicaltrialsregister.eu/ctr-search/search)

Date searched: 26 May 2023

Records retrieved: 163

Two separate searches were used, retrieving 163 records in total, which were imported into EndNote 20 and deduplicated.

1. ((“diabetic retinopathy”) AND (Aflibercept OR Eylea OR Zaltrap OR Bevacizumab OR Avastin OR Mvasi OR Alymsys OR Aybintio OR Equidacent OR Onbevzi OR Oyavas OR Zirabev OR “rhuMAB VEGF” OR Ranibizumab OR Lucentis OR rhuFab OR Pegaptanib OR Macugen)) = **113 hits**
2. ((“diabetic retinopathy”) AND ((anti OR trap or inhibitor OR antagonist) AND (VEGF OR “vascular endothelial growth factor” OR vasculotropin OR “vascular permeability factor” OR VPF))) = **50 hits**

**WHO International Clinical Trials Registry Platform**

via <https://trialsearch.who.int/>

Date searched: 26 May 2023

Records retrieved: 198

Two separate searches were used in Advanced Search, retrieving 198 records in total, which were imported into EndNote 20 and deduplicated.

1. Advanced Search

**Condition:** (diabetic retinopathy)

**Intervention:** (Aflibercept OR Eylea OR Zaltrap OR Bevacizumab OR Avastin OR Mvasi OR Alymsys OR Aybintio OR Equidacent OR Onbevzi OR Oyavas OR Zirabev OR rhuMAB VEGF OR Ranibizumab OR Lucentis OR rhuFab OR Pegaptanib OR Macugen)

Recruitment Status: ALL = 194 records for 180 trials

2. Advanced Search

**Condition:** (diabetic retinopathy)

**Intervention:** ((VEGF OR vascular endothelial growth factor OR vasculotropin OR vascular permeability factor or VPF) AND (anti OR trap or inhibitor or antagonist))

Recruitment Status: ALL = 23 records for 18 trials

**List of excluded studies****Randomised controlled trial of DME (35)**

Bayer AG. An open-label, randomized, active-controlled, parallel-group, Phase-3b study of the efficacy, safety, and tolerability of three different treatment regimens of 2mg aflibercept administered by intr.

Braimah IZ, Kenu E, Amissah-Arthur KN, Akafo S, Kwarteng KO, Amoaku WM. Safety of intravitreal ziv-aflibercept in choroido-retinal vascular diseases: a randomised double-blind intervention study. *PLOS ONE* 2019;**14**:e0223944.

Bressler SB, Qin H, Beck RW, Chalam KV, Kim JE, Melia M, Wells JA 3rd; Diabetic Retinopathy Clinical Research and Network. Factors associated with changes in visual acuity and central subfield thickness at 1 year after treatment for diabetic macular edema with ranibizumab. *Arch Ophthalmol* 2012;**130**:1153–61.

Bressler SB, Qin H, Melia M, Bressler NM, Beck RW, Chan CK, *et al.*; Diabetic Retinopathy Clinical Research and Network. Exploratory analysis of the effect of intravitreal ranibizumab or triamcinolone on worsening of diabetic retinopathy in a randomized clinical trial. *JAMA Ophthalmol* 2013;**131**:1033–40.

Bressler SB, Liu D, Glassman AR, Blodi BA, Castellarin AA, Jampol LM, *et al.*; Diabetic Retinopathy Clinical Research and Network. Change in diabetic retinopathy through 2 years: secondary analysis of a randomized clinical trial comparing aflibercept, bevacizumab, and ranibizumab. *JAMA Ophthalmol* 2017;**135**:558–68.

Department of Ophthalmology and Medical University of Vienna. A Randomized, Double-masked Study with Intracocular Bevacizumab (Avastin®) Compared with Intravitreal Ranibizumab (Lucentis®) in Patients with Persistent Diabetic Macular Edema or Persistent Active. URL: [www.clinicaltrialsregister.eu/ctr-search/search?query=eudract\\_number:2008-001469-28](http://www.clinicaltrialsregister.eu/ctr-search/search?query=eudract_number:2008-001469-28)

Dhoot D, Hill L, Tarnowski K, Stoilov I. Baseline factors associated with  $\geq$  2-step diabetic retinopathy (DR) severity improvement with ranibizumab (RBZ). *Investigative Ophthalmol Visual Sci Conference* 2018;**59**.

Dhoot DS, Hill LF, Ghanekar A, Tarnowski KW, Ali FS. Baseline factors associated with diabetic retinopathy improvement in RIDE/RISE. *Ophthalmol Retina* 2021;**5**:101–3.

Dhoot DS, Moini H, Reed K, Du W, Vitti R, Berliner AJ, Singh RP. Functional outcomes of sustained improvement on Diabetic Retinopathy Severity Scale with intravitreal aflibercept in the VISTA and VIVID trials. *Eye* 2022;**19**:19.

Dimitriou E, Theodossiadis P, Chatzirallis A, Kazantzis D, Theodossiadis G, Chatziralli E. Intravitreal ranibizumab alone or in combination with panretinal photocoagulation

for the treatment of proliferative diabetic retinopathy with coexistent macular edema: long-term outcomes in real-life data. *Investigative Ophthalmol Visual Sci Conference* 2020;**61**.

Ekinci M, Ceylan E, Cakici O, Tanyildiz B, Olcaysu O, Cagatay HH. Treatment of macular edema in diabetic retinopathy: comparison of the efficacy of intravitreal bevacizumab and ranibizumab injections. *Expert Rev Ophthalmol* 2014;**9**:139–43.

Euctr-009909-25-De. Evaluation of the Efficacy and Safety of a Macugen Monotherapy versus Combined Therapies in the Treatment of Diabetic Retinopathy - A Single Centre, Randomized, Prospective Phase II Trial. 2009.

Glassman AR, Stockdale CR, Beck RW, Baker C, Bressler NM; Diabetic Retinopathy Clinical Research and Network. Evaluation of masking study participants to intravitreal injections in a randomized clinical trial. *Arch Ophthalmol* 2012;**130**:190–4.

Gonzalez VH. Pegaptanib in diabetic retinopathy: improvements in diabetic macular edema, retinal neovascularization, and diabetic retinopathy severity. *Am Academy Ophthalmol* 2006:192.

Gonzalez VH, Wang PW, Ruiz CQ. Panretinal photocoagulation for diabetic retinopathy in the RIDE and RISE trials: not '1 and done'. *Ophthalmology* 2019;**21**:21.

Gonzalez VH, Wang PW, Ruiz CQ. Panretinal photocoagulation for diabetic retinopathy in the RIDE and RISE trials: not '1 and done'. *Ophthalmology* 2021;**128**:1448–57.

Hassan M, Sadiq MA, Halim MS, Afridi R, Nguyen NV, Sepah YJ. Short-term effects of ranibizumab on diabetic retinopathy severity and progression. *Ophthalmol Retina* 2018;**2**:749–51.

Hassan M, Sadiq MA, Halim MS, Afridi R, Nguyen NV, Sepah YJ. Short-term effects of ranibizumab on diabetic retinopathy severity and progression in the ranibizumab for edema of the macula in diabetes - Protocol 3 with high dose (READ-3) study. *Investigative Ophthalmol Visual Sci Conference* 2018;**59**.

Irct201205029617N. Efficacy of Macular laser Photocoagulation with or without Intravitreal Injection of Bevacizumab (Avastin) or Triamcinolone Acetonide for Diffuse Diabetic Macular Edema. 2012.

Mehta H, Lim LL, Nguyen V, Qatarnah D, Wickremasinghe SS, Hodgson LAB, *et al.* Development of new proliferative diabetic retinopathy in the BEVORDEX trial. *Ophthalmol Retina* 2019;**3**:286–7.

Mitchell P, McAllister I, Larsen M, Staurengi G, Korobelnik JF, Boyer DS, *et al.* Evaluating the impact of intravitreal aflibercept on diabetic retinopathy progression in the VIVID-DME and VISTA-DME studies. *Ophthalmol Retina* 2018;**2**:988–96.

NCT. *Laser-Ranibizumab-Triamcinolone for Proliferative Diabetic Retinopathy*. 2007. URL: <https://clinicaltrials.gov/show/NCT00445003>

NCT. *Anterior and Posterior Segment Vascular Changes Following Laser and Anti-Vascular Endothelial Growth Factor (VEGF) Treatment of Diabetic Retinopathy*. 2009.

NCT. *Laser Therapy Combined with Intravitreal Aflibercept vs Intravitreal Aflibercept Monotherapy (LADAMO)*. 2015. <https://clinicaltrials.gov/show/NCT02432547>

Novartis Pharma and AG. *A 12-Month, 2-Arm, Randomized, Double-Masked, Multicenter Phase III Study Assessing the Efficacy and Safety of Brolucizumab Every 4 Weeks versus Aflibercept Every 4 Weeks in Adult Patients with Vis*.

Novartis Pharma Gmb and H. *A Randomized, Single-blinded, Multicenter, Phase IV Study to Compare Systemic VEGF Protein Dynamics Following Monthly Intravitreal Injections of 0.5 mg Ranibizumab versus 2 mg Aflibercept until St*.

Novartis Pharma and AG. *A Two-year, Three-arm, Randomized, Double Masked, Multicenter, Phase III Study Assessing the Efficacy and Safety of Brolucizumab versus Aflibercept in Adult Patients with Visual Impairment due to D*.

Novartis Pharma and AG. *A Two-year, Two-arm, Randomized, Double Masked, Multicenter, Phase III Study Assessing the Efficacy and Safety of Brolucizumab versus Aflibercept in Adult Patients with Visual Impairment due to Dia*.

Oxurion NV. *A Phase 2, Randomised, Single-masked, Active-controlled, Multicentre Study to Evaluate the Efficacy and Safety of Intravitreal THR-317 Administered in Combination with Ranibizumab, for the Treatment*.

Quark Pharmaceuticals and Inc. *An Open-Label Dose Escalation Study of PF-04523655 (Stratum I) Combined With A Prospective, Randomized, Double-Masked, Multi-Center, Controlled Study (Stratum II) Evaluating The Efficacy and Safety*.

Sadiq MA, Hassan M, Soliman MK, Afridi R, Do DV, Nguyen QD, Sepah YJ. Effects of two different doses of ranibizumab on diabetic retinopathy severity. *Ophthalmol Retina* 2017;**1**:566–7.

Sameen M, Khan MS, Mukhtar A, Yaqub MA, Ishaq M. Efficacy of intravitreal bevacizumab combined with pan retinal photocoagulation versus panretinal photocoagulation alone in treatment of proliferative diabetic retinopathy. *Pakistan J Med Sci* 2017;**33**:142–5.

Sasongko MB, Rogers S, Constantinou M, Sandhu SS, Wickremasinghe SS, Al-Qureshi S, Lim LL. Diabetic retinopathy progression 6 months post-cataract surgery with intravitreal bevacizumab vs triamcinolone: a secondary analysis of the DiMECAT trial. *Clin Exp Ophthalmol* 2020;**48**:793–801.

Shahraki T, Arabi A, Nourinia R, Beheshtizadeh NF, Entezari M, Nikkiah H, *et al.* Panretinal photocoagulation versus intravitreal bevacizumab versus a proposed modified combination therapy for treatment of proliferative diabetic retinopathy: a Randomized Three-Arm Clinical Trial (CTPDR Study). *Retina* 2022;**42**:1065–76.

Yan P, Qian C, Wang W, Dong Y, Wan G, Chen Y. Clinical effects and safety of treating diabetic macular edema with intravitreal injection of ranibizumab combined with retinal photocoagulation. *Therapeutics Clin Risk Manage* 2016;**12**:527–33.

### **RCT of vitreous haemorrhage or vitrectomy (86)**

Ahmadieh H, Shoeibi N, Entezari SM. Intravitreal bevacizumab for early post-vitrectomy hemorrhage in diabetics: a randomized, doubleMasked clinical trial. *Am Academy Ophthalmol* 2008:181.

Ahmadieh H, Shoeibi N, Entezari M, Monshizadeh R. Intravitreal bevacizumab for prevention of early postvitrectomy hemorrhage in diabetic patients: a randomized clinical trial. *Ophthalmology* 2009;**116**:1943–8.

Ahn J, Woo SJ, Chung H, Park KH. The effect of adjunctive intravitreal bevacizumab for preventing postvitrectomy hemorrhage in proliferative diabetic retinopathy. *Ophthalmology* 2011;**118**:2218–26.

Albuquerque TL, Pierozzi GS, Araujo AC. C, Neto NH, Carregal TB, Martins MC, *et al.* Comparative, randomized, double blinded study of the use of anti-VEGF in patients with vitreous hemorrhage or tractional retinal detachment

secondary to diabetic retinopathy. *Investigative Ophthalmol Visual Sci* 2014;**55**:4391.

Aleman I, Velazquez JC, Rush SW, Rush RB. Ziv-aflibercept versus bevacizumab administration prior to diabetic vitrectomy: a randomised and controlled trial. *Br J Ophthalmol* 2019;**103**:1740–6.

Arevalo JF, Lasave AF, Kozak I, Al Rashaed S, Al Kahtani E, Maia M, *et al.*; Pan-American Collaborative Retina Study Group. Preoperative bevacizumab for tractional retinal detachment in proliferative diabetic retinopathy: a prospective randomized clinical trial. *Am J Ophthalmol* 2019;**207**:279–87.

Bhavsar A. A randomized trial evaluating intravitreal ranibizumab or intravitreal saline for vitreous hemorrhage from proliferative diabetic retinopathy. *Investigative Ophthalmol Visual Sci Conference* 2013;**54**.

Bhavsar AR, Torres K, Beck RW, Friedman SM, Glassman AR, Maturi RK, *et al.*; Diabetic Retinopathy Clinical Research Network. Randomized clinical trial evaluating intravitreal ranibizumab or saline for vitreous hemorrhage from Proliferative Diabetic Retinopathy Diabetic Retinopathy Clinical Research Network. *JAMA Ophthalmol* 2013;**131**:283–93.

Castillo J, Aleman I, Rush SW, Rush RB. Preoperative bevacizumab administration in proliferative diabetic retinopathy patients undergoing vitrectomy: a randomized and controlled trial comparing interval variation. *Am J Ophthalmol* 2017;**183**:1–10.

Castillo Velazquez J, Aleman I, Rush SW, Rush RB. Bevacizumab before diabetic vitrectomy: a clinical trial assessing 3 dosing amounts. *Ophthalmol Retina* 2018;**2**:1010–20.

Chelala E, Nehme J, El Rami H, Aoun R, Dirani A, Fadlallah A, Jalkh A. Efficacy of intravitreal ranibizumab injections in the treatment of vitreous hemorrhage related to proliferative diabetic retinopathy. *Retina* 2018;**38**:1127–33.

ChiCtr. *Feasibility Study of Anti-VEGF Instead of Intraoperative PRP in Proliferative Diabetic Retinopathy*. 2018.

ChiCtr. *A Prospective and Randomized Controlled Clinical Study for Pre- and After-operative Intravitreal Injection of Anti-VEGF Combined with Pars Plana Vitrectomy*. 2020.

ChiCtr. *A Prospective Randomized Controlled Study of Long-acting Dexamethasone Implant to Improve the Prognosis of PDR Patients after Vitrectomy*. 2021.

ChiCTR1800019455. *Effects of Intraocular Injection of Different Anti-VEGF Drugs on Inflammatory Factors in Aqueous Humor of Patients with Diabetic Retinopathy*. 2018.

ChiCTR2000035032. *Efficacy of Different Doses of Anti-VEGF with Vitrectomy in the Treatment of Proliferative Diabetic Retinopathy*. 2020.

Comyn O, Bainbridge JWB. A pilot randomized controlled trial of ranibizumab pre-treatment for diabetic vitrectomy (The RaDiVit study). *Investigative Ophthalmol Visual Sci* 2014;**55**:2302.

Comyn O, Wickham L, Charteris DG, Sullivan PM, Ezra E, Gregor Z, *et al.* Ranibizumab pretreatment in diabetic vitrectomy: a pilot randomised controlled trial (the RaDiVit study). *Eye* 2017;**31**:1253–8.

Comyn O, Lange C, Bainbridge JWB. Vitreous and plasma cytokine levels in subjects with advanced proliferative diabetic retinopathy in the Ranibizumab in Diabetic Vitrectomy (RaDiVit) Study. *Investigative Ophthalmol Visual Sci Conference* 2019;**60**.

Cui J, Chen H, Lu H, Dong F, Wei D, Jiao Y, *et al.* Efficacy and safety of intravitreal conbercept, ranibizumab, and triamcinolone on 23-gauge vitrectomy for patients with proliferative diabetic retinopathy. *J Ophthalmol* 2018;**2018**:4927259.

da R Lucena D, Ribeiro JA, Costa RA, Barbosa JC, Scott IU, de Figueiredo-Pontes LL, Jorge R. Intraoperative bleeding during vitrectomy for diabetic tractional retinal detachment with versus without preoperative intravitreal bevacizumab (IBeTra study). *Br J Ophthalmol* 2009;**93**:688–91.

di Lauro R, De Ruggiero P, di Lauro R, di Lauro MT; Romano MR. Intravitreal bevacizumab for surgical treatment of severe proliferative diabetic retinopathy. *Graefes Arch Clin Exp Ophthalmol* 2010;**248**:785–91.

Diabetic Retinopathy Clinical Research and Network. Randomized clinical trial evaluating intravitreal ranibizumab or saline for vitreous hemorrhage from proliferative diabetic retinopathy. *JAMA Ophthalmol* 2013;**131**:283–93.

- Dong F, Yu C, Ding H, Shen L, Lou D. Evaluation of intravitreal ranibizumab on the surgical outcome for diabetic retinopathy with tractional retinal detachment. *Medicine* 2016;**95**:e2731.
- Dong X. Effect of ranibizumab on the efficacy of vitrectomy in patients with PDR[J]. *Int Eye Sci* 2019;**19**:809812.
- Luo W. Effect of ranibizumab combined with vitrectomy on the serum VEGF-A and SDF-1 expression in patients with proliferative diabetic retinopathy. *Int Eye Sci* 2019;**19**:438–41.
- Euctr-000780-21-Gb. A Randomised, Single-masked, Phase IV Pilot Study of the Efficacy and Safety of Adjunctive Intravitreal Avastin® (Bevacizumab) in the Prevention of Early Postoperative Vitreous Haemorrhage Following Diabetic Vitrectomy – Intravitreal Avastin® in Diabetic Vitrectomy. 2007.
- Euctr-015559-25-Gb. Preoperative Intravitreal Ranibizumab for Persistent Diabetic Vitreous Haemorrhage: A Randomized, Double-masked, Controlled Study – Vitreous Haemorrhage Study. 2010. URL: [www.who.int/trialsearch/Trial2.aspx?TrialID=EUCTR2009-015559-25-GB](http://www.who.int/trialsearch/Trial2.aspx?TrialID=EUCTR2009-015559-25-GB)
- Euctr-024062-22-Gb. A Prospective, Randomised Controlled Trial of Ranibizumab Pre-treatment in Diabetic Vitrectomy – A Pilot Study – A Pilot RCT of Ranibizumab in Diabetic Vitrectomy – The RaDiVit Study. 2011.
- Farahvash MS, Majidi AR, Roohipoor R, Ghassemi F. Preoperative injection of intravitreal bevacizumab in dense diabetic vitreous hemorrhage. *Retina* 2011;**31**:1254–60.
- Ferraz DA, Morita C, Preti RC, Nascimento VP, Maia OO, de Barros AC, *et al.* Use of intravitreal bevacizumab or triamcinolone acetonide as a preoperative adjunct to vitrectomy for vitreous haemorrhage in diabetics. *Revista Brasileira De Oftalmologia* 2013;**72**:12–16.
- Gao S, Lin Z, Chen Y, Xu J, Zhang Q, Chen J, Shen X. Intravitreal conbercept injection as an adjuvant in vitrectomy with silicone oil infusion for severe proliferative diabetic retinopathy. *J Ocul Pharmacol Ther* 2020;**36**:304–10.
- Genovesi-Ebert F, Rizzo S, Di Bartolo E, Miniaci S, Vento A, Palla M, Cresti F. Injection of intravitreal Avastin before vitrectomy surgery in the treatment of severe proliferative diabetic retinopathy. *Invest Ophthalmol Vis Sci* 2007;**48**:ARVO E-Abstract 5044.
- Glassman AR, Beaulieu WT, Maguire MG, Antoszyk AN, Chow CC, Elman MJ, *et al.* Visual acuity, vitreous hemorrhage, and other ocular outcomes after vitrectomy vs aflibercept for vitreous hemorrhage due to diabetic retinopathy: a secondary analysis of a randomized clinical trial. *JAMA Ophthalmol* 2021;**139**:725–33.
- Han XX, Guo CM, Li Y, Hui YN. Effects of bevacizumab on the neovascular membrane of proliferative diabetic retinopathy: reduction of endothelial cells and expressions of VEGF and HIF-1alpha. *Mol Vision* 2012;**18**:1–9.
- Hernandez-Da Mota SE, Nunez-Solorio SM. Experience with intravitreal bevacizumab as a preoperative adjunct in 23-G vitrectomy for advanced proliferative diabetic retinopathy. *Eur J Ophthalmol* 2010;**20**:1047–52.
- Hu BJ, Zeng Q, Liu XL, Li XR, Song WJ. Influence of intravitreal avastin on the expression of cell factors in retinal proliferative membrane in proliferative diabetic retinopathy eye. [Chinese]. *Zhonghua Shiyan Yanke Zazhi/Chinese Journal of Experimental Ophthalmology* 2013;**31**:55–59.
- Hu Z, Cao X, Chen L, Su Y, Ji J, Yuan S, *et al.* Monitoring intraocular proangiogenic and profibrotic cytokines within 7 days after adjunctive anti-vascular endothelial growth factor therapy for proliferative diabetic retinopathy. *Acta Ophthalmol* 2021;**14**:14.
- Jeon S, Lee WK. Intravitreal bevacizumab increases intraocular interleukin-6 levels at 1 day after injection in patients with proliferative diabetic retinopathy. *Cytokine* 2012;**60**:535–9.
- Jiang TT, Gu JX, Zhang PJ, Chen WW, Chang Q. The effect of adjunctive intravitreal conbercept at the end of diabetic vitrectomy for the prevention of post-vitrectomy hemorrhage in patients with severe proliferative diabetic retinopathy: a prospective, randomized pilot study. *BMC Ophthalmol* 2020;**20**:9.
- Jiao C, Spee C, He S, Mullins R, Elliott D, Hinton DR, Sohn EH. Angiofibrotic response to bevacizumab on fibrovascular membranes in proliferative diabetic retinopathy. *Invest Ophthalmol Vis Sci* 2014;**55**:5821.
- Jorge DM, Tavares Neto JEDS, Poli-Neto OB, Scott IU, Jorge R. Intravitreal bevacizumab (IVB) versus IVB in combination with pars plana vitrectomy for vitreous hemorrhage secondary to proliferative diabetic retinopathy: a randomized clinical trial. *Int J Retina Vitreous* 2021;**7**:35.

- JPRN-UMIN. *Low Dose of Intravitreal Bevacizumab (Avastin) Used as Preoperative Adjunct Therapy for Proliferative Diabetic Retinopathy*. 2012.
- Kanclerz P, Raczynska K. Preoperative bevacizumab as an adjunct for vitrectomy in proliferative diabetic retinopathy patients. *Ophthalmologica J Int d'ophtalmologie* 2016;**236**:58.
- Li Q, Wang JH, Zhang MM, Wang Y. Effect of ranibizumab intravitreal injection before 23G-vitrectomy surgery in the treatment of patients with proliferative diabetic retinopathy [Chinese]. *Int Eye Sci* 2016;**16**:1959–61.
- Li B, Li MD, Ye JJ, Chen Z, Guo ZJ, Di Y. Vascular endothelial growth factor concentration in vitreous humor of patients with severe proliferative diabetic retinopathy after intravitreal injection of conbercept as an adjunctive therapy for vitrectomy. *Chine Med J* 2020;**133**:664–9.
- Manabe A, Shimada H, Hattori T, Nakashizuka H, Yuzawa M. Randomized controlled study of intravitreal bevacizumab 0.16 mg injected one day before surgery for proliferative diabetic retinopathy. *Retina* 2015;**35**:1800–7.
- Meng N, Ren BC. Effect of intravitreal injection of bevacizumab for vitreous hemorrhage in patients with proliferative diabetic retinopathy [Chinese]. *Int Eye Sci* 2016;**16**:972–4.
- Modarres M, Nazari H, Falavarjani KG, Naseripour M, Hashemi M, Parvaresh MM. Intravitreal injection of bevacizumab before vitrectomy for proliferative diabetic retinopathy. *Eur J Ophthalmol* 2009;**19**:848–52.
- NCT. *Intravitreal Bevacizumab for Proliferative Diabetic Retinopathy*. 2007. URL: <https://clinicaltrials.gov/show/NCT00423059>
- NCT. *Evaluation of Ranibizumab in Proliferative Diabetic Retinopathy (PDR) Requiring Vitrectomy*. 2007. URL: <https://clinicaltrials.gov/show/NCT00516464>
- NCT. *Preoperative Bevacizumab for Vitreous Hemorrhage*. 2008. URL: <https://clinicaltrials.gov/show/NCT00596297>
- NCT. *Safety and Efficacy of Intravitreal Ranibizumab as a Preoperative Adjunct Treatment before Vitrectomy Surgery in Proliferative Diabetic Retinopathy (PDR) Compared to Vitrectomy Alone*. 2009. URL: <https://clinicaltrials.gov/show/NCT00931125>
- NCT. *Acute Changes in Intraocular Cytokines after Intravitreal Bevacizumab*. 2011. URL: <https://clinicaltrials.gov/show/NCT01439178>
- NCT. *Ranibizumab in Diabetic Vitrectomy. A Prospective, Randomised Controlled Trial of Ranibizumab Pre-treatment in Diabetic Vitrectomy – A Pilot Study*. 2011. URL: <https://ClinicalTrials.gov/show/NCT01306981>
- NCT. *Prospective Randomized Controlled Study of Intravitreal Injection of Bevacizumab for Proliferative Diabetic Retinopathy*. 2013. URL: <https://clinicaltrials.gov/show/NCT01854593>
- NCT. *Aflibercept Injection for Proliferative Diabetic Retinopathy*. 2013. URL: <https://clinicaltrials.gov/show/NCT01805297>
- NCT. *Pre-operative Intravitreal Bevacizumab for Tractional Retinal Detachment Secondary to Proliferative Diabetic Retinopathy*. 2013. URL: <https://clinicaltrials.gov/show/NCT01976923>
- NCT. *Comparison of Interval Variation and Dosage in Preoperative Bevacizumab and Ziv-Aflibercept Administration in Proliferative Diabetic Retinopathy Undergoing Vitrectomy*. 2015. URL: <https://clinicaltrials.gov/show/NCT02590094>
- NCT. *25-G Vitrectomy with Ranibizumab or Triamcinolone Acetonide on PDR in China-Randomized Clinical Trial*. 2015. URL: <https://clinicaltrials.gov/show/NCT02447185>
- NCT. *Intravitreal Injection of Ranibizumab versus Sham before Vitrectomy in Patients with Proliferative Diabetic Retinopathy*. 2016. URL: <https://clinicaltrials.gov/show/NCT02857491>
- NCT. *Pre-vitrectomy Intravitreal Ranibizumab for Patients with Proliferative Diabetic Retinopathy Combined With Diabetic Macular Edema*. 2020. URL: <https://clinicaltrials.gov/show/NCT04464694>
- Pakzad-Vaezi K, Albani DA, Kirker AW, Merkur AB, Kertes PJ, Eng KT, et al. A randomized study comparing the efficacy of bevacizumab and ranibizumab as pre-treatment for pars plana vitrectomy in proliferative diabetic retinopathy. *Ophthalmic Surg Lasers Imaging Retina* 2014;**45**:521–4.
- Petrarca R, Soare C, Wong R, Desai R, Neffendorf J, Simpson A, Jackson TL. Intravitreal ranibizumab for persistent diabetic vitreous haemorrhage: a randomised,

- double-masked, placebo-controlled feasibility study. *Acta Ophthalmol* 2020;**98**:E960–7.
- Qi QF, Shi YW, Guo T. Clinical observation on preoperative application of bevacizumab in proliferative diabetic retinopathy [Chinese]. *Int Eye Sci* 2014;**14**:1646–8.
- Ren XJ, Bu SC, Zhang XM, Jiang YF, Tan LZ, Zhang H, Li XR. Safety and efficacy of intravitreal conbercept injection after vitrectomy for the treatment of proliferative diabetic retinopathy. *Eye* 2019;**33**:1177–83.
- Reza NM, Hosein AM, Hesamsadat H, Amir EM, Narges H, Amin N. Intravitreal tissue plasminogen activator in diabetic vitreous hemorrhage. *Int J Pharm Res* 2019;**11**:823–7.
- Sohn EH, He S, Kim LA, Salehi-Had H, Javaheri M, Spee C, et al. Angiofibrotic response to vascular endothelial growth factor inhibition in diabetic retinal detachment: report no. 1. *Arch Ophthalmol* 2012;**130**:1127–34.
- Starnes DC, Lalane R, Walia H, Farooq A, Frazier H, Marcus W, et al. Endolaserless vitrectomy with intravitreal aflibercept injection (IAI) for proliferative diabetic retinopathy (PDR)-related vitreous hemorrhage: LASER LESS TRIAL 1-year results. *Invest Ophthalmol Vis Sci Conference* 2019;**60**.
- Su L, Ren X, Wei H, Zhao L, Zhang X, Liu J, et al. Intravitreal conbercept (Kh902) for surgical treatment of severe proliferative diabetic retinopathy. *Retina* 2016;**36**:938–43.
- Sun M, Li MX. Study of anti-vascular endothelial growth factor medicine for proliferative diabetic retinopathy at perioperative period [Chinese]. *Int Eye Sci* 2015;**15**:1772–4.
- Sun L, Tao Y. Effects of bevacizumab on CTGF and PEDF in proliferative membrane in patients with PDR [Chinese]. *Int Eye Sci* 2017;**17**:1051–4.
- Tegins E, Javaheri M, Elliott D, Kim L, Salehi-Had H, Hinton D, Sohn E. 1 year clinical outcomes of a randomized clinical trial investigating pre-operative adjunctive bevacizumab for tractional retinal detachment (TRD) due to proliferative diabetic retinopathy (PDR). *Invest Ophthalmol Vis Sci Conference* 2013;**54**.
- Victor AA, Gondhowiardjo TD, Waspadji S, Wanandi SI, Bachtiar A, Suyatna FD, Muhiddin H. Effect of laser photocoagulation and bevacizumab intravitreal in proliferative diabetic retinopathy: review on biomarkers of oxidative stress. *Med J Indones* 2014;**23**:79–86.
- Wang YP, Chen MZ, Chen GC, Chen YJ. Clinical effect of vitrectomy with intravitreal ranibizumab for diabetic retinopathy [Chinese]. *Int Eye Sci* 2014;**14**:1257–9.
- Wildan A, Winarto, Kristina TN. Aflibercept and bevacizumab injection effects on visual acuity of post vitrectomy diabetic retinopathy. *Pakistan J Med Health Sci* 2019;**13**:1214–8.
- Yamaji H, Shiraga F, Shiragami C, Nomoto H, Fujita T, Fukuda K. Reduction in dose of intravitreal bevacizumab before vitrectomy for proliferative diabetic retinopathy. *Arch Ophthalmol* 2011;**129**:106–7.
- Yang XC, Xu JB, Wang RL, Mei Y, Lei H, Liu J, et al. A randomized controlled trial of conbercept pretreatment before vitrectomy in proliferative diabetic retinopathy. *J Ophthalmol* 2016;**2016**:8.
- Yao TT, Yang Y, Jin XL, Wang YX, Zhou YL, Xu AJ, et al. Intraocular pharmacokinetics of anti-vascular endothelial growth factor agents by intraoperative subretinal versus intravitreal injection in silicone oil-filled eyes of proliferative diabetic retinopathy: a randomized controlled pilot study. *Acta Ophthalmol* 2020;**98**:e795–e800.
- Yin N, Zhao S, Zhu HN. Efficacy comparison of conbercept and ranibizumab as pre-treatment for pars plana vitrectomy in proliferative diabetic retinopathy [Chinese]. *Int Eye Sci* 2017;**17**:1300–02.
- Yu XQ, Cao GP, Tang MX. Effect of vitrectomy combined medication hyperplastic on patients with diabetic retinopathy [Chinese]. *Int Eye Sci* 2015;**15**:1402–04.
- Zahaf A, Zghal I, Fekih O, Zayani M, Mahjoub A, Bouguila H. Preoperative intravitreal bevacizumab effects on the course of Pars Plana vitrectomy in diabetic vitreous hemorrhage. *Acta Ophthalmol Conference* 2015;**93**.
- Zaman Y, Rehman AU, Memon AF. Intravitreal Avastin as an adjunct in patients with proliferative diabetic retinopathy undergoing pars plana vitrectomy. *Pakistan J Med Sci* 2013;**29**:590–2.
- Zhao XL, Yang G, Yang J, Zhang JJ. Effect of intravitreal conbercept vs triamcinolone acetonide at the end of surgery on macular structure and function in patients with severe proliferative diabetic retinopathy. *Int J Clin Exp Med* 2017;**10**:14511–18.



Zhou AY, Zhou CJ, Yao J, Quan YL, Ren BC, Wang JM. Panretinal photocoagulation versus panretinal photocoagulation plus intravitreal bevacizumab for high-risk proliferative diabetic retinopathy. *Int J Ophthalmol* 2016;**9**:1772–8.

Zhou J, Liu Z, Chen M, Luo ZH, Li YQ, Qi GY, Liu T. Concentrations of VEGF and PIGF decrease in eyes after intravitreal conbercept injection. *Diabetes Ther Res Treatment Educ Diabetes Related Disorders* 2018;**9**:2393–8.

### Exclude population (others) (17)

Altaweel MM. Changes in severity of diabetic retinopathy following Pegaptanib (Macugen®) Therapy. *Invest Ophthalmol Vis Sci* 2006;**47**:ARVO E-abstract 5441.

Chae JB, Joe SG, Yang SJ, Lee JY, Sung KR, Kim JY, et al. Effect of combined cataract surgery and ranibizumab injection in postoperative macular edema in nonproliferative diabetic retinopathy. *Retina* 2014;**34**:149–56.

Cheema RA, Al-Mubarak MM, Amin YM, Cheema MA. Role of combined cataract surgery and intravitreal bevacizumab injection in preventing progression of diabetic retinopathy: prospective randomized study. *J Cataract Refract Surg* 2009;**35**:18–25.

Euctr-004648-12-Es. *This Is a Phase 3, Multicenter, Randomized, Masked, Controlled, Parallel Group Study of 12 Months Duration in Treatment Naïve Subjects with RVO*. 2017.

Department of Ophthalmology and Medical University of Vienna. *European Intravitreal Avastin® Trial 1*. [www.clinicaltrialsregister.eu/ctr-search/search?query=eudract\\_number:2005-003132-21](http://www.clinicaltrialsregister.eu/ctr-search/search?query=eudract_number:2005-003132-21)

JPRN-jRCTs031180307. *The Effect of an Anti-VEGF Drug on Proliferative Retinopathy*. 2019. URL: <https://jrct.niph.go.jp/latest-detail/jRCTs031180307>

Kodiak Sciences and Inc. *A Prospective, Randomized, Double-masked, Active Comparator-controlled, Multi-center, Two-arm, Phase 3 Study to Evaluate the Efficacy and Safety of Intravitreal KSI-301 Compared with Intravitreal A*.

NCT. *Analysis of Aqueous and Vitreous Humor*. 2017. URL: <https://ClinicalTrials.gov/show/NCT02067013>

Novartis Farmacéutica and SA. *A 12-Month, Phase IIIb, Randomized, Visual Acuity, Assessor-masked, Multicenter Study Assessing the Efficacy and Safety of Ranibizumab 0.5mg in Treat and Extend Regimen Compared to Monthly Regimen*.

Novartis Pharma Services and AG. *A 24-Month Randomized, Double-masked, Multicenter, Phase II Study Assessing Safety and Efficacy of Verteporfin (Visudyne®) Photodynamic Therapy Administered in Conjunction with Lucentis™ versus Luc*.

Novartis Pharma Services and AG. *A 24-Month, Phase IIIb, Open-label, Randomized, Active-controlled, 3-Arm, Multicenter Study Assessing the Efficacy and Safety of an Individualized, Stabilization-criteria-driven PRN Dosing Regimen W*.

Novartis Pharma Services and AG. *A 24-Month, Phase IIIb, Open-label, Single Arm, Multicenter Study Assessing the Efficacy and Safety of an Individualized, Stabilization Criteria-driven PRN Dosing Regimen with 0.5-mg Ranibizumab IN*.

Novartis Pharma Services and AG. *A 24-Month, Phase IIIb, Randomized, Double-masked, Multicenter Study Assessing the Efficacy and Safety of Two Treatment Regimens of 0.5 mg Ranibizumab Intravitreal Injections Guided by Functional A*.

Novartis Pharma and AG. *A 64-Week, Two-arm, Randomized, Double-masked, Multi-center, Phase IIIb Study Assessing the Efficacy and Safety of Brolicizumab 6mg Compared to Aflibercept 2 mg in a Treat-to-Control Regimen in PA*.

NCT. *Effects of Intravitreal Ranibizumab for Macular Edema with Nonproliferative Diabetic Retinopathy*. 2016. URL: <https://ClinicalTrials.gov/show/NCT02834663>

Opthea Limited. *A Phase 3, Multicentre, Double-masked, Randomised Study to Evaluate the Efficacy and Safety of Intravitreal OPT-302 in Combination with Ranibizumab, Compared with Ranibizumab Alone, in Participants*.

Yu B, Liu Z. The clinical efficacy of vitreous injection of ranibizumab in patients with ocular fundus disease and its effect on hemorheology. *Int J Clin Exp Med* 2019;**12**:11249–56.

### Irrelevant intervention (3)

Abadia B, CalvoP, Ferreras A, Bartol F, Verdes G, Pablo L. Clinical applications of dexamethasone for aged eyes. *Drugs Aging* 2016;**33**:639–46.

Altun A, Kanar HS, Aki SF, Arsan A, Hacisalihoglu A. Effectiveness and safety of coadministration of intravitreal dexamethasone implant and silicone oil endotamponade for proliferative diabetic retinopathy with tractional diabetic macular edema. *J Ocular Pharmacol Ther* 2021;**37**:131–7.

CTRI. *A Clinical Study to Assess and Compare the Efficacy and Safety of Hydroxychloroquine and Teneeligiptin in Type 2 Diabetes Patients with Non-proliferative Diabetic Retinopathy*. 2020.

#### **Irrelevant comparator (4)**

Antoszyk AN, Glassman AR, Beaulieu WT, Jampol LM, Jhaveri CD, Punjabi OS, *et al*. Network DRCR Retina. Effect of intravitreal aflibercept vs vitrectomy with panretinal photocoagulation on visual acuity in patients with vitreous hemorrhage from proliferative diabetic retinopathy: a randomized clinical trial. *JAMA* 2020;**324**:2383–95.

Khodabandeh A, Fadaifard S, Abdollahi A, Karkhaneh R, Roohipour R, Abdi F, *et al*. Role of combined phacoemulsification and intravitreal injection of bevacizumab in prevention of postoperative macular edema in non-proliferative diabetic retinopathy. *J Curr Ophthalmol* 2018;**30**:245–9.

Shi R, Ma Y, Wang F, Wang JP. Effects of intravitreal injection on the expression of vascular endothelial growth inhibitor in vitreous of proliferative diabetic retinopathy [Chinese]. *Int Eye Sci* 2015;**15**:985–8.

Yan P, Zhang XH, Zhang L, Li J. Effect of intravitreal injection of ranibizumab combined with voritine on hemorrhagic proliferative diabetic retinopathy and its effect on visual acuity and endothelial growth factor [Chinese]. *Chine J Pharm Biotechnol* 2019;**26**:127–30.

#### **No relevant outcomes (1)**

Khalaf H, Rostamizadeh M, Gonzalez VH. Foveal Avascular Zone in high risk proliferative diabetic retinopathy treated with intravitreal aflibercept injection (ELYSIAN). *Invest Ophthalmol Vis Sci Conference* 2018;**59**.

#### **Inappropriate trial design (27)**

Ababneh OH, Yousef YA, Gharaibeh AM, Abu Ameerh MA, Abu-Yaghi NE, Al Bdour MD. Intravitreal bevacizumab in the treatment of diabetic ocular neovascularization. *Retina* 2013;**33**:748–55.

Abdallah W, Fawzi AA. Anti-VEGF therapy in proliferative diabetic retinopathy. *Int Ophthalmol Clin* 2009;**49**:95–107.

Al-Khersan H, Hariprasad SM, Salehi-Had H. Dexamethasone and anti-VEGF combination therapy for the treatment of diabetic macular edema. *Ophthalmic Surg Lasers Imaging Retina* 2019;**50**:4–7.

Bakri SJ, Donaldson MJ, Link TP. Rapid regression of disc neovascularization in a patient with proliferative diabetic

retinopathy following adjunctive intravitreal bevacizumab. *Eye* 2006;**20**:1474–5.

Beaulieu WT, Bressler NM, Gross JG; Diabetic Retinopathy Clinical Research Network. Panretinal photocoagulation versus ranibizumab for proliferative diabetic retinopathy: patient-centered outcomes from a randomized clinical trial reply. *Am J Ophthalmol* 2017;**177**:233–233.

Bi SS, Jiang T, Chen Y, Ma XF. Effects of laser photocoagulation combined with anti-VEGF drugs at different time in the treatment of diabetic retinopathy. *Int Eye Sci* 2020;**20**:613–8.

Brown DM, Wykoff CC. Intravitreal aflibercept for proliferative diabetic retinopathy. *Lancet* 2017;**390**:2141–2141.

Browning DJ, Lee C, Stewart MW, Landers MB 3rd. Vitrectomy for center-involved diabetic macular edema. *Clin Ophthalmol* 2016;**10**:735–42.

Chen E, Park CH. Use of intravitreal bevacizumab as a preoperative adjunct for tractional retinal detachment repair in severe proliferative diabetic retinopathy. *Retina* 2006;**26**:699–700.

Chen PY, Wang TW, Wang WC, Liao JC, Yang SA, Hsu YT. Clinical Outcome of Diabetic Retinopathy with the Treatment of Photocoagulation versus Anti-VEGF. 2020.

Desapriya E, Khoshpouri P, Al-Isa A. Panretinal photocoagulation versus ranibizumab for proliferative diabetic retinopathy: patient-centered outcomes from a randomized clinical trial. *Am J Ophthalmol* 2017;**177**:232–3.

Ergur O, Bayhan HA, Kurkcuoglu P, Takmaz T, Gurdal C, Can I. Comparison of panretinal photocoagulation (PRP) with PRP plus intravitreal bevacizumab in the treatment of proliferative diabetic retinopathy. [Turkish] Proliferatif diyabetik retinopati tedavisinde tek basina panretinal fotokoagulasyon (PRF) ile PRF ve intravitreal bevacizumab kombinasyonunun karsilastirilmesi. *Retina-Vitreus* 2009;**17**:273–7.

Gibson JM, McGinnigle S. Diabetes: intravitreal ranibizumab for proliferative diabetic retinopathy. *Nature Rev Endocrinol* 2016;**12**:130–1.

Glassman AR. Results of a randomized clinical trial of aflibercept vs panretinal photocoagulation for proliferative

diabetic retinopathy: is it time to retire your laser?. *JAMA Ophthalmol* 2017;**135**:685–6.

Gross JG, Glassman AR. A novel treatment for proliferative diabetic retinopathy: anti-vascular endothelial growth factor therapy. *JAMA Ophthalmol* 2016;**134**:13–4.

Gupta MP, Kiss S, Chan RV. Reversal of retinal vascular leakage and arrest of progressive retinal nonperfusion with monthly anti-vascular endothelial growth factor therapy for proliferative diabetic retinopathy. *Retina* 2018;**38**:e74–e75.

Hershberger V, Hill LF, Tuomi LL, Ghanekar A. Ranibizumab-induced diabetic retinopathy improvement—results from patients at high risk for vision loss in ride/rise and Protocol S. *Diabetes* 2018;**67**:A158.

Krishnan R, Goverdhan S, Lochhead J. Intravitreal pegaptanib in severe proliferative diabetic retinopathy leading to the progression of tractional retinal detachment. *Eye* 2009;**23**:1238–9.

Krzystolik MG, Filippopoulos T, Ducharme JF, Loewenstein JI. Pegaptanib as an adjunctive treatment for complicated neovascular diabetic retinopathy. *Arch Ophthalmol* 2006;**124**:920–1.

Li J, Liu F. Clinical evidence on the treatment of non-proliferative diabetic retinopathy. *Chine J Evidence-Based Med* 2007;**7**:894–8.

Melia M, Edwards A, Kollman C. Interim analysis with sample size re-estimation for binary outcome in a trial of intravitreal ranibizumab versus saline injection for prevention of vitrectomy in eyes with proliferative diabetic retinopathy and vitreous hemorrhage. *Clin Trials* 2012;**9**:523–4.

Olsen TW. Anti-VEGF pharmacotherapy as an alternative to panretinal laser photocoagulation for proliferative diabetic retinopathy. *JAMA* 2015;**314**:2135–6.

Ospedale Sacro Cuore-Don and Calabria. *Evaluation of Safety and Efficacy on Visual Acuity Outcome of Intravitreal Somministration of Bevacizumab in Patients with Diabetic Retinopathy*. [www.clinicaltrialsregister.eu/ctr-search/search?query=eudract\\_number:2006-005315-10](http://www.clinicaltrialsregister.eu/ctr-search/search?query=eudract_number:2006-005315-10)

Parikakis E. Laser or anti-VEGF for proliferative diabetic retinopathy. *Acta Ophthalmol* 2018;**96**:94.

Tan TE, Sivaprasad S, Wong TY. Anti-vascular endothelial growth factor therapy for complications of diabetic retinopathy—from treatment to prevention?. *JAMA Ophthalmol* 2023;**141**:223–5.

Wise J. Lucentis offers treatment alternative for diabetic retinopathy, trial finds. *BMJ* 2015;**351**:h6145.

Zucchiatti I, Bandello F. Intravitreal ranibizumab in diabetic macular edema: long-term outcomes. *Dev Ophthalmol* 2017;**60**:63–70.

### **Not a randomised controlled trial (17)**

CHICTR-OON. *Effect of Anti VEGF on the Expression of Vitreous Ang2 in Patients with PDR* 2017. URL: [www.who.int/trialsearch/Trial2.aspx?TrialID=ChiCTR-OON-17012170](http://www.who.int/trialsearch/Trial2.aspx?TrialID=ChiCTR-OON-17012170)

Chung EJ, Kang SJ, Koo JS, Choi YJ, Grossniklaus HE, Koh HJ. Effect of intravitreal bevacizumab on vascular endothelial growth factor expression in patients with proliferative diabetic retinopathy. *Yonsei Med J* 2011;**52**:151–7.

Department of Ophthalmology and M U W. Disease-modification under treatment with aflibercept in advanced diabetic retinopathy – a pilot study.

EUCTR2006-005315-10-IT. *Evaluation of Safety and Efficacy on Visual Acuity Outcome of Intravitreal Somministration of Bevacizumab in Patients with Diabetic Retinopathy – ND*. 2006.

He F, Yu W. Longitudinal neovascular changes on optical coherence tomography angiography in proliferative diabetic retinopathy treated with panretinal photocoagulation alone versus with intravitreal conbercept plus panretinal photocoagulation: a pilot study. *Eye* 2020;**34**:1413–18.

IRCT138903314232N1. *Intravitreal Bevacizumab (Avastin) Therapy for Proliferative Diabetic Retinopathy*. 2010.

JPRN-UMIN. *Evaluate the Effect of Intravitreal Bevacizumab Injection for Ocular Proliferative Diseases*. 2016.

Kernt M, Cserhati S, Seidensticker F, Liegl R, Kampik A, Neubauer A, et al. Improvement of diabetic retinopathy with intravitreal Ranibizumab. *Diabetes Res Clin Pract* 2013;**100**:e11–3.

Lopez-Lopez F, Gomez-Ulla F, Rodriguez-Cid MJ, Arias L. Triamcinolone and bevacizumab as adjunctive therapies to panretinal photocoagulation for proliferative diabetic retinopathy. *ISRN Ophthalmol Print* 2012;**2012**:267643.

NCT. *Intravitreal Bevacizumab for Management of Active Progressive Proliferative Diabetic Retinopathy (PDR)*. 2006. URL: <https://ClinicalTrials.gov/show/NCT00370721>

NCT. *Analysis of Angiogenic Factor Levels in Eyes with Diabetic Retinopathy*. 2012. URL: <https://ClinicalTrials.gov/show/NCT02026843>

NCT. *Combined Triple Therapy in Diabetic Retinopathy (DRP)*. 2012. URL: <https://clinicaltrials.gov/study/NCT00806169>

NCT. *Effect of Macugen (Pegaptanib) on Surgical Outcomes and VEGF Levels in Diabetic Patients with PDR (Diabetic Retinopathy or CSDME (Macular Edema))*. 2012. URL: <https://ClinicalTrials.gov/show/NCT00446381>

NCT. *Ziv-aflibercept in Ocular Disease Requiring Anti-VEGF Injection*. 2015. URL: <https://ClinicalTrials.gov/show/NCT02486484>

Park YJ, Ahn J, Kim TW, Park SJ, Joo K, Park KH, Shin JY. Efficacy of bevacizumab for vitreous haemorrhage in proliferative diabetic retinopathy with prior complete panretinal photocoagulation. *Eye*:8.

Park JM, Lee SJ. The effect of panretinal photocoagulation and additive intravitreal bevacizumab injections on central retinal vessel diameters in diabetic retinopathy. *Acta Ophthalmol Conference* 2015;93.

Vidinova CN, Gouguchkova PT, Dimitrov T, Vidinov KN, Nocheva H. [Comparative clinical and ultrastructural analysis of the results from ranibizumab and aflibercept in patients with PDR]. *Klinische Monatsblätter für Augenheilkunde* 2020;237:79–84.

### **Protocols of excluded and ongoing studies (20)**

Frimley Park Hospital and NHS. Foundation Trust. A randomised controlled trial of efficacy of Pegaptanib sodium in the prevention of proliferative diabetic retinopathy.

Fakultní nemocnice Královské and Vinohrady. A randomized, 12 months, active controlled study of the efficacy of repeated doses of intravitreal aflibercept in subjects with proliferative diabetic retinopathy.

Euctr-000658-30-Ie. *Randomised Controlled Trial of Intravitreal Bevacizumab vs. Conventional Treatment for Proliferative Diabetic Retinopathy – Randomised Controlled Trial of Intravitreal Bevacizumab vs. Conventional Treatment for Proliferative*. 2007.

Euctr-001856-36-Fr. *Efficacy and Safety of Aflibercept (Eylea®) in Proliferative Diabetic Retinopathy*. 2016.

Euctr-004203-39-Cz. *Study of Effect of Intravitreal Aflibercept in Subjects with Proliferative Diabetic Retinopathy*. 2014.

Euctr-006795-10-Gb. *A Randomised Controlled Trial of Efficacy of Pegaptanib Sodium in the Prevention of Proliferative Diabetic Retinopathy – EPPDR*. 2008.

ISRCTN. *A Prospective Randomised Controlled Trial Assessing the Efficacy of Pegaptanib Sodium (Macugen®) in the Prevention of Proliferative Diabetic Retinopathy*. 2010.

NCT. *Ranibizumab for Treatment of Persistent Diabetic Neovascularization Assessed by Wide-Field Imaging*. 2008. URL: <https://clinicaltrials.gov/show/NCT00606138>

NCT. *Prospective, Randomized, Open Label, Phase II Study to Assess Efficacy and Safety of Macugen® (Pegaptanib 0.3 mg Intravitreal Injections) Plus Panretinal Photocoagulation and PRP (Monotherapy) in the Treatment with High Risk PDR*. 2011. URL: <https://clinicaltrials.gov/show/NCT01281098>

NCT. *Prevention of Macular Edema in Patients with Diabetic Retinopathy Undergoing Cataract Surgery*. 2013. URL: <https://clinicaltrials.gov/show/NCT01988246>

NCT. *Treatment with Intravitreal Aflibercept Injection for Proliferative Diabetic Retinopathy, The A.C.T Study*. 2013. URL: <https://clinicaltrials.gov/show/NCT01813773>

NCT. *Safety and Efficacy of Aflibercept in Proliferative Diabetic Retinopathy*. 2015. URL: <https://ClinicalTrials.gov/show/NCT02151695>

NCT. *Conbercept vs Panretinal Photocoagulation for the Management of Proliferative Diabetic Retinopathy*. 2016. URL: <https://clinicaltrials.gov/show/NCT02911311>

NCT. *Intravitreal Aflibercept as Indicated by Real-Time Objective Imaging to Achieve Diabetic Retinopathy Improvement*. 2018. URL: <https://clinicaltrials.gov/show/NCT03531294>

NCT. *Multicenter Clinical Study of Anti-VEGF Treatment on High Risk Diabetic Retinopathy (DR)*. 2018. URL: <https://clinicaltrials.gov/show/NCT03452657>

NCT. *A Multicenter, Randomized Study in Participants with Diabetic Retinopathy without Center-involved Diabetic*

Macular Edema to Evaluate the Efficacy, Safety, and Pharmacokinetics of Ranibizumab Delivered via the Port Delivery System Relative to the Comparator Arm. 2020. URL: <https://clinicaltrials.gov/show/NCT04503551>

NCT. Intravitreal Bevacizumab for Nonproliferative Diabetic Retinopathy. 2020. URL: <https://clinicaltrials.gov/show/NCT04511715>

NCT. Study of Efficacy and Safety of Brolocizumab versus Panretinal Photocoagulation Laser in Patients with Proliferative Diabetic Retinopathy. 2020. URL: <https://ClinicalTrials.gov/show/NCT04278417>

NCT. Intravitreal Bevacizumab vs Laser vs Combination of Bevacizumab and Modified Laser in PDR. 2021. URL: <https://clinicaltrials.gov/show/NCT04800679>

TCTR. Change of OCT Findings after Intravitreal Anti-VEGF Injection in Patients with Diabetic Tractional Retinal Detachment: A Randomized Controlled Trial. 2021.

### Irretrievable (1)

Neri Alvarez-Villalobos Humberto de León-Gutiérrez Fernando Ruiz-Hernandez. Safety and clinical effectiveness behavior of bevacizumab biosimilars in the intravitreal application.

## Trials not included in meta-analyses

TABLE 3 Trials not included in meta-analyses

Trial	Key paper(s)	Anti-VEGF	Comparator	Location	Sample size	Population
<b>No PRP arm</b>						
RECOVERY	Alagorie 2021	Aflibercept (monthly)	Aflibercept (quarterly)		40 eyes	PDR
<b>Conference abstracts</b>						
Garcia	Garcia-Aguirre 2008	Bevacizumab	PRP	Mexico	10 persons	NPDR, PDR
Ernst	Ernst 2012	Bevacizumab	PRP	USA	10 persons	NPDR, PDR
MEDICARE	Dufour 2017	Aflibercept	PRP	France	20 persons	PDR
Oh	Oh 2014 CA	Bevacizumab (+ PRP)	PRP	South Korea	125 persons	NPDR, PDR
Ramezani	Ramezani 2021	Bevacizumab (+ PRP)	PRP	Unknown	153 eyes	PDR
Tardieu	Tardieu 2022	Not stated	PRP	Unknown	40 persons	PDR
<b>Papers in Chinese</b>						
Bi	Bi 2020	Ranibizumab (+ PRP)	PRP	China	120 persons	Unclear
Meng	Meng 2019	Ranibizumab (+ PRP)	PRP	China	80 persons	PDR
Zhou	Zhou 2017	Bevacizumab (+ PRP)	PRP	China	30 persons	Unclear
<b>Trials from before 2010</b>						
Cho	Cho 2009–2010	Bevacizumab (+ PRP)	PRP + Triamcinolone	China	91 eyes	NPDR, PDR
Mirshahi	Mirshahi 2008	Bevacizumab (+ PRP)	PRP, Sham injection	Iran	80 eyes	PDR

continued

TABLE 3 Trials not included in meta-analyses (continued)

Trial	Key paper(s)	Anti-VEGF	Comparator	Location	Sample size	Population
Tonello	Tonelo 2008	Bevacizumab (+ PRP)	PRP	Brazil	30 eyes	PDR
<b>Unused or unspecified anti-VEGFs</b>						
Chen/Zhou	Chen 2017	Unclear	PRP	China	120 persons	PDR
Gonzalez	Gonzalez 2007/2009/2014	Pegaptanib sodium	PRP	USA	20 persons	PDR
He	He 2020	Conbercept (+ PRP)	PRP	China	44 eyes	PDR
Leal	Leal 2013	Pegaptanib sodium (+ PRP)	PRP	Portugal	22 persons	PDR
Wang	Wang 2019	Conbercept (+ PRP)	PRP	China	64 persons	NPDR, PDR
<b>No protocol-specified outcomes</b>						
Helmy	Helmy 2023	Ranibizumab	PRP	Egypt	50 persons	PDR
Preti	Preti 2013	Bevacizumab (+ PRP)	PRP	S. America	42 persons	PDR
Rentiya	Rentiya 2022	Ranibizumab (+ PRP)	PRP	Brazil	30 persons	PDR

## Risk-of-bias assessment

TABLE 4 Full risk-of-bias assessment – Table A

Trial	Randomisation process		Deviations from intended interventions		Missing outcome data	
	Judgement	Comments	Judgement	Comments	Judgement	Comments
Ahmad 2012 <sup>27</sup>	Some concerns	Randomised by 'simple lottery'. No further details No allocation concealment method reported No evidence of significant differences in key prognostic factors	Some concerns	No placebo States 'the physician did not know which eye has been injected', but the control group did not receive a placebo injection No CONSORT diagram, and no reporting of deviation from the intervention due to the trial context ITT/mITT not reported. The risk that the analysis was not based on ITT principles cannot be excluded	Low	All participants completed the 90 days follow-up

TABLE 4 Full risk-of-bias assessment – Table A (continued)

Trial	Randomisation process		Deviations from intended interventions		Missing outcome data	
	Judgement	Comments	Judgement	Comments	Judgement	Comments
Ali 2018 <sup>28</sup>	Some concerns	States the study is randomised, with allocation by 'simple lottery method'. No further details No information on whether allocation was concealed	Some concerns	No placebo. Contralateral design No CONSORT diagram, and no reporting of deviation from the intervention due to the trial context ITT/mITT not reported The risk that the analysis was not based on ITT principles cannot be excluded	Some concerns	No information on loss to follow-up. No evidence that the result was not biased by any possible missing outcome data Likelihood of significant missingness may be limited by relatively short follow-up duration
CLARITY <sup>23</sup>	Low	Computer generated with minimisation. Central allocation by trials unit No significant baseline imbalances	Low	No placebo. 'The treating ophthalmologists and participants were not masked' CONSORT diagram reported. No evidence of deviation from intended intervention due to the trial context Analyses conducted according to ITT principles	Low	Available for 91% (211/232) at 52 weeks Appropriate sensitivity analyses for missing BCVA data with prespecified alternative scenarios were conducted and showed no evidence of bias
Ferraz <i>et al.</i> 2015 <sup>32</sup>	Some concerns	Described as randomised. No other details  No details on allocation concealment  Contralateral design  No evidence of significant differences in key prognostic factors	Some concerns	Placebo controlled. Contralateral design  Trial registry entry described as single masked (participants)  Masking only reported for outcome assessors ('examiners' and participants), not for carers  No CONSORT diagram, and no reporting of deviation from the intervention due to the trial context  ITT/mITT not reported  The risk that the analysis was not based on ITT principles cannot be excluded	Low	3% (2/60) eyes excluded due to VH in the control arm. It appears that all other randomised eyes were analysed
Marashi 2017 <sup>26</sup>	High	Described as randomised. No other details No details on allocation concealment Eighty per cent had DME at baseline in the IVB arm vs. 20% in the control arm Although the trial is small, the difference is large and considered unlikely to be due to chance alone. No adjustments for baseline imbalance were performed	Some concerns	No placebo No CONSORT diagram, and no reporting of deviation from the intervention due to the trial context ITT/mITT not reported The risk that the analysis was not based on ITT principles cannot be excluded	Some concerns	No information on loss to follow-up. Follow-up duration means that the risk of at least some loss to follow-up is high No evidence that the result was not biased by any possible missing outcome data

continued

TABLE 4 Full risk-of-bias assessment – Table A (continued)

Trial	Randomisation process		Deviations from intended interventions		Missing outcome data	
	Judgement	Comments	Judgement	Comments	Judgement	Comments
PANORAMA <sup>25</sup>	Low	<p>Patients were randomised according to a central randomisation scheme with treatment assignments provided by an interactive voice response system/interactive web response system to the designated study pharmacist (or qualified designee)</p> <p>Some differences in sex at baseline: higher rate of males in 2q16 (56%) and 2q8 (60%) compared with control (52%), but no other imbalances in reported baseline characteristics</p>	Low	<p>Placebo controlled. Participants, outcome assessors and study personnel were masked throughout the study period, except for study drug administration which was done by an unmasked physician</p> <p>Rates of participants not assessed were higher in the control group (73%) at 100 weeks (and 52 weeks) compared with aflibercept arms (84% and 83%), although participants were masked throughout the study period, and there was no evidence of changes from the intended intervention that occurred because of the trial context</p> <p>All participants analysed. LOCF imputation method used</p>	Low	<p>Rates of participants not assessed were higher in the control group (73%) at 100 weeks compared with aflibercept arms (84% and 83%)</p> <p>Sensitivity analysis: primary efficacy analysis was also performed using all observed measurements (regardless of whether rescue treatment was given). Protocol also stated that for sensitivity analyses, only true missing values would be imputed using the LOCF procedure, and that baseline values would be carried forward if all post-baseline observations were missing or non-gradable</p> <p>Sensitivity analysis results showed similar results to main analyses for DRSS, although all are based on the LOCF principle, and sensitivity analyses were not performed for BCVA</p> <p>The risk that the higher rate of missingness in the control arm is partly due to its true value cannot be excluded. However, due to the size of the difference in missingness, any possible bias arising is likely to be small</p>



TABLE 4 Full risk-of-bias assessment – Table A (continued)

Trial	Randomisation process		Deviations from intended interventions		Missing outcome data	
	Judgement	Comments	Judgement	Comments	Judgement	Comments
PRIDE <sup>33</sup>	Some concerns	A number of differences in baseline characteristics, including key variables, although differences do not clearly favour one arm and may have occurred by chance. Differences in mean age (ranibizumab: 52.5, PRP: 53, ranibizumab + PRP: 55), age distribution (< 65 years: 86%, 86%, 72%); smoker (14%, 26%, 35%); duration of diabetes (25 years, 23 years, 21 years), Mean mm <sup>2</sup> NVD + NVE: 9.4, 5.4, 4.1; ETDRS: 83.3, 80.5, 80.0	Low	No masking Analyses conducted based on ITT principle, using LOCF	Some concerns	23% (25/108) of randomised participants not measured at 12 months No significant differences in rates of missingness across groups
PROTEUS <sup>34</sup>	Low	Computer-generated block randomisation. Central allocation implemented through electronic platform Large and statistically significant difference in mean age [ranibizumab + PRP: 58.8 years (13.3), PRP: 52.0 (11.9)]. Non-statistically significant difference in sex (31.7% vs. 41.3% female) Difference in time since diagnosis not reported In a multivariate analysis, 'age, HbA1c, and number of PRP treatments did not show a significant association with BCVA difference from baseline to month 12' Re-analysis with IPD provided by trialist suggested low concerns	Low	CONSORT diagram reported. No evidence of deviation from intended intervention due to trial context ITT-principle-based primary analysis	Some concerns	
Protocol S	Low	Permuted block randomisation. Stratification by site and presence of centrally involved DME Central allocation concealment with web-based tool from trials unit No evidence of baseline imbalances	Low	No placebo. Masking only for outcome assessors All eyes randomised received the treatment allocated Analyses conducted according to ITT principles	Low	83% (382/394) completed 2-year follow-up. Of those, 5% (18/394) died, 12% (48/394) withdrew or missed their visit For missing data at 2 years, statistical analysis plan reports 'Markov chain Monte Carlo multiple imputation-based on treatment group, the randomisation stratification factors, and all available visual acuity data from assessment visits prior to 2 years'

continued

TABLE 4 Full risk-of-bias assessment – Table A (continued)

Trial	Randomisation process		Deviations from intended interventions		Missing outcome data	
	Judgement	Comments	Judgement	Comments	Judgement	Comments
Protocol W <sup>24</sup>	Low	Central, web-based (DRCR network) randomisation, stratified by DR severity level No evidence of baseline imbalances	Low	Placebo controlled. Participants masked. Investigators and study co-ordinators unmasked CONSORT diagram reported. No evidence of deviation from intended intervention due to trial context Analyses conducted according to ITT principles	Low	68.5% (137/200, or 74.9% excluding 17 deaths) completed their 4-year visit in intervention arm, vs. 67.3% (134/199, 73.2% excluding 16 deaths) Multiple imputation (Markov model) used for missing data (assumes data are missing at random). Model included treatment group, study eye laterality, baseline DRSS, baseline visual acuity and change in visual acuity from baseline to each protocol assessment visit up to and including 4 years. Missingness documented, balanced between arms and unlikely to depend on its true value
Rebecca 2021 <sup>29</sup>	Some concerns	Described as randomised. No other details No details on allocation concealment No evidence of significant differences in key prognostic factors	Some concerns	No placebo No CONSORT diagram, and no reporting of deviation from the intervention due to the trial context ITT/mITT not reported The risk that the analysis was not based on ITT principles cannot be excluded	Some concerns	No information on loss to follow-up. Follow-up duration means that the risk of at least some loss to follow-up is high No evidence that the result was not biased by any possible missing outcome data

TABLE 4 Full risk-of-bias assessment – Table A (continued)

Trial	Randomisation process		Deviations from intended interventions		Missing outcome data	
	Judgement	Comments	Judgement	Comments	Judgement	Comments
Roohipoor 2019 <sup>30</sup>	Some concerns	Random block method, but no further details on how allocation sequence was generated. No information on allocation concealment No evidence of significant differences in key prognostic factors	Some concerns	No placebo. Contralateral design No CONSORT diagram, and no reporting of deviation from the intervention due to the trial context ITT/mITT not reported The risk that the analysis was not based on ITT principles cannot be excluded	Some concerns	Significant loss to follow-up. Only 59% (19 out of 32) completed 10 months follow-up No evidence that the result was not biased by missing outcome data Reasons for loss to follow-up were not reported. The risk that at least some missingness could be due to visual acuity outcomes cannot be excluded
Sao Paulo A <sup>36</sup>	Some concerns	Randomised based on a computer-generated sequence. No further details reported There were differences in age (mean PASCAL arm age was 7.5 years older than ranibizumab and 2.2 years older than ETDRS), although they were not statistically significant	Some concerns	No placebo No evidence of deviation from the intervention due to the trial context ITT/mITT not explicitly reported	Some concerns	13/48 (27%) withdrew. No significant difference in withdrawal between arms No evidence that the result was not biased by missing outcome data Reasons for loss to follow-up were not reported. The risk that at least some missingness could be due to visual acuity outcomes cannot be excluded
Sao Paulo B <sup>35</sup>	Some concerns	Block randomisation (blocks of 2), allocation drawn randomly by technician from one of two identical opaque envelopes. No further information on randomisation and allocation concealment No evidence of significant differences in key prognostic factors	Some concerns	No placebo No CONSORT diagram, and no reporting of deviation from the intervention due to the trial context ITT/mITT not reported The risk that the analysis was not based on ITT principles cannot be excluded	Some concerns	Only 72.5% (29/40) participants analysed at 48 weeks No evidence that the result was not biased by missing outcome data Significant loss to follow-up. Reported reasons for loss to follow-up were generally appropriate (incl. four deaths and two ocular events, four did not return for assessment, one not specified). No clear imbalances between arms

CONSORT, consolidated standards of reporting trials; DRCR, diabetic retinopathy clinical research retina network; ITT, intention to treat; IVB, intravitreal bevacizumab; LOCF, last-observation carried forward; mITT, modified intention to treat.

TABLE 5 Full risk-of-bias assessment – Table B

Trial	Measurement of the outcome		Selection of the reported result		Overall bias
	Judgement	Comments	Judgement	Comments	Judgement
Ahmad 2012 <sup>27</sup>	High	Snellen chart, converted to log-MAR Participants unmasked (no placebo). No mention of blinding of outcome assessors Participants and study personnel may have been influenced by knowledge of the intervention	Some concerns	Insufficient information about analysis plans	High
Ali 2018 <sup>28</sup>	High	Appears to be ETDRS, standard scale No placebo	Some concerns	No protocol	High
CLARITY <sup>23</sup>	Some concerns	ETDRS, standard scale The lack of blinding of participants means raises some concerns, although appropriate steps were taken to mask the optometrists assessing BCVA Optometrists 'masked to treatment allocation throughout the study. The optometrists received the participants into the visual acuity lanes with a visual acuity-specific source data worksheet that included the PIN and details of the study eye and non-study eye to be refracted, but with no previous records or case report forms by which the patient's treatment arm could be identified'	Low	A SAP 'was finalised before data lock and agreed with oversight committees'	Low
Ferraz 2015 <sup>32</sup>	Low	ETDRS Outcome assessors masked throughout the study period	Some concerns	Insufficient information about analysis plans. Outcome retrospectively reported in trial registry	Some concerns
Marashi 2017 <sup>26</sup>	High	Snellen scale, converted to log-MAR No placebo Participants and study personnel may have been influenced by knowledge of the intervention	Low	Protocol registered around time of study start, and prespecified outcome and time point were reported	High
PANORAMA <sup>25</sup>	Low	ETDRS method Outcome assessors were masked throughout the study period	Low		Low
PRIDE <sup>33</sup>	High	ETDRS, standard. No masking of outcome assessors	Low	SAP not mentioned Protocol registered before time of study start, and prespecified outcome and time point were reported	Some concerns
PROTEUS <sup>34</sup>	High	Standard ETDRS No placebo. Participants and outcome assessors were aware of the intervention Participants and study personnel may have been influenced by knowledge of the intervention	Low	No SAP Outcome and follow-up specified in prospectively registered protocol	Some concerns

TABLE 5 Full risk-of-bias assessment – Table B (continued)

Trial	Measurement of the outcome		Selection of the reported result		Overall bias
	Judgement	Comments	Judgement	Comments	Judgement
Protocol S <sup>31</sup>	Some concerns	E-ETDRS Participants unmasked (no placebo), but protocol states that 'visual acuity testers [...] will be masked to treatment group at annual visits'	Low	SAP v1.0 is dated March 2015 Protocol first published December 2011, primary completion dated January 2015 Outcome specified in prospectively registered protocol	Low
Protocol W <sup>24</sup>	Low	DRSS Outcome assessors masked	Low	SAP reported and finalised before unblinded outcome data were available for analysis	Low
Rebecca 2021 <sup>29</sup>	High	BCVA. Scale not reported, but standard outcome No placebo. Participants and outcome assessors were aware of the intervention Participants and study personnel may have been influenced by knowledge of the intervention	Some concerns	Insufficient information about analysis plans	High
Roohipoor 2019	High	BCVA measured using standard Snellen chart No placebo Participants and study personnel may have been influenced by knowledge of the intervention	Some concerns	SAP not mentioned in protocol or publication. 10 months follow-up assessment was not pre-specified (unlike 6 months)	High
Sao Paulo A	High	Standard ETDRS No placebo. Participants were aware of the intervention. No masking of outcome assessor reported	Some concerns	No SAP Outcome and follow-up specified in protocol, but unclear if prospectively registered	High
Sao Paulo B	High	ETDRS, converted to log-MAR No blinding of outcome assessor, who performed the interventions Participants and study personnel may have been influenced by knowledge of the intervention	Some concerns	Insufficient information about analysis plans	High

SAP, statistical analysis plan.

## Appendix 2 Proliferative diabetic retinopathy: all best corrected visual acuity analyses

All figures and tables relate to the trials of PDR, excluding the two trials (PANORAMA, Protocol W) of non-proliferative retinopathy. For their results, see [Appendix 4](#).




FIGURE 9 All ETDRS data (as mean change from baseline) by drug and type of intervention.




FIGURE 10 All ETDRS data (as mean change from baseline) by trial and drug type.




FIGURE 11 Mean difference in ETDRS between anti-VEGF and control arms over time.




FIGURE 12 Mean difference between anti-VEGF and control arms by ETDRS at randomisation.

## Figures and forest plots summarising best corrected visual acuity data

Note from these figures that there appears to be a possible decline in benefit to vision over time, and that the benefit

of anti-VEGF may be greater in people with poorer initial vision, but these differences may be confounded by differences between types of anti-VEGF.




FIGURE 13 Forest plot of all mean differences in ETDRS between anti-VEGF and control (right side favours anti-VEGF).





FIGURE 14 Forest plot of all mean differences in log-MAR between anti-VEGF and control (left side favours anti-VEGF).




## Forest plots of meta-analyses of best corrected visual acuity

## Up to 1 year




**FIGURE 15** Meta-analysis of mean differences in ETDRS between anti-VEGF and control up to 1 year of follow-up (right side favours anti-VEGF).




**FIGURE 16** Meta-analysis of mean differences in log-MAR between anti-VEGF and control up to 1 year of follow-up (left side favours anti-VEGF).

## One to 2 years' follow-up



**FIGURE 17** Meta-analysis of mean differences in ETDRS between anti-VEGF and control with 1–2 years' of follow-up (right side favours anti-VEGF).



**FIGURE 18** Meta-analysis of mean differences in log-MAR between anti-VEGF and control with 1–2 years' of follow-up (left side favours anti-VEGF).

Maximum follow-up in trial (up to 2 years)




FIGURE 19 Meta-analysis of mean differences in ETDRS between anti-VEGF and control at end of trial (right side favours anti-VEGF).




FIGURE 20 Meta-analysis of mean differences in log-MAR between anti-VEGF and control at end of trial (left side favours anti-VEGF).

## Network meta-analyses of best corrected visual acuity (using logarithm of the minimum angle of resolution)

Note: From this point forward on, meta-analyses of BCVA measured using log-MAR are presented. Some analyses

using ETDRS were performed but are not included here. Similarly, only random-effects analyses are presented for simplicity, as differences between random- and fixed-effect analyses were minimal.

### Analyses at up to 1 year of follow-up




FIGURE 21 Network diagram of BCVA at up to 1 year of follow-up.





FIGURE 22 All treatment comparisons for 1-year random-effects NMA of log-MAR.



**FIGURE 23** Probability of treatments for 1-year random-effects NMA of log-MAR.


**TABLE 6** Results of NMA of log-MAR up to 1 year – comparisons between treatments

Intervention	Comparator	Mean difference	95% CI	
Aflibercept	PRP	-0.084	-0.222	0.056
Bevacizumab	PRP	-0.198	-1.213	0.785
Bevacizumab + PRP	PRP	-0.172	-0.279	-0.069
Ranibizumab	PRP	-0.121	-0.233	-0.006
Ranibizumab + PRP	PRP	-0.078	-0.165	0.013
Bevacizumab	Aflibercept	-0.115	-1.142	0.853
Bevacizumab + PRP	Aflibercept	-0.088	-0.273	0.082
Ranibizumab	Aflibercept	-0.037	-0.213	0.130
Ranibizumab + PRP	Aflibercept	0.006	-0.151	0.173
Bevacizumab + PRP	Bevacizumab	0.026	-0.947	1.027
Ranibizumab	Bevacizumab	0.077	-0.913	1.098
Ranibizumab + PRP	Bevacizumab	0.121	-0.867	1.151
Ranibizumab	Bevacizumab + PRP	0.051	-0.095	0.208
Ranibizumab + PRP	Bevacizumab + PRP	0.094	-0.040	0.236
Ranibizumab + PRP	Ranibizumab	0.043	-0.067	0.160


**TABLE 7** Results of NMA of log-MAR up to 1 year – ranking probabilities

Treatment arm	Probability of ranking					
	1st (%)	2nd (%)	3rd (%)	4th (%)	5th (%)	6th (%)
PRP	0.00	0.13	0.65	4.93	37.65	56.65
Aflibercept	4.33	12.65	24.65	30.48	23.10	4.80
Bevacizumab	50.73	5.05	2.38	2.63	3.83	35.40
Bevacizumab + PRP	33.60	44.23	14.73	5.68	1.73	0.05
Ranibizumab	10.25	30.30	33.60	17.40	7.25	1.20
Ranibizumab + PRP	1.10	7.65	24.00	38.90	26.45	1.90


**Analyses at 1–2 years' follow-up**



**FIGURE 24** Network diagram of BCVA at up to 1–2 years of follow-up.



**FIGURE 25** All treatment comparisons for 1–2 year random-effects NMA of log-MAR.



**FIGURE 26** Probability of treatments for 1–2-year random-effects NMA of log-MAR.

**TABLE 8** Results of NMA of log-MAR 1–2 years – comparisons between treatments


Intervention	Comparator	Mean difference	95% CI	
Aflibercept	PRP	-0.080	-0.225	0.100
Bevacizumab	PRP	-0.182	-1.181	0.816
Ranibizumab	PRP	-0.072	-0.171	0.017
Ranibizumab + PRP	PRP	-0.068	-0.152	0.020
Bevacizumab	Aflibercept	-0.102	-1.095	0.899
Ranibizumab	Aflibercept	0.008	-0.200	0.187
Ranibizumab + PRP	Aflibercept	0.012	-0.174	0.189
Ranibizumab	Bevacizumab	0.110	-0.887	1.104
Ranibizumab + PRP	Bevacizumab	0.114	-0.885	1.112
Ranibizumab + PRP	Ranibizumab	0.004	-0.100	0.114

**TABLE 9** Results of NMA of log-MAR 1–2 years – ranking probabilities


Treatment arm	Probability of ranking				
	1st (%)	2nd (%)	3rd (%)	4th (%)	5th (%)
PRP	0.05	1.20	6.25	37.93	54.58
Aflibercept	21.38	34.58	20.98	17.00	6.08
Bevacizumab	56.20	3.15	2.88	3.35	34.43
Ranibizumab	13.13	30.28	34.13	20.15	2.33
Ranibizumab + PRP	9.25	30.80	35.78	21.58	2.60

This article should be referenced as follows:

Simmonds M, Llewellyn A, Walker R, Fulbright H, Walton M, Hodgson R, *et al.* Anti-VEGF drugs compared with laser photocoagulation for the treatment of diabetic retinopathy: a systematic review and meta-analysis. [published online ahead of print December 11 2024]. *Health Technol Assess* 2024. <https://doi.org/10.3310/PCGV5709>

*Analysis at maximum follow-up time (up to 2 years)*

**FIGURE 27** All treatment comparisons for end-of-trial random-effects NMA of log-MAR.



**FIGURE 28** Probability of treatments for end-of-trial random-effects NMA of log-MAR.



**TABLE 10** Results of NMA of log-MAR at end of trial – comparisons between treatments

Intervention	Comparator	Mean difference	95% CI	
Aflibercept	PRP	-0.087	-0.228	0.049
Bevacizumab	PRP	-0.209	-1.176	0.782
Bevacizumab + PRP	PRP	-0.171	-0.284	-0.064
Ranibizumab	PRP	-0.085	-0.177	-0.004
Ranibizumab + PRP	PRP	-0.069	-0.151	0.016
Bevacizumab	Aflibercept	-0.122	-1.117	0.881
Bevacizumab + PRP	Aflibercept	-0.085	-0.265	0.093
Ranibizumab	Aflibercept	0.002	-0.167	0.151
Ranibizumab + PRP	Aflibercept	0.017	-0.139	0.180
Bevacizumab + PRP	Bevacizumab	0.038	-0.956	1.030
Ranibizumab	Bevacizumab	0.124	-0.868	1.100
Ranibizumab + PRP	Bevacizumab	0.140	-0.830	1.116
Ranibizumab	Bevacizumab + PRP	0.087	-0.054	0.225
Ranibizumab + PRP	Bevacizumab + PRP	0.102	-0.032	0.243
Ranibizumab + PRP	Ranibizumab	0.015	-0.086	0.128

**TABLE 11** Results of NMA of log-MAR at end of trial – ranking probabilities

Treatment arm	Probability of ranking					
	1st (%)	2nd (%)	3rd (%)	4th (%)	5th (%)	6th (%)
PRP	0.00	0.63	4.80	35.88	58.70	0.00
Aflibercept	20.58	29.20	23.05	17.80	4.35	20.58
Bevacizumab	5.90	2.70	1.85	3.65	33.43	5.90
Bevacizumab + PRP	46.80	9.78	3.88	1.23	0.10	46.80
Ranibizumab	17.03	31.83	31.10	16.20	0.93	17.03
Ranibizumab + PRP	9.70	25.88	35.33	25.25	2.50	9.70

## Network meta-analyses allowing for interaction with follow-up time and best corrected visual acuity at randomisation

### Allowing for variation with follow-up time

Network meta-analyses incorporating all follow-up times

(longest in each trial), allowing varying effect of anti-VEGF with follow-up time. Time variation is assumed to be the same for all types of anti-VEGF. A selection of output plots is presented. Results are presented for the predicted effects after 1 year of follow-up.




FIGURE 29 All treatment comparisons for time-adjusted random-effects NMA of log-MAR.

TABLE 12 Results of NMA of log-MAR adjusting for time – comparisons between treatments

Intervention	Comparator	Mean difference	95% CI
Aflibercept	PRP	-0.086	-0.221 0.045
Bevacizumab	PRP	-0.199	-1.218 0.858
Bevacizumab + PRP	PRP	-0.112	-0.245 0.027
Ranibizumab	PRP	-0.119	-0.214 -0.023
Ranibizumab + PRP	PRP	-0.075	-0.153 0.001
Bevacizumab	Aflibercept	-0.112	-1.135 0.945
Bevacizumab + PRP	Aflibercept	-0.026	-0.216 0.163
Ranibizumab	Aflibercept	-0.033	-0.200 0.130
Ranibizumab + PRP	Aflibercept	0.011	-0.138 0.163
Bevacizumab + PRP	Bevacizumab	0.086	-0.957 1.127
Ranibizumab	Bevacizumab	0.080	-0.973 1.105
Ranibizumab + PRP	Bevacizumab	0.123	-0.919 1.138
Ranibizumab	Bevacizumab + PRP	-0.007	-0.197 0.184
Ranibizumab + PRP	Bevacizumab + PRP	0.037	-0.123 0.195
Ranibizumab + PRP	Ranibizumab	0.044	-0.064 0.155


**TABLE 13** Results of NMA of log-MAR adjusting for time – ranking probabilities

Treatment arm	Probability of ranking					
	1st (%)	2nd (%)	3rd (%)	4th (%)	5th (%)	6th (%)
PRP	0.00	0.05	0.65	5.13	38.10	56.08
Aflibercept	7.60	19.98	25.30	25.10	17.48	4.55
Bevacizumab	51.98	3.88	2.58	2.78	4.35	34.45
Bevacizumab + PRP	18.73	30.70	17.90	17.35	12.35	2.98
Ranibizumab	19.40	34.13	25.88	14.88	5.25	0.48
Ranibizumab + PRP	2.30	11.28	27.70	34.78	22.48	1.48

### **Allowing for variation over time and by logarithm of the minimum angle of resolution at randomisation**

Network meta-analyses incorporating all follow-up times (longest in each trial), allowing for varying effect of anti-VEGF by follow-up duration and varying effect by trial

mean log-MAR at randomisation. Time and log-MAR variation are assumed to be the same for all types of anti-VEGF. A selection of output plots is presented. Results are presented for the predicted effects after 1 year of follow-up and at mean baseline BCVA across trials.

**FIGURE 30** All treatment comparisons for time-adjusted and baseline BCVA adjusted random-effects NMA of log-MAR.

**TABLE 14** Results of NMA of log-MAR adjusting for time and baseline BCVA – comparisons between treatments

Intervention	Comparator	Mean difference	95% CI	
Aflibercept	PRP	-0.085	-0.310	0.119
Bevacizumab + PRP	PRP	-0.116	-0.281	0.050
Ranibizumab	PRP	-0.117	-0.247	0.011
Ranibizumab + PRP	PRP	-0.073	-0.187	0.041
Bevacizumab + PRP	Aflibercept	-0.031	-0.287	0.228
Ranibizumab	Aflibercept	-0.032	-0.288	0.233
Ranibizumab + PRP	Aflibercept	0.012	-0.218	0.251
Ranibizumab	Bevacizumab + PRP	-0.001	-0.239	0.225
Ranibizumab + PRP	Bevacizumab + PRP	0.043	-0.151	0.234
Ranibizumab + PRP	Ranibizumab	0.044	-0.112	0.203


**TABLE 15** Results of NMA of log-MAR adjusting for time and baseline BCVA – ranking probabilities

Treatment arm	Probability of ranking				
	1st (%)	2nd (%)	3rd (%)	4th (%)	5th (%)
PRP	0.08	0.70	4.23	19.50	75.50
Aflibercept	19.25	23.45	21.43	22.70	13.18
Bevacizumab + PRP	38.40	23.30	18.10	15.68	4.53
Ranibizumab	36.93	30.48	19.63	10.75	2.23
Ranibizumab + PRP	5.35	22.08	36.63	31.38	4.58

## Network meta-analyses of reduced networks

*Assuming anti-vascular endothelial growth factor and anti-vascular endothelial growth factor + panretinal photocoagulation are equivalent*

This analysis assumes that anti-VEGF only arms and anti-VEGF + PRP arms have equal effect. To be used to assess differences between anti-VEGF types. A model allowing effect to vary with time and baseline log-MAR was used. Results are presented for the predicted effects after 1 year of follow-up and at mean baseline BCVA across trials.



**FIGURE 31** Results from a reduced network to compare anti-VEGFs.

**TABLE 16** Results of reduced network to compare anti-VEGFs – comparisons between treatments

Intervention	Comparator	Mean difference	95% CI	
Aflibercept	PRP	-0.091	-0.245	0.063
Bevacizumab	PRP	-0.126	-0.261	0.007
Ranibizumab	PRP	-0.094	-0.173	-0.023
Bevacizumab	Aflibercept	-0.035	-0.238	0.174
Ranibizumab	Aflibercept	-0.003	-0.166	0.163
Ranibizumab	Bevacizumab	0.032	-0.142	0.200


**TABLE 17** Results of reduced network to compare anti-VEGFs – ranking probabilities

Treatment	Probability of ranking			
	1st (%)	2nd (%)	3rd (%)	4th (%)
PRP	0.00	0.80	11.48	87.73
Aflibercept	25.08	32.93	33.35	8.65
Bevacizumab	53.60	23.33	20.18	2.90
Ranibizumab	21.33	42.95	35.00	0.73

**Assuming all types of anti-vascular endothelial growth factor are equivalent**

This analysis assumes that all three anti-VEGF drugs have

equal effect. To be used to assess the overall effect of anti-VEGF. A model allowing effect to vary with time and baseline log-MAR was used.



**FIGURE 32** Results from a reduced network to compare treatment classes.

**TABLE 18** Results of reduced network to compare treatment classes – comparisons between treatments

Intervention	Comparator	Mean difference	95% CI	
Anti-VEGF	PRP	-0.089	-0.180	-0.019
Anti-VEGF + PRP	PRP	-0.108	-0.192	-0.039
Anti-VEGF + PRP	Anti-VEGF	-0.019	-0.126	0.083

**TABLE 19** Results of reduced network to compare treatment classes – ranking probabilities

Treatment	Probability of ranking		
	1st (%)	2nd (%)	3rd (%)
PRP	0.03	1.20	98.78
Anti-VEGF	33.05	65.88	1.08
Anti-VEGF + PRP	66.93	32.93	0.15

## Threshold analyses

### Up to 1 year




FIGURE 33 Threshold analyses of data up to 1 year of follow-up.

### One to 2 years




FIGURE 34 Threshold analyses of data with 1–2 years of follow-up.

### Maximum follow-up (up to 2 years)




FIGURE 35 Threshold analyses of data at end of trial (up to 2 years).

### Allowing for effect variation with time and baseline logarithm of the minimum angle of resolution




FIGURE 36 Threshold analyses of model adjusting for effect of time and baseline log-MAR.

### Reduced network (for comparing anti-vascular endothelial growth factors)

Adjusted for follow-up time and BCVA at baseline.




FIGURE 37 Threshold analysis of simplified network to compare anti-VEGF types, with time and baseline BCVA adjustment.

### Reduced network (comparing anti-vascular endothelial growth factor to panretinal photocoagulation)

Adjusted for follow-up time and BCVA at baseline.




FIGURE 38 Threshold analyses of simplified network to compare anti-VEGF to PRP, adjusted for follow-up time and baseline BCVA.

## Appendix 3 Other outcomes

This appendix presents tables and figures for all analyses, using data from publications of included RCTs for outcomes other than BCVA. These mostly consist of forest plots without meta-analysis, because the evidence was generally too limited in extent, and too diverse in intervention and follow-up times, to justify a full meta-analysis.

### Neovascularisation of the disc




FIGURE 39 Forest plot of all NVD data (left side favours anti-VEGF).

As meta-analysis was not possible for most outcomes, the forest plots without meta-analysis include trials of proliferative and non-proliferative retinopathy, to aid comparison.

### Forest plots of outcomes without meta-analysis

These forest plots show results for all anti-VEGF types, and at all follow-up times. Note that this means some trials appear more than once in a forest plot.



### Neovascularisation elsewhere




FIGURE 40 Forest plot of all NVE data (left side favours anti-VEGF).

### Diabetic macular oedema




FIGURE 41 Forest plot of DME incidence (left side favours anti-VEGF).

### Improvement in diabetic retinopathy severity score (Diabetic Retinopathy Severity Scale)




FIGURE 42 Forest plot of improvement in DRSS severity (right side favours anti-VEGF).

### Proliferative retinopathy incidence




FIGURE 43 Forest plot of proliferative DR (left side favours anti-VEGF).

### Regression of neovascularisation




FIGURE 44 Forest plot of regressive neovascularisation (left side favours anti-VEGF).

### Use of other treatments




FIGURE 45 Forest plot of use of other treatments (left side favours anti-VEGF).

## Vitrectomy




FIGURE 46 Forest plot of vitrectomy incidence (left side favours anti-VEGF).

## Vitreous haemorrhage




FIGURE 47 Forest plot of vitreous haemorrhage incidence (left side favours anti-VEGF).

## Adverse event outcomes

These forest plots show results for all anti-VEGF types, and at all follow-up times. Note that this means some trials appear

more than once in a forest plot. For simplicity, only adverse event outcomes reported in two or more studies are presented.

## Cataracts




FIGURE 48 Forest plot of cataracts data (left side favours anti-VEGF).

### Conjunctival haemorrhage




FIGURE 49 Forest plot of conjunctival haemorrhage data (left side favours anti-VEGF).

### Cardiovascular mortality




FIGURE 50 Forest plot of cardiovascular mortality data (left side favours anti-VEGF).

### Death (all-cause mortality)




FIGURE 51 Forest plot of death data (left side favours anti-VEGF).

### Myocardial infarction




FIGURE 52 Forest plot of myocardial infarction data (left side favours anti-VEGF).


**Ocular pain**

FIGURE 53 Forest plot of ocular pain data (left side favours anti-VEGF).


**Raised intraocular pressure**

FIGURE 54 Forest plot of raised intraocular pressure data (left side favours anti-VEGF).


**Retinal detachment**

FIGURE 55 Forest plot of retinal detachment data (left side favours anti-VEGF).


**Retinal tear**

FIGURE 56 Forest plot of retinal data (left side favours anti-VEGF).


**Serious adverse event (however defined)**

FIGURE 57 Forest plot of serious adverse event data (left side favours anti-VEGF).


**Stroke**

FIGURE 58 Forest plot of stroke data (left side favours anti-VEGF).

**Meta-analyses of other outcomes and adverse events**

All meta-analyses presented assumed that the impact of anti-VEGF on outcome (or adverse event) is the same for all types of anti-VEGF (in isolation or combined with

PRP), and at all follow-up times. For trials with multiple time points, the longest follow-up was used. For trial with multiple arms, only one anti-VEGF arm was used; arms using anti-VEGF alone were preferred.


**Neovascularisation elsewhere**

FIGURE 59 Meta-analysis of NVE (left side favours anti-VEGF).

### Neovascularisation of the disc




FIGURE 60 Meta-analysis of NVD (left side favours anti-VEGF).

### Other non-vision outcomes

This forest plot shows the summary results of each meta-analysis (each bar is a meta-analysis result). Meta-analyses

are restricted to trials of proliferative retinopathy. Full forest plots for each outcome are not presented.




FIGURE 61 Meta-analysis summary for non-vision outcomes in PDR trials (left side favours anti-VEGF).

### Adverse events

This forest plot shows the summary results of each meta-analysis (each bar is a meta-analysis result). Meta-analyses

are restricted to trials of proliferative retinopathy. Full forest plots for each outcome are not presented.




FIGURE 62 Meta-analysis summary for adverse events (left side favours anti-VEGF).

## Appendix 4 Non-proliferative diabetic retinopathy

This section reports the findings of the two trials in non-proliferative retinopathy. As both trials compared aflibercept to sham injection, no NMAs were performed.

PANORAMA had two aflibercept arms: injections every 8 weeks or every 16 weeks. Only the 16-week arm is analysed here, as that was the schedule used in Protocol W.

### Best corrected visual acuity




FIGURE 63 Mean difference in ETDRS after 2 years in NPDR trials.




FIGURE 64 Mean difference in log-MAR after 2 years in NPDR trials.

### Diabetic macular oedema in non-proliferative retinopathy

Diabetic macular oedema was the only outcome other than BCVA reported in both trials of NPDR.




FIGURE 65 Diabetic macular oedema incidence in NPDR trials.



## Other outcomes




FIGURE 66 Non-BCVA outcomes in NPDR trials.

## Adverse events




FIGURE 67 Adverse event outcomes in NPDR trials.