



# City Research Online

## City St George's, University of London

**Citation:** Northcott, S., Devane, N., Behn, N., Nassor, J., Nazim, I., Omar, A., O'Regan, T., Thomas, S. & Hilari, K. (2024). Emotional And Language Recovery For People With Aphasia (Ela): Co-Designing A Novel Intervention. *International Journal of Stroke*, 19(3\_SUPPL), p. 79. doi: 10.1177/17474930241300284 ISSN 1747-4930 doi: 10.1177/17474930241300284

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/34366/>

**Link to published version:** <https://doi.org/10.1177/17474930241300284>

**Copyright and Reuse:** Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

## Emotional and Language recovery for people with Aphasia (ELA): co-designing a novel intervention

Authors: Sarah Northcott, Niamh Devane, Nicholas Behn, Jamila Nassor, Iqra Nazim, Ayan Omar, Thérèse O'Regan, Shirley Thomas, Katerina Hilari

### Introduction

Having aphasia increases the risk of becoming depressed, anxious and isolated. People with aphasia want to work on communication, confidence, and wellbeing. We aimed to co-design a novel intervention that would address these priorities through integrating two evidence-based therapy approaches: elaborated semantic feature analysis (ESFA); and solution focused brief therapy (SFBT), a brief psychological therapy.

### Methods

A series of workshops were held with five people with aphasia (6 workshops); three family members (3 workshops); and four speech and language therapists (SLTs) (3 workshops). In addition, meetings were held with experts in SFBT and elaborated SFA. This information was analysed using qualitative content analysis and mapped onto a therapy manual. Consultants with aphasia and student SLTs trialled therapy components for further refinement.

### Results

Themes from the workshops and expert meetings included: important features of therapy (e.g., sense of progress; therapeutic relationship; real-life impact); managing endings; involving family (e.g., noticing their hopes and needs); choosing meaningful words to target in therapy; supporting home practice; dosage and delivery; integrating impairment and wellbeing work; therapist training and support needs. Initial feedback has been positive: student SLTs reported the intervention helped them to 'see the person'; people with aphasia found it satisfying, enjoyable and described re-engaging in valued activities post therapy.

### Conclusions

ELA is a language therapy delivered within a holistic, humanistic framework, designed to be delivered within busy clinical services. It harnesses the behavioural change components of SFBT to facilitate clients taking linguistic gains from the therapy room into their everyday life.