



# City Research Online

## City St George's, University of London

**Citation:** Powell, C., Adisa, O., Herlitz, L., Bains, S., Eyrúnardóttir Clark, S., Deighton, J., Syed, S., Gilbert, R., Feder, G. & Howarth, E. (2025). Domestic abuse, primary care and child mental health services: A systems analysis of service coordination from professionals' perspectives. *Children and Youth Services Review*, 169, 108076. doi: 10.1016/j.childyouth.2024.108076


This is the published version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/34520/>

**Link to published version:** <https://doi.org/10.1016/j.childyouth.2024.108076>

**Copyright and Reuse:** Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).



## Domestic abuse, primary care and child mental health services: A systems analysis of service coordination from professionals' perspectives

Claire Powell<sup>a,\*</sup>, Olumide Adisa<sup>b</sup>, Lauren Herlitz<sup>c</sup>, Shivi Bains<sup>d</sup>, Sigrún Eyrúnardóttir Clark<sup>e</sup>, Jessica Deighton<sup>f</sup>, Shabeer Syed<sup>g</sup>, Ruth Gilbert<sup>h</sup>, Gene Feder<sup>i</sup>, Emma Howarth<sup>j</sup>

<sup>a</sup> Senior Research Fellow, Population, Policy, and Practice Research and Teaching Department, University College London Great Ormond Street Institute of Child Health, London, UK

<sup>b</sup> Senior research fellow, Institute of Social Justice and Crime, University of Suffolk, Ipswich, UK

<sup>c</sup> Research fellow, Population, Policy, and Practice Research and Teaching Department, University College London Great Ormond Street Institute of Child Health, London, UK

<sup>d</sup> Research student, Population, Policy, and Practice Research and Teaching Department, University College London Great Ormond Street Institute of Child Health, London, UK

<sup>e</sup> Research fellow, Rapid Research Evaluation and Appraisal Lab (RREAL), Department of Targeted Intervention, University College London, London, UK

<sup>f</sup> Professor, Evidence Based Practice Unit, Anna Freud National Centre for Children and Families and University College London, London, UK

<sup>g</sup> Senior research associate, Population, Policy, and Practice Research and Teaching Department, University College London Great Ormond Street Institute of Child Health, London, UK

<sup>h</sup> Professor, Population, Policy, and Practice Research and Teaching Department, University College London Great Ormond Street Institute of Child Health, London, UK

<sup>i</sup> Professor, Centre for Academic Primary, Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

<sup>j</sup> Reader, School of Psychology, University of Sussex, Brighton, UK

### ARTICLE INFO

#### Keywords:

Domestic abuse  
Health services  
Systems  
Children  
Mental health

### ABSTRACT

**Objective:** We explored how services work together to support parents and children experiencing both parental intimate partner violence (IPV) and parental or child mental health problems by drawing on the perspectives of professionals working in primary care, children and young people's mental health services (CYPMHS), and domestic abuse services.

**Methods:** We conducted a qualitative study, interviewing 38 professionals in three geographically contrasting local authority areas in England. We carried out framework analysis using a systems approach and mapping techniques to understand the service interrelationships and boundary judgements of professionals.

**Results:** The relationships between domestic abuse services, CYPMHS, and primary care were complex, involving funders and commissioners, local authority strategic groups, and wider services such as schools and children's centres. Participants consistently identified a gap in the relationship between statutory CYPMHS and domestic abuse services. Other service gaps were for children living with ongoing or intermittent IPV and for children and parents with needs falling below or between service thresholds. There was a gap in support services for users of abusive behaviour to prevent future IPV. An overview of staff perspectives revealed differing views on treating the effects of trauma, and the co-ordination and sequencing of care.

**Conclusion:** Improving the response to children and adults experiencing mental health problems in the wake of IPV requires a systems perspective to understand the barriers to service co-ordination. Our findings indicate a particular need to address the gap between CYPMHS and domestic abuse services. Current ways of working with adults could be adapted for children, in addition to learning from examples of best practice in the study sites.

### 1. Introduction

Globally over one in four women are reported to have experienced intimate partner violence (IPV) in their lifetime (Sardinha et al., 2022)

and in the UK it is estimated that one in five children have been exposed to parental IPV. (Radford et al., 2011) Intimate partner violence refers to physical, sexual or psychological harm, including threatening, aggressive and coercive behaviours, by an intimate partner or ex-partner.

\* Corresponding author.

E-mail address: [c.powell@ucl.ac.uk](mailto:c.powell@ucl.ac.uk) (C. Powell).

<https://doi.org/10.1016/j.chilyouth.2024.108076>

Received 29 May 2024; Received in revised form 29 November 2024; Accepted 6 December 2024

Available online 18 December 2024

0190-7409/© 2024 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

(Krug et al., 2002) In the UK, children living with parental IPV are now acknowledged as primary victims of abuse. (Act, 2021).

The impact of IPV on parental and child mental health is well documented, (Bacchus et al., 2018; Devries et al., 2018; McTavish et al., 2016) as is the associated increased use of healthcare services by survivors of IPV. (Rivara et al., 2007) Findings from the UK's ALSPAC longitudinal study show that parents and their children who had experienced IPV were more likely to experience depression than families that were not exposed. Children who experienced both parental IPV and maternal depression were themselves more likely to experience depression than their peers who were exposed to only one of these parental factors. (Gondek et al., 2023) Similarly, parents with IPV recorded in English electronic healthcare records in the first 1000 days of their child's life were more likely to have recorded mental and physical health problems than parents with no recorded IPV. (Syed et al., 2023).

In recognition of the links between family health outcomes and parental IPV, UK policy emphasises the importance of partnership working between health services and domestic abuse agencies, along with clear referral pathways for victims of violence and abuse and a co-ordinated local responses. (HM Government. Call to End Violence Against Women and Girls [Internet]., 2010) National Institute for Health and Care Excellence (NICE) guidance suggests a multi-agency approach to IPV (NICE. Domestic violence and abuse [Internet]., 2016) but in general, current policy and guidance is focused on the response to adult victims with more limited advice as to how this should be operationalised for working with children.

Although multiagency approaches for parental IPV are advocated, evidence from practice shows siloed provision for adult and CYP mental health and IPV. (Allen et al., 2022) Reported factors contributing to this include: challenges in multi-agency co-operation for children with child protection concerns (including parental IPV); (Barlow et al., 2023) healthcare infrastructure; (Lewis et al., 2018) and administrative issues affecting referral pathways, funding and multi-agency miscommunication. (Lewis et al., 2018; Howarth et al., 2016).

A gap in specialist support for children exposed to IPV is well recognised. (Domestic Abuse Commissioner, 2023) Research into outcomes for children living with IPV found that most interventions for children are delivered by the voluntary (or non-profit) sector. (Howarth et al., 2016) Even where IPV services, such as IRIS (a specialist domestic violence and abuse training, support and referral programme for General Practices), (Sohal et al., 2020) are linked to UK general practice (family medicine) there are no recommended approaches for how either physical or mental health services should identify children's needs or co-ordinate support when children also have mental health problems.

Within mental health services, there has been a move towards trauma-informed approaches driven by survivors (Sweeney et al., 2018; Sweeney and Taggart, 2018) that aim to deliver services in a way that do not retraumatise people who have experienced trauma, including IPV. However an analysis of UK policy identified varying interpretations of trauma-informed care and a lack of UK or NHS-wide strategy or funding (Emsley et al., 2022) whilst an overview of trauma-informed training for mental health practitioners found that most mental health trusts (regional organisations that provide mental health services) in England have not yet implemented training for their staff. (McNally et al., 2023).

Previous research has criticised health services responding to IPV for taking a narrow, single sector perspective, suggesting a systems approach would be useful. We define a systems approach in this context as an innovative lens that considers the complex interconnections of multiple sectors or service types at more than one operational level (e.g. frontline practice and commissioning) and also identifies key systems actors who coordinate these systems. (Adisa et al., 2024) There is some evidence that research is beginning to take this wider perspective. However, recent research identifying systems factors that increase screening and identification of IPV in healthcare settings did not include children. (O'Campo et al., 2011) There is also a lack of research on the

'enabling conditions' (García-Moreno et al., 2015) or systems barriers to service co-ordination specifically for families experiencing mental health problems and IPV.

We addressed the following research questions using a systems approach: 1) How do domestic abuse, primary care and child mental health services co-ordinate services for families experiencing mental health problems and IPV? 2) What are professional perspectives of how services should support families experiencing mental health problems and IPV? 3) How do professionals perceive service boundaries for families experiencing mental health problems and IPV?

## 2. Methods

We used a qualitative study design to understand how primary health care, child mental health, and domestic abuse services work together to identify and support families experiencing parental IPV and mental health problems (in either parent or child or both). We followed the Consolidated Criteria for Reporting Qualitative Research (COREQ) (See supplement 1) and we pre-registered our protocol. (Powell et al., 2022) Changes to the protocol are detailed in supplement 2 and see supplement 3 for our glossary and definitions.

We carried out 36 semi-structured interviews between November 2022 and February 2023 with 38 professionals in three areas of England. Participants were asked to describe a family's journey through their services, including how mental health problems and IPV were identified and recorded, support provided, and the coordination of inter-service relationships.

### 2.1. Public & Patient involvement

We consulted our lived experience advisory group at the study design and data analysis stages. The group contributed questions to the interview schedule and highlighted areas of importance in the analysis. In addition we formed three professional advisory groups, one for each recruitment area, which included at least one practising clinician/professional from each of our participant groups (i.e. domestic abuse services, primary care, child mental health services). These groups input into the interview schedule design, supported with participant recruitment, and fed back on the findings.

### 2.2. Theoretical framework: Systems approach

We used Meadows' definition of a system for this study: a system typically comprises of "three kinds of things: elements, interconnections, and a function or purpose" and "is more than the sum of its parts" (Meadows, 2008). Therefore, a system can be characterised or perceived as a set of relationships in which the elements when taken together form a whole. The whole itself may be part of wider systems with their own set of elements that are linked to other systems. As our focus was on service co-ordination, we were particularly interested in the interconnections between elements (i.e. relevant health and social care services).

We also drew on Ulrich's theory of boundary critique which suggests that it is important to access a 'diverse variety of stakeholder views in defining problems and complex boundaries'. (Ulrich et al., 1996) In our study, we were interested in service boundaries, i.e. who was perceived as eligible for and able to access services.

### 2.3. Ethics

We did not ask participants about personal experiences of IPV; however we were aware the subject matter may have been difficult or that participants may themselves have experienced IPV. Participants were able to feedback on the interview process and they were offered a debrief sheet with contact numbers for support services. We collected written informed consent and participants were free to withdraw at any

point. Our study was approved by University College London Research Ethics Committee (17893/003) and NHS Health Research Authority (315188).

#### 2.4. Sampling & recruitment

The study areas were based on pre-existing contacts from an earlier study conducted in England. (Powell et al., 2022) We aimed for contrasting areas based on geography but also willingness from a key contact to provide support. Our final areas were 1) a northern, primarily rural local authority; 2) a western, city council local authority; 3) an inner-city London local authority.

In consultation with the study advisory groups we designed a maximum variation sampling framework to recruit participants from a range of professional roles (including frontline, managerial, and strategic/commissioning) across primary care (n = 6), child mental health services (including voluntary sector and statutory sector covering universal to specialist services) (n = 13), and domestic abuse services (n = 19) in each area. We used the framework to target specific professionals, but we included other professionals if they expressed an interest and met our criteria. See Table 1 in below for participant demographics. (See also Table 2 in supplement 4 for sampling framework, recruitment targets and final numbers, and supplement 3 for definitions of professional groups.)

#### 2.5. Procedure

Advisory group members emailed the study information to relevant organisations or individuals and interested participants contacted the study team directly. We also used snowball sampling, asking interview participants to suggest suitable colleagues. Once a potential participant made contact, we shared the information sheet, written informed consent form and brief demographics questionnaire in REDCap, (Harris et al., 2019) and organised an online interview in Teams. 43 professionals contacted the research team and 7 (16 %) did not take part in an interview.

We offered individual 60-minute interviews (n = 31) and 90-minute

**Table 1**  
Interview participant demographics.

	Total (%)
<i>Gender</i>	
Female	37 (97 %)
Male	1 (3 %)
<i>Age</i>	
18 to 25 years	1 (3 %)
26 to 35 years	8 (21 %)
36 to 45 years	13 (34 %)
46 to 55 years	11 (29 %)
55 + years	5 (13 %)
<i>Ethnicity*</i>	
Black, Black British, Caribbean or African (Caribbean, African, Any other Black, Black British or Caribbean background)	1 (3 %)
Mixed or Multiple ethnic groups (White and Black Caribbean, White and Black African, White and Asian, Any other Mixed or multiple ethnic background)	4 (10 %)
White (English, Welsh, Scottish, Northern Irish or British, Irish, Gypsy or Irish Traveller, Roma, Any other White background)	33 (87 %)
<i>Years in profession</i>	
Under 3 years	9 (24 %)
3 to 5 years	8 (21 %)
6 to 10 years	6 (16 %)
11 to 15 years	7 (18 %)
Over 15 years	8 (21 %)

group interviews (n = 2) where there were two or more participants from the same organisation. For GPs, we offered a shorter 30-minute individual interview (n = 3) to encourage participation. We adapted the interview schedule for each type of role and all participants were sent the interview questions in advance.

Interviews were carried out by experienced female qualitative researchers (CP, LH, SC) who met weekly throughout the interview period to discuss any concerns and track recruitment progress. Interviews took place online and were audio-recorded in Teams and sent securely to a professional transcriber for transcription. Participants were offered a shopping voucher for their time and the opportunity to review summary findings.

#### 2.6. Analysis

While data were being collected, each interview was summarised in an area-specific rapid assessment procedure (RAP) sheet (Vindrola-Padros et al., 2020) (a structured template) to facilitate initial analysis. This enabled real-time consideration of possible themes, as well as rapid cross-case comparison, and reflection on new probes or follow-up questions. The RAP sheets were used to create area-specific summaries, which were sent to all participants and advisory groups for optional feedback after data collection was completed.

Following the feedback and team discussion we carried out framework analysis. (Ritchie et al., 2003) After the initial inductive indexing and coding of the data in NVivo 12 by one researcher (CP), (Lumivero. NVivo (Version 12) [Internet]., 2017) we applied our systems lens to organise the codes and structure the analysis of themes. We used the codes to create system maps in Kumu (relationship mapping software, 2023) to visualise our interpretations related to organisational relationships. Initial stages of analysis were discussed with the interview team, the wider research team, and the lived experience group to refine the themes and the analysis.

#### 2.7. Reflexivity

Whilst we did not use an explicitly reflexive analytic approach we did reflect on our positions during the data collection and analysis stages. The data collection was carried out by three female researchers (CP, LH, SC), two of whom have extensive experience in the voluntary sector prior to entering academic research. The team reflected on their non-practice backgrounds when designing the interview schedules (e.g. explaining to participants they were not experts) and during the analysis through discussion of themes. The wider research team, with extensive experience in domestic abuse research, included two practising clinicians, a GP and a clinical psychologist, who drew on their dual expertise as researchers and practitioners when commenting on the analysis. This was in addition to our survivor advisory group who also feedback on the findings and whose comments are integrated in the paper.

### 3. Results

We explored how primary care, domestic abuse, and child mental health services in England work together and we developed three broad themes: 1) the interrelationships of organisations and cross-sector groups involved in care for children and parents experiencing mental health problems (MHP) and IPV; 2) the boundaries of care between services; and 3) tensions in perspectives on how CYP/parents should be supported.

#### 4. Interrelationships of organisations and cross-sector groups involved in care for children and parents experiencing MHP and IPV

We identified three groups of organisational interrelationships that affect how primary care, domestic abuse and child mental health

services work together to support families: A – funding-commissioning relationships; B – interface relationships (i.e. between frontline service delivery and funding/commissioning); C – inter-service relationships. For each group we present a systems map based on the interview codes and thematic summaries.

4.1. Funding and commissioning relationships

Participants highlighted the key role that funding and commissioning play in service co-ordination for families. The three core aspects of funding and commissioning relationships were: national government budget allocation; wider commissioning relationships which support or hinder service delivery; individual commissioners and what they do. See Fig. 1 for a representation of these relationships.

4.1.1. National government budget allocation

The Domestic Abuse Act (2021) was cited as a positive piece of legislation which has made provision for specific funding of children’s domestic abuse support services; this was seen as potentially funding some otherwise unavailable mental health support for children and increasing domestic abuse services’ capacity to work with other services. However, service commissioners and leaders pointed out that the use of national government funding is often restricted and does not always allow sufficient time or flexibility to create the service co-ordination needed, for example, to implement joint commissioning and contracting. Furthermore, we repeatedly heard from professionals that the current decade was one of the worst periods of funding for public services they had experienced in their careers, and this was limiting and reducing available services that children and families could access and their capacity for working together:

“All of us, schools, CAMHS, social care, [are] all very underfunded, very fractured kind of services and we’re all trying to come together and ... it’s like a broken piece of pottery and you’re trying to put all the pieces back together and all the bits aren’t really fitting” (P10, CYPMHS worker)

Other participants discussed this fragmentation and explained that lack of funding meant that only less intensive wellbeing work, often in groups, could be funded, not the in-depth one-to-one mental health support that parents and children needed.

4.2. Wider commissioning relationships

Participants discussed a variety of commissioning relationships which might support or hinder service commissioning and co-ordination. Joint commissioning arrangements with bodies such as NHS England and Integrated Care Boards (ICBs) were seen as supportive and productive, crossing the health-social care interface, and enabling relationships between services. When local councils provided support for domestic abuse as a political priority, this enabled commissioners and service leaders to create innovative services. However participants explained that regional bodies, such as Violence Reduction Units (VRUs), might be separately commissioned to deliver IPV services funded by central government. When these bodies did not communicate with local commissioners or delivered their programmes in parallel, this negatively affected local service co-ordination resulting in service gaps or duplication:

“It’s not just about receiving the money; it’s about how we’re managing those resources. You know, implement something that’s a little bit of a bigger scale, that’s got economy of scale, that will provide the same services across bigger patches. But this is the problem, when we see things being delivered pan-[region], ... commissioners, locally, have got very little say around it and that’s the problem.” (P13, Domestic Abuse)

4.2.1. Individual commissioners and what they do

Local authority service commissioners’ prioritisation (or not) of domestic abuse, their knowledge and expertise, cross-sector networks, and capacity were considered by participants to directly affect inter-service co-ordination:

“Maybe commissioners who’ve got 70 different portfolios and DAs just a little drop in the ocean, would they have the time and capacity, would they prioritise?” (P13, Domestic Abuse)

According to our participants, commissioning effective services that worked well together depended on vision, commissioning relationships across sectors, and the integration of administrative procedures such as joint contracting and shared case management systems. Without these things in place, mental health problems in the context of domestic abuse could be overlooked because it did not seem to be anyone’s specific responsibility.




Fig. 1. Map of funding and commissioning relationships.

### 4.3. Interface relationships

There were three types of relationships at the interface of service delivery and funding-commissioning that participants described where primary care, child mental health and domestic abuse services might cross paths and could help or hinder cross-service working. These were: 1) local area strategic groups; 2) safeguarding partnerships; 3) domestic homicide reviews. These were all affected by funding-commissioning relationships and directly shaped service delivery relationships. See Fig. 2 for a map of these relationships.

#### 4.3.1. Local area strategic groups

Strategic groups (including both steering and operational groups) bring together different health and social care partners and can facilitate joined-up support pathways. These groups were described as part of the commissioning process to oversee service delivery and partnership working and were seen as key to co-ordinating support between services. However, member attendance and inclusion were key to their effectiveness, as reported by interviewees. An observation primarily from the domestic abuse sector was the non-attendance of mental health services (despite being invited):

“I’d say one of the biggest gaps in all these meetings I’ve been talking about is representation from [mental health trust]. ... And certainly CAMHS... I’ve never seen them strategically round the table, at all.” (P14, Domestic Abuse)

In addition to this we heard from some domestic abuse services and relevant (particularly small) charity sector service providers about not being invited:

“At the minute we don’t have a voice there [Domestic Abuse Board] ...if you think about the 400 plus people we worked with last year, their voices are not being represented. I’m really keen for us to have a seat, so that we can have those conversations with other services, we haven’t had strong links with health, so that’s an area that we really want to develop and make some links there.” (P15, Domestic Abuse)

The exclusion of domestic abuse services, including specialist ‘by and for’ services was also highlighted in discussions around multi-agency meetings. Participants in strategic roles across health and domestic abuse suggested this might add to the communication challenges between voluntary sector domestic abuse (and related) services and statutory CYPMHS.

#### 4.3.2. Safeguarding partnerships

Safeguarding partnerships were seen to help joined up working and as an opportunity to think about the whole family, as well as encouraging multi-disciplinary working. It was a key structure for bringing awareness of domestic abuse into health services and for multi-agency referrals. Where there was limited cross-working between domestic abuse, primary care and child mental health services, safeguarding was identified by interviewees across professional groups as the only arena where these services might work together because of their clear purpose:

“It’s within safeguarding we contribute to how those services work together... safeguarding should be a thread through all those services and how they work together.” (P16, Primary Care)

However some small charity sector organisations, especially by and for services, felt they were not always seen as equal partners in the process:

“The only thing I would say is we’re not always or experiences have been at times where we don’t get the notes, so we won’t get the sort of notes or minutes like others will, we’re just sort of called in and ... not necessarily, I suppose, seen as equals in relation to other.” (P24, Domestic Abuse)

#### 4.3.3. Domestic homicide reviews (DHRs)

DHRs were also seen as an opportunity to improve service coordination, often highlighting poor service communication or challenges identifying families at risk. Some participants from the domestic abuse sector highlighted that they had only been able to develop positive relationships with mental health services after a DHR had flagged issues in cross-service communication.

### 4.4. Inter-service relationships

Participants reported that referrals were the primary relationship between services. In this section we describe the nature of relationships between domestic abuse, child mental health and primary care services, as well as the impact of wider services. Fig. 3 shows the wide eco-system of services in which families experiencing both domestic abuse and mental health might access support.

#### 4.4.1. Connecting services between domestic abuse, CYPMHS and primary care services

According to our interviewees, children’s centres (family support centres for families with children under 5 years) and Family Hubs (a




Fig. 2. Map of interface relationships.




Fig. 3. Map of inter-service relationships.

development of children's centres that can offer services for older children) seemed particularly well-integrated with domestic abuse, CYPMHS and primary care services. Early help services (for families who don't meet the social care threshold but need additional support) appeared to have good connections to both domestic abuse services and CYPMHS. Small voluntary sector organisations reported that they needed visibility from statutory services (i.e. advertising and referrals) to be well-integrated in the local service offer.

The main gap that professionals described (and found challenging) was that between domestic abuse services and statutory CYPMHS, perhaps related to the challenges identified in funding, commissioning and interface relationships described above. One participant described it as a 'wall' (P17, domestic abuse), another as 'very hit and miss with a lot of misses' (P24, domestic abuse), and in general the relationship was described as mediated by social care or schools. Participants across domestic abuse services reported that in their areas they were unable to refer to mental health services directly without the support of another local authority service (e.g. a school) or a healthcare professional.

Primary care services seemed less well-integrated with wider local authority services. However, GPs described positive relationships with health visitors which enabled them to be alerted to families with children under 5 years exposed to parental IPV. There did not seem to be a similar relationship with community-based professionals for older children. Whilst GPs described making referrals to CYPMHS for young people exposed to IPV, there was rarely any other contact between the two services. One participant described the primary care role as:

"We are the holder of the information... our important thing is to make sure it's all visible and it's reviewed and it's there for other people to see." (P18, Primary Care)

Other participants felt there previously had been more opportunity for conversations with GPs around mental health support for young people and parents, but this was no longer possible given increased caseloads in primary care.

#### 4.4.2. Support from the wider network

Participants described the impact of poor service co-ordination in the wider network (i.e. beyond primary care, domestic abuse and CYPMHS) and how this affected support for families with a range of needs. A particular issue was the lack of communication between adult mental health services and substance misuse services and the practice of excluding parents from accessing mental health services until substance misuse was addressed. This could leave parents at greater risk of harm from their partners using abusive behaviours, less likely to be able to

address their issues, as well as having a negative impact on their children.

Participants explained how disagreements between social care services and Family Hubs about categorising parental relationships as conflict or domestic abuse might mean families were referred to inappropriate support, leaving families unable to access domestic abuse or CYPMHS services when needed. Different definitions of IPV and inter-parent conflict across services seemed to underpin this, although one participant speculated that categorising parental interactions as conflictual rather than abusive enabled cases to be passed from social care to other services, which might help reduce the high caseloads in social care.

## 5. Boundaries of care between services

Participants discussed how the threshold criteria for access into CYPMHS and adult mental health services prevented families experiencing IPV from accessing timely mental health support. Four groups of children and families referred for mental health problems were identified by practitioners as particularly affected by access criteria: children who live with current/ongoing domestic abuse; children below social care safeguarding thresholds; parents/children with mental health presentations that fall between the primary and secondary mental health care thresholds; and mothers with high needs and residency of their children. Practitioners discussed the impact of exclusion from services for families and the implications for service co-ordination.

### 5.1. Children living with domestic abuse (i.e. still living with the abusing parent) interviewees perceived that many mental health service criteria exclude children who are not considered to be living in a situation of safety

"One of the ... exclusion criteria from many of the young people's mental health service response, is needs stability, they need to be safe, there cannot be ongoing abuse, and of course that excludes every child that is experiencing domestic abuse, living in a refuge, mum's in a coercive controlling relationship. They've got high level trauma symptoms, they're of course demonstrating harmful behaviours in school environments, and that is classified as not safe or stable enough to work with generic mental health services." (P17, Domestic Abuse)

Participants from domestic abuse services and both statutory and voluntary sector child mental health services reiterated this, highlighting the number of children with mental health needs who could not access support for this reason.

### 5.2. Children below social care safeguarding thresholds

Children perceived as living in an abusive situation that is not risky enough to meet social care thresholds were also considered to be less likely to access CYPMHS even with significant mental health needs. Interviewees acknowledged this was because of reduced funding and service cuts which meant that CYPMHS thresholds are only really met by children also meeting social care thresholds. As one primary care practitioner explained:

“The threshold for CAMHS is so high that a child really needs to be in quite significant distress, and if they were in that much distress and there was that much, you know, there was that much of a need, I would say nine times out of ten they’re already involved in social care.” (P11 Primary Care)

### 5.3. Parents/children who fall between the primary and secondary mental health care thresholds

In England, primary mental health care provides access to ‘NHS Talking Therapies’ (previously known as ‘Improving Access to Psychological Therapy’ or ‘IAPT’), which predominantly comprises CBT interventions. Interviewees from child mental health and primary care services were concerned that adults and children exposed to domestic abuse are considered too complex for this level of therapy and that standard CBT as opposed to trauma-focused CBT is often inappropriate for trauma survivors. However these same families did not then meet the diagnostic criteria or thresholds for higher tier or specialist services; this left them unable to access any statutory mental health support as described by this practitioner:

“I would say 50 % of my cases, I go back to the GP and say, you know, the NHS talking therapy services isn’t enough for this client, but then they don’t meet the threshold for secondary mental health services, so we’re kind of stuck because I can’t do much work while she’s in crisis, but there are no resources to help her come out of crisis.” (P11, Primary Care)

### 5.4. Mothers with high needs

The final group refers to mothers with high level needs including mental health problems and substance use problems, who have residency of their children. Whilst there is some specialist mental health and domestic abuse provision for mothers who no longer have residency of their children, mothers with children are unable to access residential domestic abuse provision with high-level mental health support.

### 5.5. The impact of children and parents being excluded from accessing services

The impact on services and families of these threshold criteria and lack of support for some children and families was described by this practitioner:

“I think the waits are just so, so severe and the child has to be so disabled, by the time they get seen by CAMHS in, sadly, you just feel that sometimes you think a stitch in time would save nine and I fear that, that there often isn’t that possibility, there isn’t that capacity in the service to see these children before they do get completely unravelled.” (P19, Primary Care)

Several participants explained that children would deteriorate on lengthy waiting lists. Parents were also more likely to return to abusive partners if they were unable to access timely and appropriate support. Wider support services described how they might avoid even making referrals to mental health services because they did not want families to be rejected:

“What we don’t want is families being bounced around, being told ... you’re not relevant for this service... People don’t like to hear that and that has ramifications down their life, to be told that they’re not complex enough or too complex.” (P20, Primary Care)

Voluntary sector services and primary care services described how they tried to fill the gap by keeping families on for longer in their own services, recommending private therapy or trying to find alternative support through schools. The boundaries of care between services seemed both to reflect funding and commissioning arrangements (and their gaps) but also practitioner perspectives on who should be supported and how in different services.

## 6. Tensions in perspectives on how families should be supported

On analysing interviewee perspectives of how care should be provided, there were three points of divergence: 1) the problem with ‘safety first’; 2) the role of child and young people mental health services; 3) who is responsible for users of abusive behaviour.

### 6.1. The problem with ‘safety first’

There was general agreement from participants that mental health interventions for parents and children were limited when environmental factors (such as poor housing) are not addressed, and that trauma responses cannot be processed while traumatic experiences are ongoing. However, participants outside of mental health services were keen to point out that it is difficult to keep people safe if they are mentally unwell or distressed:

“A really big barrier for most women to make positive decisions about relationships, you know, having the strength to make decisions to leave relationships, ... being able to prioritise their children’s needs... really whittles down to their capacity and mental health. If a mum is in complete mental health distress, then it’s really difficult for me to put all these options out ... if they’re in crisis, none of those things are really easy to take on board and to action.” (P11, Primary Care)

Domestic abuse services felt they were equipped to offer emotional support, mental health safety planning or advocacy around service access, but not therapy or formal mental health interventions. Domestic abuse services staff – in particular, advocates with a focus on practical support – described how it was difficult not to be pushed into a therapist role when providing support. Staff felt that approaches which brought in consultation and supervision from mental health specialists in domestic abuse services would be useful to negotiate some of these challenges.

### 6.2. The role of child and young people mental health services

The next diverging perspective was around the role of CYPMHS for families experiencing domestic abuse. Many domestic abuse practitioners felt CYPMHS saw domestic abuse primarily as a safeguarding issue rather than as a risk for or contributor to mental health problems:

“I guess is it the case that mental health services don’t really understand domestic abuse and sexual violence, and maybe they are only seeing it in terms of safeguarding of their own clients, rather than how could they be working in prevention, how could they be working in crisis, and how could they be working in recovery?” (P12, Domestic Abuse)

This participant felt mental health services could be part of domestic abuse prevention and recovery for children because they are already in contact with so many children who have experienced parental IPV.

There was a perception that there is a mismatch between CYPMHS thresholds and diagnostic criteria, and children’s presentations when affected by trauma (in this case, domestic abuse), i.e. that children could

experience severe mental distress and need support, but this did not ‘fit’ service criteria. Child mental health staff from both the statutory and voluntary sectors agreed that short-term interventions were unsuitable for addressing the impact of domestic abuse, and that there were not the resources in the public sector for children impacted by trauma.

However, domestic abuse staff and voluntary sector children’s mental health staff felt that there were some more fundamental differences between themselves and statutory child mental health practitioners’ perspectives on mental health. The conflict seemed to centre on whether trauma-related mental distress is qualitatively different from ‘mental illness’ – with a perception that domestic abuse services and the voluntary sector had a trauma-informed approach to mental health, whilst mental health services did not.

Some participants suggested that where statutory child mental health services were aware of domestic abuse, their role was providing support for children with historic trauma, “we’re in the game of processing past trauma” (P21, CYPMHS), whilst it was seen as the voluntary sector’s role to provide support with current trauma. In addition, there was a general perception that statutory child mental health services are less aware of the complexities of domestic abuse, particularly around coercive control, jealousy, stalking and harassment or challenges facing people trying to leave abusive relationships.

We identified from the interviews that domestic abuse services were systematically asking as part of their intake assessments about the mental health of parents and children, whilst statutory child mental health services were not systematically asking about or identifying family IPV. This was reflected in levels of training staff had received too – with all domestic abuse staff reporting some mental health training, whilst few child mental health practitioners had received IPV training, and notably most of those who had were based in the voluntary sector. (See [supplement 5 for further details](#)).

Participants agreed that there is a lack of clarity about who is responsible for what support at what time for families experiencing domestic abuse – in particular, around trauma support, mental health interventions, counselling, and referral pathways in general. Participants agreed that they wanted to address the siloed ways of working between domestic abuse services and CYPMHS. CYPMHS practitioners discussed how difficult it was for them to identify domestic abuse if children did not talk about it and accompanying parents did not provide this information.

### 6.3. Who is responsible for users of abusive behaviour?

The final diverging perspective was around support for users of abusive behaviour (usually fathers) to change their behaviour. There was general agreement that there is a gap in support for parents who use abusive behaviour and young people who may begin to in their own relationships. Some participants felt their support of children and the non-harming parent was limited if there was no support available for the parent using abusive behaviour to end their abuse. There was a sense that parents who use violence are ignored by services, in particular social services and mental health services, and that outside of specialist perpetrator programmes there is a lack of responsibility for and skills to work with this group of parents.

Some specialists described how colleagues might be frightened of parents who use abusive behaviour or worry about upsetting them (and the impact on the family). However this fear resulted in avoidance of any contact with the parents who use abusive behaviour:

“I was worried about if I said something that made this man very angry that he would then actually seriously hurt his family or kill them.” (P23, CYPMHS)

Specialist domestic abuse workers wanted mental health services to be able to hold perpetration and mental health in mind simultaneously, both for the parent who uses violence and for young people who start to display abusive behaviour:

“We see adolescent boys who are using abuse against their girlfriends but who have experienced a lot of childhood trauma and domestic violence as children and the CAMHS professionals tend to hold the boys’ trauma in mind over and above the trauma he’s causing.” (P22, domestic abuse)

Child and adult mental health services were the focus because this was often the service that users of abusive behaviour might already be in contact with, according to our participants, and they felt it was better for the work to start with a professional who was already in contact with the parent or young person. Support for people who use violence to change was seen as key to prevention of future domestic abuse both in adults but also in young victims who might go on to have their own harmful relationships.

## 7. Discussion

This qualitative study explored professional perspectives of service co-ordination for families experiencing IPV and mental health problems. The systems mapping of interrelationships identified complex barriers and enablers to service co-ordination in the wider system through to the relationships between the frontline services themselves. A key gap was identified between domestic abuse services and statutory CYPMHS. Exploration of the boundaries of care suggested that children living in households with ongoing or intermittent IPV and children and parents with mental health or social care needs below or between service thresholds might be less able to access mental health support. An overview of staff perspectives revealed differing views on addressing the effects of trauma, the co-ordination and sequencing of care, and service responsibility for parents who use abusive behaviour.

The key gap we identified between statutory CYPMHS services and voluntary sector domestic abuse services in part explains why children can be invisible in the wider domestic abuse support system. (Roy et al., 2022) This service gap has been highlighted across the voluntary sector service literature and reiterates recent calls for better integration between specialist IPV and mental health services (Oram et al., 2022) with joint commissioning as the central component. (Domestic Abuse Commissioner, 2023) Our lived experience group felt that there was a particular gap for teenage boys and that any link between domestic abuse services and CYPMHS would need to proactively address this. This gap in service provision, particularly in relation to domestic abuse refugees has been highlighted in the grey and academic literature. (Baker, 2009; Radford et al., 2011).

This gap between CYPMHS and domestic abuse services appeared to be a specific manifestation of a deeper difference in perspectives between these services relating to how trauma is recognised and addressed in mental health services, (Oram et al., 2022; Coughlan et al., 2024) and how services can support children who are not considered to be in a position of safety. This is an ongoing debate around CYPMHS and their ability to provide therapeutic support for children with unstable home lives and has been identified in ongoing research about the mental health needs of children with a social worker. (Coughlan et al., 2024) An analysis of CYPMHS policy argues that a move towards individualised understandings of mental health excludes consideration of social conditions. (Callaghan et al., 2017) Earlier research with adult survivors of IPV had similar critiques of mental health services (Humphreys and Thiara, 2003) and was echoed by our lived experience group who felt that CYPMHS can have a narrow perspective that does not relate mental health to the context a child might be living in. This professional separation between mental health and IPV is also reflected in child safeguarding policy: a recent review examined the representation of domestic abuse and found that mental health is rarely mentioned. (Russell et al., 2022).

This disciplinary and service separation between parental IPV and child mental health is also reflected in our finding that children (and parents) often fell between lower levels of mental health support and

specialist mental health services. This has been widely reported as an issue across mental health services for children with more complex needs. For example, a recent evaluation of mental health support teams in schools (classed as a lower level of mental health support) found that children who were underserved by this service included children with special educational needs, those from ethnic minority backgrounds, and children with challenging family circumstances, including parental IPV. (Ellins et al., 2023) Therapists in the study, like our participants, felt that the time-limited, low-intensity CBT on offer was not suited to these children, who also did not meet thresholds for more specialist support. (Ellins et al., 2023) This suggests a wider systemic problem with service thresholds and the support offer affecting children living with social adversities and has been highlighted in ongoing research about child mental health. (Coughlan et al., 2024; Morgan et al., 2024).

These systemic problems align with Hester's three planet model (Hester, 2011) which explains the contradictions in professional practice and discourse between child protection, domestic abuse and child contact services. CYPMHS practitioners and others in child-focused health and social care services who work with children affected by domestic abuse also bring their practices and understanding of domestic abuse. This was highlighted in our study by professionals who reported disagreements about categorising IPV as parental conflict and the implications for service referrals. Inter-professional consensus on the difference between conflict and abuse is central to providing appropriate trauma-informed support at the right time for adult and child survivors. (Hodes and Mennicke, 2019).

### 7.1. Strengths and limitations

This exploratory study used a systems lens and highlighted areas that warrant further indepth research, such as factors affecting thresholds to health and social care services for families affected by domestic abuse and their variation by region, service demand and service structures. Whilst this study incorporated the views of a range of health and domestic abuse professionals across three geographically varied areas in England, primary care was under-represented in our sample. Further work is needed to understand primary care practitioner perspectives; this is important given their central role in referring parents and children. However, resonance between our findings and national surveys (Domestic Abuse Commissioner, 2023; Standing Together Against Domestic Abuse, 2020), suggests the findings are more widely transferable beyond the three study sites. Research involving a wider range of health services and local authority services would enable further understanding of these perspectives, and in-depth work is required on the recognition of and support for IPV in CYPMHS. In addition, further work could be carried out to focus on specific geographic and regional differences.

### 7.2. Implications for policy and practice

Our findings show that system-level intervention might be needed to improve service co-ordination, consistent with the wider health and domestic abuse service literature. Evaluation of Independent Domestic Violence Advisors (IDVA) in maternity services highlighted ongoing funding and local area health commissioning and management support were needed for service co-ordination. (Forbes et al., 2023) Similarly, national reviews of services have highlighted the impact of commissioning decisions, (Domestic Abuse Commissioner, 2023; Connect Centre, 2021) funding cycles (Standing Together Against Domestic Abuse, 2020; Ava, 2023) and funding restrictions, (Juliette et al., 2023) and missing health service input in strategic meetings (Standing Together Against Domestic Abuse, 2020) on cross-sector service co-ordination. A review of health practitioner readiness to address IPV emphasised the need for supportive systems, (Hegarty et al., 2020) whilst the recent Lancet Psychiatry Commission on IPV highlighted trauma-informed support is only possible with system-wide changes. (Oram et al., 2022) Thus, attention needs to be paid to funding,

commissioning and contracting processes, and the involvement of strategic groups, particularly smaller voluntary sector organisations and by and for services.

Our findings suggest there needs to be a strengthening of the relationship between domestic abuse services and CYPMHS including the development of advice, referrals and treatment pathways, in addition to mechanisms to facilitate cross-service advice and guidance. Currently, evidence about best practice is based on work with adults but suggests that there are pre-existing ways of working that could be expanded. Early work from the extension of the IRIS programme (domestic abuse advocates in general practice) has shown the success of extending the model to consider children (Szilassy et al., 2024) and this a possible model that could be developed in or with CYPMHS. A trial in the UK that trained domestic abuse advocates to support mental health had a positive impact on women's mental health (Ferrari et al., 2018) suggesting that similarly this model could be extended to enable children accessing domestic abuse services to access psychologically informed support. Each of our study areas had an example of positive practice for children, these are described in supplement 6, Table 3.

Clinicians from mental health services consulting to the voluntary sector, suggested by our participants, might support the identification of families who fall through the service gap, in addition to strengthening the relationship between the sectors. Similarly domestic abuse organisations, such as SafeLives (SafeLives [Internet]. [cited, 2024]) could be systematically funded to provide training for health and social care professionals and commissioners to raise awareness and skills and support wider understanding of domestic abuse and the difference with conflict. Local organisations could offer advice to mental health professionals working with families with experience of abuse. Furthermore, policy initiatives in the UK, such as Family Hubs and Early Help services, where health and social care services are already working together, provide pre-existing structures for strengthening the system in addition to opportunities for prevention. (Cleaver et al., 2019).

## 8. Conclusion

Improving service co-ordination for families experiencing IPV and mental health problems requires a systems approach to address the many barriers to accessing support parents and their children face. Improving the link between CYPMHS and domestic abuse services through ensuring relevant professionals from statutory and voluntary sector attend strategic groups, jointly contracting embedded workers across services and joint care pathways could go some way in addressing the main gaps in support.

## Funding

This study is funded by the National Institute for Health and Care Research (NIHR) Policy Research Programme (funder reference: PR-PRU-1217-21301; UCL award code: 177763). GF and OA's salaries were supported by the UK Prevention Research Partnership (Violence, Health and Society; MR-VO49879/1), an initiative funded by UK Research and Innovation Councils, the Department of Health and Social Care (United Kingdom) and the UK devolved administrations, and leading health research charities. The views expressed in this article are those of the authors and not necessarily those of the UK Prevention Research Partnership or any other funder.

## Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Acknowledgements

We wish to thank all the interview participants for giving their time to take part. We are grateful to Voices, our survivor advisory group and our three professional advisory groups for supporting this study and using their expertise to improve it.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.chilgyouth.2024.108076>.

## Data availability

The authors do not have permission to share data.

## References

- Adisa O, Bond E, editors. *Tackling Domestic Abuse and Sexual Violence: A Systems Approach* [Internet]. Cham: Springer International Publishing; 2024 [cited 2024 Nov 28]. (Palgrave Studies in Victims and Victimology). Available from: <https://link.springer.com/10.1007/978-3-031-58600-2>.
- Allen, K., Melendez-Torres, G. J., Ford, T., Bonell, C., Finning, K., Fredlund, M., et al. (2022). Family focused interventions that address parental domestic violence and abuse, mental ill-health, and substance misuse in combination: A systematic review. *PLoS One*, 17(7), Article e0270894.
- AVA. Funding for Support Services for Victims and Survivors of Violence Against Women and Girls in London [Internet]. London Councils; 2023 [cited 2024 Apr 10]. Available from: <https://avaproject.org.uk/wp-content/uploads/2023/05/1716-5-LC-London-Mapping-Report-WEB-03.pdf>.
- Bacchus, L. J., Ranganathan, M., Watts, C., & Devries, K. (2018). Recent intimate partner violence against women and health: A systematic review and meta-analysis of cohort studies. *BMJ Open*, 8(7), Article e019995.
- Baker, H. (2009). 'Potentially violent men?': Teenage boys, access to refuges and constructions of men, masculinity and violence. *J Soc Welf Fam Law.*, 31(4), 435–450.
- Barlow J, McMillan AS, Bowen E. Improving outcomes for children with child protection concerns who have been exposed to domestic abuse [Internet]. 2023 [cited 2023 Nov 9] p. 196. Available from: <https://foundations.org.uk/wp-content/uploads/2023/06/improving-outcomes-for-children-with-child-protection-concerns-who-have-been-exposed-to-domestic-abuse.pdf>.
- Callaghan, J. E., Fellin, L. C., & Warner-Gale, F. (2017). A critical analysis of child and adolescent mental health services policy in England. *Clin Child Psychol Psychiatry.*, 22(1), 109–127.
- Cleaver, K., Maras, P., Oram, C., & McCallum, K. (2019). A review of UK based multi-agency approaches to early intervention in domestic abuse: Lessons to be learnt from existing evaluation studies. *Aggress Violent Behav.*, 1(46), 140–155.
- Connect Centre. Roadmap Evaluation. Final Report. [Internet]. University of Central Lancashire; 2021 [cited 2024 Apr 10]. Available from: [https://clock.uclan.ac.uk/39447/2/Roadmap\\_Report\\_280921.pdf](https://clock.uclan.ac.uk/39447/2/Roadmap_Report_280921.pdf).
- Coughlan B, Russell A, Mannes J, Denford E, Hillier A, Davis L, et al. Children's social workers' perspectives on mental health and working with child mental health teams: Systematic review and narrative synthesis [Internet]. OSF; 2024 [cited 2024 Nov 29]. Available from: <https://osf.io/6mbr4>.
- Coughlan B, Mannes J, Marshall N, Morgan T, Ford T, Crozier-Roche F, et al. Mental Health Needs of Children and Young People in England, 2016-2021: An Epidemiological Study, and Comparison of Young People Who Have Received Protective and Care Interventions and Their Peers [Internet]. OSF; 2024 [cited 2024 Nov 29]. Available from: <https://osf.io/dx8e7>.
- Devries, K., Knight, L., Petzold, M., Merrill, K. G., Maxwell, L., Williams, A., et al. (2018). Who perpetrates violence against children? A systematic analysis of age-specific and sex-specific data. *BMJ Paediatr Open.*, 2(1), Article e000180.
- Domestic Abuse Act [Internet]. King's Printer of Acts of Parliament; 2021. Available from: <https://www.legislation.gov.uk/ukpga/2021/17/contents/enacted>.
- Domestic Abuse Commissioner. A Patchwork of Provision. How to meet the needs of victims and survivors across England and Wales. [Internet]. 2023 [cited 2023 Nov 10] p. 140. Available from: [https://domesticabusecommissioner.uk/wp-content/uploads/2022/11/A-Patchwork-of-Provision\\_Technical-Report\\_Nov-2022\\_Final.pdf](https://domesticabusecommissioner.uk/wp-content/uploads/2022/11/A-Patchwork-of-Provision_Technical-Report_Nov-2022_Final.pdf).
- Ellins, J., Hocking, L., Al-Haboubi, M., Newbould, J., Fenton, S. J., Daniel, K., et al. (2023). Early evaluation of the children and young people's mental health trailblazer programme: a rapid mixed-methods study. *Health Soc Care Deliv Res.*, 1–137.
- Emsley, E., Smith, J., Martin, D., & Lewis, N. V. (2022). Trauma-informed care in the UK: Where are we? A qualitative study of health policies and professional perspectives. *BMC Health Serv Res.*, 22(1), 1164.
- Ferrari, G., Feder, G., Agnew-Davies, R., Bailey, J. E., Hollinghurst, S., Howard, L., et al. (2018). Psychological advocacy towards healing (PATH): A randomized controlled trial of a psychological intervention in a domestic violence service setting. *PLoS One*, 13(11), Article e0205485.
- Forbes, C., Alderson, H., Domoney, J., Papamichail, A., Berry, V., McGovern, R., et al. (2023). A survey and stakeholder consultation of Independent Domestic Violence Advisor (IDVA) programmes in English maternity services. *BMC Pregnancy and Childbirth.*, 23(1), 404.
- García-Moreno, C., Hegarty, K., d'Oliveira, A. F. L., Koziol-McLain, J., Colombini, M., & Feder, G. (2015). The health-systems response to violence against women. *The Lancet Lond Engl.*, 385(9977), 1567–1579.
- Gondek, D., Howe, L. D., Gilbert, R., Feder, G., Howarth, E., Deighton, J., et al. (2023). Association of interparental violence and maternal depression with depression among adolescents at the population and individual level. *Journal of the American Medical Association Netw Open.*, 6(3), Article e231175.
- Harris, P. A., Taylor, R., Minor, B. L., Elliott, V., Fernandez, M., O'Neal, L., et al. (2019). The REDCap consortium: Building an international community of software platform partners. *Journal of Biomedical Informatics.*, 1(95), Article 103208.
- Hegarty, K., McKibbin, G., Hameed, M., Koziol-McLain, J., Feder, G., Tarzia, L., et al. (2020). Health practitioners' readiness to address domestic violence and abuse: A qualitative meta-synthesis. *PLoS One*, 15(6), Article e0234067.
- Hester, M. (2011). The three planet model: towards an understanding of contradictions in approaches to women and children's safety in contexts of domestic violence. *The British Journal of Social Work.*, 41(5), 837–853.
- HM Government. Call to End Violence Against Women and Girls [Internet]. 2010 [cited 2023 Nov 9]. Available from: <https://assets.publishing.service.gov.uk/media/5a7b30d1ed915d429748d70a/vawg-paper.pdf>.
- Hodes, C., & Mennicke, A. (2019). Is it conflict or abuse? a practice note for furthering differential assessment and response. *Clinical Social Work Journal.*, 47(2), 176–184.
- Howarth, E., Moore, T. H., Welton, N. J., Lewis, N., Stanley, N., MacMillan, H., et al. (2016). IMPROVING Outcomes for children exposed to domestic Violence (IMPROVE): An evidence synthesis. *Public Health Res.*, 4(10), 1–342.
- Humphreys, C., & Thiara, R. (2003). Mental health and domestic violence: 'I Call it symptoms of abuse'. *The British Journal of Social Work.*, 33(2), 209–226.
- Juliette, C., Rodriguez, D., Walker, E., Arnott, B., Moore, I., Clarkson, M., et al. (2023). Domestic abuse programmes for children & families. Programme promise and evaluation feasibility. [Internet]. *Foundations.*, 58. Available from: [cited 2023 Nov 14] <https://foundations.org.uk/wp-content/uploads/2023/06/domestic-abuse-programmes-for-children-and-families-programme-promise-and-evaluation-feasibility.pdf>.
- Krug, E. G., Mercy, J. A., Dahlberg, L. L., & Zwi, A. B. (2002). The world report on violence and health. *The Lancet.*, 360(9339), 1083–1088.
- Lewis, N. V., Feder, G. S., Howarth, E., Szilassy, E., McTavish, J. R., MacMillan, H. L., et al. (2018). Identification and initial response to children's exposure to intimate partner violence: A qualitative synthesis of the perspectives of children, mothers and professionals. *BMJ Open.*, 8(4), Article e019761.
- Lumivero. NVivo (Version 12) [Internet]. 2017. Available from: [www.lumivero.com](http://www.lumivero.com).
- McNally, K., Ragan, K., Varese, F., & Lovell, K. (2023). The ongoing importance of the routine enquiry into trauma and abuse and trauma-informed care within mental health trusts in England. *J Psychiatr Ment Health Nurs.*, 30(6), 1130–1136.
- McTavish, J. R., MacGregor, J. C. D., Wathen, C. N., & MacMillan, H. L. (2016). Children's exposure to intimate partner violence: An overview. *International Review of Psychiatry.*, 28(5), 504–518.
- Meadows, D. H. (2008). Thinking in systems: a primer. *Chelsea Green Publishing.*, 242 p.
- Morgan, T., Crozier-Roche, F., Drayak, T., Smith, J., Graham, D., Marshall, N., et al. (2024). Addressing the mental health needs of children with a social worker. *Paediatrics & Child Health.*, 34(11), 406–411.
- NICE. Domestic violence and abuse [Internet]. 2016 p. 37. (Quality standard). Available from: <https://www.nice.org.uk/guidance/qs116/resources/domestic-violence-and-abuse-pdf-75545301469381>.
- Oram, S., Fisher, H. L., Minnis, H., Seedat, S., Walby, S., Hegarty, K., et al. (2022). The Lancet Psychiatry Commission on intimate partner violence and mental health: Advancing mental health services, research, and policy. *The Lancet Psychiatry.*, 9(6), 487–524.
- O'Campo, P., Kirst, M., Tsamis, C., Chambers, C., & Ahmad, F. (2011). Implementing successful intimate partner violence screening programs in health care settings: Evidence generated from a realist-informed systematic review. *Social science and medicine.*, 72(6), 855–866.
- Powell C, Deighton J, Gilbert R, Howarth E, Feder G. Intimate partner violence and mental health of children and parents or carers: qualitative study protocol [Internet]. OSF; 2022 [cited 2024 Dec 12]. Available from: <https://osf.io/4ynrh>.
- Powell, C., Feder, G., Gilbert, R., Paulauskaite, L., Szilassy, E., Woodman, J., et al. (2022). Child and family-focused interventions for child maltreatment and domestic abuse: Development of core outcome sets. *BMJ Open.*, 12(9), Article e064397.
- Radford, L., Corral, S., Bradley, C., Fisher, H., Bassett, C., Howat, N., et al. (2011). Child abuse and neglect in the UK today. *NSPCC.*, 205.
- Radford, L., Aitken, R., Miller, P., Ellis, J., Roberts, J., & Firkic, A. (2011). Meeting the needs of children living with domestic violence in London: Research report. *Refuge/NSPCC.*
- [relationship mapping software]. Kumu [Internet]. 2023 [cited 2023 Nov 10]. Available from: <https://kumu.io/>.
- Ritchie, J., Spencer, L., & O'Connor, W. (2003). Carrying out qualitative analysis. In J. Ritchie, & J. Lewis (Eds.), *Qualitative Research Practice*. London: Sage.
- Rivara, F. P., Anderson, M. L., Fishman, P., Bonomi, A. E., Reid, R. J., Carrell, D., et al. (2007). Healthcare utilization and costs for women with a history of intimate partner violence. *Am J Prev Med.*, 32(2), 89–96.
- Roy, J., Williamson, E., Pitt, K., Stanley, N., Man, M. S., Feder, G., et al. (2022). 'It felt like there was always someone there for us': Supporting children affected by domestic violence and abuse who are identified by general practice. *Health & Social Care in the Community.*, 30(1), 165–174.

- Russell, A., Clements, K., Duschinsky, R., Howarth, E., Mayes, T., Reisel, A., et al. (2022). Domestic violence and abuse in local child safeguarding policy: How is the problem represented? *Health Soc Care Community*, 30(6), e3871–e3884.
- SafeLives [Internet]. [cited 2024 Nov 14]. We are SafeLives - Ending domestic abuse. Available from: <http://honeycomb.demo.fatbeehive.com>.
- Sardinha, L., Maheu-Giroux, M., Stöckl, H., Meyer, S. R., & García-Moreno, C. (2022). Global, regional, and national prevalence estimates of physical or sexual, or both, intimate partner violence against women in 2018. *The Lancet*, 399(10327), 803–813.
- Sohal, A. H., Feder, G., Boomla, K., Dowrick, A., Hooper, R., Howell, A., et al. (2020). Improving the healthcare response to domestic violence and abuse in UK primary care: Interrupted time series evaluation of a system-level training and support programme. *BMC Medicine*, 18(1), 48.
- Standing Together Against Domestic Abuse. In Search of Excellence. A refreshed guide to effective domestic abuse partnership work - The Coordinated Community Response (CCR) [Internet]. Standing Together; 2020 [cited 2023 Nov 14]. Available from: <https://static1.squarespace.com/static/5ee0be2588f1e349401c832c/t/5fd78eaf72a0a65a94da967e/1607962290051/In+Search+of+Excellence+2020.pdf>.
- Sweeney, A., & Taggart, D. (2018). (Mis)understanding trauma-informed approaches in mental health. *J Ment Health*, 27(5), 383–387.
- Sweeney, A., Filson, B., Kennedy, A., Collinson, L., & Gillard, S. (2018). A paradigm shift: Relationships in trauma-informed mental health services. *BJPsych Adv*, 24(5), 319–333.
- Syed, S., Gilbert, R., Feder, G., Howe, L. D., Powell, C., Howarth, E., et al. (2023). Family adversity and health characteristics associated with intimate partner violence in children and parents presenting to health care: A population-based birth cohort study in England. *The Lancet Public Health*, 8(7), e520–e534.
- Szilassy, E., Coope, C., Emsley, E., Williamson, E., Barbosa, E. C., Johnson, M., et al. (2024). Feasibility of a reconfigured domestic violence and abuse training and support intervention responding to affected women, men, children and young people through primary care. *BMC Prim Care*, 25(1), 38.
- Ulrich W. A primer to critical systems heuristics for action researchers [Internet]. Hull, UK: University of Hull; 1996 [cited 2023 Nov 10]. Available from: [https://wulrich.com/downloads/ulrich\\_1996a.pdf](https://wulrich.com/downloads/ulrich_1996a.pdf).
- Vindrola-Padros, C., Chisnall, G., Cooper, S., Dowrick, A., Djellouli, N., Symmons, S. M., et al. (2020). Carrying out rapid qualitative research during a pandemic: Emerging lessons from COVID-19. *Qualitative Health Research*, 30(14), 2192–2204.