



City Research Online

City St George's, University of London

Citation: Tweedy, J., Dvoriashyna, M., Crawshaw, J., Overby, D., Repetto, R., Roberts, P., Spelman, T., Stewart, P. & Foss, A. (2025). A model of the mechanisms underpinning unconventional aqueous humor outflow. *Investigative Ophthalmology and Visual Science*, 66(4), 75. doi: 10.1167/iovs.66.4.75

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/34949/>

Link to published version: <https://doi.org/10.1167/iovs.66.4.75>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

A model of the mechanisms underpinning unconventional aqueous humor outflow

Jennifer H. Tweedy,¹ Mariia Dvoriashyna,² Jessica R. Crawshaw,^{3,4} Darryl R. Overby,⁵ Rodolfo Repetto,⁶ Paul A. Roberts,⁷ Tamsin A. Spelman,⁸ Peter S. Stewart,⁹ Alexander J. E. Foss¹⁰

¹ Department of Mathematical Sciences, University of Bath, UK,

email: jht55@bath.ac.uk, ORCID: 0000-0002-0084-2465;

² School of Mathematics and Maxwell Institute for Mathematical Sciences, University of Edinburgh, Edinburgh, UK,

email: m.dvoriashyna@ed.ac.uk, ORCID: 0000-0002-6057-1919;

³ Wolfson Centre for Mathematical Biology, Mathematical Institute, University of Oxford, UK,

email: Jessica.Crawshaw@maths.ox.ac.uk, ORCID: 0000-0003-0797-1144;

⁴ School of Mathematical Sciences, Queensland University of Technology, Brisbane, Australia;

⁵ Department of Bioengineering, Imperial College London, UK,

email: d.overby@imperial.ac.uk, ORCID: 0000-0001-9894-7515;

⁶ Department of Civil, Chemical and Environmental Engineering, University of Genoa, Italy,

email: rodolfo.repetto@unige.it, ORCID: 0000-0001-9332-5063;

⁷ Department of Optometry and Visual Sciences, City St George's, University of London, UK,

email: p.a.roberts@univ.oxon.org, ORCID: 0000-0001-5293-6431;

⁸ Sainsbury Laboratory, University of Cambridge, 47 Bateman Street, Cambridge, CB2 1LR

email: Tamsin.Spelman@slcu.cam.ac.uk, ORCID: 0000-0002-4991-7931;

⁹ School of Mathematics & Statistics, University Place, University of Glasgow, UK,

email: Peter.Stewart@glasgow.ac.uk, ORCID: 0000-0002-0971-8057;

¹⁰ Department of Ophthalmology, Nottingham University Hospitals NHS Trust, UK,
email: Alexander.Foss@nottingham.ac.uk, ORCID: 0000-0001-9649-0072.

Correspondence: Jennifer H. Tweedy Department of Mathematical Sciences, University of Bath,
UK; jht55@bath.ac.uk

Received: March 11, 2024, with revisions on October 1, 2024 and January 24, 2025

Word count: 10,875 (7,487 without appendix)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Abstract

Purpose To develop a mathematical model of the unconventional outflow pathway.

Methods The unconventional pathway is modeled as having two key components: the uveo-vortex and the trans-scleral pathways. The uveo-vortex pathway is modeled using Starling's law and the trans-scleral flow using predominately hydrostatic forces. We include transcytosis from the choriocapillaris (CC) and collapsibility of the suprachoroidal space (SCS) as particular features. There is considerable uncertainty in a number of model parameter values, and we identify the most significant ones using sensitivity analysis.

Results The model successfully generates a fluid flow from anterior to posterior in the choroidal tissue and the SCS, which also demonstrates many of the known physiological features, including the insensitivity of the unconventional flow to fluctuations in the intraocular pressure (IOP), albumin removal by the trans-scleral flow and the CC as a net absorber of fluid from, and supplier of albumin to, the choroidal tissue. The model supports the two previously proposed mechanisms of the action of prostaglandin $F_{2\alpha}$ analogues.

Conclusions We have developed a theoretical model of the unconventional aqueous outflow pathway that successfully captures its physiological features and elucidates the actions of prostaglandin $F_{2\alpha}$ analogues and other drugs.

INTRODUCTION

It has long been well-known that most of the aqueous humor drains from the eye via the trabecular meshwork and the canal of Schlemm, the so-called conventional pathway. The presence of a second outflow pathway, termed the unconventional flow (also known as the uveo-scleral flow), was first reported in 1903 by Leber; he observed that tracers introduced into the anterior chamber of the eye could traverse into the suprachoroidal space (SCS) between the choroid and sclera.¹ However, it was not until the late 1960's that a clear description of the unconventional outflow pathway emerged, following a series of pioneering experiments by Bill *et al.*^{2,3,4,5,6,7} In particular, it was noted that tracer particles travel from the anterior chamber to the SCS at a rate 200-fold greater than the reverse movement in the cynomolgus monkey,⁸ and the rate is independent of particle size,⁴ indicating a bulk flow of fluid.

A schematic diagram of aqueous flow is shown in Figure 1. The ciliary epithelium, which overlies the ciliary processes, produces aqueous, from where it flows through the posterior chamber, anteriorly through the pupil (cyan arrows) into the anterior chamber and exits the eye via two distinct pathways: the conventional (green arrow) and unconventional routes (black arrows). The conventional flow traverses the trabecular meshwork, enters Schlemm's canal, and exits via the episcleral veins. The unconventional flow, which accounts for a sizable fraction (5 to 40%) of the total outflow,⁹ continues posteriorly through the choroidal tissue and leaves the eye via either the trans-scleral or uveo-vortex routes. As a point of terminology, we avoid the term uveo-scleral flow, as this has variably been used to describe either the unconventional pathway as a whole or just the trans-scleral pathway.

The trans-scleral flow is important as it removes choroidal interstitial albumin. In other tissues,

1
2 typically, the lymphatic systems clears interstitial albumin, but this is absent in the choroid (though
3
4 the presence of lymphatics in the ciliary body has been reported,¹⁰ but their existence remains con-
5
6 troversial¹¹). The concentration of albumin in the aqueous within the anterior chamber is estimated
7
8 at around only 1% of the plasma level,^{12,13} but, on its passage through the tissue of the iris root,
9
10 the interstitial fluid (IF) accumulates albumin from the fenestrated capillaries of the ciliary body
11
12 stroma,^{14,15} meaning that when it reaches the ciliary body, the level of albumin is much higher,
13
14 with levels as high as 74% of that in plasma being reported.^{16,17} In other experiments, researchers
15
16 have measured the albumin concentration in the uveal IF as around 10% of the plasma level in
17
18 monkeys, 30% in rabbits and 19-35% in humans.^{18,19} Albumin from the plasma in the vessels of
19
20 the choriocapillaris (CC) also accumulates in the IF. This is predominantly via transcytosis²⁰ (see
21
22 also the large pore system²¹), although transport through fenestrae and active transport play a mi-
23
24 nor role.²¹ The trans-scleral flow is dominated by direct (pressure-driven) flow across the sclera,
25
26 although there is also a small amount of drainage through the perivascular spaces (by mechanisms
27
28 still not well understood).^{22,23,24} Indeed, the hydraulic conductivity of sclera is quite sufficient to
29
30 account for a trans-scleral flow without invoking special routes.²⁵

31
32 Starling forces arise due to a combination of local hydrostatic and osmotic pressure differences
33
34 between the capillaries and the surrounding IF. In most capillaries the balance of these forces favors
35
36 net exudation from the vessels,²⁶ due to the low interstitial hydrostatic pressure. In the CC, how-
37
38 ever, the balance favors net absorption due to the high intraocular pressure (IOP). This uveo-vortex
39
40 flow into the CC accounts for a large fraction of the total unconventional flow.^{27,28}

41
42 Interest in the unconventional outflow increased following the discovery that prostaglandin $F_{2\alpha}$
43
44 ($PGF_{2\alpha}$) analogues, such as latanoprost, act by enhancing this flow, thus reducing the IOP. These are
45
46 the most frequently prescribed class of drugs used to treat glaucoma, and there are two hypothesized
47

1
2 mechanisms of their action. The first postulates a reduced pathway resistance through the ciliary
3
4 muscle due to either or both of relaxing it²⁹ and opening intramuscular spaces.³⁰ The opening of
5
6 the spaces by $\text{PGF}_{2\alpha}$ analogues is thought to be due to release of matrix metalloproteases (MMPs)
7
8 from the ciliary muscle, resulting in a degree of autodigestion of the ciliary body extracellular
9
10 matrix.³¹ This results in the appearance of extracellular spaces in the ciliary muscular tissue, which
11
12 have been observed in histological samples taken within a few days of commencing treatment.³⁰
13
14 The second mechanism also involves increased MMP expression, this time acting on the sclera to
15
16 increase its hydraulic conductivity, which also increases the flow.^{32,33}

17
18 However, we note that there are other potential mechanisms that could play a contributory
19
20 role. For example, changes are reported in the vasculature when using $\text{PGF}_{2\alpha}$ analogues. These
21
22 include acting as a vasoconstrictor or a vasodilator, but, for the eye, one of the commonest side
23
24 effects is hyperemia, which is a vasodilatory response.³⁴ In addition, there is increased hydraulic
25
26 conductance of the capillary walls, and increases of up to three-fold have been reported;^{35,36} we also
27
28 note that MMPs are involved in regulation of the endothelial hydraulic conductance.³⁷ Furthermore,
29
30 latanoprost causes choroidal thickening, but the reason for this is not clear.³⁸

31
32 Other drugs also have an effect on the unconventional pathway: atropine,^{39,40} tropicamide⁴¹
33
34 (both antagonists to the acetylcholine receptor) and epinephrine⁴² all enhance the flow, while pilo-
35
36 carpine^{43,40} (an agonist to the acetylcholine receptor) reduces the flow. It has been proposed that
37
38 atropine and pilocarpine act by affecting the resistance to flow through the ciliary muscle. However,
39
40 these drugs can also act on the vasculature: pilocarpine increases blood vessel hydraulic conduc-
41
42 tance,⁴⁴ tropicamide reduces vessel albumin conductance and epinephrine decreases blood vessel
43
44 hydraulic conductance. In this paper, we explore how changes in the vasculature affect unconven-
45
46 tional flow.
47

1
2 A large increase in unconventional flow is required to reduce the IOP significantly, for the
3
4 following reasons. To reduce the IOP by $r\%$, we need to reduce the hydrostatic pressure drop
5
6 across the trabecular meshwork and Schlemm's canal, $IOP - p_{\varepsilon}$, (where p_{ε} is episcleral venous
7
8 pressure) by a larger proportion $r_d\%$, given by $r_d = IOP \cdot r / (IOP - p_{\varepsilon})$. The conventional
9
10 outflow is proportional to this pressure drop and thus also reduces by $r_d\%$.⁴⁵ The remainder of the
11
12 aqueous flow must be taken up by the unconventional pathway, meaning that its flow must grow
13
14 by $Q_c r_d / Q_u\%$, where Q_c and Q_u are the flows (prior to IOP reduction) in the conventional and
15
16 unconventional pathways, respectively. Thus to achieve a moderate (say $r = 20\%$) reduction in the
17
18 IOP requires a percentage increase in the unconventional flow that is many times greater, both as
19
20 Q_c / Q_u is many times greater than unity and also as $r_d > r$. A further aim of this work is therefore
21
22 to explore in general the mechanisms by which unconventional flow could be enhanced.

23
24 The unconventional drainage pathway has the counter-intuitive property of being relatively un-
25
26 affected by the IOP.^{3,5,46} This is in contrast to the conventional flow, which increases in proportion
27
28 to the difference between the IOP and the episcleral venous pressure. Bill,⁵ working on living
29
30 cynomolgus monkeys, found that at a physiological IOP (11 mmHg) the conventional and uncon-
31
32 ventional flow rates were $0.80 \pm 0.11 \mu\text{l}/\text{min}$ and $0.44 \pm 0.06 \mu\text{l}/\text{min}$, respectively, but when IOP was
33
34 artificially raised to 22 mmHg, these rates increased to $4.18 \pm 0.12 \mu\text{l}/\text{min}$ and $0.63 \pm 0.08 \mu\text{l}/\text{min}$,
35
36 respectively. This represents a fivefold increase in the conventional flow and less than 50% increase
37
38 in the unconventional flow (see also Figure 5A below).

39
40 Various explanations have been proposed to explain the insensitivity of the flow to the IOP,⁴⁶
41
42 which involve both the trans-scleral and uveo-vortex flows. One possibility is collapsibility of the
43
44 fluid domain: the collapse of the SCS and/or ciliary muscle will increase the hydraulic resistance,
45
46 thereby impacting the change in flow rate,⁴⁶ which would affect the trans-scleral component of the
47

1
2 unconventional flow. Another plausible mechanism is that the pressure in the CC mirrors the IOP,⁴⁷
3
4 meaning that, as the IOP varies, there is little change in the difference in hydrostatic plus oncotic
5
6 pressure between the IF and CC, and, as this drives the uveo-vortex flow, this flow also does not
7
8 increase.

9
10 There are numerous theoretical models in the literature that have been successfully developed
11
12 to describe aqueous production,^{48,49} flow^{50,51,52,53} and drainage,⁵⁴ and fluid flows more generally in
13
14 the eye.^{55,56} However, the mechanics of the unconventional outflow pathway represents an under-
15
16 studied aspect of ocular physiology, and there remains a notable absence of mathematical models
17
18 in this area. In this paper, we develop a novel mathematical model of this pathway, which allows
19
20 us to test rigorously the mechanisms described above and theoretically assess the effectiveness of
21
22 potential pharmacological therapies.

23 24 25 26 **MATHEMATICAL MODEL**

27
28
29 In this Section, we give an overview of the mathematical model for a general audience. Full details
30
31 of the mathematical model and its derivation can be found in the Supplementary Material, while the
32
33 section ‘Details of mathematical model’ in the Appendix contains a shorter summary of the same
34
35 information, consisting of statements of the key modeling assumptions and the equations that are
36
37 used in the model development, as well as the full governing equations that are used to obtain the
38
39 results presented in this paper.

General description of the model

We model IF flow and transport of albumin in the choroidal tissue and the SCS, a potential space between the choroid and sclera, which is illustrated in Figure 2. IF, modeled as a Newtonian incompressible fluid with dynamic viscosity μ and uniform absolute temperature T , flows posteriorly from the anterior chamber, where it has hydrostatic pressure IOP, through the iris root, ciliary muscle and into the choroidal tissue and SCS. From there it flows out of the eye through the sclera or via the CC. Albumin flows into the choroidal tissue and SCS through the ‘inlet’ at the anterior choroid and leaves via advection across the sclera. There is also exchange with the CC via both advection and diffusion.

We treat the choroidal tissue as a perfused porous layer of tissue, permeable to IF, with (Darcy) permeability k_C (proportional to the hydraulic conductivity). Albumin diffuses in this tissue with diffusion coefficient D , and is advected with the IF flow. The choroid has a prescribed thickness, h_C , which varies from anterior to posterior, and h_{C0} is the average thickness.

The SCS is a potential space that exists between the choroid and sclera, which, although narrow in comparison to both of these tissues, represents a relatively low resistance pathway for IF flow.^{57,58,59} We assume (for simplicity and modeling expedience) that this space maintains a uniform prescribed thickness, h_{P0} , under physiological conditions. It is expected that the hydrostatic pressure in the SCS will change from its physiological value during non-physiological conditions, such as changes in the IOP or after pharmacological treatment. It is straightforward to separate the sclera and the choroid during surgery, and, furthermore, the observations of Croft *et al.*⁶⁰ show choroidal movement of between 0.1 and 1 mm parallel to the sclera during accommodation, both of which suggest there is little or no adhesion between the choroid and sclera. For simplicity in this

1
2 work, we assume there is no adhesion between these surfaces. Hence, any variation in IF pressure
3
4 in the SCS or choroidal tissue could cause the choroidal tissue to deform (the scleral tissue is much
5
6 stiffer), altering the thickness of the SCS.

7
8 As such, we assume that the actual thickness of the SCS in non-physiological conditions, h_P ,
9
10 is dependent on the departure from the physiological value of the difference between the pressure
11
12 in the SCS itself and that on the apical retinal pigment epithelium (RPE), p_R (see Equation (3)
13
14 in the Appendix for a precise definition), and we assume p_R equals the IOP (but interrogate the
15
16 effect of lower values in Figure 7B in the Appendix). If this pressure difference is positive and
17
18 large compared to λ_P (a parameter characterizing the elastic stiffness of the SCS), such as might
19
20 occur during conditions of raised IOP, we model the thickness of the SCS as approximately linearly
21
22 dependent on this pressure difference. To account for small and/or negative pressure differences (as
23
24 is expected in normal physiological conditions), we modify the dependence to ensure that the SCS
25
26 never fully closes. With this approach the SCS can, in principle, become arbitrarily narrow, and
27
28 one could apply a suitable threshold to demarcate regions in which the SCS can be considered to be
29
30 collapsed (an extension of this approach could be used to facilitate understanding of experimental
31
32 observations⁶¹). Further details on our choice of constitutive law for the mechanical response of
33
34 the SCS can be found in the Supplementary Material.

35
36 The scleral surface of the SCS exhibits both cellular components and extracellular matrix.⁶²
37
38 However, it is not clear to what extent this structure inhibits the IF flow along this space and so for
39
40 this reason we adopt two approaches: in the main text we assume that IF can flow freely within
41
42 the SCS, while in the Supplementary Material we also consider the case of the IF flowing in the
43
44 gaps between material present in the space (flow through a porous medium). We find that the
45
46 predictions of the model are qualitatively insensitive to this choice. We assume that the IF does not
47

1
2 slip at the scleral and choroidal bounding surfaces of the SCS; the use of a more general condition
3
4 is discussed in the Supplementary Material, where we show that the corresponding correction to
5
6 our results would be very small.

7
8 The ‘inlet’ of the model incorporates the posterior end of the ciliary body. We model the short
9
10 region (about 3 mm) of tissue from the anterior chamber to the inlet, which comprises the iris
11
12 root and ciliary body, as flow through a resistor with fixed resistance, R_{icm} , driven by a pressure
13
14 difference. Thus, the hydrostatic pressure difference between the IOP in the anterior chamber and
15
16 that at the inlet of the model (lower) equals R_{icm} multiplied by the volume flow rate. Albumin in
17
18 the ciliary body region is assumed to be well mixed and we prescribe zero diffusive flux at the inlet.

19
20 The inner surface of the choroid is bounded by the retinal pigment epithelium (RPE), across
21
22 which IF is pumped at a prescribed rate, $q_{\mathcal{R}}$. We assume there is no transport of albumin across the
23
24 RPE, owing to the presence of tight junctions connecting its cells.

25
26 We model the sclera as a rigid layer with thickness h_S . The inner surface of the sclera is
27
28 assumed to be spherical with radius R_0 , and the region under consideration in the mathematical
29
30 model spans up to an angle θ_0 from the posterior, corresponding to a distance of approximately
31
32 28.1 mm. On its outer surface we assume a uniform prescribed orbital pressure, $p_{\mathcal{O}}$, and albumin
33
34 concentration, $c_{\mathcal{O}}$. IF transport across the sclera is modeled using Starling’s law on the assumption
35
36 that albumin is the only solute contributing to the local osmotic pressure difference between the
37
38 IF in the choroidal tissue and that in the orbit. The hydraulic conductance is $k_S/(h_S\mu)$, where
39
40 k_S is the (Darcy) permeability of the scleral tissue, osmotic reflection coefficient is σ_S , and ideal
41
42 gas constant is R_g . Albumin transport across the sclera is modeled by the Patlak equation⁶³ with
43
44 permeability coefficient β_S .

45
46 Within the choroidal tissue, IF and albumin pass to and from the fenestrated CC. We describe
47

1
 2 the CC as uniformly distributed within the choroidal tissue, with surface area per unit volume
 3
 4 of tissue S_{CC} , uniform hydrostatic pressure p_{CC} and albumin concentration c_{CC} . The hydrostatic
 5
 6 pressure in the CC is assumed to be uniformly $p_{CC} = \text{IOP} + 5 \text{ mmHg}$ (and accordingly varies with
 7
 8 the IOP, see also Page 50).⁴⁷ IF transport across the CC is modeled by Starling's equation, again on
 9
 10 the assumption that the local osmotic pressure difference only depends on differences in albumin
 11
 12 concentration between plasma and IF. In this case, the hydrostatic pressure difference tends to force
 13
 14 fluid out of the CC, while the osmotic pressure difference favours absorption into the CC. The
 15
 16 hydraulic conductance is $L_{p,CC}$ and reflection coefficient is σ_{CC} . Albumin is primarily transported
 17
 18 by transcytosis,²⁰ which for simplicity we model as a diffusion-like term with albumin conductance
 19
 20 $P_{a,CC}$. The reflection coefficient σ_{CC} is near unity;⁶⁴ therefore, there is also little advective transport
 21
 22 of albumin into the CC.

23
 24 The unconventional flow rate predicted by the model is defined as the total rate of IF flow
 25
 26 through the inlet of the model.

29 **Simplification of the model**

30
 31
 32 We focus on the transport along a fixed anterior-posterior slice bisecting the eye, and we assume all
 33
 34 variables are independent of their circumferential position around the eye, see Figure 2. Choroidal
 35
 36 thickness is around 2% of its anterior–posterior length, with the thickness of the SCS being much
 37
 38 smaller. As such, it is appropriate to assume variations in all variables across the thickness of
 39
 40 the SCS are much smaller than those along the length ($\sim 28.1 \text{ mm}$) of the SCS. For a given
 41
 42 distance posteriorly, it emerges that the IF pressures in the choroidal tissue and SCS vary negligibly
 43
 44 in the direction perpendicular to the sclera, and so, by continuity, these are equal, meaning that
 45
 46 we can work in terms of a single pressure variable, p , in both spaces, which depends only on
 47

1
2 the distance from the inlet. Similarly, a single albumin concentration, c , may be defined in both
3
4 spaces. After simplification, the resulting system of equations, consisting of four coupled ordinary
5
6 differential equations, governs the IF pressure, p , IF flow rate, Q ($\mu\text{l}/\text{min}$), albumin concentration,
7
8 c , and albumin flow rate, J (mol/s). These are stated in Equations (19)–(22) in the Appendix, with
9
10 boundary conditions below. We solve them using MATLAB R2024a (Mathworks), employing the
11
12 `bvp4c` solver, and present the results herein.

13 14 15 **Parameter values** 16

17
18 The model developed here is dependent on 28 parameters, whose typical physiological values are
19
20 listed in Table 1.

21
22 Three parameter values for which we have no information are the hydraulic resistance of the
23
24 iris root and ciliary muscle, R_{icm} , the (Darcy) permeability of the choroidal tissue, k_C , and the
25
26 physiological thickness of the SCS, h_{p0} . To set these three, we assume that the pressures at the
27
28 inlet and posterior pole match those in the literature,⁵⁸ and that the flow rates of IF through the
29
30 choroidal tissue and the SCS are equal at the inlet, see also Page 49. The latter choice is explored
31
32 in Figure 7A.

33
34 Furthermore, we could not find any value for the elasticity parameter quantifying changes in
35
36 the thickness of the SCS in non-physiological conditions, λ_P . We choose a value for this such that
37
38 there is significant expansion or collapse of the SCS over the range of IOP considered in this paper
39
40 (5 to 25 mmHg), which in turn allows us to assess whether collapsibility of the SCS is a possible
41
42 mechanism explaining the observed pressure insensitivity of the unconventional flow to the IOP.

43
44 Moreover, there are several other values listed in the Table that were either obtained indirectly
45
46 or for which values reported in the literature vary significantly. Details of all of these choices are
47

1
2 given in the Appendix (see references in the Table), and we have also indicated those parameter
3
4 values whose values have a degree of uncertainty ('*') or are very uncertain (**').

5 6 7 **Sensitivity analysis** 8

9
10 To quantify the effect of the uncertainty in the parameter values on the model output, we perform a
11
12 global sensitivity analysis using the extended Fourier amplitude sensitivity test, eFAST.⁶⁵ eFAST is
13
14 a variance-based approach that dissects output variance in the predictions using a spectral analysis
15
16 to calculate sensitivity indices (SI) for each parameter under consideration. These indices measure
17
18 to what degree each parameter value affects the model output. The first-order SI describes the
19
20 'isolated effect' of each parameter, as it quantifies the reduction in variance of the model output if
21
22 that parameter were kept fixed. The total SI captures both this first-order effect and the interaction
23
24 of the given parameter with the other parameters, as it measures the expected variance that would
25
26 remain if all but the given parameter were fixed. The sensitivity analysis was carried out using the
27
28 MATLAB codes developed by the group of Dr Kirschner.⁶⁶ More details on the implementation
29
30 are reported in the Appendix.

Table 1: Baseline physiological values of the model parameters, with references, where appropriate. Parameters whose values have a significant degree of uncertainty are denoted with ‘*’, while those for which we have no, or very little, information with ‘***’. For more details on the choices of parameter values, please see the Appendix (where appropriate, page references are provided).

Parameter value	Description
IF	
$\mu = 0.7 \cdot 10^{-3} \text{ Pa s}$	dynamic viscosity of IF (equal to that of water at 37°C)
$T = 37 + 273^\circ\text{K}$	absolute temperature
IOP = 15 mmHg	physiological intraocular pressure
Choroidal tissue	
$k_C = 2000k_S^{**}$	Darcy permeability of choroidal tissue, see Page 49
$D = 61 \cdot 10^{-12} \text{ m}^2\text{s}^{-1}$	albumin diffusion coefficient ^{67,68}
$h_C = (7.77\theta^3 - 45.4\theta^2 - 57.4\theta + 397) \mu\text{m}$	thickness profile (θ is the angle in radians subtended at the center of the eye between the point and the posterior pole), see Page 48
$h_{C0} = 266 \mu\text{m}$	typical choroidal thickness, ⁶⁹ see Page 48
SCS	
$h_{P0} = 2.4 \mu\text{m}^{**}$	thickness of SCS in physiological conditions, see Page 48

Table 1: (continued)

Parameter value	Description
$p_{\mathcal{R}} = \text{IOP}^*$	pressure on apical RPE, see Page 51
$\lambda_{\mathcal{P}} = 100 \text{ Pa}^{**}$	elasticity parameter, see Page 50
Inlet	
$R_{icm} = 1.27 \text{ mmHg}/(\mu\text{l}/\text{min})^{**}$	resistance of iris root and ciliary muscle, see Page 49
RPE	
$q_{\mathcal{R}} = 3 \cdot 10^{-8} \text{ m/s}^*$	flow across RPE, ^{70,71} see Page 50
Sclera	
$h_S = 0.5 \text{ mm}^*$	thickness of sclera, ⁷² see Page 48
$R_0 = 1.15 \cdot 10^{-2} \text{ m}$	radius of inner scleral surface, ⁷³ see Page 48
$\theta_0 = 140^\circ$	angle subtended at center of eye between anterior limit of choroid and posterior pole of eye, see Figure 2 and Page 48
$p_{\mathcal{O}} = 3 \text{ mmHg}^*$	orbital pressure, ⁷⁴ see Page 50
$c_{\mathcal{O}} = 0.19c_{cc}^*$	albumin concentration in orbit, ¹⁹ see Page 51
$k_S = 5.85 \cdot 10^{-18} \text{ m}^2$	Darcy permeability of scleral tissue, ⁷⁵ see Page 49
$\sigma_S = 0.38$	albumin reflection coefficient of sclera, ⁷⁵ see Page 53
$R_g = 8.314 \text{ J}/(\text{mol } ^\circ\text{K})$	ideal gas constant
$\beta_S = 8.3 \cdot 10^{-9} \text{ m/s}$	albumin conductance of sclera, ⁷⁵ see Page 53

Table 1: (continued)

Parameter value	Description
CC	
$S_{cc} = 12000 \text{ m}^{-1} *$	surface area of CC per unit volume of choroid, see Page 48
$p_{cc} = \text{IOP} + 5 \text{ mmHg} *$	hydrostatic pressure in CC, ⁴⁷ see Page 50
$c_{cc} = 0.66 \text{ mM}$	albumin concentration in blood vessels, ⁷⁶ see Page 51
$L_{p,cc} = 8 \cdot 10^{-11} \text{ m/s/Pa} **$	hydraulic conductance of the CC, ⁶⁴ see Page 50
$\sigma_{cc} = 0.95 *$	albumin reflection coefficient of CC, ⁶⁴ see Page 53
$P_{d,cc} = 8 \cdot 10^{-10} \text{ m/s} **$	albumin conductance of CC, ⁷⁷ see Page 53

RESULTS

In this Section, we start by presenting the model results with the baseline parameter set listed in Table 1, which we refer to as the ‘reference physiological case’; we follow this with the results of the sensitivity analysis of this solution. We will then discuss non-physiological cases, to examine the response of the unconventional flow to changing conditions: firstly, we impose a change in the IOP to determine how this influences the unconventional flow; secondly, we change the model parameters to investigate the postulated action of $\text{PGF}_{2\alpha}$ analogues, as well as comparing alternative explanations of the action of these drugs.

Reference physiological case

We use the parameter values listed in Table 1, and solve the governing equations to obtain the IF pressure and flow and the albumin concentration, with results shown in Figure 3.

The green curve in Figure 3A shows the pressure distribution in the IF, p , along the tissue of the choroid and the SCS, which we hereinafter refer to as the ‘choroid–SCS’, and, for comparison, the uniform pressures in the CC, p_{CC} (red), the orbit, p_O (cyan), and the IOP (magenta) are also shown. As prescribed by our choice of model parameters (R_{icm} , k_C and h_{p0} , see p. 49), the pressure is 14.2 mmHg at the inlet (posterior ciliary body / anterior choroid, slightly below the IOP of 15 mmHg), and 11.3 mmHg at the posterior pole. Both values are well above the orbital pressure of 3 mmHg due to the high hydraulic resistance of the sclera. Our results show that the pressure rapidly decreases over the peripheral choroid–SCS near to the inlet, and has a near uniform pressure distribution in the posterior part of the choroid–SCS.

Figure 3B shows albumin concentrations, with the green curve showing the concentration in the choroid–SCS, c , with the uniform albumin concentrations imposed in the CC, c_{CC} (red), and orbit, c_O (cyan), plotted for comparison. The concentration in the choroid–SCS is everywhere higher than the orbital concentration and lower than that in the CC, and it exhibits only a slight decrease over the peripheral choroid–SCS, with a near uniform concentration in the posterior. The overall mean concentration is 28.8% of that in plasma (0.66 mM, see Table 1), which is within the range 10–35% reported in the literature.^{18,19}

In Figures 3C and D we show the IF and albumin flow rates, respectively, along the choroid–SCS. The total IF flow rate (purple solid line), which is the sum of flows through the choroidal tissue (dashed line) and the SCS (dotted line), decreases towards the posterior pole. At the inlet,

1
2 the overall IF flow rate in the choroidal tissue and that in the SCS are equal by our choice of
3 parameters, and the unconventional flow rate equals the sum of these two rates. The value predicted
4 by the model has the right order of magnitude: assuming an aqueous production rate of $2.5 \mu\text{l}/\text{min}$,
5
6 the model predicts that the unconventional pathway accounts for approximately 25% of the total
7
8 outflow.⁹ The albumin flow rate (Figure 3D) follows the same pattern as that of the IF, decreasing
9
10 as it progresses through the domain.
11
12

13 Figure 3E details the IF exchange between the choroid–SCS and the surrounding compartments,
14 specifically the CC (red); sclera (blue); and RPE (yellow). Positive values indicate flows into the
15
16 choroid–SCS, while negative values indicate flows out. Table 2 gives the corresponding flow rates
17
18 of IF between the compartments. IF enters the choroid–SCS from both the inlet ($0.63 \mu\text{l}/\text{min}$, 19%)
19
20 and the RPE ($2.64 \mu\text{l}/\text{min}$, 81%), and leaves through the sclera ($1.67 \mu\text{l}/\text{min}$, 51%); in addition,
21
22 there is a net flow into the CC of $1.67 \mu\text{l}/\text{min}$ (49%).
23
24

25 Figure 3F and Table 2 show the corresponding exchanges for albumin (there is no albumin flow
26
27 across the RPE, as it is assumed to be impermeable to albumin). Thus, albumin enters the choroid–
28
29 SCS by a combination of flow through the inlet (61%) and transcytosis from the CC (39%), while
30
31 it leaves the choroid–SCS via advection with the IF flow into the CC (14%) and through the sclera
32
33 (86%). We note that the rate of albumin loss from the CC is $10.6 \cdot 10^{-13} \text{ mol/s}$ per eye, in good
34
35 agreement with measurements of Bill in a rabbit eye.⁷⁸ Also, note that the crystalloid component
36
37 of IF flows into the CC, but the net flow of albumin is out of the CC.
38
39
40

41 **Sensitivity analysis**

42
43
44 As mentioned, there is a degree of uncertainty in some of the parameter values, and to understand
45
46 the potential effect of this on the model results, we perform a global sensitivity analysis, whose
47

Table 2: IF

and albumin flow rates predicted by the model between the choroid–SCS and other compartments

(negative values indicate a flow out of the choroid–SCS).

Compartment	IF flow rate ($\mu\text{l}/\text{min}$)	Albumin flow rate (10^{-12} mol/s)
From inlet	0.63	2.60
Into/from CC	-1.61	1.06 (-0.57 enters CC by advection 1.63 leaves CC by transcytosis)
Through sclera	-1.66	-3.66
From RPE	2.64	0

output is two sensitivity indices (SI). These are plotted in Figure 4, which shows the effect of the choice of parameter values upon the resulting unconventional flow rate (A) and the average albumin concentration (B). The heights of the bars indicate the first-order (blue) and total-order (red) SI for each model parameter. In each case, the values of the first and total SI have similar magnitudes, suggesting that there is little interaction between the model parameters. For those parameter values with the highest SIs (for which we assume that a change in their value has the most significant effect on the output), the small black arrows show the direction of influence; an upward-pointing arrow indicates that increasing the parameter value increases the corresponding model output.

In Figure 4A, we see that the IF flow rate is most sensitive to the thickness of the SCS, h_{p0} . This is because the resistance of the SCS decreases as $1/h_{p0}^3$ when this thickness is increased, resulting in higher flow rates. Scleral (Darcy) permeability, k_S , is another important parameter: increasing k_S results in a higher trans-scleral outflow, and thus in a larger unconventional flow rate. Pressure

1
2 in the CC, p_{CC} , also plays an important role; increasing p_{CC} reduces the uveo-vortex flow into the
3
4 CC, and thus the unconventional flow rate.

5
6 Figure 4B shows that the average albumin concentration in the choroidal tissue is most sensitive
7
8 to the pressure in the CC, p_{CC} ; increasing this value results in decreasing albumin concentration.

9
10 Other important parameters are the albumin conductance of the CC, $P_{d,CC}$, and the flow across the
11
12 RPE, q_R . Increasing q_R results in a decrease of albumin in the tissue, as the fluid that comes from
13
14 the RPE is free of albumin. Increasing $P_{d,CC}$ increases the release of albumin from the CC and
15
16 thus its concentration in the choroid–SCS. In comparison to the key parameters mentioned here,
17
18 the remaining parameters have little influence on IF flow or albumin concentration.

19 20 21 **Dependence of the unconventional flow on the IOP**

22
23
24 In Figure 5A, we report with dots the experimental measurements by Bill⁵ on living monkeys
25
26 for the conventional (green) and unconventional (black) routes (with dashed straight lines drawn
27
28 through these points for clarity). Our predictions of unconventional outflow in the baseline case
29
30 are also reported (thick blue curve). The model predicts that unconventional flow rate increases in
31
32 a nearly linear fashion with the IOP, and that the relationship is slightly nonlinear at low values of
33
34 the IOP, with a steeper gradient at low IOP.

35
36 Our model includes various mechanisms that control the response of the model to a change of
37
38 IOP. In particular, (1) osmotic pressure differences due to local differences in albumin concentration
39
40 between the plasma in the CC and the IF in the choroidal tissue, which affects fluid exchange across
41
42 vessel walls, and osmotic pressure differences across the sclera, which affects the trans-scleral flow,
43
44 and (2) the existence of the collapsible SCS. A strength of mathematical modeling is that these
45
46 effects can be individually switched on or off, and this is done in Figure 5B, as indicated in the
47

1
2 table in the legend.

3
4 If we remove the effect of differences in albumin concentration, both across the vessel walls of
5
6 the CC and across the sclera, on the IF flow (which we do by setting the reflection coefficients for
7
8 the CC and sclera (σ_{CC} and σ_S , respectively) to zero, see dashed lines in Figure 5B), the direction of
9
10 the flow is towards the iris root, opposite to what is observed physiologically. This is because there
11
12 is no uveo-vortex flow in this case as the hydrostatic pressure is higher in the CC than it is in the
13
14 choroidal interstitial tissue. Since the resistance to outflow through the sclera is much higher than
15
16 the resistance to flow of the choroid–SCS, the IF partly escapes through the inlet towards the ciliary
17
18 body. We note that this direction of the unconventional flow is due to the choice of parameters in
19
20 Table 1, and it can be modified by lowering the resistance of the sclera, for instance. If the SCS and
21
22 oncotic pressure are both absent (red dashed line), the model predicts a perfectly linear dependence
23
24 of the IF flow rate on the IOP. With the SCS present (blue dashed line), the magnitude of the
25
26 flow is larger, as the combined resistance of the choroid–SCS is lower than that of the choroidal
27
28 tissue alone. The scenario changes significantly when the effect of osmotic pressure differences are
29
30 included (solid curves) and the IF flows posteriorly through the inlet in the correct direction.

31
32 We note that the predicted values of the unconventional flow rates with no SCS (red curve, errors
33
34 of 12 and 3% at 11 and 22 mmHg, respectively) fit the absolute values given by Bill's data slightly
35
36 better than those with the SCS present (blue curve, errors of 27 and 18%). However, the slope
37
38 of our predicted values with increasing IOP is noticeably closer to that of Bill's data in the case
39
40 with the SCS present (between 11 and 22 mmHg the slope of Bill's data is $0.0173 \mu\text{l}/\text{min}/\text{mmHg}$,
41
42 and the corresponding slopes of the blue (with SCS) and red (without SCS) curves are 0.0165 and
43
44 $0.202 \mu\text{l}/\text{min}/\text{mmHg}$, respectively). Given that Bill's data was measured in monkeys (which have
45
46 a very different baseline IOP, 11 vs. 15 mmHg in humans), we feel that this slope is a more robust
47

comparator between the approaches, since it quantifies the unconventional flow rate in terms of an IOP increase from a baseline value (which could vary from individual to individual within a species).

The use of prostaglandin $F_{2\alpha}$ analogues

In this Section, we will test various possible mechanisms of unconventional flow increase when $PGF_{2\alpha}$ analogues are used. The mechanisms we will consider include: lowering the resistance of the iris root and ciliary muscle, R_{icm} , increasing the scleral hydraulic conductivity, k_S , lowering the CC pressure, p_{CC} , and increasing the capillary hydraulic conductance, $L_{p,CC}$. In all cases, we will consider the departure from the reference physiological case, while keeping the IOP fixed at 15 mmHg.

The standard explanation for the increase in the unconventional flow rate when using $PGF_{2\alpha}$ analogues is that there is a reduction in the resistance to the flow across the ciliary muscular tissue, corresponding to a drop in the parameter R_{icm} in our model.^{29,30} We show this effect in Figure 6A (red curve and top axis). As R_{icm} reduces, the flow rate increases, by up to 20% when R_{icm} is reduced by a factor of 2. However, given the large increase in unconventional flow that is required for significant IOP reduction, this increase is unlikely to be sufficient to explain the action of the drug. Accordingly, we seek other contributory factors to explain the action of $PGF_{2\alpha}$ analogues.

A second possible explanation is that there is an increase in scleral hydraulic conductivity, k_S . The dependence of the unconventional flow rate on k_S is shown in Figure 6A (blue curve and bottom axis), indicating a 20% increase in the flow rate if the hydraulic conductivity is doubled.

Next, we explore the effects of venodilation, which is an alternative explanation for the increase in unconventional flow, in Figure 6B. We show the increase of the unconventional outflow rate

1
2 when CC hydrostatic pressure is reduced (red dashed and top axis). The pressure is reduced from
3
4 its physiological value of IOP + 5 mmHg (left) to IOP + 2.5 mmHg (right), which increases the
5
6 unconventional flow by only about 10%. We also show that the flow rate decreases with increasing
7
8 hydraulic conductance of the CC, $L_{p,CC}$ (blue curve and bottom axis). Increasing $L_{p,CC}$ twofold
9
10 decreases the flow rate by about 2%.

11
12 A possible explanation for the reported appearance of fluid spaces in the ciliary muscular tissue
13
14 is that an increase of albumin in the uveal interstitial tissue results in tissue edema. In Figure 6C,
15
16 we show the corresponding concentration of albumin at both the inlet and the average over the
17
18 choroid–SCS domain as hydraulic conductance of the CC, $L_{p,CC}$, increases. The concentration
19
20 increases with $L_{p,CC}$, because the exchange of albumin between the IF and blood is enhanced, and
21
22 this effect is more pronounced at the inlet (20% for a two-fold increase and 27% for a three-fold
23
24 increase) than the average over the choroid–SCS domain (9% and 11%, respectively). This is
25
26 because the trans-scleral flow removes the excess albumin along the length of the choroid–SCS,
27
28 but this effect is cumulative from the inlet.

29
30 This pronounced increase in albumin concentration explains the small decrease in unconven-
31
32 tional flow that is observed when increasing the hydraulic conductance, $L_{p,CC}$ (blue curve in Fig-
33
34 ure 6B), since the increased albumin concentration increases the osmotic pressure in the choroidal
35
36 tissue, reducing the flow from the choroidal tissue into the CC. This is despite the fact that resis-
37
38 tance to flow of the uveo-vortex pathway is reduced.

39
40 Figure 6D summarises the mechanisms discussed in this Section by showing the relative in-
41
42 crease of the unconventional flow due to the four different effects considered in Figures 6A,B.

DISCUSSION AND CONCLUSIONS

In this paper, we have presented a new mechanistic mathematical model of the unconventional flow pathway in the human eye. This model encodes the known anatomical features of unconventional flow, as described in the Section ‘Mathematical Model’, and it is encouraging that it succeeds in capturing some of the key physiological features.

Like many mathematical models of biological systems, our model depends on several parameters, some of which are uncertain. To understand the impact of our choice of parameters on the model outputs, we conducted a global sensitivity analysis. We found that, while the quantitative predictions are influenced by the specific parameter set used, the qualitative behaviour remains consistent. Importantly, the model offers a plausible explanation for the physics underlying the unconventional flow pathway in terms of the mechanisms of insensitivity of the flow rate to the IOP and the mechanisms of the action of $\text{PGF}_{2\alpha}$ analogues.

Pressure insensitivity

A key feature of the unconventional flow is that it is relatively insensitive to the IOP,⁷⁹ and our model similarly predicts only a weak dependency on the IOP, as well as predicting a flow of the correct magnitude and direction. Inspection of Figure 5A indicates that the relationship between the unconventional flow rate and the IOP is fairly close to linear (though the graph does show a slight curvature); the conventional flow has a linear dependence. However, the unconventional flow is only weakly dependent on the IOP, in the sense that increasing the IOP from 11 to 22 mmHg induces a relatively small percentage increase on the unconventional flow ($\approx 32\%$). This is not the case for the conventional flow, which increases in proportion to the difference from episcleral

1
2 venous pressure (corresponding to a 450% increase over the same range). This large difference is a
3
4 consequence of the fact that the IOP for which the unconventional flow rate is zero is < 5 mmHg,
5
6 whereas the corresponding pressure where the conventional flow rate is zero is the episcleral ve-
7
8 nous pressure of 8.4 mmHg. This significant difference in the pressures where the two flow rates
9
10 are zero, rather than any nonlinearity in the flow rate–IOP relationship, is the primary reason why
11
12 the observed change in the unconventional flow rate when the IOP is changed is so small com-
13
14 pared to the corresponding change in the conventional flow rate. This agrees with and explains the
15
16 experimental observations of Bill.⁵

17
18 To explain the pressure insensitivity mechanically, we note that there are two features of the
19
20 model that, in combination, reduce the dependence of the flow rate on the IOP, as follows. Firstly,
21
22 the uveo-vortex flow (the flow into the venous capillaries) is affected by the albumin concentration
23
24 in the IF, which is much lower than that in plasma, meaning the resulting osmotic pressure differ-
25
26 ence acts to absorb IF into the venous capillaries. Since the CC pressure is set at 5 mmHg higher
27
28 than the IOP, the hydrostatic pressure difference between the IF and blood does not change with
29
30 the IOP, as the IOP is varied, and thus the difference in both the hydrostatic and osmotic pressures
31
32 between IF and blood do not depend strongly on the IOP, with the result that the uveo-vortex flow
33
34 does not depend strongly on the IOP also.⁸⁰

35
36 Secondly, the SCS is compressible, resulting in it having a variable hydraulic resistance. When
37
38 the pressure in the SCS is low, the SCS compresses, raising its resistance to IF flow and tending to
39
40 reduce the flow. Conversely, the large hydrostatic pressure drop between the IOP and the relatively
41
42 low SCS pressure drives increased IF flow. When the pressure in the SCS is high, the SCS expands,
43
44 tending to increase flow, but there is a small pressure drop, which tends to reduce the flow. The
45
46 result of these opposing mechanisms is that the flow rate in this space is relatively weakly dependent
47

1
2 upon the IOP, although our results suggest that this effect is relatively small.

3
4 In contrast, the other components governing the unconventional flow in the model, including the
5
6 flow through the inlet resistor, choroidal tissue and the trans-scleral flow,⁸¹ are all dependent on the
7
8 IOP. However, the relative insensitivity of the uveo-vortex flow on the IOP and the compressibility
9
10 of the SCS in combination with the other components results in a reduced overall sensitivity of the
11
12 unconventional flow rate to the IOP compared with that of the conventional flow rate, as observed
13
14 in experiments and predicted by the model.

15 16 17 **IOP-lowering action of PGF_{2α} analogues**

18
19
20 This model sheds light on the mechanisms of action of PGF_{2α} analogues used to treat glaucoma,
21
22 which work by increasing the unconventional outflow. Our model shows that reducing the resis-
23
24 tance of the iris root and ciliary muscle does have some effect on the unconventional flow rate, but
25
26 this is not enough to explain the expected pressure drop fully, suggesting that additional mecha-
27
28 nisms are required. These drugs are also reported to increase scleral hydraulic conductivity, and
29
30 our model predicts a modest increase in unconventional flow under this condition.

31
32 We also tested two further possible mechanisms. If the choroidal vessels dilate, the CC pressure
33
34 will be reduced, and in these conditions, our results indicate a relatively small increase in the
35
36 unconventional flow. Furthermore, increasing the hydraulic conductivity of the CC has little effect
37
38 on the flow (it decreases very slightly).

39
40 It is likely that the observed IOP-lowering action of PGF_{2α} analogues is due to a combination
41
42 of the first three mechanisms. We do note, however, that the current investigation is preliminary.
43
44 In particular, we assumed a normal IOP of 15 mmHg, but in reality the drug is usually given to
45
46 patients with primary open-angle glaucoma who have a much higher IOP. For these patients, the
47

1
2 unconventional outflow would be a much higher proportion of total outflow (as the conventional
3
4 outflow is impaired). Further research is required to understand fully the role of $\text{PGF}_{2\alpha}$ analogues
5
6 in pathological conditions.

7 8 9 **Other actions of $\text{PGF}_{2\alpha}$ analogues**

10
11
12 The observation that latanoprost causes choroidal thickening mentioned in the Introduction could
13
14 be explained due to direct action of the drug, but our model gives a possible alternative explanation
15
16 for this, as follows. Since MMPs are involved in regulation of endothelial hydraulic conductance,
17
18 it could be that they increase the hydraulic conductance of the CC walls sufficiently to raise the
19
20 interstitial albumin level, particularly near the inlet, see Figure 6C, and such an increase is known to
21
22 cause tissue edema. The observed appearance of fluid-filled spaces in the ciliary muscle following
23
24 the use of $\text{PGF}_{2\alpha}$ analogues could be due to a similar mechanism.

25
26 It is also known that latanoprost stimulates accumulation of tracer from the anterior chamber
27
28 in the cervical lymph nodes,⁸² and that the pressure-lowering effects of latanoprost are compro-
29
30 mised in patients who have had their cervical lymph nodes removed.⁸³ We anticipate that use of
31
32 the drug would increase the trans-scleral albumin flux. For example, the percentage increases in
33
34 the trans-scleral albumin flux corresponding to the conditions of the four bars in Figure 6D are, re-
35
36 spectively, 17%, 40%, 34% and 10%; these are bigger increases even than the increases in IF flow,
37
38 suggesting that the albumin flux could increase significantly when the drug is used. This would
39
40 result in increased orbital albumin, which enters the orbital lymphatics *en route* to the destination
41
42 in the cervical lymph nodes. Failure to clear this washed-out albumin in patients with no cervical
43
44 lymph nodes could result in increased albumin concentration in the choroid–SCS. High albumin
45
46 concentrations in the choroid–SCS would reduce the uveo-vortex flow due to the reduced oncotic
47

1
2 pressure, and this could explain the poor results of latanoprost treatment in these patients.
3

4 5 **The role of lymphatics** 6

7
8 Although not the original goal of the model, our model does provide some insight into the role of
9
10 lymphatics. Our model shows that the IOP-dependent trans-scleral component of the unconven-
11
12 tional flow effectively removes albumin from the choroid–SCS. In this way, it takes on the role that
13
14 is undertaken by lymphatics in most other tissues (although there have been reports of lymphatics
15
16 present in the ciliary body,¹⁰ but this has not been confirmed by others⁸⁴). None have been reported
17
18 in the choroid, and, although tracers from the anterior chamber have been recovered from cervical
19
20 lymph nodes,⁸² this does not mean that the site of entry into the lymphatic system had to be in
21
22 the eye. The data is consistent with the tracers entering the lymphatic system in the orbit or in
23
24 the conjunctiva after exiting the eye.⁴⁶ The fact that our model produces accurate predictions with-
25
26 out incorporating lymphatics is in line with the consensus view that the choroid lacks a lymphatic
27
28 supply.
29

30 31 **Future potential** 32

33
34 There are several effects that could be worth further investigation, which we mention in this sec-
35
36 tion. First, as mentioned in the introduction, a number of other drugs can affect the unconventional
37
38 pathway, and extensions of this work could have implications for our understanding of their actions.
39
40 Second, increasing vascular hydraulic conductance or reducing venous pressure is expected to en-
41
42 hance the unconventional flow. There are drugs that can do this, and it would be worth exploring
43
44 their use as a potential alternative glaucoma treatments.
45

1
2 Third, the unconventional flow has been shown to carry particulate tracers from the SCS to the
3
4 posterior retina,⁸⁵ raising the possibility of a noninvasive route for drug delivery to the back of the
5
6 eye from the anterior chamber. Any agent that can enter the anterior chamber can potentially be
7
8 delivered to the posterior regions of the retina or the choroid by the unconventional flow. Although
9
10 this is currently considered unlikely,⁸⁶ it is worth further consideration, and using drugs such as
11
12 those mentioned in this paper to enhance the unconventional flow would also enhance the drug
13
14 delivery along this route. A model based on the one presented herein could be used as a starting
15
16 point to understand the distribution of the agent.

17
18 Fourth, the complex physiology represented by the resistor at the inlet of the model is highly
19
20 simplified, and could be explored in an extension of the model. A fraction of the unconventional
21
22 flow will leave via the vessels in the ciliary body, as well as across the sclera in this region, and we
23
24 do not account for this.

25
26 Fifth, the pressure at the inlet might not be uniform as there could be ‘hotspots’ of outflow in
27
28 the anterior chamber around the collector channels, which might perturb the pressure field there.

29
30 Sixth, in reality, the CC has a spatial distribution of pressures, with higher pressures near the
31
32 arterioles supplying blood to the CC and lower pressures near the venules that drain it, which are
33
34 spaced on a typical lengthscale of 400 to 800 μm , with pressure variation between these points.⁸⁷
35
36 Given that the vessels of the CC have large diameters of about 20 μm compared with typical cap-
37
38 illary vessels, and that the pressure drop needed to drive channel flow scales as the third power of
39
40 the channel width, we would expect the pressure variation within the CC to be small in comparison
41
42 with that in other capillary beds.

43
44 Seventh, saccades of the eye would affect the fluid pressures therein due to the acceleration,
45
46 and we have not accounted for these in our model.
47

Acknowledgements

None of the authors has conflicts of interest to disclose.

JHT acknowledges support from a Daphne Jackson Fellowship sponsored by the Royal Society and an Emmy Noether Fellowship from the London Mathematical Society, and is grateful to both these organisations for supporting her. She started work on this problem while based at Imperial College London.

DRO acknowledges support from the BrightFocus Foundation (G2015145).

TAS and PSS acknowledge funding from EPSRC grant number EP/P024270/1. PSS acknowledges funding from EPSRC grant numbers EP/N014642/1, EP/S030875/1 and EP/T017899/1.

The work presented herein arose from problems presented at a study group funded by the Macular Society, and a study group funded by the EPSRC-funded UK Fluids Network (grant numbers EP/N032861/1, EP/N032934/1, EP/P000851/1, EP/N032411/1, EP/N032152/1). We are grateful to both the Macular Society and the UK Fluids Network for supporting this research.

The work has also been supported by the Isaac Newton Institute (funded by EPSRC grant number EP/V521929/1).

For the purpose of open access, the author has applied a Creative Commons Attribution (CC-BY) licence to any Author Accepted Manuscript version arising.

Figures




Figure 1: Left: Schematic of the anterior segment of the eye. Aqueous humour flow in the posterior and anterior chambers is shown with cyan arrows. The conventional outflow is indicated by the green arrow, and the unconventional pathway is marked by black arrows. Right: Zoom of the choroid (based on Figure 4 by Wajer *et al.*⁸⁸). Bottom: cross-section, showing the layers of different types of vessels; top: view of apical surface of the CC.




Figure 2: Schematic diagram of the eye highlighting the set up and parameters used in the mathematical model. The anterior of the eye is at the top of the diagram and the model has rotational symmetry about the vertical axis $\theta = 0$. The inset zooms into the region we focus on in the model, which is the outer part of the posterior eye.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47




Figure 3: *Caption on the next page*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Figure 3: *Continued* Model results in the reference physiological case, showing the IF flow and albumin concentration plotted against distance from the inlet. (A) Pressure in the choroid–SCS, p , CC, p_{CC} , orbit, p_O and anterior chamber, IOP. (B) Albumin concentration in the choroid–SCS, c , CC, c_{CC} , and orbit, c_O . (C) IF flow rate posteriorly in the choroid–SCS, showing also the components of this flow in the choroidal tissue and SCS. (D) As (C) but for albumin flow rate. (E) IF flow rate per unit area of surface into (positive) and out of (negative) the choroid–SCS. (F) As (E) but for albumin (with the same colors). The flow of albumin out of the CC is the sum of the contributions due to transcytosis (out of the CC, positive) and to advection by the crystalloid fluid (into the CC, negative).




Figure 4: Results of sensitivity analysis showing the first-order (blue) and total-order (red) sensitivities for (A) the unconventional flow rate, and (B) the average albumin concentration across the choroid-SCS. The directions of influence for the most sensitive parameters are indicated by the black arrows above the bars, *i.e.* an upward (downward) arrow indicates that the flow rate/albumin concentration increases (decreases) as the given parameter is increased. The label d is the dummy variable, which is a parameter that does not appear in the model and hence represents a ‘negative control’ for the sensitivity analysis. In the simulations, k_S is varied independently of k_C .




Figure 5: Dependence of the unconventional flow rate on the IOP. (A) Comparison of conventional and unconventional flow rates. Solid dots: measurements of the conventional (green) and unconventional (black) flow rates by Bill on living monkeys with best fit dashed lines added.⁵ Blue curve: predictions of the model. (B) Detailed graph of predictions of the model accounting for various mechanisms. In the table in the figure inset, the column ‘Albumin’ indicates the presence or absence of albumin exchange with the vessels: ‘+’ indicates that there is exchange; ‘-’ that there is not (we set $\sigma_{CC} = \sigma_S = 0$ in the model). The column ‘SCS’ refers to the second mechanism: the symbols ‘+’ and ‘-’ indicate the presence or absence of a collapsible SCS, respectively. The black dashed line and blue solid curve are the same as those appearing in (A). In both figures, the black vertical line corresponds to the episcleral venous pressure of 8.4 mmHg and the black horizontal line marks zero flux. Note that, as the IOP is varied, we also vary the capillary pressure $p_{CC} = \text{IOP} + 5$ mmHg and apical RPE pressure $p_R = \text{IOP}$.




Figure 6: Figures (A) and (B) investigate four possible mechanisms of unconventional flow increase with the use of $\text{PGF}_{2\alpha}$ analogues, (C) shows albumin concentration in one case (increasing $L_{p,cc}$), and (D) summarises the four mechanisms. In all figures, we consider a departure from the reference physiological case (which is shown on the left sides of figure A-C), *i.e.* the SCS height is allowed to deform. (A) Top axis (red dashed): Increase in unconventional flow rate with decreasing iris root resistance, R_{icm} . The x -axis shows the factor by which R_{icm} is reduced, with 1 being the reference physiological case and 0.5 corresponding to the reduction of R_{icm} by a factor of 2. Bottom axis (blue solid): Unconventional flow rate increase with increasing scleral hydraulic conductivity, k_S . The x -axis is the factor by which k_S is increased. *Continued on the next page*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Figure 6: *Continued* (B) Top axis (red dashed): Unconventional flow rate increase as pressure in the capillaries, p_{CC} , is reduced (extreme left is the reference physiological case, $p_{CC} = \text{IOP} + 5 \text{ mmHg}$; extreme right is $p_{CC} = \text{IOP} + 2.5 \text{ mmHg}$). Bottom axis (blue solid): Unconventional flow rate increase with increasing hydraulic conductance of the vessel walls, $L_{p,CC}$. The x -axis is the factor by which we increase $L_{p,CC}$ from the reference physiological case. (C) Albumin concentration with increasing hydraulic conductance (x -axis is the same as the bottom one in B). The solid line is the albumin concentration at the inlet and the punctured line is the spatially averaged albumin concentration in the choroid–SCS. (D) Summary of all four mechanisms of relative increase of the flow rate: decreasing R_{icm} by a factor of 2, doubling k_S , reducing p_{CC} by 2.5 mmHg and doubling $L_{p,CC}$. The colors of the bars correspond to the colors of the lines in A and B.

Appendix

Details of mathematical model

Governing equations for the IF flow

We consider an axisymmetric model of the choroidal tissue, SCS and sclera. We assume the inner surface of the sclera is perfectly spherical with radius R_0 , and work in spherical coordinates (r, θ, ϕ) centered on the sphere with $\theta = 0$ in the direction of the posterior pole of the eye. The thickness of the SCS and choroid are given by $h_{\mathcal{P}}(\theta)$ and $h_{\mathcal{C}}(\theta)$, respectively, with $0 \leq \theta \leq \theta_0$ where $\theta = \theta_0$ gives the anterior limit of the choroid at the ciliary body region. As such, assuming axisymmetry, the SCS is defined over $R_0 - h_{\mathcal{P}}(\theta) < r < R_0$ and the choroid over $R_0 - h_{\mathcal{P}}(\theta) - h_{\mathcal{C}}(\theta) < r < R_0 - h_{\mathcal{P}}(\theta)$. We assume that $\epsilon_{\mathcal{C}} = h_{\mathcal{C}0}/R_0 \ll 1$, $\epsilon_{\mathcal{P}} = h_{\mathcal{P}0}/R_0 \ll 1$, where $h_{\mathcal{C}0}$ and $h_{\mathcal{P}0}$ are typical values of $h_{\mathcal{C}}$ and $h_{\mathcal{P}}$, respectively. We model the flow of IF and albumin transport in both the choroidal tissue and in the SCS, assuming that the IF is incompressible and Newtonian. The trans-scleral flow is driven by the difference between the IOP at the iris root and the orbital pressure (outside the sclera). In addition, within the choroid, there is IF flow and mass transport between the CC and the tissue. For simplicity, we describe the CC as uniformly distributed, with uniform properties throughout the layer.

In the tissue of the choroid, we assume incompressible Darcy flow, with $\mathbf{u}_{\mathcal{C}} = (u_{\mathcal{C}r}, u_{\mathcal{C}\theta}, 0)$ being the Darcy velocity (flow per unit area):

$$\nabla \cdot \mathbf{u}_{\mathcal{C}} = q_{\mathcal{C}}, \quad \mathbf{u}_{\mathcal{C}} = -\frac{k_{\mathcal{C}}}{\mu} \nabla p_{\mathcal{C}}, \quad (1)$$

where $k_{\mathcal{C}}$ is the Darcy permeability of the tissue, μ is the dynamic viscosity of the IF and $p_{\mathcal{C}}$ is the IF pressure. Within the choroidal tissue, the volume of IF entering the domain from the blood

1
2 vessels per unit time per unit volume of tissue is given by

$$\frac{1}{2} q_C = L_{CC} (p_{CC} - p_C - \sigma_{CC} R_g T (c_{CC} - c_C)). \quad (2)$$

3
4
5
6
7 Here L_{CC} represents the conductance of the CC, given by $L_{CC} = S_{CC} L_{p,CC}$, and p_{CC} denotes the
8 prescribed local blood pressure within these vessels. The albumin concentrations in the choroidal
9 tissue and CC are given by c_C and c_{CC} , respectively, with c_{CC} fixed. The symbol R_g denotes the
10 universal gas constant, T denotes the absolute temperature and σ_{CC} is the osmotic reflection coef-
11 ficient.
12
13
14
15

16
17 We assume the sclera is rigid, and the thickness of the SCS, h_P , is determined by the hydrostatic
18 pressure difference between the pressure there, p_P , and the pressure on the apical RPE, p_R . We
19 assume the constitutive law
20
21

$$\frac{1}{2} p_P = p_R + \lambda_P \left(\frac{h_P}{h_{P,\text{nat}}} - \left(\frac{h_P}{h_{P,\text{nat}}} \right)^{-n} \right), \quad (3)$$

22
23 in which we set the exponent n equal to unity, where λ_P is the prescribed elastic stiffness. The
24 variable $h_{P,\text{nat}}$ is the thickness that the SCS would adopt if there were no difference between the
25 IF pressure in the SCS and the pressure in the retina; note that this scenario is not physiological,
26 and that we will calculate values of $h_{P,\text{nat}}$ from the model. This law is similar to tube laws used to
27 model arteries and veins^{89,90,91,92} and the response is linearly elastic if $(p_P - p_R)/\lambda_P$ is large, but
28 the space strongly resists collapse if p_P is small, preventing complete closure.
29
30
31
32
33
34
35
36
37

38
39 In the SCS, flow is governed by the Stokes equations:

$$\frac{1}{2} \nabla \cdot \mathbf{u}_P = 0, \quad \nabla p_P = \mu \nabla^2 \mathbf{u}_P, \quad (4)$$

40
41 where $\mathbf{u}_P = (u_{Pr}, u_{P\theta}, 0)$ is IF velocity.
42
43
44
45
46
47

1
2 Treating the tissue of the sclera as a membrane, we assume that IF crosses due to the hydrostatic
3 and osmotic pressure differences across it:
4

$$5 \quad \hat{\mathbf{n}}_{\mathcal{P},\mathcal{S}} \cdot \mathbf{u}_{\mathcal{P}}|_{r=R_0} = -u_{\mathcal{P}r}|_{r=R_0} = -q_{\mathcal{S}} = -\frac{k_{\mathcal{S}}}{\mu h_{\mathcal{S}}} \left((p_{\mathcal{P}}|_{r=R_0} - p_{\mathcal{O}}) - \sigma_{\mathcal{S}} R_g T (c_{\mathcal{P}}|_{r=R_0} - c_{\mathcal{O}}) \right) \quad (5)$$

6
7
8
9 where $\hat{\mathbf{n}}_{\mathcal{P},\mathcal{S}} = -\hat{\mathbf{e}}_r$ is the inward pointing unit normal vector, $q_{\mathcal{S}}$ is the volume of IF crossing
10 the sclera per unit surface area per unit time, $k_{\mathcal{S}}$ is the Darcy permeability of the sclera, $h_{\mathcal{S}}$ is the
11 thickness of the sclera, $p_{\mathcal{O}}$ and $c_{\mathcal{O}}$ are the orbital pressure and albumin concentration, respectively
12 and $\sigma_{\mathcal{S}}$ is the reflection coefficient. We assume that IF enters the choroidal tissue from across the
13 RPE at a prescribed rate:
14
15
16
17
18

$$19 \quad \hat{\mathbf{n}}_{\mathcal{R},\mathcal{C}} \cdot \mathbf{u}_{\mathcal{C}}|_{r=R_0-h_{\mathcal{P}}-h_{\mathcal{C}}} = -q_{\mathcal{R}}, \quad (6)$$

20
21 where $q_{\mathcal{R}}$ is the flow per unit surface area and $\hat{\mathbf{n}}_{\mathcal{R},\mathcal{C}}$ is the inward pointing unit normal vector at the
22 RPE–choroid interface $r = R_0 - h_{\mathcal{P}} - h_{\mathcal{C}}$. Note that, the flow rate $q_{\mathcal{R}}$ is assumed to be uniform and
23 constant, as it is mainly driven by active ion transport⁵⁶ and only weakly depends on the hydrostatic
24 and oncotic pressure differences between the choroidal tissue and the sub-retinal space.⁹³
25
26
27
28

29 At the interface between the choroid and SCS, $r = R_0 - h_{\mathcal{P}}$, we have continuity of normal stress.
30 We assume the normal stresses of both the IF in the SCS and the fluid–porous medium complex are
31 dominated by the IF hydrostatic pressures (this assumption will be checked *a posteriori*), giving
32 the balance
33
34
35
36

$$37 \quad p_{\mathcal{C}}|_{r=R_0-h_{\mathcal{P}}} = p_{\mathcal{P}}|_{r=R_0-h_{\mathcal{P}}}. \quad (7)$$

38
39 Furthermore, mass conservation requires continuity of normal flux:
40
41

$$42 \quad \hat{\mathbf{n}}_{\mathcal{C},\mathcal{P}} \cdot \mathbf{u}_{\mathcal{C}}|_{r=R_0-h_{\mathcal{P}}} = \hat{\mathbf{n}}_{\mathcal{C},\mathcal{P}} \cdot \mathbf{u}_{\mathcal{P}}|_{r=R_0-h_{\mathcal{P}}}, \quad (8)$$

43
44 where $\hat{\mathbf{n}}_{\mathcal{C},\mathcal{P}}$ is the inward pointing unit normal vector at the choroid–SCS interface.
45
46
47

In the SCS, we need additional boundary conditions to model the physics of the tangential component of the flow, which are provided by the no-slip boundary condition:

$$\hat{\mathbf{t}}_{\mathcal{P},S} \cdot \mathbf{u}_{\mathcal{P}}|_{r=R_0} = 0, \quad \hat{\mathbf{t}}_{\mathcal{C},\mathcal{P}} \cdot \mathbf{u}|_{r=R_0-h_{\mathcal{P}}} = 0, \quad (9)$$

where $\hat{\mathbf{t}}_{\mathcal{P},S}$ and $\hat{\mathbf{t}}_{\mathcal{C},\mathcal{P}}$ are the corresponding unit tangential vectors.

Finally, we assume that the tissue between the anterior chamber and anterior choroid (comprising the tissue of the iris root and ciliary muscle) has resistance to fluid flow R_{icm} , which we assume is a constant parameter and independent of the pressure. This leads to the boundary condition

$$\text{IOP} - p = R_{icm}Q, \quad (10)$$

at $\theta = \theta_0$, where IOP is intraocular pressure, p is the mean pressure in the choroid–SCS, and Q is the volume flow through the tissue. Symmetry at the posterior pole $\theta = 0$ requires

$$\mathbf{u}_{\mathcal{C}} \cdot \hat{\mathbf{e}}_{\theta} = \mathbf{u}_{\mathcal{P}} \cdot \hat{\mathbf{e}}_{\theta} = 0. \quad (11)$$

Governing equations for albumin transport

We assume a dilute solution of albumin and use the steady diffusion–convection equation to model its transport:

$$\nabla \cdot \mathbf{j}_{\mathcal{C}} = s_{\mathcal{C}}, \quad \mathbf{j}_{\mathcal{C}} = \mathbf{u}_{\mathcal{C}}c_{\mathcal{C}} - D_{\mathcal{C}}\nabla c_{\mathcal{C}} \quad \text{in choroidal tissue}, \quad (12)$$

$$\nabla \cdot \mathbf{j}_{\mathcal{P}} = 0, \quad \mathbf{j}_{\mathcal{P}} = \mathbf{u}_{\mathcal{P}}c_{\mathcal{P}} - D_{\mathcal{P}}\nabla c_{\mathcal{P}} \quad \text{in SCS}. \quad (13)$$

In these equations, $\mathbf{j}_{\mathcal{C}}$ and $\mathbf{j}_{\mathcal{P}}$ are the fluxes of albumin in the choroidal tissue and SCS, respectively, $c_{\mathcal{C}}$ and $c_{\mathcal{P}}$ are concentrations in the choroidal tissue and SCS, respectively, $D_{\mathcal{C}}$ and $D_{\mathcal{P}}$ are the diffusion coefficients of albumin in the choroidal tissue and SCS, respectively; we assume $D_{\mathcal{C}} = D_{\mathcal{P}} = D$.

In the choroidal tissue, to model exchange of albumin with the vessels of the CC, we adapt equations based on those proposed by Kedem and Katchalsky.⁹⁴ The net rate of albumin entering the choroidal tissue per unit volume from the blood vessels locally reads:

$$s_c = \beta_{cc}(c_{cc} - c_c) + \frac{c_c + c_{cc}}{2}(1 - \sigma_{cc})q_c, \quad (14)$$

where, β_{cc} is the permeation coefficients of the CC per unit volume multiplied by R_gT , which is given by $\beta_{cc} = S_{cc}P_{a,cc}$, where $P_{a,cc}$ is the albumin conductance. The first term is a simplified description of transcytosis plus any other diffusive processes present, while the second term accounts for small advective backflow of albumin into the CC.

We assume that a flux s_S of albumin passes through the sclera (which has both advective and diffusive components and thus depends on the local fluid velocity and albumin concentrations), and that albumin is not allowed to pass through the RPE, giving the respective boundary conditions

$$\hat{\mathbf{n}}_{\mathcal{P},S} \cdot \mathbf{j}_{\mathcal{P}}|_{r=R_0} = s_S, \quad \hat{\mathbf{n}}_{\mathcal{R},C} \cdot \mathbf{j}_{\mathcal{C}}|_{r=R_0-h_{\mathcal{P}}-h_C} = 0, \quad (15)$$

with $\hat{\mathbf{n}}_{\mathcal{P},S}$, $\hat{\mathbf{n}}_{\mathcal{R},C}$ the outward pointing unit normals. At the choroid–SCS interface, $r = R_0 - h_{\mathcal{P}}$, we have continuity of concentration and normal flux:

$$c_{\mathcal{C}}|_{r=R_0-h_{\mathcal{P}}} = c_{\mathcal{P}}|_{r=R_0-h_{\mathcal{P}}}, \quad \hat{\mathbf{n}}_{\mathcal{C},\mathcal{P}} \cdot \mathbf{j}_{\mathcal{C}}|_{r=R_0-h_{\mathcal{P}}} = \hat{\mathbf{n}}_{\mathcal{C},\mathcal{P}} \cdot \mathbf{j}_{\mathcal{P}}|_{r=R_0-h_{\mathcal{P}}}. \quad (16)$$

To model the transport of albumin across the sclera we use the Patlak equation⁶³ which describes flow and species transport across a membrane, and write

$$s_S = q_S(1 - \sigma_S) \frac{c_{\mathcal{O}} - c_{\mathcal{P}}|_{r=R_0} e^{Pe_S}}{1 - e^{Pe_S}}, \quad (17)$$

where the Péclet number $Pe_S = q_S(1 - \sigma_S)/\beta_S$. Here, β_S is the permeation coefficient per unit area of the scleral tissue multiplied by R_gT , which we term the albumin conductance, σ_S is the

reflection coefficient of the scleral tissue and c_O is the orbital concentration, and q_S is the IF Darcy velocity across the sclera, given by Equation (5).

Finally in this Section, we use the boundary conditions

$$\left. \frac{\partial c_C}{\partial \theta} \right|_{\theta=\theta_0} = \left. \frac{\partial c_P}{\partial \theta} \right|_{\theta=\theta_0} = 0, \quad \left. \frac{\partial c_C}{\partial \theta} \right|_{\theta=0} = \left. \frac{\partial c_P}{\partial \theta} \right|_{\theta=0} = 0, \quad (18)$$

where (18a) follows from the assumption that the IF in the iris root and ciliary muscle is well mixed and (18b) is required for continuity of the diffusive flux of albumin.

Simplification of equations

We take advantage of the fact that the lengthscales of the fluid domain in the radial and azimuthal directions are very different, with the aspect ratio being $\epsilon_C = h_{C0}/R_0 \ll 1$. We thus scale the equations using appropriate different lengthscales in the two coordinate directions. In the choroidal tissue the radial lengthscale is h_{C0} and we scale the θ -velocity on $U_C = Q_0/(10 \times 2\pi R_0 h_{C0})$, where Q_0 is the aqueous humour production rate and the factor 10 is included so that the typical unconventional flow is around 10% of total aqueous flow. The corresponding pressure scale is $P_0 = \mu R_0 U_C / k_C$ and radial velocity scale is $\epsilon_C U_C$, and we scale the other variables accordingly. The radial component of Equation (1b) shows that the pressure is independent of the radial coordinate to leading order in ϵ_C .

In the SCS, we note that the pressure drop over a nominal length R_0 of choroidal tissue is $\mu R_0 U_C / k_C$. Using this as the scale for the pressure and with the corresponding θ -velocity scale $h_{P0}^2 U_C / k_C$, the pressure p_P is again independent of the radial coordinate to leading order. We solve for the radial dependence of the components of \mathbf{u}_C in the choroidal tissue in terms of p_C and its derivatives and in the SCS we solve for \mathbf{u}_P in terms of p_P and its derivatives, in each case finding

explicit expressions for the radial dependence. Matching the pressures in the choroidal tissue and SCS yields $p_C = p_R = p(\theta)$, along with first-order ordinary differential equations governing the IF pressure, p , in and flow rate, Q , along the choroid–SCS.

We scale the albumin concentration on c_0 , the albumin concentration in the CC ($c_0 = c_{CC}$), and scale the other variables correspondingly. We find that c_C and c_P are independent of the radial coordinate to leading order, and, after some work, can show that $c_C = c_P = c(\theta)$, and find first-order ordinary differential equations governing c and J , where J is the albumin flow rate along the choroidal tissue.

The resulting governing equations and boundary conditions for the four variables p , Q , c , J represent a great simplification over the full model (1)–(18), and they can be written:

$$\frac{dp}{d\theta} = - \frac{\mu Q}{2\pi \sin \theta (k_C h_C + h_P^3/12)}, \quad (19)$$

$$\frac{dQ}{d\theta} = 2\pi R_0^2 \sin \theta (h_C q_C + q_R - q_S), \quad (20)$$

$$\frac{dc}{d\theta} = \frac{cQ - J}{2\pi \sin \theta (D_C h_C + D_P h_P)}, \quad (21)$$

$$\frac{dJ}{d\theta} = 2\pi R_0^2 \sin \theta (h_C s_C - s_S). \quad (22)$$

Equations (20) and (22) express conservation of fluid and albumin, respectively. Equation (19) states that the pressure gradient is proportional to volumetric flow rate times the resistance to the flow, and is derived from integrating the θ -velocity from Darcy's law in the choroidal tissue (Equation (1)) and Stokes flow in the SCS (Equation (4)) over the thicknesses of these layers (multiplied by $2\pi R_0 \sin \theta$) to obtain an expression for Q that can be rearranged. Equation (21) gives an expression for the concentration gradient, and is derived from integrating Fick's law (Equations (12b) and (13b)) over the thickness of the choroid–SCS (also multiplied by $2\pi R_0 \sin \theta$) to obtain an expression for J that can be rearranged. The boundary conditions are $Q = J = 0$ at $\theta = 0$ and

Symbol	Description	Reference equation
$h_{\mathcal{P}}$	Thickness of SCS	(3)
$q_{\mathcal{C}}$	Volumetric flow rate out of vessels	(2)
$q_{\mathcal{R}}$	Volumetric flow rate across RPE	Prescribed
$q_{\mathcal{S}}$	Volumetric flow rate across sclera	(5)
$s_{\mathcal{C}}$	Species flow rate out of vessels	(14)
$s_{\mathcal{S}}$	Species flow rate across sclera	(17)

Table 3: Table giving references for expressions appearing in Equations (19)–(22).

IOP – $p = R_{icm}Q$, $J = cQ$ at $\theta = \theta_0$, and references for definitions of the symbols appearing in Equations (19)–(22) are provided in Table 3.

Regularity at $\theta = 0$ means the functions $h_{\mathcal{C}}$, $h_{\mathcal{P}}$, p , c , Q and J are even functions of θ , and we avoided numerical singularities at $\theta = 0$ (arising from division by $\sin \theta$) by solving on the domain $[\delta\theta, \theta_0]$, where $\delta\theta = \theta_0/n_{\text{pts}}$ is the grid spacing and n_{pts} is the number of grid points in θ . Note that during the solution process, the solver `bvp4c` chooses a mesh adaptively, based on the residual of the continuous solution.

This formulation has a major advantage over the equations before simplification in that it may readily be rapidly solved using a simple numerical scheme. When solving these equations with `bvp4c`, it was our experience that the choice of initial condition could be important for convergence of the program. Where possible, we changed parameter values in small steps, using the solution from the previous step as the initial condition for the current step.

1 2 **Estimation of parameter values**

3
4 The model relies on a number of parameters, some of which are difficult to estimate, owing to the
5
6 sparse availability of experimental measurements. The model parameters are reported in Table 1,
7
8 and, for those parameters requiring explanation, we report our reasoning in this Section.
9

10 11 **Geometry**

12
13 We model the eye internal to the sclera as a perfect sphere. To find an estimate for the outer scleral
14
15 radius R_0 , we note that typical healthy eye diameters in adult humans are 24.2 mm (transverse) and
16
17 23.7 mm (sagittal).⁹⁵ We average these and subtract a typical scleral thickness of $h_S = 0.5$ mm
18
19 (measured as between $[0.37, 0.86]$ mm).⁷² Our expression for the choroidal thickness profile was
20
21 derived by fitting a cubic polynomial to measurements in human emmetropes;⁶⁹ details are given
22
23 in the Supplementary Material.
24

25
26 The thickness of the SCS is small and difficult to estimate experimentally. We choose a value
27
28 of h_{P0} to provide an equal flow through the choroidal tissue and SCS, so that both flow rates are
29
30 comparable, and we include the dependency of the flow rate on this value in Figure 7A (we discuss
31
32 this Figure in more detail later).
33

34
35 In order to convert values of hydraulic and albumin conductances of the CC from the values in
36
37 individual vessels (given in the literature) to a value per unit volume of choroidal tissue (required
38
39 by the model), we need to multiply by S_{CC} , the surface area of the CC per unit volume of tissue.
40
41 We examine two ways to estimate S_{CC} . Firstly, modeling the CC as being composed of vessels
42
43 of circular cross-section, we estimate $S_{CC} = 4\phi/d$, where ϕ is the volume fraction of the CC and
44
45 d are their diameters. Denoting Φ as the area fraction occupied by the septae (the voids between
46
47

capillaries), we obtain the volume fraction of the CC as $(1 - \Phi)d/h_c$, and hence $S_{CC} = 4(1 - \Phi)/h_c$.

Secondly, the model of Zouache *et al.*⁸⁷ (ignoring the surface areas of the feeding arterioles and draining venules) gives a surface area to volume ratio of $S_{CC} = 2(1 + (2d/d_s - 1)\Phi)/h_c$, where d_s is the septal diameter, and the authors quote $d = 20 \mu\text{m}$, $d_s = 3$ to $24 \mu\text{m}$ and $\Phi = 0.25$ to 0.5 . These two methods give values of S_{CC} in the ranges $10,000$ to $15,000 \text{ m}^{-1}$ and $12,000$ to $72,000 \text{ m}^{-1}$, respectively; we take $S_{CC} = 12,000 \text{ m}^{-1}$.

Fluid

The Darcy permeability of human scleral tissue was estimated as $19 \cdot 10^{-13} \text{ cm}^4 \text{ dyn}^{-1} \text{ s}^{-1}$,⁹⁶ and, multiplying by the dynamic viscosity, this gives an estimate of k_S as $1.33 \cdot 10^{-18} \text{ m}^2$; k_S was also estimated as $5.85 \cdot 10^{-18} \text{ m}^2$.²⁵ We use the latter value, as those measurements are more recent, and study the effect of this parameter in the sensitivity analysis.

Little is known about the Darcy permeability of the choroidal tissue, k_C , except that it is larger than that of the sclera,⁹⁷ and little is known about the resistance to flow of the tissue of the iris root and ciliary muscle, R_{icm} . In addition, little is known about the SCS thickness h_{P0} ; for example, in their experiments on rabbits, Chiang *et al.*⁹⁸ assumed that the physiological SCS thickness was ‘less than $25 \mu\text{m}$ ’, which is not specific enough for our model. To set these three unknown model parameters for the reference physiological case, we (1) make the model match the IF pressures at the inlet and the pole to the values measured by Emi *et al.*, which (for physiological values of the IOP) are $0.8 \pm 0.2 \text{ mmHg}$ and $3.7 \pm 0.5 \text{ mmHg}$ lower than the IOP at the anterior and posterior of the SCS, respectively,⁵⁸ and (2) set the proportions of the unconventional flow that pass through the choroidal tissue and the SCS at the inlet. This means that if we change this proportion, we also need to adjust the parameters k_C and R_{icm} to reproduce the pressures at the inlet and the pole.

1
2 In Figure 7A we show how changing this proportion (and correspondingly changing k_C and R_{icm})
3
4 affects the unconventional flow. Even over the range 5 to 100%, the unconventional flow only
5
6 changes by around 10%, suggesting that the model is not very sensitive to the chosen value of this
7
8 proportion.

9
10 For the hydraulic conductivity of the CC, $L_{p,CC}$, we choose the value $L_{p,CC} = 8 \cdot 10^{-11}$ m/s/Pa,
11
12 which is of the same order of magnitude as that for other fenestrated capillaries (*e.g.* renal per-
13
14 itubular capillaries $4.4 \cdot 10^{-10}$ m/s/Pa and intestinal mucosa $3.8 \cdot 10^{-11}$ m/s/Pa),⁷⁷ and investigate
15
16 its effect with our sensitivity analysis.

17
18 Measured flow through the RPE in humans is (261 ± 130) $\mu\text{l}/\text{cm}^2/\text{day} \approx (3.0 \pm 1.5) \cdot 10^{-8}$ m/s,⁹⁹
19
20 in monkeys is 0.14 or 0.36 $\mu\text{l}/\text{mm}^2/\text{h} \approx 3.9 \cdot 10^{-8}$ or $1.0 \cdot 10^{-7}$ m/s,^{100,101} and in human fetal retinal
21
22 pigment epithelia *in vitro* is 10 $\mu\text{l}/\text{cm}^2/\text{h} = 2.8 \cdot 10^{-8}$ m/s.⁷¹ We use the value $q_{\mathcal{R}} = 3 \cdot 10^{-8}$ m/s,
23
24 which is similar to these values, and investigate the effect of this choice in the sensitivity analysis.

25 26 27 **Pressures**

28
29
30 In rabbit studies, for IOPs greater than around 10–15 mmHg and up to over 100 mmHg, the
31
32 choroidal venous pressure was shown to be slightly above the IOP.⁸⁰ Later work confirmed these
33
34 findings, showing the choroidal venous pressure is around 3 cmH₂O above the IOP (IOP + 2 mmHg)
35
36 and that the CC pressures are 7 to 13 cmH₂O above the IOP (IOP + 5 to IOP + 10 mmHg).⁴⁷ In
37
38 our model we set pressure in the CC to be $p_{CC} = \text{IOP} + 5$ mmHg, which is the lower boundary of
39
40 the measured values. We study the effect of this parameter in the sensitivity analysis.

41
42 We take the orbital pressure to be $p_{\mathcal{O}} = 3$ mmHg, which is within the typical range in the liter-
43
44 ature of 2 to 6 mmHg,⁷⁴ and investigate the effect of the uncertainty in the value in the sensitivity
45
46 analysis.

1
2 As discussed in the Section ‘General description of the model’ starting on Page 9, we assume
3
4 that under normal physiological conditions the SCS has uniform prescribed thickness $h_{\mathcal{P}0}$. The
5
6 SCS thickness responds elastically if the IOP changes. We choose the parameter characterising
7
8 the elasticity, see Equation (3), as $\lambda_{\mathcal{P}} = 100$ Pa. As mentioned in the Section ‘Parameter values’,
9
10 this value was chosen so that there is significant expansion or collapse of the SCS over the range
11
12 of the IOP considered in this paper (5 to 25 mmHg). The role of this parameter is investigated in
13
14 Figure 7B, where the dependency of flow rate on the IOP is shown for three values of $\lambda_{\mathcal{P}}$. The blue
15
16 curve ($\lambda_{\mathcal{P}} = 100$ Pa) is the same as that in Figure 5. Figure 7B shows that for $\lambda_{\mathcal{P}} = 1000$ Pa, the
17
18 SCS is stiff and the response to the IOP increase is almost linear. For $\lambda_{\mathcal{P}} = 10$ Pa, the SCS is more
19
20 flexible and the flow exhibits more non-linearity for small IOP, and shows a slower increase with
21
22 the IOP at high values of the IOP than in the stiff case.

23
24 We assume that the retinal pressure $p_{\mathcal{R}} = \text{IOP}$; however, there is some variation in pressure
25
26 between the anterior and posterior of the eye due to the slow flow of fluid through the vitreous
27
28 humor, requiring a pressure drop to drive it. In addition, there is a flow of IF outward through the
29
30 retina, causing an additional pressure drop over the thickness of the retina. We compare the flow
31
32 with $p_{\mathcal{R}} = \text{IOP}$ with that for $p_{\mathcal{R}} = \text{IOP} - 0.5$ mmHg in Figure 7B, which suggests the choice of
33
34 $p_{\mathcal{R}}$ has little effect.

35 36 37 **Albumin**

38
39 Albumin concentration in blood vessels is around 44 g/l,⁷⁶ giving $c_{CC} = 0.66$ mM (molecular
40
41 weight of albumin is 66.5 kDa). We could not find a reference for albumin concentration in the
42
43 orbit, but the concentration in human choroidal tissue was given as 35% of the plasma value (though
44
45 measurements in only three eyes were reported, and these varied very widely) and that in the scleral
46
47




Figure 7: (A) Flow rate through inlet as the proportion of flow through the choroidal tissue is varied. Instead of requiring that the flow rates through the choroidal tissue and SCS are equal at the inlet, we demand that the flow rate through the choroidal tissue is a fixed percentage of the total unconventional flow rate. For each value of this percentage, we calculate values of R_{icm} , k_C and h_{P0} that also fix the correct pressures at the inlet and posterior pole (more details in text). In the Figure, we show the flow rates through the choroidal tissue and SCS as well as the total unconventional flow rate. (B) Dependence of the unconventional flow rate on the IOP for three different values of λ_P (legend, with $\lambda_P = 100$ Pa (blue curve) being the baseline physiological value). The pressures $p_{CC} = IOP + 5$ mmHg and $p_R = IOP$ vary with the IOP. The blue dashed line shows the case $p_R = IOP - 0.5$ mmHg (with $p_{CC} = IOP + 5$ mmHg, $\lambda_P = 100$ Pa).

1
2 tissue as 19% (in this case, four measurements were given, which varied widely),¹⁹ while in the
3
4 monkey the choroidal tissue concentration was given as 10% of that in plasma.¹⁸ In this paper, we
5
6 fix the orbital concentration at 19% of the blood plasma value and vary it in the sensitivity analysis
7
8 (Figure 4).

9
10 The albumin conductance of a fenestrated capillary is taken as $P_{d,CC} = 8 \cdot 10^{-10}$ m/s, which
11
12 is larger than the value $4 \cdot 10^{-10}$ m/s reported by Levick²¹ for fenestrated capillaries. We choose
13
14 a higher value since there are a vast number of caveolae present in the CC,²⁰ which will enhance
15
16 albumin transport by transcytosis. With this value of $P_{d,CC}$, we calculate an albumin flux from the
17
18 CC of $10.6 \cdot 10^{-13}$ mol/s, which is in good agreement with $9.35 \cdot 10^{-13}$ mol/s found by Bill in a
19
20 rabbit eye (0.0847 μ l/min of blood plasma albumin).⁷⁸ We check the dependency of the model on
21
22 $P_{d,CC}$ with the sensitivity analysis.

23
24 The osmotic reflection coefficient for transport across the capillary walls is taken as $\sigma_{CC} =$
25
26 0.95, which is in line with the results of Michel and collaborators,⁶⁴ who report values of the
27
28 reflection coefficient for albumin at least 0.9 for vessels with high hydraulic permeability (over
29
30 about $6 \cdot 10^{-11}$ m/s/Pa).

31
32 Anderson *et al.*⁷⁵ reported the partition coefficient for the sclera as 0.62, which was found
33
34 by leaving the scleral tissue in a solution containing albumin; the partition coefficient was the
35
36 concentration of albumin in the tissue divided by that in the solution. This implies a reflection
37
38 coefficient $\sigma_S = 0.38$. The albumin conductance across human sclera was measured as $\beta_S =$
39
40 $8.3 \cdot 10^{-9}$ m/s.⁷⁵

Sensitivity analysis

The details of the eFAST method are not reported here for brevity, as they are clearly described by Saltelli *et al*⁶⁵ and Marino *et al*.⁶⁶ In this method, each considered parameter is sampled from a uniform distribution within its corresponding range; the ranges are reported in Table 4. The sampling of each parameter follows a periodic curve with a certain frequency. For each parameter, five independent curves are considered, each with a random phase shift. Each curve samples 1000 points. We computed the sensitivity of the unconventional flow rate and the average albumin concentration to variations in selected parameters.

We will outline here a rigorous definition of the SIs. Consider the input parameters $X = (X_1, X_2, \dots, X_n)$ and the output Y , so that $Y = f(X)$, where f is the model. The first-order SI for parameter i is defined as

$$S_i = \frac{V_i}{V}. \quad (23)$$

In this expression $V = \text{Var}(Y)$ is the total variance of Y , and V_i is the first-order contribution of parameter i in the variance decomposition. This is defined as $V_i = \text{Var}(E(Y|X_i))$,⁶⁵ where, according to Bayesian notation, $E(Y|X_i)$ is the expected value of Y conditional on a fixed value of X_i . Since $V = \text{Var}(E(Y|X_i)) + E(\text{Var}(Y|X_i))$, the quantity V_i describes the expected reduction in the variance of the model output if parameter i was fixed.

The total SI for parameter i is defined as

$$S_{Ti} = \frac{E(\text{Var}(Y|X_{\sim i}))}{V}, \quad (24)$$

where $X_{\sim i}$ denotes a set of all parameters except i . The numerator therefore represents the expected variance that would remain if all parameters but parameter i are kept fixed.

The dummy parameter is a variable that does not appear in the model. It is included as a form of negative control.

Table 4: Ranges considered for each parameter in the sensitivity analysis. The dummy parameter d is varied between 0 and 1.

Parameter	Range	Units	Parameter	Range	Units
$P_{d,CC}$	$[4, 12] \cdot 10^{-10}$	m^{-1}	k_S	$[2.93, 8.78] \cdot 10^{-18}$	m^2
k_C	$[5.85, 17.6] \cdot 10^{-15}$	m^2	$L_{p,CC}$	$[4, 12] \cdot 10^{-11}$	m/s/Pa
q_R	$[1.5, 4.5] \cdot 10^{-8}$	m/s	R_{icm}	$[0.63, 1.9]$	$\text{mmHg}/(\mu\text{l}/\text{min})$
p_O	$[3, 6]$	mmHg	c_O	$[0.076, 0.176]$	mM
p_{CC}	$[18, 22]$	mmHg	h_{pO}	$[1.2, 3.6]$	μm

References

- [1] A. P. Nesterov, “13. The future for surgery in the glaucomas by increasing uveoscleral out-flow,” in *Glaucoma* (J. E. Cairns, ed.), pp. 257–273, Grune & Stratton, 1986.
- [2] A. Bill, “The drainage of blood from the uvea and the elimination of aqueous humour in rabbits,” *Exp. Eye Res.*, vol. 1, pp. 200–205, 1962.
- [3] A. Bill, “The aqueous humor drainage mechanism in the cynomolgus monkey (*Macaca irus*) with evidence for unconventional routes,” *Invest. Ophthalmol. Vis. Sci.*, vol. 4, pp. 911–919, 1965.

- 1
2 [4] A. Bill and K. Hellsing, “Production and drainage of aqueous humor in the cynomolgus
3
4 money (Macaca irus),” *Invest. Ophthalmol. Vis. Sci.*, vol. 4, pp. 920–926, 1965.
5
- 6 [5] A. Bill, “Conventional and uveo-scleral drainage of aqueous humour in the cynomolgus
7
8 monkey (Macaca irus) at normal and high intraocular pressures,” *Exp. Eye Res.*, vol. 5, no. 1,
9
10 pp. 45–54, 1966.
11
- 12 [6] A. Bill, “The routes for bulk drainage of aqueous humour in rabbits with and without cyclo-
13
14 dialysis,” *Doc. Ophthalmol.*, vol. 20, pp. 157–169, 1966.
15
- 16 [7] A. Bill, “The routes for bulk drainage of aqueous humour in the vervet monkey (Cercopithe-
17
18 cus ethiops),” *Exp. Eye Res.*, vol. 5, pp. 55–57, 1966.
19
- 20 [8] J. E. Pederson and C. B. Toris, “Uveoscleral outflow: diffusion or flow?,” *Invest. Ophthal-*
21
22 *mol. Vis. Sci.*, vol. 28, no. 6, pp. 1022–1024, 1987.
23
- 24 [9] A. Alm and S. F. E. Nilsson, “Uveoscleral outflow – A review,” *Exp. Eye Res.*, vol. 88,
25
26 pp. 760–768, 2009.
27
- 28 [10] Y. H. Yücel, M. G. Johnston, T. Ly, and *et al.*, “Identification of lymphatics in the ciliary
29
30 body of the human eye: A novel “uveolymphatic” outflow pathway,” *Exp. Eye Res.*, vol. 89,
31
32 pp. 810–819, 2009.
33
- 34 [11] F. Schroedl, A. Kaser-Eichberger, S. L. Schlereth, and *et al.*, “Consensus statement on the
35
36 immunohistochemical detection of ocular lymphatic vessels,” *Invest. Ophthalmol. Vis. Sci.*,
37
38 vol. 55, no. 10, pp. 6440–6442, 2014.
39
40
41
42
43
44
45
46
47

- 1
2 [12] U. R. Chowdhury, B. J. Madden, M. C. Charlesworth, and M. P. Fautsch, “Proteome analysis
3 of human aqueous humor,” *Invest. Ophthalmol. Vis. Sci.*, vol. 51, pp. 4921–4931, 2010.
4
5
- 6 [13] R. C. Tripathi, C. B. Millard, and B. J. Tripathi, “Protein composition of human aqueous
7 humor: SDS-PAGE analysis of surgical and post-mortem samples,” *Exp. Eye Res.*, vol. 48,
8 pp. 117–130, 1989.
9
10
- 11 [14] T. F. Freddo, S. P. Dartels, M. F. Darsotti, and R. D. Kamm, “The source of proteins in the
12 aqueous humor of the normal rabbit,” *Invest. Ophthalmol. Vis. Sci.*, vol. 31, no. 1, pp. 125–
13 137, 1990.
14
15
- 16 [15] G. Raviola, “The structural basis of the blood–ocular barriers,” *Exp. Eye Res.*, vol. 25,
17 pp. 27–63, 1977.
18
19
- 20 [16] A. Bill, “Capillary permeability to and extravascular dynamics of myoglobin, albumin and
21 gammaglobulin in the uvea,” *Acta. Physiol. Scand.*, vol. 73, no. 1, pp. 204–219, 1968.
22
23
- 24 [17] T. F. Freddo, “A contemporary concept of the blood-aqueous barrier,” *Prog. Retin. Eye Res.*,
25 vol. 32, pp. 181–195, 2013.
26
27
- 28 [18] C. B. Toris, J. E. Pederson, S. Tsuboi, D. S. Gregerson, and T. J. Rice, “Extravascular al-
29 bumin concentration of the uvea,” *Invest. Ophthalmol. Vis. Sci.*, vol. 31, no. 1, pp. 43–53,
30 1990.
31
32
- 33 [19] M. R. Allansmith, C. R. Whitney, B. H. McClellan, and L. P. Newman, “Immunoglobulins
34 in the human eye: location, type, and amount,” *Arch. Ophthalmol.*, vol. 89, no. 1, pp. 36–45,
35 1973.
36
37
38
39
40
41
42
43
44
45
46
47

- 1
2 [20] M. Nakanishi, G. R. I. A. Bhutto, M. Edwards, D. S. McLeod, and G. A. Luttly, “Albumen
3
4 transport to bruch’s membrane and rpe by choriocapillaris caveolae,” *Invest. Ophthalmol.*
5
6 *Vis. Sci.*, vol. 57, no. 4, pp. 2213–2224, 2016.
- 7
8 [21] N. Herring and D. J. Paterson, *Levick’s Introduction to Cardiovascular Physiology*. CRC
9
10 Press, 6 ed., 2018.
- 11
12 [22] J. Krohn and T. Bertelsen, “Corrosion casts of the suprachoroidal space and uveoscleral
13
14 drainage routes in the human eye,” *Acta. Physiol. Scand.*, vol. 75, no. 1, pp. 32–35, 1997.
- 15
16 [23] J. Krohn and T. Bertelsen, “Light microscopy of uveoscleral drainage routes after gelatine
17
18 injections into the suprachoroidal space,” *Acta. Physiol. Scand.*, vol. 76, no. 5, pp. 521–527,
19
20 1998.
- 21
22 [24] W. Deng, C. Liu, C. Parra, J. R. Sims, M. A. Faiq, A. Sainulabdeen, H. Song, and K. C.
23
24 Chan, “Quantitative imaging of the clearance systems in the eye and the brain,” *Quant.*
25
26 *Imaging Med. Surg.*, vol. 10, no. 1, p. 1, 2020.
- 27
28 [25] T. L. Jackson, A. Hussain, A. Hodgetts, A. M. S. Morley, J. Hillenkamp, P. M. Sullivan,
29
30 and J. Marshall, “Human scleral hydraulic conductivity: age-related changes, topographical
31
32 variation, and potential scleral outflow facility,” *Invest. Ophthalmol. Vis. Sci.*, vol. 47, no. 11,
33
34 pp. 4942–4946, 2006.
- 35
36 [26] J. R. Levick, “Capillary filtration-absorption balance reconsidered in light of dynamic ex-
37
38 travascular factors,” *Exp. Physiol.: Translation and Integration*, vol. 76, no. 6, pp. 825–857,
39
40 1991.
- 41
42
43
44
45
46
47

- 1
2 [27] J. E. Pederson, D. E. Gaasterland, and H. M. MacLellan, “Uveoscleral aqueous outflow in
3
4 the rhesus monkey: importance of uveal reabsorption,” *Invest. Ophthalmol. Vis. Sci.*, vol. 16,
5
6 no. 11, pp. 1008–1017, 1977.
7
- 8 [28] S. H. Sherman, K. Green, and A. M. Laties, “The fate of anterior chamber fluorescein in
9
10 the monkey eye. 1. The anterior chamber outflow pathways,” *Exp. Eye Res.*, vol. 27, no. 2,
11
12 pp. 159–173, 1978.
13
- 14 [29] J. F. Poyer, C. Millar, and P. L. Kaufman, “Prostaglandin F_{2α} effects on isolated rhesus
15
16 monkey ciliary muscle,” *Invest. Ophthalmol. Vis. Sci.*, vol. 36, no. 12, pp. 2461–2465, 1995.
17
18
- 19 [30] E. Lütjen-Drecoll and E. Tamm, “Morphological study of the anterior segment of cynomol-
20
21 gus monkey eyes following treatment with prostaglandin F_{2α},” *Exp. Eye Res.*, vol. 47,
22
23 pp. 761–769, 1988.
24
- 25 [31] R. N. Weinreb, K. Kashiwagi, F. Kashiwagi, S. Tsukahara, and J. D. Lindsey,
26
27 “Prostaglandins increase matrix metalloproteinase release from human ciliary smooth mus-
28
29 cle cells,” *Invest. Ophthalmol. Vis. Sci.*, vol. 38, no. 13, pp. 2772–2780, 1997.
30
31
- 32 [32] J.-W. Kim, J. D. Lindsey, N. Wang, and R. N. Weinreb, “Increased human scleral permeabil-
33
34 ity with prostaglandin exposure,” *Invest. Ophthalmol. Vis. Sci.*, vol. 42, no. 7, pp. 1514–1521,
35
36 2001.
37
- 38 [33] D. D. Gaton, T. Sagara, J. D. Lindsey, and R. N. Weinreb, “Matrix metalloproteinase-1
39
40 localization in the normal human uveoscleral outflow pathway,” *Invest. Ophthalmol. Vis.*
41
42 *Sci.*, vol. 40, no. 2, pp. 363–369, 1999.
43
44
45
46
47

- 1
2 [34] J. Stjernschantz, G. Selén, M. Astin, and B. Resul, “Microvascular effects of selective
3 prostaglandin analogues in the eye with special reference to latanoprost and glaucoma treat-
4 ment,” *Prog. Retin. Eye Res.*, vol. 19, pp. 459–496, 2000.
5
6
7
8 [35] D. K. Payne, J. W. Fuseler, and M. W. Owens, “Modulation of endothelial cell permeability
9 by lung carcinoma cells: A potential mechanism of malignant pleural effusion formation,”
10 *Inflammation*, vol. 18, no. 4, pp. 407–417, 1994.
11
12
13
14 [36] E. Svensjö, “Bradykinin and prostaglandin E1, E2 and F2 α -induced macromolecular leakage
15 in the hamster cheek pouch,” *Prostaglandins Med.*, vol. 1, no. 5, pp. 397–410, 1978.
16
17
18
19 [37] J. S. Alexander and J. W. Elrod, “Extracellular matrix, junctional integrity and matrix met-
20 alloproteinase interactions in endothelial permeability regulation,” *J. Anat.*, vol. 200, no. 6,
21 pp. 561–574, 2002.
22
23
24
25 [38] Z. Duru, C. Özsaygılı, D. M. Ulusoy, A. A. Demirtaş, A. Çiçek, and N. Duru, “Does using
26 topical latanoprost affect subfoveal choroidal thickness?,” *Cutan. Ocul. Toxicol.*, vol. 38,
27 pp. 370–374, 2019.
28
29
30
31
32 [39] K. Crawford and P. L. Kaufman, “Pilocarpine antagonizes prostaglandin F2 α -induced ocular
33 hypotension in monkeys: Evidence for enhancement of uveoscleral outflow by prostaglandin
34 F2 α ,” *Arch. Ophthalmol.*, vol. 105, pp. 1112–1116, 1987.
35
36
37
38
39 [40] A. Bill, “Effects of atropine and pilocarpine on aqueous humour dynamics in cynomoigus
40 monkeys (*macaca irus*),” *Exp. Eye Res.*, vol. 6, no. 2, pp. 120–125, 1967.
41
42
43
44 [41] N. R. Taylor, A. J. Zele, A. J. Vingrys, and R. G. Stanley, “Variation in intraocular pressure
45
46
47

1
2 following application of tropicamide in three different dog breeds,” *Vet. Ophthalmol.*, vol. 10,
3
4 no. s1, pp. 8–11, 2007.

5
6 [42] A. Bill, “Early effects of epinephrine on aqueous humor dynamics in vervet monkeys (*Cer-*
7
8 *copithecus ethiops*),” *Exp. Eye Res.*, vol. 8, pp. 35–43, 1969.

9
10
11 [43] A. Bill and P. E. Walinder, “The effects of pilocarpine on the dynamics of aqueous humor in
12
13 a primate (*Macaca irus*),” *Invest. Ophthalmol. Vis. Sci.*, vol. 5, pp. 170–175, 1966.

14
15
16 [44] M. Mori, M. Araie, M. Sakurai, and T. Oshika, “Effects of pilocarpine and tropicamide on
17
18 blood-aqueous barrier permeability in man,” *Invest. Ophthalmol. Vis. Sci.*, vol. 33, pp. 416–
19
20 423, 1992.

21
22 [45] R. F. Brubaker, “Goldmann’s equation and clinical measures of aqueous dynamics,” *Exp.*
23
24 *Eye Res.*, vol. 78, pp. 633–637, 2004.

25
26
27 [46] M. Johnson, J. W. McLaren, and D. R. Overby, “Unconventional aqueous humor outflow: A
28
29 review,” *Exp. Eye Res.*, vol. 158, pp. 94–111, 2017.

30
31
32 [47] O. Mäepea, “Pressures in the anterior ciliary arteries, choroidal veins and choriocapillaris,”
33
34 *Exp. Eye Res.*, vol. 54, no. 5, pp. 731–736, 1992.

35
36 [48] R. Sacco, G. Guidoboni, J. W. Jerome, G. Bonifazi, N. M. Marazzi, A. C. Verticchio Ver-
37
38 cellin, M. S. Lang, and A. Harris, “A theoretical approach for the electrochemical character-
39
40 ization of ciliary epithelium,” *Life*, vol. 10, no. 2, p. 8, 2020.

41
42
43 [49] M. Dvoriashyna, A. J. E. Foss, E. A. Gaffney, and R. Repetto, “A mathematical model of
44
45
46
47

1
2 aqueous humor production and composition,” *Invest. Ophthalmol. Vis. Sci.*, vol. 63, p. 1, Aug.
3
4 2022.

5
6 [50] J. J. Heys, V. H. Barocas, and M. J. Taravella, “Modeling passive mechanical interaction
7
8 between aqueous humor and iris,” *Journal of Biomechanical Engineering*, vol. 123, no. 6,
9
10 pp. 540–547, 2001.

11
12
13 [51] A. D. Fitt and G. Gonzalez, “Fluid mechanics of the human eye: aqueous humour flow in
14
15 the anterior chamber,” *Bull. Math. Biol.*, vol. 68, no. 1, pp. 53–71, 2006.

16
17
18 [52] M. Dvoriashyna, R. Repetto, M. R. Romano, and J. H. Tweedy, “Aqueous humour flow in
19
20 the posterior chamber of the eye and its modifications due to pupillary block and iridotomy,”
21
22 *Math. Med. Biol.*, vol. 35, no. 4, pp. 447–467, 2018.

23
24
25 [53] M. Dvoriashyna, R. Repetto, and J. H. Tweedy, “Oscillatory and steady streaming flow in
26
27 the anterior chamber of the moving eye,” *J. Fluid Mech.*, vol. 863, pp. 904–926, 2019.

28
29
30 [54] M. C. Johnson and R. D. Kamm, “The role of Schlemm’s canal in aqueous outflow from the
31
32 human eye,” *Invest. Ophthalmol. Vis. Sci.*, vol. 24, pp. 320–325, 1983.

33
34
35 [55] J. H. Siggers and C. R. Ethier, “Fluid mechanics of the eye,” *Annu. Rev. Fluid Mech.*, vol. 44,
36
37 pp. 347–372, 2012.

38
39
40 [56] M. Dvoriashyna, A. J. E. Foss, E. A. Gaffney, and R. Repetto, “Fluid and solute transport
41
42 across the retinal pigment epithelium: a theoretical model,” *J. Roy. Soc. Interface*, vol. 17,
43
44 p. 20190735, 2020.

- 1
2 [57] H. Inomata and A. Bill, “Exit sites of uveoscleral flow of aqueous humor in cynomolgus
3
4 monkey eyes,” *Exp. Eye Res.*, vol. 25, pp. 113–118, 1977.
5
- 6 [58] K. Emi, J. E. Pederson, and C. B. Toris, “Hydrostatic pressure of the suprachoroidal space,”
7
8 *Invest. Ophthalmol. Vis. Sci.*, vol. 30, no. 2, pp. 233–238, 1989.
9
- 10
11 [59] B. Chiang, J. H. Jung, and M. R. Prausnitz, “The suprachoroidal space as a route of admin-
12
13 istration to the posterior segment of the eye,” *Adv. Drug Deliver. Rev.*, vol. 126, pp. 58–66,
14
15 2018.
16
- 17
18 [60] M. A. Croft, E. Lütjen-Drecoll, and P. L. Kaufman, “Age-related posterior ciliary muscle
19
20 restriction – A link between trabecular meshwork and optic nerve head pathophysiology,”
21
22 *Exp. Eye Res.*, vol. 158, pp. 187–189, 2017.
23
- 24
25 [61] H.-L. Li, R. Ren, and H. Gong, “Segmental Unconventional Outflow in Mouse Eyes,” *In-*
26
27 *vest. Ophthalmol. Vis. Sci.*, vol. 64, p. 26, 2023.
28
- 29
30 [62] C. Platzl, A. Kaser-Eichberger, A. Benavente-Perez, and F. Schroedl, “The choroid-sclera
31
32 interface: An ultrastructural study,” *Heliyon*, vol. 8, no. 5, p. e09408, 2022.
33
- 34
35 [63] G. A. Truskey, F. Yuan, and D. F. Katz, *Transport Phenomena in Biological Systems*. Pearson
36
37 Education, 2004.
38
- 39
40 [64] C. C. Michel, “Capillary permeability and how it may change,” *J. Physiol.*, vol. 404, no. 1,
41
42 pp. 1–29, 1988.
43
- 44
45 [65] A. Saltelli, S. Tarantola, and K. P.-S. Chan, “A quantitative model-independent method for
46
47 global sensitivity analysis of model output,” *Technometrics*, vol. 41, no. 1, pp. 39–56, 1999.

- 1
2 [66] S. Marino, I. B. Hogue, C. J. Ray, and D. E. Kirschner, “A methodology for performing
3
4 global uncertainty and sensitivity analysis in systems biology,” *J. Theor. Biol.*, vol. 254, no. 1,
5
6 pp. 178–196, 2008.
- 7
8 [67] S. Vogel, *Life’s devices: the physical world of animals and plants*. Princeton University
9
10 Press, 1988.
- 11
12 [68] N. Meechai, A. M. Jamieson, and J. Blackwell, “Translational diffusion coefficients of
13
14 bovine serum albumin in aqueous solution at high ionic strength,” *J. Colloid Interface Sci.*,
15
16 vol. 218, no. 1, pp. 167–175, 1999.
- 17
18 [69] M. Hoseini-Yazdi, S. J. Vincent, M. J. Collins, S. A. Read, and D. Alonso-Caneiro, “Wide-
19
20 field choroidal thickness in myopes and emmetropes,” *Sci. Rep.*, vol. 9, p. 3474, 2019.
- 21
22 [70] B. A. Hughes, S. S. Miller, and T. E. Machen, “Effects of cyclic AMP on fluid absorption
23
24 and ion transport across frog retinal pigment epithelium. Measurements in the open-circuit
25
26 state,” *J. Gen. Physiol.*, vol. 83, no. 6, pp. 875–899, 1984.
- 27
28 [71] G. Shi, A. Maminishkis, T. Banzon, S. Jalickee, R. Li, J. Hammer, and S. S. Miller, “Control
29
30 of chemokine gradients by the retinal pigment epithelium,” *Invest. Ophthalmol. Vis. Sci.*,
31
32 vol. 49, no. 10, pp. 4620–4630, 2008.
- 33
34 [72] S. Vurgese, S. Panda-Jonas, and J. B. Jonas, “Scleral thickness in human eyes,” *PLoS ONE*,
35
36 vol. 7, no. 1, p. e29692, 2012.
- 37
38 [73] I. Bekerman, P. Gottlieb, and M. Vaiman, “Variations in eyeball diameters of the healthy
39
40 adults,” *J. Ophthalmol.*, vol. 2014, p. 503645, 2014.
- 41
42
43
44
45
46
47

- 1
2 [74] T. J. Enz and M. Tschopp, “Assessment of orbital compartment pressure: A comprehensive
3
4 review,” *Diagnostics*, vol. 12, no. 6, p. 1481, 2022.
5
- 6 [75] O. A. Anderson, T. L. Jackson, J. K. Singh, A. A. Hussain, and J. Marshall, “Human transs-
7
8 cleral albumin permeability and the effect of topographical location and donor age,” *Invest.*
9
10 *Ophthalmol. Vis. Sci.*, vol. 49, no. 9, pp. 4041–4045, 2008.
11
- 12 [76] R. Narang, D. Ridout, C. Nonis, and J. S. Kooner, “Serum calcium, phosphorus and albumin
13
14 levels in relation to the angiographic severity of coronary artery disease,” *Int. J. Cardiol.*,
15
16 vol. 60, no. 1, pp. 73–79, 1997.
17
- 18 [77] J. R. Levick and L. H. Smaje, “An analysis of the permeability of a fenestra,” *Microvascular*
19
20 *research*, vol. 33, no. 2, pp. 233–256, 1987.
21
- 22 [78] A. Bill, “The albumin exchange in the rabbit eye,” *Acta. Physiol. Scand.*, vol. 60, no. 1–2,
23
24 pp. 18–29, 1964.
25
- 26 [79] S. F. E. Nilsson and A. Bill, “Part i – normal: Anterior segment. Physiology and neuro-
27
28 physiology of aqueous humor inflow and outflow,” in *Glaucoma* (P. L. Kaufman and T. W.
29
30 Mittag, eds.), vol. 7 of *Textbook of Ophthalmology*, Mosby-Year Book Europe Ltd., 1994.
31
- 32 [80] A. Bill, “The uveal venous pressure,” *Arch. Ophthalmol.*, vol. 69, no. 6, pp. 780–782, 1963.
33
- 34 [81] R. N. Kleinstein and I. Fatt, “Pressure dependency of transcleral flow,” *Exp. Eye Res.*, vol. 24,
35
36 pp. 335–340, 1977.
37
- 38 [82] A. L. C. Tam, N. Gupta, Z. Zhang, and Y. H. Yücel, “Quantum dots trace lymphatic drainage
39
40 from the mouse eye,” *Nanotechnology*, vol. 22, p. 425101, 2011.
41
42
43
44
45
46
47

- 1
2 [83] Y. K. Kim, K. I. Na, J. W. Jeoung, and K. H. Park, “Intraocular pressure-lowering effect
3 of latanoprost is hampered by defective cervical lymphatic drainage,” *PLoS ONE*, vol. 12,
4 p. e0169683, 2017.
5
6
7
- 8 [84] A. Kaser-Eichberger, F. Schrödl, A. Trost, and *et al.*, “Topography of lymphatic markers in
9 human iris and ciliary body,” *Invest. Ophthalmol. Vis. Sci.*, vol. 56, pp. 4943–4953, 2015.
10
11
12
- 13 [85] C. L. Rowe-Rendleman, S. A. Durazo, U. B. Kompella, and *et al.*, “Drug and gene delivery
14 to the back of the eye: from bench to bedside,” *Invest. Ophthalmol. Vis. Sci.*, vol. 55, no. 4,
15 pp. 2714–2730, 2014.
16
17
18
- 19 [86] A. Alm, “Uveoscleral outflow,” *Eye*, vol. 14, p. 488491, 2000.
20
21
- 22 [87] M. A. Zouache, I. Eames, and P. J. Luthert, “Blood flow in the choriocapillaris,” *J. Fluid*
23 *Mech.*, vol. 774, pp. 37–66, 2015.
24
25
26
- 27 [88] S. D. Wajer, M. Taomoto, D. S. McLeod, R. L. McCally, H. Nishiwaki, M. E. Fabry, R. L.
28 Nagel, and G. A. Lutty, “Velocity measurements of normal and sickle red blood cells in the
29 rat retinal and choroidal vasculatures,” *Microvascular research*, vol. 60, no. 3, pp. 281–293,
30 2000.
31
32
33
34
35
- 36 [89] C. Cancelli and T. J. Pedley, “A separated-flow model for collapsible-tube oscillations,” *J.*
37 *Fluid Mech.*, vol. 157, pp. 375–404, 1985.
38
39
40
- 41 [90] B. S. Brook, S. A. E. G. Falle, and T. J. Pedley, “Numerical solutions for unsteady gravity-
42 driven flows in collapsible tubes: evolution and roll-wave instability of a steady state,” *J.*
43 *Fluid Mech.*, vol. 396, pp. 223–256, 1999.
44
45
46
47

- 1
2 [91] E. F. Toro and A. Siviglia, “Simplified blood flow model with discontinuous vessel proper-
3 ties: analysis and exact solutions,” in *Modeling of Physiological Flows: Modeling, Simula-*
4 *tion and Applications* (G. Ambrosi, A. Quarteroni, and G. Rozza, eds.), vol. 5, pp. 19–39,
5 Milano: Springer, 2012.
6
7
8
9
10 [92] E. F. Toro, “Brain venous haemodynamics, neurological diseases and mathematical mod-
11 elling: A review,” *Appl. Math. Comput.*, vol. 272, pp. 542–579, 2016.
12
13
14
15 [93] S. Hamann, “Molecular mechanisms of water transport in the eye,” *Int. Rev. Cytol.*, vol. 215,
16 pp. 395–431, 2002.
17
18
19 [94] O. Kedem and A. Katchalsky, “Thermodynamic analysis of the permeability of biological
20 membranes to nonelectrolytes,” *Biochim. Biophys. Acta*, vol. 27, pp. 229—246, 1958.
21
22
23 [95] I. Bekerman, P. Gottlieb, and M. Vaiman, “Variations in eyeball diameters of the healthy
24 adults,” *J. Ophthalmol.*, vol. 2014, 2014.
25
26
27
28 [96] I. Fatt and B. O. Hedbys, “Flow of water in the sclera,” *Exp. Eye Res.*, vol. 10, no. 2, pp. 243–
29 249, 1970.
30
31
32
33 [97] V.-P. Ranta and A. Urtti, “Transscleral drug delivery to the posterior eye: prospects of phar-
34 macokinetic modeling,” *Adv. Drug Deliv. Rev.*, vol. 58, no. 11, pp. 1164–1181, 2006.
35
36
37
38 [98] B. Chiang, N. Venugopal, H. E. Grossniklaus, J. H. Jung, H. F. Edelhauser, and M. R. Praus-
39 nitz, “Thickness and closure kinetics of the suprachoroidal space following microneedle
40 injection of liquid formulations,” *Invest. Ophthalmol. Vis. Sci.*, vol. 58, pp. 555–564, 2017.
41
42
43
44
45
46
47

1
2 [99] E. Chihara and N. Nao-i, “Resorption of subretinal fluid by transepithelial flow of the retinal
3
4 pigment epithelium,” *Graef. Arch. Clin. Exp.*, vol. 223, pp. 202–204, 1985.

5
6 [100] J. E. Pederson and H. L. Cantrill, “Experimental retinal detachment: V. Fluid movement
7
8 through the retinal hole,” *Arch. Ophthalmol.*, vol. 102, no. 1, pp. 136–139, 1984.

9
10 [101] H. L. Cantrill and J. E. Pederson, “Experimental retinal detachment: VI. The permeability
11
12 of the blood-retinal barrier,” *Arch. Ophthalmol.*, vol. 102, no. 5, pp. 747–751, 1984.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47