



City Research Online

City St George's, University of London

Citation: Linckelmann, M. (2025). The operad of Latin hypercubes. *Glasgow Mathematical Journal*, 67(3), pp. 475-486. doi: 10.1017/s0017089525000047

This is the published version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/34996/>

Link to published version: <https://doi.org/10.1017/s0017089525000047>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

RESEARCH ARTICLE

The operad of Latin hypercubes

Markus Linckelmann

School of Science & Technology, Department of Mathematics, City St George's, University of London, London, EC1V 0HB, UK

Email: markus.linckelmann.1@city.ac.uk

Received: 25 July 2024; Accepted: 15 January 2025

Keywords: Latin hypercube; operad

2020 Mathematics Subject Classification: Primary - 18M60, 18M05; Secondary - 05B15

Abstract

We show that the sets of d -dimensional Latin hypercubes over a non-empty set X , with d running over the positive integers, determine an operad which is isomorphic to a sub-operad of the endomorphism operad of X . We generalise this to categories with finite products, and then further to internal versions for certain Cartesian closed monoidal categories with pullbacks.

1. Introduction

There are several variants of the definition of Latin hypercubes in the literature – see the discussions and references in [1, §4] and [8, Section 1]. In order to define the version, we will be considering here, we fix the following notation and conventions. Categories are assumed to be essentially small. Let X be an object in a category \mathcal{C} with finite products and let d be a positive integer. The product X^d of d copies of X in \mathcal{C} is equipped with d canonical projections $\pi_i^d: X^d \rightarrow X$, characterised by the universal property that for any object Y and morphisms $\eta_i: Y \rightarrow X$, with $1 \leq i \leq d$, there is a unique morphism $\eta: Y \rightarrow X^d$ satisfying $\eta_i = \pi_i^d \circ \eta$ for $1 \leq i \leq d$. In that case we write $\eta = (\eta_i)_{1 \leq i \leq d}$. Applied to X^{d+1} instead of Y and any subset of d of the $d+1$ canonical projections π_j^{d+1} yields for every s such that $1 \leq s \leq d+1$ a unique morphism $\tau_s^{d+1}: X^{d+1} \rightarrow X^d$ satisfying $\pi_i^{d+1} = \pi_i^d \circ \tau_s^{d+1}$ for $1 \leq i \leq s-1$, and $\pi_i^{d+1} = \pi_{i-1}^d \circ \tau_s^{d+1}$ for $s+1 \leq i \leq d+1$. With the notation above, this is equivalent to

$$\tau_s^{d+1} = (\pi_1^{d+1}, \dots, \pi_{s-1}^{d+1}, \pi_{s+1}^{d+1}, \dots, \pi_{d+1}^{d+1}).$$

If X is a non-empty set, then X^{d+1} is the Cartesian product of $d+1$ copies of X , and we have

$$\tau_s^{d+1}(x_1, x_2, \dots, x_{d+1}) = (x_1, \dots, x_{s-1}, x_{s+1}, \dots, x_{d+1}),$$

where $x_i \in X$ for $1 \leq i \leq d+1$. That is, τ_s^{d+1} is the projection from X^{d+1} to X^d which discards the coordinate s in X^{d+1} .

Definition 1.1. Let \mathcal{C} be a category with finite products, let X be an object in \mathcal{C} , and let d be a positive integer. A Latin hypercube of dimension d over X is a morphism $\lambda: L \rightarrow X^{d+1}$ in \mathcal{C} such that $\tau_s^{d+1} \circ \lambda: L \rightarrow X^d$ is an isomorphism in \mathcal{C} , for $1 \leq s \leq d+1$.

The morphism λ in this Definition is necessarily a monomorphism. Two Latin hypercubes $\lambda: L \rightarrow X^{d+1}$ and $\lambda': L' \rightarrow X^{d+1}$ are called *isomorphic* if there is an isomorphism $\alpha: L \rightarrow L'$ such that $\lambda = \lambda' \circ \alpha$. In that case α is unique since λ' is a monomorphism.

A Latin hypercube of dimension d over a non-empty set X is uniquely isomorphic to a Latin hypercube given by a subset L of X^{d+1} with the property that for any choice of d of the $d+1$ coordinates of an element in X^{d+1} there is a unique element in the remaining coordinate such that the



resulting element belongs to L . In particular, we can identify a Latin hypercube of dimension d over a non-empty set uniquely, up to unique isomorphism, as the graph of the map $f: X^d \rightarrow X$ satisfying $(x_1, x_2, \dots, x_d, f(x_1, x_2, \dots, x_d)) \in L$ for all $(x_1, x_2, \dots, x_d) \in X^d$. In this way, Latin hypercubes of dimension d can be identified with a subset, denoted $\mathcal{L}(X^d, X)$, of the set $\text{Map}(X^d, X)$ of all maps from X^d to X . Any map in $\mathcal{L}(X^d, X)$ is clearly surjective. The following result shows that the subsets $\mathcal{L}(X^d, X)$ of $\text{Map}(X^d, X)$ with $d \in \mathbb{N}$ determine an operad. We refer to [6] or [7, Part II, §1.2] for basic terminology on operads.

Theorem 1.2. *Let X be a non-empty set. There is a sub-operad \mathcal{L} of the endomorphism operad \mathcal{E} of X such that $\mathcal{L}(d) = \mathcal{L}(X^d, X)$, for all $d \in \mathbb{N}$.*

This is not the most general version in which this result can be stated. Theorem 1.2 admits a generalisation to categories with finite products, which we will describe in Theorem 3.4, and a further generalisation, in Theorem 4.5, to certain Cartesian closed monoidal categories in which \mathcal{L} can be identified with a sub-operad of an internal endomorphism operad. We have chosen to state and prove this result first in the context of non-empty sets in order to not distract from the elementary nature of the proof. Operads were first introduced for topological spaces, and Theorem 1.2 holds verbatim for non-empty compactly generated topological spaces (this is a special case of Theorem 4.5; see Remark 4.8).

Remark 1.3. As pointed out in [8], if $d = 1$, then a Latin hypercube over a non-empty set X is a subset of X^2 of the form $\{(x, \sigma(x))\}_{x \in X}$ for some permutation σ of X , and hence Latin hypercubes of dimension 1 over X correspond to the elements of the symmetric group \mathfrak{S}_X of permutations of X , so they form themselves a group. This group structure is encoded as the structural map $- \circ_1 -$ on $\mathcal{L}(1)$ of the operad \mathcal{L} .

Remark 1.4. Let X be an object in a category \mathcal{C} with finite products and let d be a positive integer. The group $\text{Aut}_{\mathcal{C}}(X) \wr \mathfrak{S}_{d+1}$ acts on X^{d+1} , hence on the class of Latin hypercubes of dimension d over X , with the base group of $d + 1$ copies of $\text{Aut}_{\mathcal{C}}(X)$ acting by composing λ with a $(d + 1)$ -tuple of automorphisms of X^{d+1} and \mathfrak{S}_{d+1} acting on X^{d+1} by permuting the $d + 1$ canonical projections $\pi_i^{d+1}: X^{d+1} \rightarrow X$. This action induces an action of $\text{Aut}_{\mathcal{C}}(X) \wr \mathfrak{S}_{d+1}$ on the isomorphism classes of Latin hypercubes of dimension d over X , which for \mathcal{C} the category of sets is the standard notion of paratopism.

Remark 1.5. Let X be an object in a category \mathcal{C} with finite products and let d be a positive integer. The canonical projections $\pi_i^d: X^d \rightarrow X$ are split surjective, with section the diagonal morphism $\delta: X \rightarrow X^d$ defined as the unique morphism such that $\pi_i^d \circ \delta = \text{Id}_X$, for $1 \leq i \leq d$. Let $\lambda: L \rightarrow X^{d+1}$ be a Latin hypercube. The $d + 1$ components $\lambda_i = \pi_i^{d+1} \circ \lambda: L \rightarrow X$ of λ are split epimorphisms. Indeed, if we choose $s \neq i$, with $1 \leq i, s \leq d + 1$, then π_i^{d+1} factors through τ_s^{d+1} ; more precisely, $\pi_i^{d+1} \circ \lambda = \pi_j^d \circ \tau_s^{d+1} \circ \lambda$ where $j = i$ if $i < s$, and $j = i - 1$ if $i > s$. Since $\tau_s^{d+1} \circ \lambda$ is an isomorphism and π_j^d a split epimorphism, it follows that their composition is a split epimorphism, and hence so is $\pi_i^{d+1} \circ \lambda$.

2. Proof of Theorem 1.2

The endomorphism operad \mathcal{E} of a non-empty set X consists of the sets $\mathcal{E}(n) = \text{Map}(X^n, X)$ for any positive integer n , together with strutral maps

$$- \circ_i -: \text{Map}(X^n, X) \times \text{Map}(X^m, X) \rightarrow \text{Map}(X^{n+m-1}, X)$$

given by

$$(f \circ_i g)(x_1, x_2, \dots, x_{n+m-1}) = f(x_1, \dots, x_{i-1}, g(x_i, \dots, x_{i+m-1}), x_{i+m}, \dots, x_{n+m-1})$$

for all positive integers n, m, i , such that $1 \leq i \leq n$, all $x_1, x_2, \dots, x_{n+m-1} \in X$, and all maps $f \in \text{Map}(X^n, X)$ and $g \in \text{Map}(X^m, X)$. The sets $\mathcal{E}(n) = \text{Map}(X^n, X)$ are equipped with the action of \mathfrak{S}_n on the n coordinates of X^n , and the identity map $\text{Id}_X \in \mathcal{E}(1) = \text{Map}(X, X)$ is the unit element of this operad. For the

associativity properties of the maps $- \circ_i -$ and their compatibility with the symmetric group actions on the sets $\text{Map}(X^n, X)$, see for instance [6, Definition 1.2] or [7, Part II, §1.2]. The main step for the proof of Theorem 1.2 is the following Lemma.

Lemma 2.1. *Let X be a non-empty set, and let d, e, i be positive integers such that $1 \leq i \leq d$. Let $f \in \mathcal{L}(X^d, X)$ and $g \in \mathcal{L}(X^e, X)$. Then $f \circ_i g \in \mathcal{L}(X^{d+e-1}, X)$.*

Proof. In order to show that $f \circ_i g$ belongs to $\mathcal{L}(X^{d+e-1}, X)$ we need to show that for any $c \in X$ and an arbitrary choice of $d + e - 2$ of the $d + e - 1$ elements $x_1, x_2, \dots, x_{d+e-1} \in X$ the remaining of these elements is uniquely determined by the equation

$$f(x_1, \dots, x_{i-1}, g(x_i, \dots, x_{i+e-1}), x_{i+e}, \dots, x_{d+e-1}) = c. \tag{1}$$

Let s be an integer such that $1 \leq s \leq d + e - 1$. Fix elements $x_1, x_2, \dots, x_{s-1}, x_{s+1}, \dots, x_{d+e-1}, c \in X$.

Consider first that case where $s \leq i - 1$ or $s \geq i + e$. Then, setting $y = g(x_i, \dots, x_{i+e-1})$, the Equation 1 becomes

$$f(x_1, \dots, x_{i-1}, y, x_{i+e}, \dots, x_{d+e-1}) = c. \tag{2}$$

All entries but the entry x_s in this equation are fixed. Since $f \in \mathcal{L}(X^d, X)$ it follows that there is a unique choice for $x_s \in X$ such that Equation 2 holds, and hence a unique choice for $x_s \in X$ such that Equation 1 holds.

Consider the remaining case where $i \leq s \leq i + e - 1$. Then in particular the elements $x_1, \dots, x_{i-1}, x_{i+e}, \dots, x_{d+e-1}$ are fixed in X . Since $f \in \mathcal{L}(X^d, X)$ it follows that there is a unique $y \in X$ such that Equation 2 holds. Thus for Equation 1 to hold, we must have

$$g(x_i, \dots, x_{i+e-1}) = y. \tag{3}$$

In this equation all but x_s have been chosen. Since $g \in \mathcal{L}(X^e, X)$ it follows that there is a unique choice $x_s \in X$ such that Equation 3 holds. In all cases, there is a unique choice of x_s such that Equation 1 holds. This shows that $f \circ_i g$ belongs to $\mathcal{L}(X^{d+e-1}, X)$ and completes the proof. \square

Proof of Theorem 1.2. Let $d \in \mathbb{N}$. A map $f \in \mathcal{E}(d) = \text{Map}(X^d, X)$ belongs to $\mathcal{L}(d) = \mathcal{L}(X^d, X)$ if and only if the set

$$L = \{(x_1, x_2, \dots, x_d, f(x_1, x_2, \dots, x_d)) \mid x_1, x_2, \dots, x_d \in X\}$$

is a Latin hypercube in X^{d+1} . Equivalently, f belongs to $\mathcal{L}(d)$ if and only for every $c \in X$ and an arbitrary choice of $d - 1$ of the d entries $x_1, \dots, x_d \in X$ the remaining entry is uniquely determined by the equation $f(x_1, x_2, \dots, x_d) = c$. The action of \mathfrak{S}_d on $\text{Map}(X^d, X)$ by permuting the d coordinates of X^d clearly preserves the subset $\mathcal{L}(d)$, and Id_X belongs to $\mathcal{L}(1)$. In order to prove Theorem 1.2, it remains to show that the sets $\mathcal{L}(d)$ are closed under the operations $- \circ_i -$. This is done in Lemma 2.1 above, and this concludes the proof.

Remark 2.2. We could have very slightly simplified the proof of Lemma 2.1 by observing that thanks to the symmetric group actions on the coordinates of Latin hypercubes (cf. Remark 1.4) it would have been sufficient in the proof of Lemma 2.1 to consider the map $f \circ_d g$ and a single s in the distinction into the two cases for s . That is, it would have been sufficient in the last part of the proof of Lemma 2.1 to consider the cases where either $1 = s \leq d - 1$ or $s = d$. We will make use of this observation in the proof of the more general Theorem 3.4 below.

Remark 2.3. The proof of Theorem 1.2, as written, involves choices of elements in the set X . In Section 3 below we will rewrite this proof in such a way that it extends to categories with finite products.

3. On Latin hypercubes in Cartesian monoidal categories

In this section, we extend Theorem 1.2 to categories with finite products. Given an object X in a category \mathcal{C} with finite products, a positive integer n , and a morphism $\lambda:L \rightarrow X^n$ in \mathcal{C} , we denote as at the beginning by $\lambda_i = \pi_i^n \circ \lambda$ the composition of λ with the i -th canonical projection $\pi_i^n: X^n \rightarrow X$, where $1 \leq i \leq n$. The morphism λ is uniquely determined by the λ_i , and we will write abusively $\lambda = (\lambda_i)_{1 \leq i \leq n}$ whenever convenient. If $\sigma \in \mathfrak{S}_n$, then σ induces an automorphism $\hat{\sigma}$ on X^n given by $\hat{\sigma}_i = \pi_{\sigma^{-1}(i)}^n$; that is, $\hat{\sigma}$ permutes the coordinates of X^n . This yields a group homomorphism $\mathfrak{S}_n \rightarrow \text{Aut}_{\mathcal{C}}(X^n)$. We write ${}^\sigma \lambda = \hat{\sigma} \circ \lambda$. We extend the earlier notation $\mathcal{L}_{\mathcal{C}}(X^d, X)$ in the obvious way.

Definition 3.1. Let \mathcal{C} be a category with finite products. Let d be a positive integer. We denote by $\mathcal{L}_{\mathcal{C}}(X^d, X)$ the subset of $\text{Hom}_{\mathcal{C}}(X^d, X)$ consisting of all morphisms $f: X^d \rightarrow X$ such that the morphism $(\text{Id}_{X^d}, f): X^d \rightarrow X^d \times X = X^{d+1}$ is a Latin hypercube.

We first identify canonical representatives in isomorphism classes of Latin hypercubes.

Proposition 3.2. Let \mathcal{C} be a category with finite products, let X be an object in \mathcal{C} , and let d, s be positive integers such that $1 \leq s \leq d + 1$. Let $\lambda: L \rightarrow X^{d+1}$ be a Latin hypercube. Then there is a unique Latin hypercube $\iota: X^d \rightarrow X^{d+1}$ such that $\tau_s^{d+1} \circ \iota = \text{Id}_{X^d}$ and such that $\iota \circ \alpha = \lambda$ for some isomorphism $\alpha: L \rightarrow X^d$. In that case we have $\alpha = \tau_s^{d+1} \circ \lambda: L \rightarrow X^d$.

Proof. By the definition of Latin hypercubes, the morphism $\alpha = \tau_s^{d+1} \circ \lambda: L \rightarrow X^d$ is an isomorphism. Then setting $\iota = \lambda \circ \alpha^{-1}$ implies immediately that α determines an isomorphism between the Latin hypercubes $\lambda: L \rightarrow X^{d+1}$ and $\iota: X^d \rightarrow X^{d+1}$. We need to show that α and ι are unique subject to these properties. Let $\iota': X^d \rightarrow X^{d+1}$ a Latin hypercube and $\alpha': L \rightarrow X^d$ an isomorphism such that $\tau_s^{d+1} \circ \iota' = \text{Id}_{X^d}$ and such that $\iota' \circ \alpha' = \lambda$. Composing this equality with τ_s^{d+1} yields

$$\alpha' = \text{Id}_{X^d} \circ \alpha' = \tau_s^{d+1} \circ \iota' \circ \alpha' = \tau_s^{d+1} \circ \lambda = \alpha$$

This implies $\iota' = \lambda \circ \alpha^{-1} = \iota$, whence the uniqueness of ι and α as stated. The result follows. □

Applied with $s = d + 1$, Proposition 3.2 implies that – as earlier in the category of sets – the morphisms in $\mathcal{L}_{\mathcal{C}}(X^d, X)$ parametrise the isomorphism classes of d -dimensional Latin hypercubes over X .

Corollary 3.3. Let \mathcal{C} be a category with finite products, let X be an object in \mathcal{C} , and let d be a positive integer. Any Latin hypercube of dimension d over X is uniquely isomorphic to a Latin hypercube of the form $\iota: X^d \rightarrow X^{d+1}$ such that ι is a section of the morphism τ_{d+1}^{d+1} discarding the coordinate $d + 1$. Any such morphism ι is then uniquely determined by its last component $f = \iota_{d+1} = \pi_{d+1}^{d+1} \circ \iota: X^d \rightarrow X$. In particular, the set $\mathcal{L}_{\mathcal{C}}(X^d, X)$ parametrises the isomorphism classes of d -dimensional Latin hypercubes over X .

We show now that the sets $\mathcal{L}_{\mathcal{C}}(X^d, X)$ form an operad, together with the structural maps – \circ_i – defined as follows. Let $f \in \mathcal{L}_{\mathcal{C}}(X^d, X)$ and $g \in \mathcal{L}_{\mathcal{C}}(X^e, X)$, where d, e are positive integers. For $1 \leq i \leq d$, the structural map

$$- \circ_i -: \text{Hom}_{\mathcal{C}}(X^d, X) \times \text{Hom}_{\mathcal{C}}(X^e, X) \rightarrow \text{Hom}_{\mathcal{C}}(X^{d+e-1}, X)$$

sends (f, g) to the morphism

$$f \circ (\text{Id}_{X^{i-1}} \times g \times \text{Id}_{X^{d-i}})$$

where we identify $X^{i-1} \times X^e \times X^{d-i} = X^{d+e-1}$ for the domain of this morphism and where we identify $X^{i-1} \times X \times X^{d-i} = X^d$ for the codomain of $\text{Id}_{X^{i-1}} \times g \times \text{Id}_{X^{d-i}}$. One checks that if X is a set, this coincides with the earlier definition of $f \circ_i g$.

Theorem 3.4. *Let X be an object in a category \mathcal{C} with finite products. There is a sub-operad \mathcal{L} of the endomorphism set operad \mathcal{E} of X such that $\mathcal{L}(d) = \mathcal{L}_{\mathcal{C}}(X^d, X)$, for all $d \in \mathbb{N}$. In particular, for any positive integers d, e, i such that $1 \leq i \leq d$, and any morphisms $f \in \mathcal{L}_{\mathcal{C}}(X^d, X)$ and $g \in \mathcal{L}_{\mathcal{C}}(X^e, X)$ we have $f \circ_i g \in \mathcal{L}_{\mathcal{C}}(X^{d+e-1}, X)$.*

Proof. The proof amounts to rewriting the proof of Theorem 1.2, including the statement and proof of Lemma 2.1, in such a way that all steps remain valid for the Cartesian products in \mathcal{C} . In order to keep this readable, we mention in each step what this corresponds to in the case where X is a non-empty set (and by considering coordinates, one easily translates this to statements in \mathcal{C}).

Let $f \in \mathcal{L}_{\mathcal{C}}(X^d, X)$ and $g \in \mathcal{L}_{\mathcal{C}}(X^e, X)$, where d, e are positive integers. That is, the morphisms

$$(\text{Id}_{X^d}, f) : X^d \rightarrow X^{d+1},$$

$$(\text{Id}_{X^e}, g) : X^e \rightarrow X^{e+1}$$

are Latin hypercubes. As in the proof of Theorem 1.2, the unitality and symmetric group actions are obvious, and we only need to show, analogously to Lemma 2.1, that $(\text{Id}_{X^{d+e-1}}, f \circ_i g)$ is a Latin hypercube. That is, we need to show that for $1 \leq s \leq d + e - 1$ and $1 \leq i \leq d$, the composition $\tau_s^{d+e} \circ (\text{Id}_{X^{d+e-1}}, f \circ_i g)$ is an automorphism of X^{d+e-1} . As pointed out in Remark 2.2, since we may permute coordinates, it suffices to do this for $i = d$ and in the two cases where either $1 = s \leq d - 1$ or $s = d$.

We consider first the case $1 = s \leq d - 1$, so $d \geq 2$. We need to show that the morphism

$$\tau_1^{d+e} \circ (\text{Id}_{X^{d+e-1}}, f \circ_d g)$$

is an automorphism of X^{d+e-1} . If X is a set, then this automorphism is given by the assignment

$$(x_1, x_2, \dots, x_{d+e-1}) \mapsto (x_2, \dots, x_{d+e-1}, f(x_1, \dots, x_{d-1}, g(x_d, \dots, x_{d+e-1}))).$$

First, the morphism $\tau_1^d \circ (\text{Id}_{X^d}, f)$ is an automorphism of X^d because (Id_{X^d}, f) is a Latin hypercube. We note that if X is a non-empty set, then the morphism $\tau_1^d \circ (\text{Id}_{X^d}, f)$ is given by the assignment

$$(x_1, \dots, x_d) \mapsto (x_2, \dots, x_d, f(x_1, \dots, x_d)).$$

Compose this with the automorphism $\hat{\sigma}$ induced by the cyclic permutation $\sigma = (1, 2, \dots, d)$ on coordinates. The resulting automorphism

$$\hat{\sigma} \circ \tau_1^d \circ (\text{Id}_{X^d}, f)$$

is, for X a set, given by the assignment

$$(x_1, \dots, x_d) \mapsto (f(x_1, \dots, x_d), x_2, \dots, x_d)$$

The Cartesian product of this automorphism with $-\times \text{Id}_{X^{e-1}}$ yields an automorphism of X^{d+e-1} , which for X a set corresponds to

$$(x_1, \dots, x_{d+e-1}) \mapsto (f(x_1, \dots, x_d), x_2, \dots, x_{d+e-1})$$

Again permuting cyclically, the coordinates yields an automorphism of X^{d+e-1} which we will denote by α , which, if X is a set, corresponds to

$$(x_1, \dots, x_{d+e-1}) \mapsto (x_2, \dots, x_{d+e-1}, f(x_1, \dots, x_d)).$$

Using the fact that $\tau_1^e \circ (\text{Id}_{X^e}, g)$ is an automorphism of X^e , combined with cyclically permuting coordinates, we obtain an automorphism γ of X^e , which corresponds to the assignment

$$(x_d, \dots, x_{d+e-1}) \mapsto (g(x_d, \dots, x_{d+e-1}), x_{d+1}, \dots, x_{d+e-1}).$$

Then $\beta = \text{Id}_{X^{d-1}} \times \eta$ is the automorphism of X^{d+e-1} which corresponds to

$$(x_1, \dots, x_{d+e-1}) \mapsto (x_1, \dots, x_{d-1}, g(x_d, \dots, x_{d+e-1}), x_{d+1}, \dots, x_{d+e-1}).$$

Similarly, $\gamma = \text{Id}_{X^{d-1}} \times \eta \times \text{Id}_X$ is an automorphism of X^{d+e-1} . Now $\alpha \circ \beta$ is an automorphism of X^{d+e-1} corresponding to

$$(x_1, \dots, x_{d+e-1}) \mapsto (x_2, \dots, x_{d-1}, g(x_d, \dots, x_{d+e-1}), x_{d+1}, \dots, x_{d+e-1}, f(x_1, \dots, x_{d-1}, g(x_d, \dots, x_{d+e-1}))).$$

Thus the automorphism $\gamma^{-1} \circ \alpha \circ \beta$ of X^{d+e-1} coincides with the morphism $\tau_1^{d+e} \circ (\text{Id}_{X^{d+e-1}}, f \circ_d g)$. This proves the result in the case $1 = s \leq d - 1$.

Consider next the case $s = d$. We need to show that the morphism $\tau_d^{d+e} \circ (\text{Id}_{X^{d+e-1}}, f \circ_d g)$ is an automorphism of X^{d+e-1} . If X is a set, then this automorphism is given by the assignment

$$(x_1, \dots, x_{d+e-1}) \mapsto (x_1, \dots, x_{d-1}, x_{d+1}, \dots, x_{d+e-1}, f(x_1, \dots, x_{d-1}, g(x_d, \dots, x_{d+e-1}))).$$

As before, by using the automorphism $\tau_1^e \circ (\text{Id}_{X^e}, g)$, cyclically permuting the coordinates and then applying $\text{Id}_{X^{d-1}} \times -$ we obtain an automorphism δ of X^{d+e-1} , which for X a set corresponds to

$$(x_1, \dots, x_{d+e-1}) \mapsto (x_1, \dots, x_{d-1}, g(x_d, \dots, x_{d+e-1}), x_{d+1}, \dots, x_{d+e-1}).$$

Similarly, applying $- \times \text{Id}_{X^{e-1}}$ to the automorphism $\tau_d^d \circ (\text{Id}_{X^d}, f)$ yields an automorphism ϵ of X^{d+e-1} , which for X a set corresponds to

$$(x_1, \dots, x_{d+e-1}) \mapsto (x_1, \dots, x_{d-1}, f(x_1, \dots, x_d), x_{d+1}, \dots, x_{d+e-1}).$$

Thus $\epsilon \circ \delta$ is the automorphism of X^{d+e-1} which for X a set corresponds to

$$(x_1, \dots, x_{d+e-1}) \mapsto (x_1, \dots, x_{d-1}, x_{d+1}, \dots, x_{d+e-1}, (f \circ_d g)(x_1, \dots, x_{d+e-1})),$$

and this is indeed the automorphism $\tau_d^{d+e-1} \circ (\text{Id}_{X^{d+e-1}}, f \circ_d g)$. This proves the second case, and the result follows. □

4. On Latin hypercubes in closed Cartesian monoidal categories

A monoidal category \mathcal{C} with unit object $\mathbb{1}$ is closed if \mathcal{C} has an internal Hom, denoted **Hom**. That is, **Hom**: $\mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{C}$ is a bifunctor such that for any object X in \mathcal{C} the functor $X \times -$ on \mathcal{C} is left adjoint to the functor **Hom**($X, -$). This adjunction yields in particular natural bijections $\text{Hom}_{\mathcal{C}}(\mathbb{1}, \text{Hom}(X, Y)) \cong \text{Hom}_{\mathcal{C}}(X, Y)$ and natural isomorphisms $\text{Hom}(\mathbb{1}, X) \cong X$; see Kelly [3] and [5] for more background material. Following [4], endomorphism operads can be defined over objects in certain closed symmetric monoidal categories. For the definition of Latin hypercubes we need in addition that \mathcal{C} is Cartesian monoidal; that is, the monoidal product is a product in the category \mathcal{C} . In that case the unit object $\mathbb{1}$ is a terminal object in \mathcal{C} . In what follows we say that a morphism $\hat{\alpha}$ between Hom objects in \mathcal{C} lifts a map α if α is the image of $\hat{\alpha}$ under the functor $\text{Hom}_{\mathcal{C}}(\mathbb{1}, -)$ modulo canonical identifications.

In order to show that the morphism sets $\mathcal{L}_{\mathcal{C}}(X^d, X)$ lift to internal objects whenever \mathcal{C} has pullbacks, we will need, from [5, Exercise 5, page 213], the fact that automorphism groups of objects lift to internal objects. We have a pullback diagram

$$\begin{array}{ccc} \text{Aut}_{\mathcal{C}}(X) & \xrightarrow{\hspace{10em}} & \{\text{Id}_{\mathbb{1}}\} \\ \delta \downarrow & & \downarrow \\ \text{Hom}_{\mathcal{C}}(X, X) \times \text{Hom}_{\mathcal{C}}(X, X) & \xrightarrow{\hspace{2em} \mu \hspace{2em}} & \text{Hom}_{\mathcal{C}}(X, X) \times \text{Hom}_{\mathcal{C}}(X, X) \end{array}$$

where δ sends $\sigma \in \text{Aut}_{\mathcal{C}}(X)$ to (σ, σ^{-1}) , where $\mu(\alpha, \beta) = (\beta \circ \alpha, \alpha \circ \beta)$ for any $\alpha, \beta \in \text{Hom}_{\mathcal{C}}(X, X)$, and where the right vertical map sends $\text{Id}_{\mathbb{1}}$ to $(\text{Id}_X, \text{Id}_X)$. The lower horizontal map μ commutes with the involution on $\text{Hom}_{\mathcal{C}}(X, X) \times \text{Hom}_{\mathcal{C}}(X, X)$ given by exchanging coordinates. The map μ and the right vertical map lift to maps on internal Hom objects, and hence, if \mathcal{C} has pullbacks, then the above diagram lifts to a pullback diagram in \mathcal{C} of the form

4.1.

$$\begin{array}{ccc}
 \mathbf{Aut}(X) & \xrightarrow{\quad\quad\quad} & \mathbb{1} = \mathbb{1} \times \mathbb{1} \\
 (\gamma, \gamma') \downarrow & & \downarrow \iota \\
 \mathbf{Hom}(X, X) \times \mathbf{Hom}(X, X) & \xrightarrow{\quad\quad\quad \nu \quad\quad\quad} & \mathbf{Hom}(X, X) \times \mathbf{Hom}(X, X)
 \end{array}$$

Composing ι with the canonical involution on $\mathbf{Hom}(X, X) \times \mathbf{Hom}(X, X)$ commutes with ν , does not change ι , while it changes (γ, γ') to (γ', γ) . The universal property of pullbacks implies that there is a unique automorphism ϵ of the object $\mathbf{Aut}(X)$ of order 2 with the property that $(\gamma', \gamma) = (\gamma, \gamma') \circ \epsilon$. This automorphism lifts the bijection given by taking inverses in the group $\mathbf{Aut}_C(X)$. Since ι is trivially a monomorphism, it follows that (γ, γ') is a monomorphism. If $\mathbf{Hom}_C(\mathbb{1}, -)$ is faithful, hence reflects monomorphism, then both γ and γ' are monomorphisms, since they lift inclusion maps. The following result shows that there are internal objects lifting the morphism sets $\mathcal{L}_C(X^d, X)$.

Theorem 4.2. *Let \mathcal{C} be a Cartesian closed monoidal category with pullbacks. Let X be an object in \mathcal{C} , and let d be a positive integer. There is an object $\mathbf{L}(X^d, X)$ in \mathcal{C} , determined uniquely up to unique isomorphism, such that we have a canonical isomorphism $\mathbf{Hom}_C(\mathbb{1}, \mathbf{L}(X^d, X)) \cong \mathcal{L}_C(X^d, X)$, and such that there is a canonical morphism $\mathbf{L}(X^d, X) \rightarrow \mathbf{Hom}(X^d, X)$ in \mathcal{C} which lifts the inclusion $\mathcal{L}_C(X^d, X) \rightarrow \mathbf{Hom}_C(X^d, X)$. If in addition the functor $\mathbf{Hom}_C(\mathbb{1}, -)$ is faithful, then the canonical morphism $\mathbf{L}(X^d, X) \rightarrow \mathbf{Hom}(X^d, X)$ is a monomorphism.*

We will need the following characterisation of the morphism sets $\mathcal{L}_C(X^d, X)$ in a category with finite products.

Lemma 4.3. *Let \mathcal{C} be a category with finite products, let X be an object in \mathcal{C} , and let d be a positive integer. We have a pullback diagram of sets*

$$\begin{array}{ccc}
 \mathcal{L}_C(X^d, X) & \xrightarrow{\quad\quad\quad} & \prod_{i=1}^d \mathbf{Aut}_C(X^d) \\
 \downarrow & & \downarrow \iota \\
 \mathbf{Hom}_C(X^d, X) & \xrightarrow{\quad\quad\quad \gamma \quad\quad\quad} & \prod_{i=1}^d \mathbf{Hom}_C(X^d, X^d)
 \end{array}$$

where $\gamma = (\gamma_i: \mathbf{Hom}_C(X^d, X) \rightarrow \mathbf{Hom}_C(X^d, X^d))_{1 \leq i \leq d}$ is defined by

$$\gamma_i(\lambda) = \tau_i^{d+1} \circ (\mathbf{Id}_{X^d}, \lambda)$$

for $1 \leq i \leq d$ and $\lambda \in \mathbf{Hom}_C(X^d, X)$, and where ι is the product of d copies of the inclusion $\mathbf{Aut}_C(X^d) \rightarrow \mathbf{Hom}_C(X^d, X^d)$.

Proof. Let $\lambda \in \mathbf{Hom}_C(X^d, X)$. By definition, we have $\lambda \in \mathcal{L}_C(X^d, X)$ if and only if $\gamma_i \in \mathbf{Aut}_C(X^d)$ for $1 \leq i \leq d$. This is clearly equivalent to the assertion that the diagram in the statement is a pullback diagram. □

Proof of Theorem 4.2. As described in the diagram 4.1, applied with X^d instead of X , there is a morphism $\mathbf{Aut}(X^d) \rightarrow \mathbf{Hom}(X^d, X^d)$ which lifts the inclusion $\mathbf{Aut}_C(X^d) \rightarrow \mathbf{Hom}_C(X^d, X^d)$. Both maps γ and ι in the diagram from Lemma 4.3 lift to morphisms $\hat{\gamma}$ and $\hat{\iota}$ between the relevant internal Hom objects, and hence, by the assumptions on \mathcal{C} , there is a pullback diagram in \mathcal{C} of the form

4.4.

$$\begin{array}{ccc}
 \mathbf{L}(X^d, X) & \longrightarrow & \prod_{i=1}^d \mathbf{Aut}(X^d) \\
 \downarrow & & \downarrow \hat{\iota} \\
 \mathbf{Hom}(X^d, X) & \xrightarrow{\hat{\gamma}} & \prod_{i=1}^d \mathbf{Hom}(X^d, X^d)
 \end{array}$$

The functor $\text{Hom}_{\mathcal{C}}(\mathbb{1}, -)$ from \mathcal{C} to the category of sets preserves pullbacks, hence sends this pullback diagram to a diagram isomorphic to that in Lemma 4.3. It follows in particular that $\text{Hom}_{\mathcal{C}}(\mathbb{1}, \mathbf{L}(X^d, X)) \cong \mathcal{L}_{\mathcal{C}}(X^d, X)$. The uniqueness statement follows from the fact that pullbacks are unique up to unique isomorphism. If the functor $\text{Hom}_{\mathcal{C}}(\mathbb{1}, -)$ is faithful, then this functor reflects monomorphisms, whence the last statement follows. \square

Theorem 4.5. *Let \mathcal{C} be a Cartesian closed monoidal category with pullbacks. Suppose that the functor $\text{Hom}_{\mathcal{C}}(\mathbb{1}, -)$ is faithful. Suppose in addition that for any two objects Y, Z in \mathcal{C} and any morphism $\zeta: Z \rightarrow \mathbf{Hom}(Y, Y)$, if the map $\text{Hom}_{\mathcal{C}}(\mathbb{1}, \zeta)$ factors through the inclusion $\mathbf{Aut}_{\mathcal{C}}(Y) \rightarrow \mathbf{Hom}_{\mathcal{C}}(Y, Y)$, then the morphism ζ factors through the morphism $\mathbf{Aut}(Y) \rightarrow \mathbf{Hom}(Y, Y)$. Let X be an object in \mathcal{C} . For any positive integer d the morphism $\mathbf{L}(X^d, X) \rightarrow \mathbf{Hom}(X^d, X)$ is a monomorphism, and, with d running over \mathbb{N} , these monomorphisms form a sub-operad of the internal endomorphism operad of X in \mathcal{C} .*

Proof. Note that since we assume $\text{Hom}_{\mathcal{C}}(\mathbb{1}, -)$ to be faithful (hence reflecting monomorphisms), it follows that the morphism $\mathbf{L}(X^d, X) \rightarrow \mathbf{Hom}(X^d, X)$ from Theorem 4.2 is a monomorphism, for any positive integer d . Furthermore, as in the proofs of Theorems 1.2, 3.4, showing the unitality and compatibility with symmetric group actions is straightforward. What remains to be proved is that the maps $- \circ_i -$ of the endomorphism operad induce maps on the subobjects $\mathbf{L}(X^d, X)$ of the internal Hom objects $\mathbf{Hom}(X^d, X)$. Let d, e, i be positive integers such that $1 \leq i \leq d$. The map

$$- \circ_i -: \text{Hom}_{\mathcal{C}}(X^d, X) \times \text{Hom}_{\mathcal{C}}(X^e, X) \rightarrow \text{Hom}_{\mathcal{C}}(X^{d+e-1}, X)$$

sends (f, g) to $f \circ (\text{Id}_{X^{i-1}} \times g \times \text{Id}_{X^{d-i}})$. Since this involves composition and products only, this map lifts to a map of internal Hom objects

$$\mathbf{Hom}(X^d, X) \times \mathbf{Hom}(X^e, X) \rightarrow \mathbf{Hom}(X^{d+e-1}, X).$$

At the level of morphism sets it follows from Theorem 3.4 that we have a commutative diagram

4.6.

$$\begin{array}{ccc}
 \text{Hom}_{\mathcal{C}}(X^d, X) \times \text{Hom}_{\mathcal{C}}(X^e, X) & \xrightarrow{- \circ_i -} & \text{Hom}_{\mathcal{C}}(X^{d+e-1}, X) , \\
 \uparrow & & \uparrow \\
 \mathcal{L}_{\mathcal{C}}(X^d, X) \times \mathcal{L}_{\mathcal{C}}(X^e, X) & \longrightarrow & \mathcal{L}_{\mathcal{C}}(X^{d+e-1}, X)
 \end{array}$$

where the vertical maps are inclusions. We need to show that this diagram lifts to the internal Hom objects and relevant subobjects. We note that the vertical maps in the diagram 4.6 lift by Theorem 4.2, and the top horizontal map lifts by the discussion preceding the diagram 4.6. What we need to show is that the bottom horizontal map in diagram 4.6 lifts as well.

Since $\mathbf{L}(X^{d+e-1}, X)$ is defined via a pullback diagram 4.4 (with $d + e - 1$ instead of d), we need to show that there is a commutative diagram of the form

4.7.

$$\begin{array}{ccc}
 \mathbf{Hom}(X^d, X) \times \mathbf{Hom}(X^e, X) & \xrightarrow{-\circ_i-} & \mathbf{Hom}(X^{d+e-1}, X) \longrightarrow \prod_{i=1}^{d+e-1} \mathbf{Hom}(X^{d+e-1}, X^{d+e-1}) \\
 \uparrow & & \uparrow \\
 \mathbf{L}(X^d, X) \times \mathbf{L}(X^e, X) & \longrightarrow & \prod_{i=1}^{d+e-1} \mathbf{Aut}(X^{d+e-1})
 \end{array}$$

Combining Lemma 4.3 (with $d + e - 1$ instead of d) and diagram 4.6 yields a commutative diagram

$$\begin{array}{ccc}
 \mathbf{Hom}_C(X^d, X) \times \mathbf{Hom}_C(X^e, X) & \xrightarrow{-\circ_i-} & \mathbf{Hom}_C(X^{d+e-1}, X) \longrightarrow \prod_{i=1}^{d+e-1} \mathbf{Hom}_C(X^{d+e-1}, X^{d+e-1}) \\
 \uparrow & & \uparrow \\
 \mathcal{L}_C(X^d, X) \times \mathcal{L}_C(X^e, X) & \longrightarrow & \prod_{i=1}^{d+e-1} \mathbf{Aut}_C(X^{d+e-1})
 \end{array}$$

The top horizontal and two vertical maps in this diagram lift to maps as in the diagram 4.7. The hypothesis on lifting maps through morphisms of the form $\mathbf{Aut}(Y) \rightarrow \mathbf{Hom}(Y, Y)$ applied to the $d + e - 1$ components on the right side of the diagram 4.7 shows the existence of the lower horizontal map making the diagram 4.7 commutative. The uniqueness of such a map follows from the fact that the right vertical map is a monomorphism, where we use that the functor $\mathbf{Hom}_C(\mathbb{1}, -)$ is faithful. \square

Remark 4.8. We do not know whether the hypothesis on lifting morphisms $Z \rightarrow \mathbf{Hom}(Y, Y)$ through $\mathbf{Aut}(Y) \rightarrow \mathbf{Hom}(Y, Y)$ is indeed needed for Theorem 4.5 to hold. This hypothesis holds in the category of compactly generated topological spaces. It is easy to see that this hypothesis holds if $\mathbf{Aut}(Y) \rightarrow \mathbf{Hom}(Y, Y)$ is a regular monomorphism (these are monomorphisms which are an equaliser of a pair of parallel morphisms), assuming as before that $\mathbf{Hom}_C(\mathbb{1}, -)$ is faithful.

5. Latin hypercubes in terms of pullback diagrams and further remarks

Definition 1.1 describes Latin hypercubes of dimension d over a non-empty set X as subsets of X^{d+1} instead as the graph of a function $X^d \rightarrow X$. We describe the composition maps $- \circ_i -$ in terms of these subsets as pullbacks.

Proposition 5.1. *Let X be a non-empty set, and let d, e, i be positive integers such that $1 \leq i \leq d$. Let $L \subset X^{d+1}$ and $M \subseteq X^{e+1}$ be Latin hypercubes over X of dimension d and e , respectively. Let $f: X^d \rightarrow X$ and $g: X^e \rightarrow X$ be the maps whose graphs are L and M , respectively. Denote by $L \circ_i M \subseteq X^{d+e}$ the Latin hypercube over X of dimension $d + e - 1$ which is the graph of the map $f \circ_i g: X^{d+e-1} \rightarrow X$.*

An element $(x_1, x_2, \dots, x_{d+e}) \in X^{d+e}$ belongs to $L \circ_i M$ if and only if there is an element $z \in X$ such that $(x_1, \dots, x_{i-1}, z, x_{e+i}, \dots, x_{d+e}) \in L$ and such that $(x_i, \dots, x_{e+i-1}, z) \in M$. Then z is uniquely determined by the elements x_1, x_2, \dots, x_{d+e} . Equivalently, we have a pullback diagram of the form

$$\begin{array}{ccc}
 L \circ_i M & \xrightarrow{\beta} & M \\
 \alpha \downarrow & & \downarrow \mu_{e+1} \\
 L & \xrightarrow{\lambda_i} & X
 \end{array}$$

where λ_i is the restriction to L of the canonical projection $\pi_i^{d+1}: X^{d+1} \rightarrow X$ onto the coordinate i , where μ_{e+1} is the restriction to M of the canonical projection $\pi_{e+1}^{e+1}: X^{e+1} \rightarrow X$ onto the coordinate $e + 1$, and

where

$$\alpha(x_1, x_2, \dots, x_{d+e}) = (x_1, \dots, x_{i-1}, z, x_{e+i}, \dots, x_{d+e}),$$

$$\beta(x_1, x_2, \dots, x_{d+e}) = (x_i, \dots, x_{i+e-1}, z).$$

Proof. Note that $L \circ_i M$ is indeed a Latin hypercube by Theorem 1.2. We have

$$(x_1, x_2, \dots, x_{d+e}) \in L \circ_i M$$

if and only if

$$x_{d+e} = (f \circ_i g)(x_1, x_2, \dots, x_{d+e-1}) = f(x_1, \dots, x_{i-1}, g(x_i, \dots, x_{i+e-1}), x_{i+e}, \dots, x_{d+e-1}).$$

Since $z = g(x_i, \dots, x_{e+i-1})$ is the unique element in X such that $(x_i, \dots, x_{i+e-1}, z) \in M$ and the unique element such that $(x_1, \dots, x_{i-1}, z, x_{i+e}, \dots, x_{d+e}) \in L$, the first statement follows. The second statement follows from the fact that z is uniquely determined by the coordinates $x_i, 1 \leq i \leq d + e$ □

It is well known that for a Latin hypercube of dimension $d \geq 2$ over a non-empty set X , fixing one of the coordinates in X^{d+1} yields a Latin hypercube of dimension $d - 1$ (this was implicitly used in the proof of Theorem 1.2). This can be extended to Cartesian monoidal categories with pullbacks. We will need the following Lemma to show this.

Lemma 5.2. *Let \mathcal{C} be a Cartesian monoidal category. Let X be an object in \mathcal{C} and let d, s be positive integers such that $s \leq d$ and $d \geq 2$. Let $c: \mathbb{1} \rightarrow X$ be a morphism in \mathcal{C} . Denote by $\text{Id}_{X^d} \times c: X^d \rightarrow X^{d+1}$ the unique morphism satisfying $\tau_{d+1}^{d+1} \circ (\text{Id}_{X^d} \times c) = \text{Id}_{X^d}$ and $\pi_{d+1}^{d+1} \circ (\text{Id}_{X^d} \times c) = t \times c$, where t is the unique morphism $X^d \rightarrow \mathbb{1}$, and where we identify $X^d = X^d \times \mathbb{1}$ and $X = \mathbb{1} \times X$. The diagram*

$$\begin{array}{ccc} X^d & \xrightarrow{\tau_s^d} & X^{d-1} \\ \text{Id}_{X^d} \times c \downarrow & & \downarrow \text{Id}_{X^{d-1}} \times c \\ X^{d+1} & \xrightarrow{\tau_s^{d+1}} & X^d \end{array}$$

is a pullback diagram.

Proof. By permuting the coordinates it suffices to show this for $s = d$. Let Z be an object in \mathcal{C} , and let $u: Z \rightarrow X^{d+1}$ and $v: Z \rightarrow X^{d-1}$ be morphisms such that

$$\tau_d^{d+1} \circ u = (\text{Id}_{X^{d-1}} \circ c) \circ v.$$

We need to show that there is a unique morphism $w: Z \rightarrow X^d$ satisfying $u = (\text{Id}_{X^d} \times c) \circ w$ and $v = \tau_d^d \circ w$. By checking on coordinates one sees that $w = (v, u_d): Z \rightarrow X^{d-1} \times X = X^d$ is the unique morphism with this property, where as before $u_d = \pi_d^{d+1} \circ u$. □

Proposition 5.3. *Let \mathcal{C} be a Cartesian monoidal category with pullbacks. Let d be an integer such that $d \geq 2$. Let $\lambda: L \rightarrow X^{d+1}$ be a morphism in \mathcal{C} . Suppose that the morphism $\lambda: L \rightarrow X^{d+1}$ is a Latin hypercube. Then the morphism $\tau_{d+1}^{d+1} \circ \lambda: L \rightarrow X^d$ is an isomorphism, and for every morphism $c: \mathbb{1} \rightarrow X$ and every pullback diagram of the form*

$$\begin{array}{ccc}
 L_c & \xrightarrow{\lambda_c} & X^d \\
 \downarrow & & \downarrow (\text{Id}_{X^d}) \times c \\
 L & \xrightarrow{\lambda} & X^{d+1}
 \end{array}$$

the morphism $\lambda_c: L_c \rightarrow X^d$ is a Latin hypercube.

Proof. The morphism $\tau_{d+1}^{d+1} \circ \lambda$ is an isomorphism as part of the definition of a Latin hypercube. Let $c: \mathbb{1} \rightarrow X$ be a morphism, and let s be an integer such that $1 \leq s \leq d$. Consider a pullback diagram as in the statement. We need to show that $\tau_s^d \circ \lambda_c: L_c \rightarrow X^{d-1}$ is an isomorphism. Since $s \leq d$ and $d \geq 2$ we can complete the diagram in the statement to a commutative diagram

$$\begin{array}{ccccc}
 L_c & \xrightarrow{\lambda_c} & X^d & \xrightarrow{\tau_s^d} & X^{d-1} \\
 \downarrow & & \downarrow \text{Id}_{X^d} \times c & & \downarrow \text{Id}_{X^{d-1}} \times c \\
 L & \xrightarrow{\lambda} & X^{d+1} & \xrightarrow{\tau_s^{d+1}} & X^d
 \end{array}$$

The right square is a pullback diagram by Lemma 5.2. Since the left square is a pullback diagram, the pasting law for pullbacks implies that the outer rectangle is a pullback diagram as well. Since the lower horizontal morphism in the outer rectangle is an isomorphism, so is the upper horizontal morphism (we use here that pullbacks preserve isomorphisms). \square

We conclude this note with some further remarks.

Remark 5.4. The notion of transversals can be adapted to the situation of Definition 1.1 as follows. Given an object X in a category with finite products and a positive integer d , a transversal in X^{d+1} is a morphism $\sigma: X \rightarrow X^{d+1}$ such that $\sigma_i = \pi_i^{d+1} \circ \sigma$ is an automorphism of X , for $1 \leq i \leq d + 1$. The morphism σ is obviously a monomorphism. We say that such a transversal is contained in a Latin hypercube $\lambda: L \rightarrow X^{d+1}$ if there is a morphism $\iota: X \rightarrow L$ such that $\lambda \circ \iota = \sigma$. In that case, ι is necessarily a monomorphism as well. If X is a non-empty set, then a transversal σ can be identified with the subset $\sigma(X)$ of X^{d+1} . If X is finite and has an abelian group structure, and if $T \subseteq X^{d+1}$ is a transversal, then we have a version of the Δ -Lemma [9, Lemma 2.1] as follows. Denote by $\alpha: X^{d+1} \rightarrow X$ the alternating sum map $\alpha(x_1, x_2, \dots, x_{d+1}) = \sum_{i=1}^{d+1} (-1)^{i-1} x_i$ and by t the sum of all involutions in X . Since the sum of all elements in X is equal to the sum t of all involutions in X , an elementary calculation shows that

$$\sum_{x \in T} \alpha(x) = \begin{cases} 0 & \text{if } d \text{ is odd} \\ t & \text{if } d \text{ is even.} \end{cases}$$

If $X = \mathbb{Z}/n\mathbb{Z}$ for some positive integer n , then $t = 0$ if n is odd, and $t = \frac{n}{2} + n\mathbb{Z}$ if n is even.

Remark 5.5. Adapting another well-known notion for Latin squares going back to Bose [2], the graph of a Latin hypercube $L \subseteq X^{d+1}$ of dimension $d \geq 1$ over a non-empty set X is the simple graph $\Gamma(L)$ with vertex set L , with an edge between two elements in L if the two elements have $d - 1$ coordinates at which they coincide. Being a Latin hypercube implies that if one fixes $d - 1$ coordinates of an element in L , then the remaining two coordinates determine each other. In particular, if two elements of L coincide at d coordinates, then these two elements are equal. If $|X| = n$ is finite, then $\Gamma(L)$ has n^d vertices and valency $\binom{d+1}{2}(n - 1)$; indeed, the neighbours of $(x_1, x_2, \dots, x_{d+1}) \in L$ are obtained by first choosing a two-element subset $\{i_1, i_2\}$ of $\{1, 2, \dots, d + 1\}$ and then replacing x_{i_1} by any of the $n - 1$ values

different from x_{i_1} and replacing x_{i_2} by the unique element such that the resulting $d + 1$ -tuple belongs to L . Adjacent vertices have $n - 2 + 2(d - 1) = n + 2d - 4$ common neighbours; indeed, two distinct elements of the form $(x_1, x_2, x_3, \dots, x_{d+1})$, $(x'_1, x'_2, x_3, \dots, x_{d+1})$ in L have the $n - 2$ common neighbours of the form $(y_1, y_2, x_3, \dots, x_{d+1})$ with $y_1 \neq x_1, x'_1$ (so necessarily $y_2 \neq x_2, x'_2$), and in addition the $2(d - 1)$ common neighbours of the form $(x_1, x'_2, x_3, \dots, y_j, \dots, x_{d+1})$, $(x'_1, x_2, x_3, \dots, z_j, \dots, x_{d+1})$ in L where $3 \leq j \leq d + 1$ and y_j, z_j are the uniquely determined elements by the remaining coordinates. If $d \geq 3$, then the length of the shortest path between non-adjacent vertices depends on the number of common coordinates.

Remark 5.6. Latin hypercubes of a fixed dimension d over objects in a category \mathcal{C} with finite products form themselves a category. Let X, Y be objects in \mathcal{C} . Let $\lambda: X^d \rightarrow X^{d+1}$ and $\mu: Y^d \rightarrow Y^{d+1}$ be Latin hypercubes. A morphism from μ to λ is a morphism $\iota: Y \rightarrow X$ such that $\lambda \circ \iota^{\times d} = \iota^{\times(d+1)} \circ \mu$. Here $\iota^{\times d}: Y^d \rightarrow X^d$ is the morphism obtained by taking d times the product of ι ; similarly for $d + 1$. Note though that the notion of isomorphism in this category is different from the notion of isomorphism considered in the Introduction, where we regard Latin hypercubes of dimension d over X as objects in the under-category of X^{d+1} . For a Latin hypercube $\lambda: X^d \rightarrow X^{d+1}$ of the form $\lambda = (\text{Id}_{X^d}, f)$ for some morphism $f: X^d \rightarrow X$, one checks easily that an automorphism of this Latin hypercube in the category defined here is an automorphism $\iota \in \text{Aut}_{\mathcal{C}}(X)$ satisfying $\iota \circ f = f \circ \iota^{\times d}$.

Acknowledgements. The present paper was partially supported by EPSRC grant EP/X035328/1.

Competing interests. None.

References

- [1] R. A. Bailey, P. J. Cameron, C. E. Praeger and C. Schneider, The geometry of diagonal groups, *Trans. Amer. Math. Soc.* **375**(8) (2022), 5259–5311.
- [2] R. C. Bose, Strongly regular graphs, partial geometries and partially balanced designs, *Pacific J. Math* **13**(2) (1963), 389–419.
- [3] G. M. Kelly, *Basic concepts of enriched category theory*, Mathematical Society Lecture Notes Series **64**, (Cambridge University Press, 1982), 245.
- [4] G. M. Kelly, On the operads of J. P. May, *Repr. Theory Appl. Categ.* **13** (2005), 1–13.
- [5] S. Mac Lane and I. Moerdijk, *Sheaves in geometry and logic* (Springer-Verlag New York, 1992), xii+629.
- [6] P. May, The geometry of iterated loop spaces, *Lectures notes in mathematics*, vol. 271 (Springer-Verlag, 1972), viii+175.
- [7] M. Markl, S. Shnider and J. Stasheff, Operads in Algebra, topology, physics, *Math. Surveys and monographs*, vol. 96 (Amer. Math. Soc, 2002), x+349.
- [8] B. D. McKay and I. M. Wanless, A census of small Latin hypercubes, *SIAM J. Discrete Math.* **22**(2) (2008), 719–736.
- [9] I. M. Wanless, Transversals in Latin squares: a survey, in *Surveys in combinatorics 2011*, London Math. Soc. Lecture Note Ser., vol. 392 (Cambridge University Press, 2011), 403–437,