



City Research Online

City, University of London Institutional Repository

Citation: Ansari, R., Herman, R. & Cartwright, M. (2025). Designing a digital intervention for children with developmental language disorder: Aligning theory, research and clinical practice. *Digital Health*, 11, doi: 10.1177/20552076251324442

This is the published version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/35285/>

Link to published version: <https://doi.org/10.1177/20552076251324442>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Designing a digital intervention for children with developmental language disorder: Aligning theory, research and clinical practice

DIGITAL HEALTH
Volume 11: 1–22
© The Author(s) 2025
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/20552076251324442
journals.sagepub.com/home/dhj



Rafiah Ansari¹ , Ros Herman¹ and Martin Cartwright²

Abstract

Introduction: Developmental language disorder (DLD) is a neurodevelopmental condition characterised by persistent and pervasive challenges in daily communication. This paper describes the process of designing a digital word-learning intervention for this population that aligns with routine clinical practice, is evidence-based, and has strong theoretical rationale.

Method: The intervention design process involved research collaboration between academics, clinicians and software professionals.

The Medical Research Council's framework for developing complex interventions provided preliminary guidance; greater design depth was introduced by adding recommended elements from a synthesis of intervention design literature. The resulting intervention design was refined with focus group feedback from speech and language therapists, which underwent reflexive thematic analysis.

Results: Intervention design features were generated by aligning word-learning traits characteristic of the disorder with evidence-based vocabulary intervention strategies and clinician requirements for embedding digital practice in routine service-delivery.

Mock ups for a mobile intervention app were developed from the design features and shared for refinement with practicing speech and language therapists who had day-to-day experience of supporting children with DLD.

Conclusion: This paper outlines the process of defining the therapeutic components of a digital intervention, guided by empirical literature and clinician insights. These initial findings will serve as a foundation for co-designing with children with DLD to shape the final design, particularly in terms of motivation and engagement features.

Investing at the design stage in the determinants of successful translation of research into practice, increases the likelihood that the resulting intervention will demonstrate both research effectiveness and clinical uptake.

Keywords

Mobile interventions, digital clinical practice, complex intervention design, neurodevelopmental disorders, developmental language disorder, speech and language therapy, translational research

Submission date: 25 April 2024; Acceptance date: 11 February 2025

¹School of Health and Psychological Sciences, Department of Language and Communication Science, City St George's, University of London, London, UK

²School of Health and Psychological Sciences, Department of Health Services Research and Management, City St George's, University of London, London, UK

Corresponding author:

Rafiah Ansari, School of Health and Psychological Sciences, Department of Language and Communication Science, City St George's, University of London, London EC1V 0HB, UK.

Email: rafiah.ansari.2@citystgeorges.ac.uk



Introduction

Developmental language disorder (DLD) is a neurodevelopmental condition that results in significant, ongoing challenges in everyday communication.¹ It is estimated to affect 7% of children,² and is a heterogeneous disorder with nature and severity varying not only between children but also within a child influenced by internal physiological and psychological mechanisms and external social and environmental factors.^{3–5} DLD can occur in isolation or alongside other neurodevelopmental disorders with broad aetiology such as Attention–Deficit/Hyperactivity Disorder and Dyslexia.¹ Difficulty with word learning is characteristic of almost half of all children with DLD,⁶ and is predictive of limitations in wider areas of language, literacy, academic attainment, social skills, and general well-being.^{7–9}

Empirical literature suggests that children with DLD struggle to associate the social, perceptual, cognitive and/or linguistic signals that help match sounds to meaning during word learning.¹⁰ For example, a child with DLD who is shown a bottle of antibiotics and a stethoscope for the first time may struggle to differentiate the words when told one is worn by a doctor. This is because they must determine the answer by combining their knowledge of object properties with what they are able to see (i.e., only one can be ‘worn’). Difficulties processing signals for language acquisition may present as immature, as divergent or both.

The role of technology

The World Health Organisation reports significantly increased interest and adoption of digitally delivered health and care in many countries since widespread remote working measures were introduced during the COVID-19 pandemic.¹¹ In particular, this has led to greater exploration of interventions being introduced through mobile devices given their ubiquitous nature, convenience and affordability.^{12,13} With regards to word learning, user-friendly mobile devices present the opportunity for children with DLD to create and access multimedia content relating to a target word, i.e., through text, images, audio and video. It is hypothesised that this would support the cross-situational, multisensory cue integration required for optimum vocabulary acquisition.

There are some indications of the potential of digital vocabulary interventions in existing research involving school-aged children with DLD when technology-enhanced vocabulary interventions have been compared with traditional paper-based approaches. For example, one study¹⁴ targeted word learning by introducing relevant video content on mobile devices as well as through books, and reported significantly greater gains in the understanding, naming and defining of target words with the use of multimedia compared to reading alone. A limitation was that the children played a passive role due to the target words and video content being pre-selected by the researchers.

Another study¹⁵ introduced a custom-made digital aid which enabled children to independently access the sound of initial letters of target words as needed during word learning, alongside strategies introduced by a speech and language therapist (SLT). The study reported significantly greater gains in the accurate naming of words when compared to SLT input alone. Whilst this finding is promising, the technology was limited to providing audio cues only.

It must be noted that the response to digital stimulus during word learning may be dependent on the child’s presenting profile. One study which explored the use of images and videos to support word learning, reported a significant correlation between severity of language difficulties in children with DLD and the extent to which background music interfered with word learning.¹⁶ This indicates that any digital vocabulary intervention would need to be adaptable to individual language and learning needs.

In addition to the digital content of an effective vocabulary intervention, research should focus on current digital clinical practice to identify core design requirements. Here the needs of the intervention provider are fundamental for translating empirical efficacy into practice.^{17,18} The provider is responsible for interpreting research findings, planning how the results can enhance practice, and evaluating meaningful impact. Despite this pivotal role, a systematic review of digital interventions for children with communication difficulties identified that the requirements for the intervention provider, most commonly the SLT, to effectively deliver the intervention tended to be overlooked.¹⁹ To overcome this gap in the literature, much can be learned from the numerous profession-wide surveys of paediatric SLTs which have explored digital service delivery during and since the remote-working measures of the COVID-19 pandemic, including work by the primary author of this paper.^{20–24} A consistent theme from the surveys is that a key influencer of technology uptake is its consideration as routine practice across the profession. As a result, identifying positive drivers to the adoption of technology in care which are used consistently across paediatric SLT services and incorporating these within the intervention design is of great importance.

These service-level requirements can then be refined by sharing early-stage intervention prototype designs during in-depth focus groups with SLTs who specifically support word learning in children with DLD. This aligns with empirical recommendations that considering provider requirements which are both profession-wide and client-specific will lead to a more sustained and effective approach to the design, development and adoption of digital solutions.^{19,25}

Developing a complex intervention

Interventions for children with DLD are complex, not only because they must cater to the heterogeneity of the

population, but also due to the range of possible delivery models with variability of place, person, resources, and activities.²⁶

Intervention programmes are predominantly led by SLTs who deliver evidence-based therapy in partnership with parents, educators and psychologists. Given the pervading nature of communication, therapy may be provided in a range of settings such as the home, school or clinic. Words targeted by such interventions should be relevant to the child, such as their interests, school curriculum and family routine.²⁷

The UK Medical Research Council (MRC) provides guidance on the development and evaluation of complex interventions used in health and care services.^{28,29} The guidance is presented over four phases: development, feasibility/piloting, evaluation and implementation. For the development phase, the MRC framework advocates the use of evidence and theory to model the processes and outcomes that constitute a complex intervention.

Bleijenberg et al.³⁰ extended the MRC framework by adding intervention design guidance with the aim to improve clinical uptake. Recommended additional steps include clearly defining the presenting condition and its implications, determining the needs to be addressed by the proposed intervention, and understanding current practice (Figure 1). The resulting coordinated list of intervention requirements can guide features for the novel intervention. As highlighted by SLTs with experience of designing digital interventions for children with disabilities, this stage requires input from specialist software designers and developers to generate appropriate technical requirements.³¹ The final output of the design process is generally a proposed prototype of the intervention.³⁰

The research described in the current paper combined the MRC complex intervention development guidance^{28,29} with recommended elements from the intervention design literature as defined by Bleijenberg et al.³⁰ to generate required content for a digital word-learning intervention for children with DLD. Through collaboration with specialist software designers and developers, design mockups were then created which were used to produce an intervention app prototype. This was shared with paediatric SLTs actively supporting children who have DLD with word learning and the therapist feedback used to refine the intervention design.

In line with best practice guidance when developing digital therapy tools for children with additional needs,^{32–34} the intervention design will also incorporate specifications obtained from children with DLD. An integrative approach is planned whereby clinical- and evidence-based practice will define the therapeutic framework for the intervention as reported in this paper. These intervention mechanics will serve as a skeleton to which contributions from children with DLD around making the therapy motivating and engaging will be added. As such, work subsequent to this

paper will involve co-design sessions with children who have DLD to understand their views and preferences around word learning. This will provide necessary content to form a complete intervention design.

This paper therefore aims to report on two areas:

- Phase 1 – The design of a digital word-learning intervention informed by existing theory and research.
- Phase 2 – Clinician feedback on the novel intervention design and how this led to further refinement.

Method

The Guidance for Reporting for Intervention Development Studies in Health Research (GUIDED)³⁵ was followed to comprehensively describe the intervention design process and to ensure completeness of reporting (Appendix A).

Phase 1 – Intervention design

The MRC intervention development guidance^{28,29} and synthesised intervention design guidance³⁰ helped generate design ideas for a digital word-learning intervention for children with DLD. This phase was undertaken by the authors who in addition to being researchers were qualified paediatric SLTs (RA and RH) and a health psychologist (MC). The work was completed over a period of 18 months and involved desktop research and remote team calls to work through the steps below.

Profiling the disorder. Word learning traits common in children with DLD were identified and related to corresponding models of language learning theory. The profile of presentation was drawn from a multinational and multidisciplinary consensus study of criteria and terminology for children with language difficulties¹ and related literature.^{4,5,10,36–40} Contributors to the consensus included SLTs, educational psychologists, psychiatrists, paediatricians and specialist teachers.

Identifying the evidence. A systematic review of articles published between 1990 and 2023 reporting on vocabulary intervention outcomes for primary-school-aged children (5–11 years) with DLD was completed.⁴¹ The review identified intervention strategies with indications of effectiveness and considered implementation factors that may influence efficacy (e.g. nature of targeted words, dosage, provider and location).

Examining existing practice. A review of existing profession-wide surveys of paediatric SLT digital practice was conducted. The review considered national surveys conducted during the COVID-19 pandemic (2020–2021) to allow learning from the wide-scale adoption of technology during the pandemic.




Figure 1. Complex intervention design process by Bleijenberg et al. (p. 89, CC BY-NC-ND 4.0 DEED)³⁰ who extended the original Medical Research Council (MRC) Framework for developing complex interventions. Starred elements in the adaption by Bleijenberg et al.³⁰ refer to the original MRC Framework.²⁸

Findings from the surveys were synthesised and analysed using the Capability, Opportunity, Motivation and Behaviour (COM-B) model.⁴² This provided a theoretical framework to identify behaviour change drivers to support the digital intervention being adopted as part of standard practice.

Modelling the process and expected outcomes. Strategies to target specific areas of need and anticipated change processes were identified. These presumed causal pathways for the intervention were synthesised in a logic model to demonstrate the relationship between intervention activities for word learning and expected vocabulary outcomes. Contextual factors which could influence intervention response were also described.

Defining intervention features. This stage involved collaboration between the research team (authors of the paper: RA, RH, MC) and a private software company, to consider the digital intervention features required to achieve word-learning efficacy and intervention uptake as informed by existing empirical literature.

Specifications included the prompts, interactions and interfaces needed for the digital intervention to achieve optimum effectiveness. The design scope extended beyond technological requirements to identify intervention design principles such as accessibility and usability standards.

Phase 2 – Clinician feedback and refinement

In this phase, feedback on a word-learning intervention prototype created from design ideas generated in Phase 1 was obtained from the professionals who would implement the intervention in practice, paediatric SLTs. Reference was made to the Reflexive Thematic Analysis Reporting Guidelines⁴³ (RTARG) to provide conceptual and methodological coherence in the analysis and reporting of findings (Supplement 1).

As the work was necessarily exploratory, a focus group was conducted with results of interest being views that supported or opposed existing intervention features. In line with RTARG recommendation, rather than seeking consensus, an inductive interpretivist paradigm was adopted, which assumed multiple subjective and separately

constructed responses to the intervention prototype that would require interpretation.^{44,45} The positionality of the researchers within the fields of SLT and health psychology was balanced by having the lead software designer facilitate the running of the focus group and support data analysis in order to provide a technical lens. The goal was for the SLT feedback to be interpreted by the researchers and software designer using their professional knowledge and findings from existing literature to guide intervention refinement.

Target population and sample size. The participant eligibility criteria comprised qualified paediatric SLTs with a minimum of 6 months of experience, currently supporting word learning in school-aged children with DLD.

Based on established guidance for focus group research,^{46,47} this study aimed to recruit between 6 and 8 participants for the group in order to have enough people to generate varying opinions and perspectives and small enough to allow each individual to participate fully and be heard.

Consent and ethical considerations. Approval for this study was obtained from the School of Health and Psychological Sciences Research Ethics Committee, City St George's, University of London, UK (ETH2122-0815).

All participants received detailed information about the purpose of the study beforehand and gave written and verbal consent. Participants were informed that their participation in the study was voluntary and that all responses would be anonymised for confidentiality. They were advised of their right to withdraw from the study at any time up to analysis without any negative implications.

Recruitment. Purposeful sampling was used by approaching a school-based SLT service within one London borough. With the permission of the service manager, an email with a cover letter and information sheet was sent to relevant team members. Written consent was obtained from six paediatric SLTs actively supporting word learning in school-aged children with DLD who expressed an interest to participate in the study.

Procedure. A focus group discussion guide was used based on a template for novel intervention approaches developed and validated by Krueger and Casey^{46,47} who are considered experts in the field of focus group research (Supplement 2).

The focus group began with an introduction by the primary author (RA) who highlighted the purpose and aims of the research. This was followed by a demonstration of the digital word-learning intervention prototype by the lead software designer. After this, SLT feedback was gained using questions from the discussion guide read by the primary author.

The focus group session lasted approximately 90 min and was concluded when it was agreed by all six participants that relevant areas had been fully explored.

Setting. The focus group took place in a meeting room at the community base for the participating clinicians in Wandsworth, London. The rationale was guidance from Krueger and Casey⁴⁶ that familiar settings are likely to elicit more open discussions.

Data collection and analysis. Clinician views were audio recorded during the focus group and then later transcribed with identifying information removed. The transcription underwent reflexive thematic analysis using explicit semantic coding based on expert guidance by Braun and Clarke.⁴⁴ In keeping with the purpose of the research, i.e., to refine intervention design requirement, this involved seeking and coding text interpreted as relating to the strengths of the intervention design or areas requiring refinement. The codes were then collated into prominent themes to provide greater understanding of participant feedback.

Transcripts were coded by the primary author in consultation with the lead software designer. Themes were then developed and refined by the research team (authors of the paper: RA, RH, MC) and the lead software designer using direct quotations to capture the reflections and recommendations of the participants. The analysis process concluded when it was agreed by the research team that saturation had been reached, i.e., all relevant codes and themes had been generated from the data.

The findings guided targeted refinement of the intervention design as reported in the results section of this paper.

Results

Phase 1 – Intervention design

Profiling the disorder. A comprehensive description of the disorder helped define the multifaceted profile of vocabulary difficulties in children with DLD.

Details of presentation (Table 1, Column 1: DLD Profile) extended beyond types of word errors,¹ to include underlying processing mechanisms and environmental factors that may be influencing observed difficulties as this would impact the intervention approach. This included the effect of attention,¹ memory,^{1,36} literacy,³⁷ social responsiveness^{38,39}, and self-management,^{1,40} on word learning as well as the influence of the saliency, frequency and consistency of sound and meaning cues.^{1,10,39}

To guide intervention targets, theoretical models were identified which corresponded to the presenting profile and provided possible explanation for the characteristics described (Table 1, Column 2: Corresponding Theory^{3,48–62}).

Table 1. Defining the clinical population, evidence-based support strategies, and the intervention features to elicit desired changes.

DLD Profile	Corresponding Theory	Recommended Strategies	Empirical Evidence for Strategy	Intervention Design Features	Change Processes
Difficulty organising and retrieving word knowledge ¹	Connectionist model ⁴⁸	Visually organise and associate information during word learning – word web approach	Improved expressive vocabulary – correct use of words ^{63,64}	Word cues clustered visually into ‘webs’	Child accurately organises, stores and retrieves word information
Sound inaccuracies in word production ¹ . Underlying processing difficulties affecting sounds, syllables, rhymes ¹	Phonological processing deficit model ^{49,50}	Explicitly teach sound properties of target word (sounds, syllables, rhymes)	Improved expressive vocabulary ^{64,71,72} Improved awareness of word sounds ⁷²	Sound cues: initial sound, final sound, syllables and and rhyme	Child develops accurate representation and retrieval of the sound components of a word
Difficulty naming and defining word properties: function, location, category, description ¹	Semantic processing deficit model ^{51,52}	Explicitly teach target word meaning (function, location, category, description)	Improved expressive vocabulary ^{64,72} Improved receptive vocabulary – correct understanding of words ⁶⁶	Meaning cues: function, location, category, description	Child develops accurate representation and retrieval of the meaning components of a word
Difficulty relating word sound and word meaning ¹	Fast mapping theory ⁵³ Phonological-semantic interactional deficits model ⁵⁴	Target word sound and meaning simultaneously	Improved expressive vocabulary ^{63,68,69,72–74} Improved receptive vocabulary ⁶⁷ Improved awareness of word sounds ⁷²	Concurrent viewing and of sound and meaning cues.	Child connects word sound to word meaning during word learning
Weak short-term memory for sounds ¹	Phonological memory deficit model ⁵⁵	Record and store word sounds during word learning.	Improved expressive vocabulary ^{14,15} Improved receptive vocabulary ¹⁴	Creates audio recordings of word	Child improves their short-term retention of word sound for more accurate word learning
Relative strengths with visuospatial memory ³⁶	Visuo-linguistic memory interaction theory ⁵⁶	Visuospatial stimuli to reinforce verbal and auditory word learning.	Improved expressive vocabulary ^{14,16} Improved receptive vocabulary ¹⁴	Uses multimedia – text, audio, image, video	Child uses multisensory processing aligned to their individual learning profile

(continued)

Table 1. Continued.

DLD Profile	Corresponding Theory	Recommended Strategies	Empirical Evidence for Strategy	Intervention Design Features	Change Processes
Relative strength with reading words (orthography) ³⁷	Simple model of reading: linguistic comprehension theory ⁵⁷ Interactive connectionist model linking phonology, semantics, and orthography ⁵⁸	Opportunity to store and access typed letters and words during word learning	Improved expressive vocabulary ^{15,69}	Enables writing and/or typing of words.	Child uses letters and written cues to support word learning
Responsive to social cuing ^{38,39}	Social interactionist theory ⁵⁹	Adult to support word learning.	Improved expressive vocabulary ^{65,69,70}	Prompts for adult to facilitate child during word learning.	Opportunity for adult to support word learning
Heterogeneous learning profiles: unique presentations between children and within child over time ^{1,4,5}	Bronfenbrenner's bioecological systems theory ³ – influence of development, environment and context.	Word learning customised to child's profile and preferences	Improved expressive vocabulary ⁶⁴	Personalisation and customisation options	Child uses learning strategies aligned to individual learning profile and preferences
Reduced central executive functioning – coordinating word knowledge with emerging cues ^{1,40}	Executive function and self-regulation interaction theory ⁶⁰	Strategies for self-management.	Improved expressive vocabulary ^{78,79}	Child-friendly interface	Child able to self-direct their learning.
Reduced statistical learning: associating word cues through co-occurrences ^{1,10}	Cross-situational word-learning model ⁶¹ Retrieval-based learning: Episodic context account ⁶²	Multiple exposure to word cues across social and physical environments.	Improved long-term vocabulary recall ^{81,82}	Learning enabled across time, place, people with prompts to reinforce learning.	Child develops robust word representation by integrating existing and emerging multisensory word cues

Identifying the evidence. Existing vocabulary intervention studies involving children with DLD, as synthesised in a systematic review,⁴¹ helped collate recommended strategies for the proposed intervention.

The review⁴¹ identified twelve interventions delivered in English (UK,^{15, 63–68} US,^{14, 69–71}, New Zealand⁷²), two in Dutch (set in the Netherlands¹⁶), one in French (set in Switzerland⁷³) and one in German (set in Germany⁷⁴); providing collective data for 288 participants (167 males, 78 females, 43 unknown).

Most studies were of medium design quality, and this, coupled with the small number of studies identified, means that practitioners should be guided by response to treatment as well as the research findings when applying the intervention in practice.

Word-learning strategies with the greatest indication of effectiveness were extracted from the review and are reported along with the study source and the targeted area of difficulty (Table 1, Column 3: Empirical Strategies and Column 4: Empirical Evidence for Strategy).

One approach, known as word-web therapy, stood out in the systematic review due to being described as popular in clinical practice as well as demonstrating significant word-learning gains during clinical trials appraised as having good quality.^{63,64} Word-webs resemble an approach commonly known as mind maps⁷⁵ and involve using diagrams which arrange sound and meaning properties around a target word to visually represent networks of association.

A key feature of word webs which was also found to effectively improve word-learning in several other studies, was the simultaneous and explicit targeting of the sound and meaning properties of words.^{63,67–69,72–74} Additional word-learning gains were reported when contextual cues were provided in the form of written text, narratives and during social interactions.^{65,69,70} Effectiveness also improved when strategies aligned to the child's presenting profile of word-learning difficulties (sound errors, meaning errors, both) and corresponding needs.⁶⁴

Three of the reviewed papers used multimedia (audio, images and videos)^{14–16} to effectively support word learning. Two of the three multimedia intervention papers^{14,16} referred to Mayer's Cognitive Theory of Multimedia Learning⁷⁶ and corresponding instructional design principles to guide intervention development.⁷⁷ The design principles optimise cognitive processing during multimedia learning by (i) minimising extraneous processing demands such as background noise that may be distracting, (ii) facilitating intrinsic processing through tasks that are accessible and manageable, and (iii) optimising novel processing using cues to align acquired and existing knowledge.

An important finding from the systematic review was specificity in intervention response, gains generally did not transfer to non-targeted words^{15,63–65,73} and showed depreciation once therapy ended.^{15,64,65,68,70,73} This

finding guided requirements for developing the novel intervention, for example, 'self-management' was included as a recommended word-learning strategy to avoid ongoing dependency on therapists.^{78,79} The metalinguistic awareness required for this self-cuing is unexpected to develop before age seven,⁸⁰ implying that the novel intervention will be most appropriate for children of upper primary school (age 7–11 years). A further strategy was the opportunity for retrieval practice across social and physical environments to reduce depreciation of learning and improve long-term vocabulary recall.^{81,82}

The review summarised the factors which may affect intervention response, including relevancy of target words, dosage, provider and location. These have been recorded as influencing factors in the novel intervention design (see Moderating Factors in Figure 2).

Examining existing practice. A review of existing surveys identified five studies that collectively reported over fifteen hundred survey responses from paediatric SLTs regarding their views and experiences of digital practice during and since the extensive remote working measures introduced in response to the COVID-19 pandemic. The surveys spanned the UK ($n = 424$; $n = 438$),^{20,21} US ($n = 269$; $n = 259$),^{22,23} and Hong Kong ($n = 135$).²⁴

Where measured, the surveys consistently indicated an increase in clinician's perception of their frequency, convenience and confidence with digital practice following COVID-19 compared to before the pandemic.^{20–23} However, the surveys also indicated a range of challenges to digital uptake, as well as measures that would facilitate overcoming the obstacles to embedding the use of technology in routine care.

A series of requirements were identified to embed the mobile word-learning intervention into regular service delivery by paediatric SLTs (Table 2, Column 1: Requirements for Routine Digital Practice). These requirements were based on areas of consensus for factors facilitating routine digital practice as reported in the reviewed surveys. Examples included ensuring that the digital word-learning intervention enables involvement of families and partner professionals so that there is alignment with standard paediatric SLT practice.

The COM-B⁴² framework of behaviour change drivers was applied to organise the requirements by type: Capability – clinician skills and knowledge; Opportunity – convenience, suitability, accessibility; and Motivation – effectiveness, quality, continuity (Table 2, Column 2: COM-B Drivers for Behaviour Change in Clinicians). Corresponding strategies were then identified to facilitate the behaviour change required for the adoption of the digital word-learning intervention as part of regular service delivery (Table 2, Column 3: Recommended Strategies). For example, multi-device crossover functionality to ensure that the intervention can be accessed by carers and partner professionals in a range of settings.




Figure 2. Logic model for a digital word-learning intervention to support children with DLD.

Modelling the process and expected outcomes. Expected change processes were modelled to link each area of need with recommended strategies (Table 1, Column 5: Change Processes; Table 2, Column 4: Change Processes).

This informed a logic model to illustrate the relationship between the components of the intervention (Figure 2). The model defines the target group, intervention approach, intervention resources, expected outputs and desired outcomes. Change mechanisms within the model propose how and why the components of the intervention may work. Factors influencing change processes are listed to highlight contextual moderators.

Defining intervention features. This stage considered the digital design features required for intervention efficacy and clinician acceptability. Table 1 (Column 6: Intervention Design Features) charts how technology could create an interactive word-web tool which, through transition from adult-guiding to self-cuing, would support a child with mapping word sounds and meaning using

multimedia text, audio, image, and videos created and viewed across place, and time.

Table 2 (Column 5: Intervention Design Features) focuses on features for implementation of the intervention in routine practice, such as embedded guidance, interoperability, app performance, and alignment with care pathways. Best practice compliance requirements were generated from Shaban and Pearson's Learning Design Framework⁸⁴ which aligns Mayer's Cognitive Theory of Multimedia Learning,⁷⁶ as utilised by existing vocabulary intervention studies,^{14,16} with user-centred principles specifically for children with learning disabilities (Table 3).⁸³ The framework and its application to intervention design have been validated and evaluated empirically.^{84,85}

Phase 2 – Clinician feedback and refinement

Intervention prototype. Figure 3 presents mock-ups of the key intervention components created using a collaborative app prototype design tool known as Figma Design.⁸⁶

Table 2. Clinician requirements to support adoption of the mobile word-learning intervention in routine digital clinical practice.

Clinician Requirements for Adopting Digital Practice as Standard ^{20–24}	COM-B ^a Drivers for Behaviour Change in Clinicians ⁴²	Recommended Strategies	Intervention Design Features	Change Processes
Practitioner knowledge and expertise	Psychological capability: A person's mental functioning such as understanding and memory.	Access to guidelines and procedures.	User guidelines embedded in app	Clinician gains knowledge on how to implement intervention
Digital tools being fit for purpose	Physical Opportunity: Involving inanimate parts, e.g., financial and material resources	Appropriate software and hardware Multi-device crossover	App compatible with core device functions such as camera, microphone, and video recorder.	Clinician can utilise the technology
Digital solution enabling holistic care	Social opportunity: Involves other people and organisations (e.g., cultural and social norms)	Family involvement and multidisciplinary care	App functions across devices and platforms. App has secure sharing capabilities	Clinician can involve carers and partner professionals
Accessibility		Equitable service	App incorporates usability design features – see Table 3 ⁸³	Clinician can use app across their caseload
Evidence of effectiveness	Reflective motivation: Involves conscious thought processes, e.g., plans and evaluations	Effective	App displays child's progress	Clinician can monitor effectiveness of intervention
Quality and performance		Reliable	Streamlined app maintenance (updates, security, bugs etc.)	Clinician experiences consistency in app performance
Aligned with existing practice		Integrated with clinical practice	App design features to be reviewed and refined by practicing clinicians	Clinician experiences continuity of care
Enables personalised care		Choice and appropriateness	App enables customisation	Clinician able to customise app to meet individual care needs
Not disruptive	Automatic motivation: Involves habitual, instinctive, affective processes (desires and habits)	Part of routine/daily situations	App incorporates usability design features – see Table 3 ⁸³	Clinician's use of app does not create disruption or add burden to them or others
Un-burdensome		Ease of use		

^aCOM-B: Capability, Opportunity, Motivation – Behaviour change framework.⁴²

These were shared for feedback during a focus group with paediatric SLTs who routinely employ word-learning strategies to support children with DLD.

The mobile app is designed for SLTs to introduce to children with DLD when supporting word learning during

therapy. Therapy sessions at school are routinely attended by support staff such as teaching assistants who can then reinforce app use during the school day as and when word-learning support is needed. Similarly, therapists can model the app to parents who can support the child at

Table 3. Usability principles and guidelines when designing applications for children with learning disabilities from Shaban and Pearson's Learning Design Framework,^{83(pp4–6)} (with permission).

Principle	Guidelines
1. The learner:	
Place user in control	Allow the child to be in control of the application Ensure easy navigation throughout the application.
Motivate the learner	Use of affirmative language with familiar vocabulary. Graphics and aesthetics (e.g., colours, prompts, background, and illustrations) are relevant to the activities' goals.
2. The learning materials:	
Modality principle	App relies on spoken and visual activities. Content is attractive, simple, and meaningful for users.
Guided activity principle	Users are given essential information to carry out each activity. The goals of the app and each activity are clear.
Segmenting principle	Tasks consist of discrete activities. Each activity has its own level
Signalling principle	Most important elements are highlighted. Verbal and pictorial cues are used sparingly in activities.
Coherence principle	All activities are relevant to the app goals. A summary of the learner's achievement is provided for each activity.
3. The learning environment:	
Consistency	Elements are consistent throughout the app (e.g., characters, colours, backgrounds, buttons, icons, etc.).
Aesthetic and minimalist design	No distracting elements (e.g., movement, blinking, scrolling). Max of 3–4 choices (e.g., drop-down menu items). Contrast between text and background. Fonts are clear and readable. Icons are simple, labelled, and familiar for users.
Help	Users can access help at any time throughout the app. Users can skip instructions.

home. The aim is for the child to move from adult facilitation to independent app use so that they can self-cue during word learning.

In terms of technical features, the intervention is developed as a web-based app that is accessible across multiple platforms, browsers, and devices, but optimised for tablets, as children are more likely to access these both at school and at home. Since the app is delivered through a browser, it does not need to be downloaded or installed, saving storage space. Additionally, app maintenance - including updates, security management, troubleshooting, and bug fixes - can be managed in real time.

The prototype mock-ups were not intended to cover all design requirements generated in Phase 1, but rather to provide a scaffold from which therapist feedback, based on their experience supporting word learning in children with DLD, could guide further design iterations. Below is a summary of the features presented to the therapists:

- A word-learning intervention app prototype in the form of interactive word-webs, which create networks of sound and meaning cues for target words by uploading text, images, audio and videos in a structured and synchronised manner.
- On the home screen, children start by typing a personally chosen word and creating or uploading an image they associate with that word (Figure 3(a)). This leads to a screen of word cues: sound (first sound, last sound, syllables, rhymes) and meaning (category, adjectives, verbs, location). The child selects the cue they wish to explore (Figure 3(b)).
- The next screen allows the child to create, upload and replay multimedia content (text, image, audio, and video) related to the selected sound and meaning feature of the personally relevant word (Figure 3(c)). Content can be built up over time (Figure 3(d)). A summary screen shows the learning for each cue (Figure 3(e)).




Figure 3. (a–e) Intervention Mockup. From clockwise top left: (a) Home screen. (b) Word cues screen. (c) Content upload. (d) Range of uploaded content. (e) Summary of learning.

Table 4. Participant information (paediatric speech and language therapists).

Level of specialism	Experience supporting DLD
Generalist (<i>n</i> = 2)	6 months–1 year (<i>n</i> = 2)
Specialist (<i>n</i> = 3)	2–4 years (<i>n</i> = 2)
Highly specialist (<i>n</i> = 1)	5+ years (<i>n</i> = 2)

- The app is designed to enable children with DLD to build up a bank of meaningful words. It is organised logically so that it can be accessed, added to, and shared across different time points and environments.

Participant characteristics. Table 4 summarises key participant demographics and demonstrates that the paediatric SLTs provided a varied sample in terms of specialism and experience.

Summary of thematic analysis. The focus group data was categorised into three overarching sections in line with the purpose of the study, i.e., to inform the intervention: design strengths, design weaknesses, design suggestions. Within each category prominent themes were defined, and

these have been reported below with direct quotations to allow greater understanding of the clinician feedback.

Category 1: Design strengths. The therapist’s first impressions were positive. There was praise for the visual structure of the app (Theme 1), the use of features to enable customisation (Theme 2), and the opportunity to create and save resources more conveniently than with paper-based interventions (Theme 3).

Theme 1: A clear app layout is valued because it means easier navigation.

The therapists found that the logical, minimalist design helped make the app intuitive to use.

‘I like the graphics. I think it’s quite nicely laid out and quite easy to use’ (Participant 5).

‘What you need to click on and what you can fill is really clear’ (Participant 2).

The use of muted background colours was also seen as a positive as it was ‘not overwhelming or distracting’ (Participant 3).

Theme 2: Word-learning apps that are customisable to children’s needs and preferences are likely to be engaging.

The therapists welcomed that the app enabled children to add personalised content relating to a target word using their preferred modalities (audio, images, video).

‘They (the children) can easily capture what’s around them at school and just use’ (Participant 2).

'I think children would be more motivated because they love working with tablets. "Oh, look that's our voices" and that's probably more memorable for them. It's not the Teaching Assistant writing things down on a piece of paper. They're creating it and experiencing the learning' (Participant 6).

'When you look at the style of learning of the young people we work with, it is so much more visual, multi-sensory based, and what this does is allow you to access that multi-sensory experience really quickly and easily' (Participant 1).

Theme 3: Apps can make therapy easier compared to paper-based approaches.

The ways in which a mobile app enables more efficient and convenient implementation of word-learning activities compared to traditional paper approaches were discussed.

'The resources can be produced without going to too much effort with preparing. So, if you want action pictures, you have to go in and get pictures from the school ... whereas on a tablet you can just quickly take them and then talk about it' (Participant 2).

'It's good ... practically as well because you don't have to worry about printing resources in advance ... you've got it all on the iPad' (Participant 4).

'The Teachers can have it out whenever the child's doing their writing or they can go back to it if they've forgotten about the word so it's like a prompt ... always there if needed which makes things a lot easier' (Participant 3).

Category 2: Design weaknesses. Areas requiring improvement included greater clarity for the user (child and supporting adult) on the purpose of the app (Theme 1), better alignment to existing language learning approaches (Theme 2), and more comprehensive monitoring and reviewing of progress (Theme 3).

Theme 1: App prompts that are unclear and ambiguous can disengage the user.

For example, therapists felt that certain wording used to describe activities was open to interpretation.

'It is better if things are explicit, rather than expecting the adult or children to infer what they should do. So instead of "Write something here" if you want the child to type a word on the first page, put "Tap to type a word" as the prompt' (Participant 4).

In addition, an illustration to represent word cues was considered too abstract for the children to understand or relate to which could lead to disengagement.

'Pictures as cue icons I think are really good, but the image that's been chosen for hand clapping to represent syllables, it's quite abstract compared to photos of actual hands' (Participant 6).

Theme 2: Therapists are hesitant to use app design features that they do not recognise from established paper-based therapy approaches.

The app design presented to the therapists organised word sounds and meaning cues separately. Whilst the therapists felt that this provided visual balance and symmetry, they commented that it did not accurately represent therapeutic practice where cues circle the central target word.

'I think I anticipated looking like what our word learning usually look like. Obviously, it's a different format, but I wonder if our kids are maybe not going to be using the app all the time, if they're using paper word webs in class, then it'll be confusing' (Participant 3).

Therapists also felt that if the app intervention only focused on individual words rather than also considering the use of those words in sentences then this would restrict the strength of gain and benefits.

'I think using well in a sentence is often the biggest teller of whether they've understood the word' (Participant 5).

Theme 3: Intervention apps that do not effectively monitor progress miss an important part of the therapy process.

Therapists highlighted the influential role that progress markers play in therapy as they 'motivate the children by allowing them to track their achievements' (Participant 5).

Therapists also stated that being able to monitor progress is important for planning purposes and for providing updates to parents/carers and other professionals.

'I'll usually get vocabulary lists from teachers, and rate them with the children to determine what they've already looked at, what they already feel confident with, then decide which we're going to be looking at together. I then have a bank of words to which I add ratings and progress to share with the children and with teachers' (Participant 3).

In addition, a mechanism for reviewing and reinforcing gains was called for.

'I think being able to revise the words is key. I really liked the ways you can explore words, but I worry that once you walk away from it, you weren't prompted to return' (Participant 1).

A need for a range of summaries of learning was also commented on. Whilst a summary screen showing the complete word web for a particular word was available, the app extended to all words targeted by an individual child and across all children on a certain therapist's caseload or attending a certain school.

Category 3: Design suggestions. A range of features were proposed to enhance the app. The suggestions aimed to better explain to users what the app is for (Theme 1), to motivate the child (Theme 2), and to assist professionals with caseload management (Theme 3):

Theme 1: Intervention apps should clearly define their purpose and intended use.

The use of prompts to highlight key features that were both written and verbal using a text-to-speech feature was advised. This is integrated into many familiar technologies and enables text to be converted into speech sounds imitative of the human voice.

There were calls for embedded tutorials to clarify app purpose and use.

‘An explanation for everybody who uses the app would be really helpful ... reduce confusion and inconsistency’ (Participant 3).

Theme 2: Apps designed for children need embedded reward systems based on activities and stages completed to motivate and show achievement.

The use of games, trophies and more child-centred features were suggested.

‘They’re so used to video games; they might want to see some of what they enjoy already to make them want to use this’ (Participant 2).

However, the importance of a balance between engaging and distracting was discussed.

‘Just thinking about all the stuff kids like to play with online. How can we add some of the motivators but without it being confusing or too much’ (Participant 6).

Theme 3: Intervention apps should have features to assist caseload management by enabling professionals to record and share progress.

Therapists suggested embedding a vocabulary measure within the app which could be completed at baseline and repeated as required with the date stamped.

Summary screens and dashboards were requested to demonstrate progress made at the level of caseloads, schools and individual children. These should be easy to share with parents/carers and other professionals both electronically and through print outs.

‘Because we’re in multiple schools we need more just one page with all the users. Instead, perhaps having per school, so opening a school folder and then having all the users that would be easier to navigate or filtering all users by school’ (Participant 4).

Design refinements following clinician feedback

Findings from the clinician feedback have led to the following updates which are summarised visually in Figures 4–6:

App prompts

- Onboarding tutorials for parents and professionals have been added (Figure 4(a): Background information, Figure 4(b): Step-by-step guide)
- Text-to-speech features have been added enabling the children to click and hear written words.
- In-app prompts (images and text) have been updated to be more explicit and understandable (for example Figure 5(a)).

Monitoring progress

- An empirically advocated baseline vocabulary assessment which can be repeated at any time has been included (Figure 5(b)).^{87–89}
- A therapist dashboard has been added which includes summary information per child (Figures 6(a)–(b)).

Therapy approach

- The word-learning cues have been arranged in a circle around the target word to align with the paper-based word web format commonly used in current practice (Figure 5(c)).⁶⁴
- Prompts to practice the target word in sentences have been added.

Child engagement

- Achievement pop-ups have been added which appear following completion of activities (Figure 5(d)). Design ideas for further engagement features will be sought through separate co-design sessions with children who have DLD as described in the Introduction section of this paper.

Discussion

This paper describes how the design of a digital word-learning intervention for children with DLD has been informed by empirical evidence and practitioner experience.

Interventions for this population are complex due to diversity in presentation and variability of intervention contexts.²⁶ This creates a tension in intervention design research between fidelity of replication and the need to tailor interventions to be sensitive to different delivery models.

To address this challenge, the MRC complex intervention development framework^{28,29} and complimentary design guidelines³⁰ were adopted to create a digital word-learning intervention based on theory, research, and existing surveys of professional practice. The empirically informed intervention design was then refined using feedback from therapists actively supports word learning in children with DLD. This approach has led to an intervention design that combines robust research literature with the day-to-day implementation experience of therapists as discussed further below.

A customised intervention

DLD represents a heterogeneous condition with presentation influenced by multiple biopsychosocial factors.^{4,5} To account for this, the intervention has been designed to align strategies according to types of word-learning difficulties (sound errors, meaning errors, both) and preferred learning styles. Practitioners will therefore be able to personalise the therapy according to the individual needs of the children being supported. This feature makes it unique from digital word-learning interventions in existing studies^{14–16} and ensures agreement with best practice guidance for DLD vocabulary interventions^{48,64} and mobile health and care applications.^{12,13}




Figure 4. (a and b) Updated intervention mock-up: onboarding tutorials for parents and professionals. From top down: (a) Background information. (b) Step-by-step guide.

Evidence-based design

Word-learning strategies for the intervention were derived from a systematic review of intervention efficacy studies and related work involving children with DLD.⁴¹ This ensured that the design was grounded in research evidence.

However, as reported by Utianski et al.,⁹⁰ practicing SLTs consistently cite lack of contextual considerations as a barrier to translating findings from field studies to clinical care. To address this, factors influencing the implementation of evidence-based word-learning strategies such as location, provider, intensity, and opportunities for reinforcement, were considered as part of the intervention design. This replicates the inclusion of contextual moderators when defining evidence-based strategies for novel interventions that have been reported, albeit sparingly, in the existing literature. For example, McKean et al.⁹¹ worked with paediatric SLTs and

health visitors to develop a hybrid face-to-face and digital intervention to promote children's language development within the home environment. Evidence-based techniques were selected for the intervention by carefully considering which strategies from systematic and scoping reviews could be applied routinely by caregivers. This was done to increase the likelihood of intervention uptake.

Routine service delivery

To ensure relevancy to the clinical landscape in which the intervention would be delivered, the surveyed experiences of paediatric SLTs following the rapid, widespread adoption of technology in response to COVID-19 remote working measures were considered.

A unique research perspective emerged from these existing surveys, as therapists were able to reflect on their




Figure 5. (a-d) Updated intervention mockup. From clockwise top left: (a) Home screen. (b) Word test. (c) Arrangement of word cues. (d) Achievement pop-up.

experience of digital practice at scale as opposed to following specific clinical trials which would have generally been the norm for digital intervention research previously. For example, therapists extended their evaluation of digital solutions beyond individual episodes of care to broader implications such as barriers to extending digital care models to families and partner professionals.

These wider requirements helped identify design features for the digital word-learning intervention to not only support initial uptake amongst clinicians but also effective and sustained adoption within complex care pathways. This is important as estimates indicate that less than half of clinical innovations ever make it into standard service delivery despite demonstrating research effectiveness.⁹²

Mapping change processes

In keeping with the complexity of the population and their needs, several theoretical models and frameworks informed the change mechanisms underpinning the intervention design.

For example, rather than referring to a single language learning model, a range of language processing theories were selected which correspond to the range of word finding needs and presentation observed in children with DLD. This is in keeping with current empirical thinking around multiple, interacting etiological pathways influencing vocabulary development in children with DLD.^{64,93,94}

In addition, a behaviour change framework was applied to the surveys of professional digital practice to understand the capability, opportunity and motivation that paediatric SLTs require to embed technology within routine service

delivery. This approach replicates work by Stringer et al.⁹⁵ who used a behaviour change framework to understand the needs of paediatric SLTs in order to transform their regular service delivery model. In that case, the goal was the introduction of collaborative interventions with teachers rather than adoption of technology in care. With both studies, a list of intervention requirements were identified which were anticipated to facilitate change in routine practitioner practice.

The theoretical assumption of how and why the intervention would be effective were synthesised to inform a logic model of intervention processes which coordinated the therapeutic content, the recipients, the context, the outputs and expected outcomes. Existing digital therapeutic research has demonstrated that this structured approach to modelling intervention components makes it easier to identify potential gaps, inconsistencies, and areas of improvement during later evaluation and implementation stages.^{96,97}

Bridging the gap between clinicians and software professionals.

This study uniquely combined clinical and technical thinking to generate app design features which met word-learning efficacy requirements. As explored by informatics researchers Du and Tekinbas,³¹ growth in technology adoption by SLTs has generally not been matched by increased collaboration with software experts. This has resulted in therapists often attempting and failing to adapt mainstream applications for use during specialist interventions.⁹⁸ A gap in shared epistemology between clinical and computer sciences has been proposed as a barrier to




Figure 6. (a–b) Updated intervention mockup. From top down: (a) Therapist dashboard. (b) Summary per child.

jointly designing digital interventions.³¹ The research in this paper was able to overcome this issue by synthesising frameworks and guidance from both fields as well as ensuring that clinical academics and software professionals worked together to interpret data relating to intervention design features.

Clinician feedback. Gaining the views of therapists who regularly implement word-learning interventions whilst still at the ideation stage proved to be a valuable and informative step. Areas not yet considered were added to the design, increasing the likelihood of adoption in practice. These design amendments were made easily without

substantial time or resource implications, which is significant as lengthy and expensive delays in the journey from research to practice has been attributed partly to inefficient research methodologies.^{90,99} Furthermore, in research that has included SLTs among other health care workers, clinician contribution to digital intervention design has been identified as an influencing factor for care providers when selecting a preferred solution.¹⁰⁰

Limitations

As the intervention design is heavily informed by the evidence on current interventions and practice, the process

inevitably suffers from the same limitations as these sources of information. With regards to vocabulary intervention effects, the greatest weaknesses identified in the systematic review were limitations with the quantity and quality of studies.⁴¹ As such, the evidence of efficacy should be treated with caution and intervention outcomes monitored closely during delivery. Similarly, the findings from the surveys of clinical digital practice which have guided the intervention design, provide a snapshot of service delivery by a sample of the profession at particular points in time.

Another important consideration is that the work in this paper provides partial design features for the final intervention. The therapeutic elements will be further shaped and enhanced by the design views of children with DLD. End-user contributions will predominantly guide motivational and engagement aspects such as the design of child-friendly app prompts, symbols for word-learning cues, and progress markers. As discussed by researchers who have undertaken similar work designing interventions for neurodiverse children, the merging of empirical, clinical and end-user perspectives is complex as it requires careful orchestration between different disciplines, generations and contributions.³³ Equality of influence may not always be achieved, and in previous studies, it is often the views of the children that are marginalised.^{32,33} As an alternative to therapy-driven approaches when creating digital products to facilitate neurodiverse children, some researchers have called for experience-driven design where the child's preferences drive technical specifications.¹⁰¹ Given the dearth of research in the intervention design space, there is learning to be gained from having a range of methodologies and as the field becomes more established future work could focus on comparing approaches.

Conclusion and next steps

This paper describes how design ideas for a digital word-learning intervention were generated from theory, research and practice through collaboration between academics, clinicians and software professionals.

A subsequent study will identify design requirements generated by children with DLD to ensure that the resulting intervention is motivational and engaging. This will be followed by usability testing as a precursor to clinical evaluation trials.

These research stages will create new knowledge to iteratively influence the design, development, evaluation and adoption in routine practice of the digital word-learning intervention.

Contributorship: RA researched literature and conceived the study. RA, RH and MC were involved in gaining ethical approval, and data analysis. RA wrote the first draft of the manuscript. All authors reviewed and edited the manuscript and approved the final version of the manuscript.

A private software company, Big Lemon[®], provided technical expertise.

Declaration of conflicting interests: The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Ethical approval: Approval for this study was obtained from the School of Health and Psychological Sciences Research Ethics Committee, City St George's, University of London, UK (ETH2122-0815).

Funding: The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The primary author, RA (Clinical Doctoral Research Fellow, ICA-CDRF-2018-04-ST2-029), was funded by the Health Education England/National Institute for Health Research (NIHR) for this research project. The views expressed in this publication are those of the authors and not necessarily those of the NIHR, NHS or the UK Department of Health and Social Care.

Guarantor: RH.

ORCID iD: Rafiah Ansari  <https://orcid.org/0000-0003-4163-1100>

Supplemental material: Supplemental material for this article is available online.

References

1. Bishop DV, Snowling MJ, Thompson PA, et al. Phase 2 of CATALISE: a multinational and multidisciplinary Delphi consensus study of problems with language development: terminology. *J Child Psychol Psychiatry* 2017; 58: 1068–1080.
2. Norbury CF, Gooch D, Wray C, et al. The impact of non-verbal ability on prevalence and clinical presentation of language disorder: evidence from a population study. *J Child Psychol Psychiatry* 2016; 57: 1247–1257.
3. Bronfenbrenner U and Morris P. The bioecological model of human development. In: Lerner RM and Damon W (eds) *Handbook of child psychology: theoretical models of human development*. 6th ed. New York: John Wiley & Sons, 2006, pp.793–828.
4. Law J, Reilly S and McKean C. (eds). *Language development: Individual differences in a social context*. Cambridge: Cambridge University Press, 2022.
5. Boerma T, Ter Haar S, Ganga R, et al. What risk factors for developmental language disorder can tell us about the neurobiological mechanisms of language development. *Neurosci Biobehav Rev* 2023; 154: 105398.
6. Rice ML and Hoffman L. Predicting vocabulary growth in children with and without specific language impairment: a longitudinal study from 2; 6 to 21 years of age. *J Speech Lang Hear Res* 2015; 58: 345–359.

7. Bleses D, Makransky G, Dale PS, et al. Early productive vocabulary predicts academic achievement 10 years later. *Appl Psycholinguist* 2016; 37: 1461–1476.
8. Girard LC, Pingault JB, Doyle O, et al. Expressive language and prosocial behaviour in early childhood: Longitudinal associations in the UK Millennium Cohort Study. *Eur J Dev Psychol* 2017 Jul 4; 14: 381–398.
9. Oslund EL, Clemens NH, Simmons DC, et al. The direct and indirect effects of word reading and vocabulary on adolescents' Reading comprehension: comparing struggling and adequate comprehenders. *Read Writ* 2018; 31: 355–379.
10. McGregor KK, Smolak E, Jones M, et al. What children with developmental language disorder teach us about cross-situational word learning. *Cogn Sci* 2022; 46: e13094.
11. World Health Organization. COVID-19 and digital health: what can digital health offer for COVID-19. WHO, Geneva. Available online: www.who.int/china/news/feature-stories/detail/Covid-19-and-digital-health-what-can-digital-health-offer-for-Covid-19 (accessed 13 September 2022). 2020.
12. Brahmhatt DH, Ross HJ and Moayed Y. Digital technology application for improved responses to health care challenges: lessons learned from COVID-19. *Can J Cardiol* 2022; 38: 279–291.
13. Jat AS, Grønli TM and Ghinea G. Navigating Design Science Research in mHealth Applications: A Guide to Best Practices. *IEEE Trans Eng Manage* 2024; 71: 14472–1448.
14. Lowman JJ and Dressler EV. Effects of explicit vocabulary videos delivered through iPods on students with language impairments. *J Spec Educ Technol* 2016; 31: 195–206.
15. Best W. Investigation of a new intervention for children with word-finding problems. *Int J Lang Commun Disord* 2005; 40: 279–318.
16. Smeets DJ, Van Dijken MJ and Bus AG. Using electronic storybooks to support word learning in children with severe language impairments. *J Learn Disabil* 2014; 47: 435–449.
17. Damschroder LJ, Aron DC, Keith RE, et al. Fostering implementation of health services research findings into practice: a consolidated framework for advancing implementation science. *Implement Sci* 2009; 4: 1–5.
18. Turner KM, Rousseau N, Croot L, et al. Understanding successful development of complex health and healthcare interventions and its drivers from the perspective of developers and wider stakeholders: an international qualitative interview study. *BMJ Open* 2019; 9: e028756.
19. Law J, Dornstauder M, Charlton J, et al. Tele-practice for children and young people with communication disabilities: employing the COM-B model to review the intervention literature and inform guidance for practitioners. *Int J Lang Commun Disord* 2021; 56: 415–434.
20. Ansari R, Loraine E and Gréaux M. Impact of COVID-19 on digital practice in UK paediatric speech and language therapy and implications for the future: a national survey. *Int J Lang Commun Disord* 2022; 57: 1112–1129.
21. Charlton J, Gréaux M, Kulkarni A, et al. UK Paediatric speech and language therapists' perceptions on the use of telehealth in current and future clinical practice: an application of the APEASE criteria. *Int J Lang Commun Disord* 2024; 59: 1163–1179.
22. Campbell DR and Goldstein H. Genesis of a new generation of telepractitioners: the COVID-19 pandemic and pediatric speech-language pathology services. *Am J Speech Lang Pathol* 2021; 30: 2143–2154.
23. Campbell DR and Goldstein H. Evolution of telehealth technology, evaluations, and therapy: effects of the COVID-19 pandemic on pediatric speech-language pathology services. *Am J Speech Lang Pathol* 2022; 31: 271–286.
24. Fong R, Tsai CF and Yiu OY. The implementation of telepractice in speech language pathology in Hong Kong during the COVID-19 pandemic. *Telemed e-Health* 2021; 27: 30–38.
25. Greenhalgh T, Wherton J, Papoutsis C, et al. Beyond adoption: a new framework for theorizing and evaluating non-adoption, abandonment, and challenges to the scale-up, spread, and sustainability of health and care technologies. *J Med Internet Res* 2017 Nov 1; 19(11): e8775.
26. Law J, McKean C, Murphy CA, et al. (eds). *Managing children with developmental language disorder: Theory and practice across Europe and beyond*. London: Routledge, 2019.
27. Dennis J, Law J and Charlton J. Speech and language therapy interventions for children with primary speech and/or language disorders:(Protocol). *Cochrane Database of Systematic Reviews* 2017(1): 1–21.
28. Craig P, Dieppe P, Macintyre S, et al. Developing and evaluating complex interventions: the new Medical Research Council guidance. *Br Med J* 2008; 337: a1655.
29. Skivington K, Matthews L, Simpson SA, et al. A new framework for developing and evaluating complex interventions: update of Medical Research Council guidance. *Br Med J* 2021; 374: n2061.
30. Bleijenbergh N, de Man-van Ginkel JM, Trappenburg JC, et al. Increasing value and reducing waste by optimizing the development of complex interventions: enriching the development phase of the medical research council (MRC) framework. *Int J Nurs Stud* 2018; 79: 86–93.
31. Du Y and Tekinbas KS. Bridging the gap in mobile interaction design for children with disabilities: Perspectives from a pediatric speech language pathologist. *Int J Child Comput Interact* 2020; 23–24; 100152.
32. Malinverni L, Mora-Guiard J, Padillo V, et al. An inclusive design approach for developing video games for children with Autism Spectrum Disorder. *Comput Human Behav* 2017 Jun 1; 71: 535–549.
33. Mora Guiard J, Malinverni L and Pares N. Narrative-based elicitation: orchestrating contributions from experts and children. In: *CHI'14 extended abstracts on human factors in computing systems*. Toronto: CHI ACM, 2014, pp.1159–1164.
34. Moltrecht B, Patalay P, Bear HA, et al. A transdiagnostic, emotion regulation app (Eda) for children: design, development, and lessons learned. *JMIR Form Res* 2022; 6: e28300.
35. Duncan E, O'Cathain A, Rousseau N, et al. Guidance for reporting intervention development studies in health research (GUIDED): an evidence-based consensus study. *BMJ Open* 2020; 10: e033516.
36. Brylka M and Cygan HB. Selective short-term memory impairment for verbalizable visual objects in children with developmental language disorder. *Res Dev Disabil* 2024; 144: 104637.

37. Clark GT and Reuterskiöld C. Orthographic support for word learning in clinical populations: a systematic review. *Lang Speech Hear Serv Sch* 2021; 52: 937–948.
38. Toseeb U and St Clair MC. Trajectories of prosociality from early to middle childhood in children at risk of developmental language disorder. *J Commun Disord* 2020; 85: 105984.
39. McGregor KK, Ohlmann N, Eden N, et al. Abilities and disabilities among children with developmental language disorder. *Lang Speech Hear Serv Sch* 2023; 54: 927–951.
40. Baron LS and Arbel Y. Inner speech and executive function in children with developmental language disorder: implications for assessment and intervention. *Perspect ASHA Spec Interest Groups* 2022; 7: 1645–1659.
41. Ansari R, Chiat S, Cartwright M and Herman R. Vocabulary interventions for children with Developmental Language Disorder: A systematic review. *Front Psychol* 2025; 16: 1517311.
42. Michie S, Van Stralen MM and West R. The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implement Sci* 2011; 6: 1–2.
43. Braun V and Clarke V. Supporting best practice in reflexive thematic analysis reporting in palliative medicine: a review of published research and introduction to the reflexive thematic analysis reporting guidelines (RTARG). *Palliat Med* 2024; 38: 608–616.
44. Braun V and Clarke V. One size fits all? What counts as quality practice in (reflexive) thematic analysis? *Qual Res Psychol* 2021; 18: 328–352.
45. Braun V and Clarke V. How do you solve a problem like COREQ? A critique of consolidated criteria for reporting qualitative research. *Methods Psychol* 2024; 11: 100155.
46. Krueger RA and Casey MA. Designing and conducting focus group interviews. In: Krueger RA, Casey MA, Donner J, Kirsch S and Maack JN (eds) *Social development papers: social analysis – selected tools and techniques*. Washington, DC: The World Bank, 2001, pp.4–23.
47. Krueger RA. *Focus groups: A practical guide for applied research*. California: Sage Publications, 2014.
48. Best W, Fedor A, Hughes L, et al. Intervening to alleviate word-finding difficulties in children: case series data and a computational modelling foundation. *Cogn Neuropsychol* 2015; 32: 133–168.
49. Constable A, Stackhouse J and Wells B. Developmental word-finding difficulties and phonological processing: the case of the missing handcuffs. *Appl Psycholinguist* 1997; 18: 507–536.
50. German DJ and Newman RS. The impact of lexical factors on children's word-finding errors. *J Speech Lang Hear Res* 2004; 47: 624–636.
51. McGregor KK and Appel A. On the relation between mental representation and naming in a child with specific language impairment. *Clin Linguist Phon* 2002; 16: 1–20.
52. Messer D and Dockrell JE. Children with word finding difficulties: continuities and profiles of abilities. *First Lang* 2013; 33: 433–448.
53. Carey S. Acquiring a single new word. *Pap Rep Child Lang Dev* 1978; 15: 17–29.
54. Chiat S. Mapping theories of developmental language impairment: premises, predictions and evidence. *Lang Cognit Process* 2001; 16: 113–142.
55. Gathercole SE and Baddeley AD. Phonological memory deficits in language disordered children: is there a causal connection? *J Mem Lang* 1990; 29: 336–360.
56. Baddeley AD, Hitch GJ and Allen R. A multicomponent model of working memory. In: Logie R, Camos V and Cowan N (eds) *Working Memory: The State of the Science*. Oxford: Oxford University Press, 2021, pp.10–43.
57. Hoover WA and Gough PB. The simple view of reading. *Reading and Writing: An Interdisciplinary Journal* 1990; 2: 127–160.
58. Snowling MJ. Language and literacy skills: who is at risk and why? In: Bishop DVM and Leonard L (eds) *Speech and language impairments in children*. Hove: Psychology Press, 2014, pp.259–274.
59. Bruner JS. The ontogenesis of speech acts. *J Child Lang* 1975; 2: 1–9.
60. Barkley RA. The executive functions and self-regulation: an evolutionary neuropsychological perspective. *Neuropsychol Rev* 2001; 11: 1–29.
61. Yu C and Ballard DH. A unified model of early word learning: integrating statistical and social cues. *Neurocomputing* 2007; 70: 2149–2165.
62. Karpicke JD, Lehman M and Aue WR. Retrieval-based learning: an episodic context account. In: Ross B (ed) *Psychology of learning and motivation*, Vol. 61. New York: Academic Press Inc, 2014, pp.237–284.
63. Best W, Hughes LM, Masterson J, et al. Intervention for children with word-finding difficulties: a parallel group randomised control trial. *Int J Speech Lang Pathol* 2018; 20: 708–719.
64. Best W, Hughes L, Masterson J, et al. Understanding differing outcomes from semantic and phonological interventions with children with word-finding difficulties: a group and case series study. *Cortex* 2021; 134: 145–161.
65. Marks I and Stokes SF. Narrative-based intervention for word-finding difficulties: a case study. *Int J Lang Commun Disord* 2010; 45: 586–599.
66. Nash M and Donaldson ML. Word learning in children with vocabulary deficits. *J Speech Lang Hear Res* 2005; 48: 439.
67. Parsons S, Law J and Gascoigne M. Teaching receptive vocabulary to children with specific language impairment: a curriculum-based approach. *Child Lang Teach Ther* 2005; 21: 39–59.
68. Wright SH. Teaching word-finding strategies to severely language-impaired children. *Eur J Disord Commun* 1993; 28: 165–175.
69. Steele SC, Willoughby LM and Mills MT. Learning word meanings during reading: effects of phonological and semantic cues on children with language impairment. *Int J Speech Lang Pathol* 2013; 15: 184–197.
70. Storkel HL, Komesidou R, Fleming KK, et al. Interactive book reading to accelerate word learning by kindergarten children with specific language impairment: identifying adequate progress and successful learning patterns. *Lang Speech Hear Serv Sch* 2017; 48: 108–124.

71. Wing CS. A preliminary investigation of generalization to untrained words following two treatments of children's word-finding problems. *Lang Speech Hear Serv Sch* 1990; 21: 151–156.
72. Zens NK, Gillon GT and Moran C. Effects of phonological awareness and semantic intervention on word-learning in children with SLI. *Int J Speech Lang Pathol* 2009; 11: 509–524.
73. Ardanouy E, Delage H and Zesiger P. Effectiveness of a group intervention for lexical enrichment in 6-to-10-year-old children with developmental language disorder. *Child Lang Teach Ther* 2023; 39: 218–233.
74. Motsch HJ and Marks DK. Efficacy of the lexicon pirate strategy therapy for improving lexical learning in school-age children: a randomized controlled trial. *Child Lang Teach Ther* 2015; 31: 237–255.
75. Buzan T and Buzan B. *The mind map book*. Essex, UK: Educational Publishers LLP, 2006.
76. Mayer RE. *Multimedia learning (2nd ed.)*. New York, NY: Cambridge University Press, 2009.
77. Mayer RE. Using multimedia for e-learning. *J Comput Assisted Learn* 2017; 33: 403–423.
78. Wittmann S. A case study in word finding. *Child Lang Teach Ther* 1996; 12: 300–313.
79. Senter R, Chow JC and Willis EC. Speech-language pathology interventions for children with executive function deficits: a systematic literature review. *Lang Speech Hear Serv Sch* 2023; 54: 336–354.
80. Edwards HT and Kirkpatrick AG. Metalinguistic awareness in children: a developmental progression. *J Psycholinguist Res* 1999; 28: 313–329.
81. Kueser JB, Leonard LB, Deevy P, et al. Word-learning trajectories influence long-term recall in children with developmental language disorder and typical development. *J Commun Disord* 2021; 94: 106160.
82. Leonard LB, Deevy P, Karpicke JD, et al. After initial retrieval practice, more retrieval produces better retention than more study in the word learning of children with developmental language disorder. *J Speech Lang Hear Res* 2020; 63: 2763–2776.
83. Shaban A and Pearson E. A learning design framework to support children with learning disabilities incorporating gamification techniques. In *Extended Abstracts of the 2019 CHI Conference on Human Factors in Computing Systems* 2019 May 2 (pp. 1–6).
84. Pearson E and Shaban A. Evaluation of a prototype interactive working memory application for children with learning disabilities. In *Proceedings of the 16th International Web for All Conference* 2019 May 13 (pp. 1–2).
85. Shaban A, Pearson E and Chang V. Evaluation of user experience, cognitive load, and training performance of a gamified cognitive training application for children with learning disabilities. *Front Comput Sci* 2021; 3: 617056.
86. Figma Inc. (Version Released 2022). Figma Design Tool Figma: The Collaborative Interface Design Tool.
87. Dale E. Vocabulary measurement: techniques and major findings. *Elem Engl* 1965; 42: 895–948.
88. Steele SC and Mills MT. Vocabulary intervention for school-age children with language impairment: a review of evidence and good practice. *Child Lang Teach Ther* 2011; 27: 354–370.
89. Westby C. Assessing vocabulary breadth and depth. *Word Mouth* 2024; 35: 7–9.
90. Utianski RL, Spencer TD and Wallace SE. Clinical impact requires clinical practice research. *Perspect ASHA Spec Interest Groups* 2022; 7: 651–662.
91. McKean C, Watson R, Charlton J, et al. 'Making the most of together time': development of a health visitor-led intervention to support children's early language and communication development at the 2–2½-year-old review. *Pilot Feasibility Stud* 2022; 8: 35.
92. Bauer MS and Kirchner J. Implementation science: what is it and why should I care? *Psychiatry Res* 2020; 283: 112376.
93. Bédard VB, Trudeau N and Macleod AA. Profiles of word-finding difficulties in school-aged children. *J Child Lang* 2024; 51: 1268–1289.
94. Raymer AM. Word finding and lexical-semantic disorders. In: Brown GG, King TZ, Haaland KY and Crosson B (eds) *APA Handbook of neuropsychology: neurobehavioral disorders and conditions: accepted science and open questions*. US: American Psychological Association, 2023, pp.23–41
95. Stringer H, Nicholson D, Hope K, et al. Using the Behaviour Change Wheel to discover barriers and enablers for engaging Early Childhood Educators in partnership working with Speech and language therapy Services. Poster Presentation, Community for Allied Health Professional Research (CAHPR). 2020. Accessed at: <https://cahpr.csp.org.uk/system/files/documents/2023-01/ID%203.6.pdf>.
96. Berry A, McClellan C, Wanless B, et al. A tailored app for the self-management of musculoskeletal conditions: evidencing a logic model of behavior change. *JMIR Format Res* 2022; 6: e32669.
97. Indraratna P, Biswas U, Liu H, et al. Process evaluation of a randomised controlled trial for TeleClinical care, a smartphone-app based model of care. *Front Med* 2022; 8: 780882.
98. Du Y, Lubniewski K, Price L, et al. "They can't believe they're a tiger": insights from pediatric speech-language pathologist mobile app users and app designers. *Int J Lang Commun Disord* 2023; 58: 1717–1737.
99. Morris ZS, Wooding S and Grant J. The answer is 17 years, what is the question: understanding time lags in translational research. *J R Soc Med* 2011; 104: 510–520.
100. Leigh S and Ashall-Payne L. The role of health-care providers in mHealth adoption. *Lancet Digit Health* 2019; 1: e58–e59.
101. Frauenberger C, Spiel K and Makhaeva J. Thinking outside the box-designing smart things with autistic children. *Int J Human-Computer Interact* 2019; 35: 666–678.

APPENDIX A. GUIDED – A guideline for intervention development studies³⁵

Item description	Where item is located in manuscript
1. Report the context for which the intervention was developed.	RESULTS SECTION Phase 1: Examination of current practice and context to define scope of intervention. Figure 2: Logic Model which includes contextual factors influencing the intervention (moderators)
2. Report the purpose of the intervention development process.	RESULTS SECTION Table 1, Column 6 and Table 2, Column 5: Intervention change processes
3. Report the target population for the intervention development process.	RESULTS SECTION Table 1, Column 1: Developmental Language Disorder profile
4. Report how any published intervention development approach contributed to the development process	MRC Framework for Developing and Evaluating Complex Interventions ^{28,29} Additional design steps from synthesis of design literature ³⁰
5. Report how evidence from different sources informed the intervention development process	RESULTS SECTION Table 1: Recommended strategy > Empirical evidence Intervention design feature > Change process
6. Report how/if published theory informed the intervention development process.	RESULTS SECTION: Table 1, Column 2: Theories underlying DLD profile. Table 2, Column 2: COM-B model of clinician's behaviour-change drivers ⁴²
7. Report any use of components from an existing intervention in the current intervention development process.	RESULTS SECTION: Table 1: Recommended strategy > Empirical evidence > Intervention design feature > Change process
8. Report any guiding principles, people or factors that were prioritised when making decisions during the intervention development process.	RESULTS SECTION: Table 1, Column 3 and Table 2, Column 3: Recommended strategies Clinician feedback and refinement.
9. Report how stakeholders contributed to the intervention development process.	RESULTS SECTION: Phase 2: Clinician feedback and refinement.
10. Report how the intervention changed in content and format from the start of the intervention development process.	RESULTS SECTION: Phase 2: Design refinements following clinician feedback.
11. Report any changes to interventions required or likely to be required for subgroups.	RESULTS SECTION Figure 2: Logic Model which includes contextual factors influencing the intervention (moderators)
12. Report important uncertainties at the end of the intervention development process.	LIMITATIONS SECTION
13. Follow TIDieR guidance when describing the developed intervention.	Planned
14. Report the intervention development process in an open access format.	Planned