



City Research Online

City, University of London Institutional Repository

Citation: Macchia, L., Okafor, C. N., Breedlove, T., Shiba, K., Piper, A., Johnson, B. & VanderWeele, T. J. (2025). A cross-national analysis of childhood predictors of physical pain. *Commun Med (Lond)*, 5(1), 337. doi: 10.1038/s43856-025-00997-2

This is the published version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/35644/>

Link to published version: <https://doi.org/10.1038/s43856-025-00997-2>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

<https://doi.org/10.1038/s43856-025-00997-2>

A cross-national analysis of childhood predictors of physical pain



Lucía Macchia¹✉, Chukwuemeka N. Okafor², Thomas Breedlove³, Koichiro Shiba⁴, Alan Piper⁵, Byron Johnson³ & Tyler J. VanderWeele⁶

Abstract

Background The socioeconomic, psychosocial, and behavioural factors that influence physical pain have been largely explored. However, evidence on the childhood circumstances that shape physical pain in adulthood is scarce.

Methods Using a nationally representative dataset of 202,898 individuals (Age range: 18–99, 49% female, 51% male, 0.03% other) from 22 countries and a random effects meta-analysis, we examine 13 potential childhood predictors of physical pain in adulthood.

Results We find that childhood experiences, personal attributes, and familial and social circumstances have meaningful and varied associations with adulthood levels of physical pain. Specifically, we find that people whose parents were divorced, single, or died (vs married) when they were children, those who reported that their family found it very difficult to live with their family's household income when they were growing up (vs got by), those who experienced abuse (vs not), felt like an outsider (vs not), reported poor physical health while growing up (vs good) and reported more frequent religious attendance at age 12 (vs never) have a greater risk of experiencing pain later in life. The strength of these associations differs by country reflecting diverse societal influences.

Conclusions Our study provides valuable insights on the early-life experiences that shape physical pain in adulthood. These findings enhance our understanding of early-life predictors of adult wellbeing and offer comprehensive evidence for designing interventions to reduce physical pain levels.

Plain language summary

Knowing the childhood predictors of physical pain is crucial for the development of approaches to reducing pain later in life. In this study, we used rigorous statistical techniques to examine the relationship between 13 childhood circumstances and physical pain in adulthood across 22 countries from all over the world. We found that people who had disadvantageous situations during childhood, like difficulties with family income, abuse, or poor physical health, had a higher risk of experiencing physical pain later in life. Other circumstances, like frequent religious attendance at age 12, were also linked to greater physical pain in adulthood. We also found that the link between childhood circumstances and pain in adulthood varied by country.

A large body of work suggests that many diseases and conditions that occur in adulthood may have their origins in childhood^{1–3}. For instance, Slopen et al.⁴ have shown that childhood experiences like parental love, two-parent family structure, and high socioeconomic status were linked to good cardiovascular health in middle adulthood. Previous work also found that parental warmth during childhood was negatively associated with harmful health behaviours like drug use and smoking in mid-life⁵, and that social and emotional support during childhood can be protective against later depression⁶. Related work has shown that frequent early-life religious attendance was linked to better physical health in later life⁷. While the association between childhood circumstances and health in adulthood has been studied in detail, less is known about the childhood experiences that shape physical pain in adulthood.

Prior work has attempted to address this empirical question. For instance, using data from the 1958 British Cohort Study, Jones et al.⁸ found that childhood adversity, represented by familial financial hardship, living in institutional care, and maternal death, was linked to greater chronic pain in adulthood. Other related studies also used cohort data from the United Kingdom and found that low socio-economic status in childhood was linked to greater pain in adulthood^{9–11}. Mustard et al.¹² have shown the same association using Canadian data. Focusing on other childhood predictors and using data from the United States, prior work has found that maternal depression¹³, physical abuse¹⁴, and an accumulation of different adverse events¹⁵ were associated with greater physical pain in adulthood. A strong link between physical abuse during childhood and pain later in life was also found in Japan¹⁶. The stress that results from

¹School of Health and Medical Sciences, City St George's, University of London, London, UK. ²Long School of Medicine, University of Texas Health Science Center San Antonio, San Antonio, USA. ³Institute for Studies of Religion, Baylor University, Waco, TX, USA. ⁴Boston University School of Public Health, Boston, MA, USA. ⁵Leeds University Business School, Leeds, UK. ⁶Human Flourishing Program, Harvard University, Cambridge, MA, USA.

✉ e-mail: lucia.macchia@citystgeorges.ac.uk

exposure to negative events seems to have lifelong consequences for physical pain¹⁷.

Although this body of work sheds light on a very important question, most of the conclusions were drawn based on samples from specific countries and only a limited number of childhood circumstances were explored. This scenario might have its roots in the challenge of collecting relevant data in different regions at the same time. Here, we use data from a diverse and international sample of 202,898 individuals across 22 countries to explore 13 potential childhood predictors, assessed retrospectively, of physical pain in adulthood. These predictors include age, gender, immigration status, marital status or family structure, religious service attendance and religious affiliation at age 12, relationship with mother and father, feeling like an outsider, abuse, self-rated health, immigration status, and subjective financial status of family while growing up. The selection of these childhood circumstances is based on the previously described body of work that documented strong relationships between these factors and mental and physical health in adulthood.

Examining childhood experiences is crucial because they have been associated with important behavioural outcomes in adulthood. For instance, childhood adversity has been found to be linked to harmful health behaviours like smoking and heavy drinking¹⁸ and weaker interpersonal relationships¹⁹. Prior work has also shown that people who experienced adversity while growing up died 20 years earlier than those who did not experience adversity while growing up²⁰.

The present research is motivated by the Ecological Systems Theory (EST)²¹ which suggests that one key predictor of people's circumstances in adulthood, including physical and mental health is the set of immediate environmental systems during childhood (e.g. family structure and relationships, socioeconomic status). Building on this theory that highlights the importance of an individual's environment while growing up, we examine the following hypotheses. Hypothesis 1 suggests that childhood experiences, personal attributes, and familial or social circumstances will have varied associations with adulthood levels of physical pain. Hypothesis 2 proposes that the strength of these associations will differ by country, reflecting diverse societal influences that are specific to each country. Hypothesis 3 suggests that some of the observed associations between the 13 childhood predictors and an individual's physical pain in adulthood will be robust against potential unmeasured confounding assessed through E-values.

To explore these hypotheses we (1) calculated descriptive statistics for the various childhood predictors across the 22 countries (Table 1), (2) conducted the meta-analysis of regression estimates for each of the childhood predictors using a random effects meta-analysis (Table 2), (3) estimated E-values to assess the robustness of the meta-analytic effect estimates in Table 2 to potential unmeasured confounding (Table 3), and (4) presented analyses by country and graphs that visually display the childhood predictor effect estimates in each of the 22 countries in the Supplementary Information.

Our study aims to provide comprehensive insights into early-life experiences that contribute to the levels of physical pain in adulthood. The study shows that childhood experiences, personal attributes, and familial and social circumstances have meaningful and varied associations with adulthood pain. These findings enhance our understanding of early-life predictors of adult outcomes and provide valuable evidence for developing approaches to reducing physical pain levels.

Methods

All analyses were pre-registered with COS prior to data access (https://osf.io/zshpy/?view_only=34700935c63449968221300d51b91350).

The description of the methods below has been adapted from VanderWeele et al.²². Further methodological detail is available elsewhere^{23–29}.

Data

The Global Flourishing Study (GFS) is a study of 202,898 participants from 22 geographically and culturally diverse countries, with nationally representative sampling within each country, concerning the distribution of determinants of wellbeing. Wave 1 of the data included the following

countries and territories: Argentina, Australia, Brazil, Egypt, Germany, Hong Kong, India, Indonesia, Israel, Japan, Kenya, Mexico, Nigeria, the Philippines, Poland, South Africa, Spain, Sweden, Tanzania, Turkey, the United Kingdom, and the United States. The countries were selected to (a) maximise coverage of the world's population, (b) ensure geographic, cultural, and religious diversity, and (c) prioritise feasibility and existing data collection infrastructure. Data collection was carried out by Gallup Inc. Data for Wave 1 were collected principally during 2023, with some countries beginning data collection in 2022 and exact dates varying by country²⁷. Four additional waves of panel data on the participants will be collected annually from 2024 to 2027. The precise sampling design to ensure nationally representative samples varied by country and further details are available in Ritter et al.²⁷. Survey items included aspects of well-being such as happiness, health, meaning, character, relationships, and financial stability³⁰, along with other demographic, social, economic, political, religious, personality, childhood, community, health, and wellbeing variables. The data are publicly available through the Center for Open Science (<https://www.cos.io/gfs>). During the translation process, Gallup adhered to the TRAPD model (translation, review, adjudication, pretesting, and documentation) for cross-cultural survey research (ccsg.isr.umich.edu/chapters/translation/overview). Additional details about methodology and survey development can be found in the GFS Questionnaire Development Report²³, the GFS Methodology²⁷, the GFS Codebook, and the GFS Translations documents²⁴.

This project was ruled EXEMPT for Institutional Review Board (IRB) review by the Baylor University IRB (#1841317-2). Gallup Inc. IRB approved the study on November 16, 2021 (#2021-11-02). All data collection was performed in accordance with the ethical standards of Gallup and with the 1964 Helsinki Declaration and its later amendments. Informed consent was obtained during the respondent recruitment stage of fieldwork. Consent was also obtained at the start of the survey. The exact wording varies across countries depending on the local laws and regulations governing data protection. All personally identifiable information (PII) was removed from the data used in this study by Gallup Inc.

Measures

Childhood Demographic Variables: Relationship with mother during childhood was assessed with the question: 'Please think about your relationship with your mother when you were growing up. In general, would you say that relationship was very good, somewhat good, somewhat bad, or very bad?' Responses were dichotomised to very/somewhat good versus very/somewhat bad. An analogous variable was used for relationship with father. We dichotomised these two variables to avoid a multi-collinearity issue because these two variables were highly correlated. 'Does not apply' was treated as a dichotomous control variable for respondents who did not have a mother or father due to death or absence. Parental marital status during childhood was assessed with responses of married, divorced, never married, and one or both had died. Financial status was measured with: 'Which one of these phrases comes closest to your own feelings about your family's household income when you were growing up, such as when YOU were around 12 years old?' Responses were lived comfortably, got by, found it difficult, and found it very difficult. Abuse was assessed with yes/no responses to 'Were you ever physically or sexually abused when you were growing up?' Participants were also asked: 'When you were growing up, did you feel like an outsider in your family?' Childhood health was assessed by: 'In general, how was your health when you were growing up? Was it excellent, very good, good, fair, or poor?' Immigration status was assessed with: 'Were you born in this country, or not?' Religious attendance during childhood was assessed with: 'How often did YOU attend religious services or worship at a temple, mosque, shrine, church, or other religious building when YOU were around 12 years old?' with responses of at least once/week, one-to-three times/month, less than once/month, or never. Gender was assessed as male, female, or other. The choice of age 12 reflects a balance between ensuring adequate recall ability and capturing a key stage in developmental transitions. Continuous age (year of birth) was classified as 18–24, 25–29, 30–39, 40–49, 50–59, 60–69, 70–79, and 80 or older.

Childhood religious tradition/affiliation had response categories of Christianity, Islam, Hinduism, Buddhism, Judaism, Sikhism, Baha'i, Jainism, Shinto, Taoism, Confucianism, Primal/Animist/Folk religion, Spiritism, African-Derived, some other religion, or no religion/atheist/agnostic; precise response categories varied by country³¹. Racial/ethnic identity was assessed in some, but not all, countries, with response categories varying by country. In some cases, the reference category was chosen based on the category with the largest number of respondents (e.g. subjective financial status of family growing up—got by) or the mid-category (e.g. health when you were growing up—good). For additional details on the assessments see the COS GFS codebook or Crabtree et al.²³.

Outcome Variable: Our pain measure comes from the following question: 'How much bodily pain have you had during the past 4 weeks?' Respondents could answer a lot, some, not very much, or none at all. In the main analyses, this variable was dichotomised as a lot/some (1) vs. not very much/none at all (0). As a post-hoc sensitivity analysis we provide the equivalent results of Table 2 with a different dichotomisation point where the outcome is treated as lot/some/not very much (1) vs. none at all (0).

Statistics and reproducibility

Descriptive statistics for the observed sample, weighted to be nationally representative within the country, were estimated for each childhood demographic category. A weighted modified Poisson regression model with complex survey-adjusted standard errors was fit within each country for physical pain on all of the aforementioned childhood predictor variables simultaneously. In the primary analyses, random effects meta-analyses of the regression coefficients^{32,33} along with confidence intervals, lower and upper limits with 95% prediction intervals, and I² for evidence concerning variation within a given demographic category across countries³⁴. Plots of estimates are available in the online supplement. Religious affiliation/tradition and race/ethnicity were used within country as control variables, when available, but these coefficients themselves were not included in the meta-analyses since categories/responses varied by country. All meta-analyses were conducted in R³⁵ using the metafor package³⁶. Within each country, a global test of association of each childhood predictor variable group with physical pain was conducted, and a pooled *p*-value³⁷ across countries reported concerning evidence for association within any country. Bonferroni corrected *p*-value thresholds are provided based on the number of childhood demographic variables^{38,39}. For each childhood predictor, we calculated *E*-values to evaluate the sensitivity of results to unmeasured confounding. The *E*-value informs whether the results can be influenced by factors not included in the analyses. For instance, large *E*-values suggest that the unmeasured confounding can be strongly associated with both the exposure and outcome whereas small *E*-values suggest weak unmeasured confounding. An *E*-value is the minimum strength of the association an unmeasured confounder must have with both the outcome and the predictor, above and beyond all measured covariates, for an unmeasured confounder to explain away an association⁴⁰. As a supplementary analysis, population-weighted meta-analyses of the regression coefficients were estimated. All analyses were pre-registered with COS prior to data access, with only slight subsequent modification in the regression analyses due to multicollinearity (https://osf.io/zshpy/?view_only=34700935c63449968221300d51b91350, see also Text S.1 in the Supplementary Information); all code to reproduce analyses are openly available in an online repository²⁶.

Missing data. Missing data on all variables was imputed using multi-variate imputation by chained equations, and five imputed datasets were used^{41–44}. To account for variation in the assessment of certain variables across countries (e.g. religious affiliation/tradition and race/ethnicity), the imputation process was conducted separately in each country. This within-country imputation approach ensured that the imputation models accurately reflected country-specific contexts and assessment methods. Sampling weights were included in the imputation model to account for missingness to be related to probability of inclusion.

Accounting for complex sampling design. The GFS used different sampling schemes across countries based on the availability of existing panels and recruitment needs²⁷. All analyses accounted for the complex survey design components by including weights, primary sampling units, and strata. Additional methodological detail, including accounting for the complex sampling design, is provided elsewhere²⁵.

Reporting summary

Further information on research design is available in the Nature Portfolio Reporting Summary linked to this article.

Results

Descriptive statistics

Table 1 shows the number and percentage of people across each childhood predictor in the observed sample: 63% had a very good relationship with their mother, 53% had a very good relationship with their father, 75% had parents who were married, 41% reported that they could get by with their family income while they were growing up, 82% had not suffered from abuse, 84% had not felt like an outsider while growing up, 33% reported to have had excellent health while growing up, 94% were born in the country in which the survey was conducted, and 41% attended religious services at least once a week when they were 12 years of age. Most individuals were middle-aged (30–39 years old (20%), 40–49 years old (17%), 50–59 (16%)), 51% of respondents were women, and 49% were men. These descriptive statistics for each country separately can be found in Tables S1–S22 in the Supplementary Information (SI).

Random effects meta-analysis

Table 2 provides a summary estimate from a random effects meta-analysis of the multivariate logistic regression estimates across countries for each of the childhood predictors. The table shows that the risk ratios for having had a good or somewhat good relationship with mother and with father were 1.02 and 1.00, respectively. This suggests that there is virtually no difference in the risk of feeling pain between people who had a good or somewhat good relationship and those who had a bad or somewhat bad relationship with both mother and father. The table also shows that people whose parents were divorced (RR = 1.06, *p* < 0.001, 95% CI [1.02, 1.10]), single (RR = 1.11, *p* < 0.001, 95% CI [1.01, 1.21]), or died (RR = 1.07, *p* < 0.001, 95% CI [1.02, 1.13]) had a greater risk of experiencing physical pain in comparison to those whose parents were married. Participants who reported that their family found it very difficult to live with their household income when they were growing up (vs got by, RR = 1.07, *p* < 0.001, 95% CI [1.03, 1.11]), those who experienced abuse (vs no, RR = 1.25, *p* < 0.001, 95% CI [1.22, 1.29]), those who felt like an outsider (vs no, RR = 1.16, *p* < 0.001, 95% CI [1.11, 1.20]), and those who reported poor physical health while growing up (vs good, RR = 1.20, *p* < 0.001, 95% CI [1.07, 1.35]) had a greater risk of experiencing pain. People who were born in a different country to the one in which the survey was conducted (RR = 0.94, *p* < 0.001, 95% CI [0.89, 1.00]) had a lower risk of experiencing physical pain in comparison to those who were born in the country in which the survey was conducted. Religious attendance at age 12 was positively linked to physical pain: People who attended religious services at least once a week (RR = 1.04, *p* < 0.001, 95% CI [1.00, 1.08]), 1–3 times a month (RR = 1.07, *p* < 0.001, 95% CI [1.03, 1.13]) or less than once a month (RR = 1.02, *p* < 0.001, 95% CI [0.98, 1.07]) had a greater risk of experiencing pain than those who never attended a religious service at age 12. Regarding key demographic factors, the analysis shows that older people (vs younger) and women (vs men) had a greater risk of experiencing physical pain. The findings shown in Table 2 confirm *Hypothesis 1* of this study: childhood experiences, personal attributes, and familial or social circumstances have varied associations with adulthood levels of physical pain.

Country-specific estimates of these regression models can be found in Tables S23–S244 in the Supplementary Information. Figures S1–S27 show graphs for each of the childhood predictor estimates in each of the 22 countries. These figures show the information presented in Tables S23–S44

Table 1 | Nationally representative childhood descriptive statistics of the observed sample

Characteristic	N = 202,898 ^a	Characteristic	N = 202,898 ^a
Relationship with mother		Age group	
Very good	127,836 (63%)	1998–2005; age: 18–24	27,007 (13%)
Somewhat good	52,439 (26%)	1993–1998; age: 25–29	20,700 (10%)
Somewhat bad	11,060 (5.5%)	1983–1993; age: 30–39	40,256 (20%)
Very bad	4,642 (2.3%)	1973–1983; age 40–49	34,464 (17%)
Does not apply	5,965 (2.9%)	1963–1973; age 50–59	31,793 (16%)
Missing	956 (0.5%)	1953–1963; age 60–69	27,763 (14%)
Relationship with father		1943–1953; age 70–79	16,776 (8.3%)
Very good	107,742 (53%)	1943 or earlier; age 80+	4119 (2.0%)
Somewhat good	55,714 (27%)	Missing	20 (<0.1%)
Somewhat bad	15,807 (7.8%)	Gender	
Very bad	8278 (4.1%)	Male	98,411 (49%)
Does not apply	13,985 (6.9%)	Female	103,488 (51%)
Missing	1372 (0.7%)	Other	602 (0.3%)
Parent marital status		Missing	397 (0.2%)
Yes, married	152,001 (75%)	Country	
No, divorced	17,726 (8.7%)	Argentina	6724 (3.3%)
Never married	15,534 (7.7%)	Australia	3844 (1.9%)
No, one or both of them had died	7794 (3.8%)	Brazil	13,204 (6.5%)
Missing	9843 (4.9%)	Egypt	4729 (2.3%)
Subjective financial status of family growing up		Germany	9506 (4.7%)
Lived comfortably	70,861 (35%)	India	12,765 (6.3%)
Got by	82,905 (41%)	Indonesia	6992 (3.4%)
Found it difficult	35,852 (18%)	Israel	3669 (1.8%)
Found it very difficult	12,606 (6.2%)	Japan	20,543 (10%)
Missing	674 (0.3%)	Kenya	11,389 (5.6%)
Abuse		Mexico	5776 (2.8%)
Yes	29,139 (14%)	Nigeria	6827 (3.4%)
No	167,279 (82%)	Philippines	5292 (2.6%)
Missing	6479 (3.2%)	Poland	10,389 (5.1%)
Outsider growing up		South Africa	2651 (1.3%)
Yes	28,732 (14%)	Spain	6290 (3.1%)
No	170,577 (84%)	Tanzania	9075 (4.5%)
Missing	3589 (1.8%)	Turkey	1473 (0.7%)
Self-rated health growing up		United Kingdom	5368 (2.6%)
Excellent	67,121 (33%)	United States	38,312 (19%)
Very good	63,086 (31%)	Sweden	15,068 (7.4%)
Good	47,378 (23%)	Hong Kong	3012 (1.5%)
Fair	19,877 (9.8%)		
Poor	4906 (2.4%)		
Missing	530 (0.3%)		
Immigration status			

Table 1 (continued) | Nationally representative childhood descriptive statistics of the observed sample

Characteristic	N = 202,898 ^a	Characteristic	N = 202,898 ^a
Born in this country	190,998 (94%)		
Born in another country	9791 (4.8%)		
Missing	2110 (1.0%)		
Age 12 rel. service attendance			
At least 1/week	83,237 (41%)		
1–3/month	33,308 (16%)		
<1/month	36,928 (18%)		
Never	47,445 (23%)		
Missing	1980 (1.0%)		

^an (%)

but organised by childhood predictor, rather than by country. These country-specific findings show that some aspects like having experienced abused were significantly positively associated with adult pain in all the countries of our sample (Fig. S9). However, the link between adult pain and other aspects, like the subjective financial status of family growing up varied by country. For instance, finding it very difficult to live with the family's household income while growing up was significantly positively associated with adult pain in Poland, Turkey, Egypt, and Brazil, whereas that link was insignificant in the other countries (Fig. S8). The results shown in Tables S23–S44, and Figs. S1–S27 confirm *Hypothesis 2*: The strength of these associations differ by country, reflecting diverse societal influences that are specific to each country.

Unmeasured confounding

The E-values shown in Table 3 suggest that some of the observed associations were moderately robust to unmeasured confounding. For example, for having experienced abuse, an unmeasured confounder that was associated with both having experienced abuse and physical pain by risk ratios of 1.82 each (above and beyond the covariates already adjusted for) could explain away the association, but weaker joint confounder associations could not. To shift the confidence interval to include the null, an unmeasured confounder associated with both having experienced abuse and physical pain by risk ratios of 1.73 each could suffice, but weaker joint confounder associations could not.

Country-specific E-values can be found in Tables S45–S66 in the Supplementary Information. Overall, the analyses presented in Table 3, and Tables S45–S66 confirm *hypothesis 3* of this study: the observed associations between the 13 childhood predictors and an individual's physical pain in adulthood are robust against potential unmeasured confounding assessed through E-values.

Alternative meta-analyses in which each country's results were weighted by the *actual* 2023 population size are presented in Table S67. These results are consistent with those presented in Tables 2 and 3.

Discussion

In this study, we used a dataset of 202,898 individuals from 22 countries and a random effects meta-analysis to examine 13 childhood predictors of physical pain in adulthood. Overall, we found that people whose parents were divorced, single, or died (vs married) when they were children, those who reported that their family found it very difficult to live with their household income when they were growing up (vs got by), those who experienced abuse (vs not), felt like an outsider (vs not), reported poor physical health while growing up (vs good) and reported more frequent religious attendance at age 12 (vs never) had a greater risk of experiencing pain later in life. These findings are in line with the EST²¹ which supports the idea that environmental factors, including childhood circumstances, can influence people's development, experiences, and wellbeing.

Table 2 | Random effects meta-analysis of regression of physical pain on childhood predictors

Variable	Category	RR	95% CI	Estimated proportion of effects by threshold		I ²	Global p-value
				<0.90	>1.10		
Relationship with mother	(Ref: Very bad/somewhat bad)	1.02	(0.98, 1.05)	0.00	0.00	21.9	6.52e-06
	Very good/somewhat good						
Relationship with father	(Ref: Very bad/somewhat bad)						1.89e-05
	Very good/somewhat good	1.00	(0.98, 1.02)	0.00	0.00	<0.1#	
Parent marital status	(Ref: Parents married)						3.96e-06
	No, divorced	1.06	(1.02, 1.10)	0.00	0.27	46.2	
	Single, never married	1.11	(1.01, 1.21)	0.00	0.32	89.7	
	No, one or both of them had died	1.07	(1.02, 1.13)	0.00	0.36	57.0	
Subjective financial status of family growing up	(Ref: Got by)						2.19e-06
	Lived comfortably	0.97	(0.93, 1.00)	0.18	0.09	74.4	
	Found it difficult	1.08	(1.05, 1.10)	0.00	0.23	38.1	
	Found it very difficult	1.07	(1.03, 1.11)	0.00	0.18	34.8	
Abuse	(Ref: No)						1.05e-06
	Yes	1.25	(1.22, 1.29)	0.00	1.00	58.8	
Outsider growing up	(Ref: No)						1.98e-06
	Yes	1.16	(1.11, 1.20)	0.00	0.77	71.8	
Self-rated health growing up	(Ref: Good)						1.55e-06
	Excellent	0.85	(0.81, 0.90)	0.64	0.00	86.5	
	Very good	0.91	(0.89, 0.93)	0.36	0.00	22.7	
	Fair	1.08	(1.04, 1.12)	0.00	0.23	56.7	
	Poor	1.20	(1.07, 1.35)	0.00	0.59	91.2	
Immigration status	(Ref: Born in this country)						6.40e-06
	Born in another country	0.94	(0.89, 1.00)	0.36	0.00	44.2	
Age 12 religious service attendance	(Ref: Never)						3.46e-06
	At least 1/week	1.04	(1.00, 1.08)	0.05	0.14	52.0	
	1–3/month	1.07	(1.03, 1.13)	0.00	0.32	62.2	
	<1/month	1.02	(0.98, 1.07)	0.00	0.18	60.8	
Year of birth	(Ref: 1998–2005; age 18–24)						1.37e-06
	1993–1998; age 25–29	1.03	(1.00, 1.06)	0.00	0.00	<0.1#	
	1983–1993; age 30–39	1.06	(1.00, 1.11)	0.09	0.36	75.9	
	1973–1983; age 40–49	1.12	(1.06, 1.18)	0.05	0.73	76.1	
	1963–1973; age 50–59	1.25	(1.14, 1.36)	0.05	0.77	89.8	
	1953–1963; age 60–69	1.28	(1.14, 1.43)	0.05	0.77	91.9	
	1943–1953; age 70–79	1.41	(1.22, 1.63)	0.00	0.77	92.7	
	1943 or earlier; age 80+	1.56	(1.32, 1.83)	0.05	0.95	84.9	
Gender	(Ref: Male)						1.18e-06
	Female	1.14	(1.10, 1.19)	0.00	0.73	84.8	
	Other	0.25	(0.04, 1.52)	0.22	0.56	99.6	

Note. #estimate of heterogeneity is likely unstable, please see Figs. S1–S27 for more detail on heterogeneity of effects. 95% CI: The 95% confidence interval for that estimate.

Estimated proportion of effects by threshold: The estimated proportion of effects across the implied distribution of effects sizes within the 22 countries that are either less than a RR of 0.9 or above a RR of 1.1.

I²: An estimate of how much of the variability in effect sizes is due to heterogeneity across countries vs. sampling variability. Global p-value: Test of the null hypothesis that the RR is 1 in all of the 22 countries. RR risk ratio.

Our analyses show that the strength of these associations differs by country. For instance, while abuse was significantly positively linked with adult pain in all countries, finding it very difficult to live with family's household income while growing up was significantly negatively associated with pain in specific countries like Poland, Turkey, Egypt, and Brazil. Historical, macroeconomic, and cultural factors might help to explain these differences. For instance, stronger social safety nets and the development of resilience that help people to cope with financial difficulties can explain why finding it very difficult to live with family's household income can have

different links with adult pain across countries. More research is needed to understand the specific historical, macroeconomic, and cultural aspects that might influence cross-country findings.

The general findings are in line with prior work that showed that children who grew up with married parents had better physical health and psychological wellbeing than those who grew up with single or separated parents⁴⁵. Socioeconomic status^{9–12} and abuse^{14,16} during childhood have been found to be associated with greater physical pain in adulthood in the United States, Canada, and Japan. Discrimination has also been found to be

Table 3 | Sensitivity of meta-analyzed childhood predictors to unmeasured confounding

Variable	Predictor (level)	E-value for Estimate	E-value for 95% CI
Relationship with mother	(Ref: Very bad/somewhat bad)		
	Very good/somewhat good	1.14	1.00
Relationship with father	(Ref: Very bad/somewhat bad)		
	Very good/somewhat good	1.05	1.00
Parent marital status	(Ref: Parents married)		
	No, divorced	1.31	1.18
	Single, never married	1.45	1.13
	No, one or both of them had died	1.35	1.15
Subjective financial status of family growing up	(Ref: Got by)		
	Lived comfortably	1.22	1.00
	Found it difficult	1.36	1.28
	Found it very difficult	1.35	1.22
Abuse	(Ref: No)		
	Yes	1.82	1.73
Outsider growing up	(Ref: No)		
	Yes	1.59	1.47
Self-rated health growing up	(Ref: Good)		
	Excellent	1.63	1.47
	Very good	1.43	1.36
	Fair	1.38	1.26
	Poor	1.69	1.34
Immigration status	(Ref: Born in this country)		
	Born in another country	1.31	1.03
Age 12 religious service attendance	(Ref: Never)		
	At least 1/week	1.24	1.03
	1–3/month	1.36	1.19
	<1/month	1.17	1.00
Year of birth	(Ref: 1998–2005; age 18–24)		
	1993–1998; age 25–29	1.20	1.00
	1983–1993; age 30–39	1.30	1.05
	1973–1983; age 40–49	1.49	1.32
	1963–1973; age 50–59	1.80	1.55
	1953–1963; age 60–69	1.87	1.54
	1943–1953; age 70–79	2.17	1.74
	1943 or earlier; age 80+	2.49	1.98
Gender	(Ref: Male)		
	Female	1.55	1.44
	Other	7.46	1.00

a strong predictor of physical pain over time in Australia⁴⁶. Poor physical health in childhood tends to be strongly correlated with poor physical health in adulthood⁴⁷. The psychological distress that results from childhood disadvantages might explain the findings of this study^{17,48} as prior work has shown that negative emotions like distress and anxiety can create new pain and exacerbate existing pain⁴⁹.

It is possible that other childhood circumstances not explored here play a role in adult pain. For instance, Sheffler et al.⁵⁰ found that coping strategies mediated the link between adverse childhood experiences and mental and physical health outcomes later in life. Moreover, prior work has suggested that positive childhood experiences like engaging in prosocial behaviour, or having a large number of friends at school, can mitigate the effect of childhood negative experiences in adult health, pain, and wellbeing⁵¹. Future research should explore these possibilities.

Although recent work has shown that frequent early-life religious service attendance was linked to better physical health in later life⁷, in this study, we found that more frequent religious service attendance during childhood was linked to greater pain in adulthood which is consistent with some other studies⁵². Although religious participation is often associated with better health, pain may be an exception⁵². It might be the case that children with poorer physical health sought religious attendance for relief and that their poor health during childhood predicted their greater pain in adulthood. However, we attempted to control for this possibility by adjusting for self-rated health in childhood. More research is needed to fully understand the dynamics between religious service attendance in childhood and physical pain later in life.

The potential multicollinearity among childhood predictors also deserves special attention. For instance, our results show that the risk of

experiencing pain for people who had a good relationship with their mother or father does not differ from the risk of those who had a bad relationship with their mother or father. However, these results should be interpreted with caution due to the potential correlation among these two variables and marital status. It might be the case that people's relationship with their parents during childhood is captured by marital status, which, out of these three variables, is the one that yielded the most reliable estimate of heterogeneity (see the note of Table 2). It is also possible that prior studies relating pain to relationship with mother and father may have been confounded by marital status.

Our study has some limitations. First, our pain measure does not allow us to explore the intensity or the type of pain (e.g. back pain, neck pain), or whether pain interferes with daily activities. Second, because the childhood predictors were reported retrospectively, they may be subject to recall bias. However, for recall bias to completely explain the observed associations of the childhood predictors with adult physical pain, the effect of adult physical pain on the retrospective assessments of the childhood predictors would have to be at least as strong as the observed associations themselves⁵³. Many of these observed associations were quite substantial. Third, our analyses might be subject to remaining unmeasured confounding, which we tried to address with the E-value sensitivity analysis reported in Table 3.

In the last decades, physical pain has become one of the most important public health challenges⁵⁴. The health and economic benefits of childhood policy have, moreover, received extensive recent attention⁵⁵. The findings of this study link these two important domains of enquiry and might inform the design of early life interventions that would contribute to reducing physical pain in adulthood.

Data availability

All the data that support the findings of this article are openly available on the Open Science Framework (<https://www.cos.io/gfs>). The specific dataset used was Wave 1 non-sensitive Global data <https://osf.io/sm4cd/> available February 2024–March 2026 via pre-registration and publicly from then onwards. The source data for Figs. S1–S27 is in Supplementary Data that can be found in https://osf.io/j2rh6/?view_only=8ba067c103544a188e106bd9ed9a5293. All analyses were pre-registered with COS prior to data access (https://osf.io/zshpy/?view_only=34700935c63449968221300d51b91350).

Code availability

All code to reproduce analyses are openly available in an online repository²⁶.

Received: 15 June 2024; Accepted: 26 June 2025;

Published online: 07 August 2025

References

1. Elstad, J. I. Childhood adversities and health variations among middle-aged men: A retrospective lifecourse study. *Eur. J. Public Health* **15**, 51–58 (2005).
2. Barboza et al. Adverse childhood experiences and physiological wear-and-tear in midlife: Findings from the 1958 British birth cohort. *Proc. Natl. Acad. Sci. USA* **112**, E738–E746 (2015).
3. Berkman, L. F. Social epidemiology: social determinants of health in the United States: are we losing ground? *Annu. Rev. Public Health* **30**, 27–41 (2009).
4. Slopen, N., Chen, Y., Guida, J. L., Albert, M. A. & Williams, D. R. Positive childhood experiences and ideal cardiovascular health in midlife: Associations and mediators. *Prev. Med.* **97**, 72–79 (2017).
5. Chen, Y., Kubzansky, L. D. & VanderWeele, T. J. Parental warmth and flourishing in mid-life. *Soc. Sci. Med.* **220**, 65–72 (2019).
6. Brinker, J. & Cheruvu, V. K. Social and emotional support as a protective factor against current depression among individuals with adverse childhood experiences. *Prev. Med. Rep.* **5**, 127–133 (2017).
7. Upenieks, L. & Schafer, M. H. Religious attendance and physical health in later life: a life course approach. *J. Health Soc. Behav.* **61**, 486–502 (2020).
8. Jones, G. T., Power, C. & Macfarlane, G. J. Adverse events in childhood and chronic widespread pain in adult life: results from the 1958 British Birth Cohort Study. *Pain* **143**, 92–96 (2009).
9. Power, C. et al. Life-course influences on health in British adults: effects of socio-economic position in childhood and adulthood. *Int. J. Epidemiol.* **36**, 532–539 (2007).
10. Macfarlane, G. J., Norrie, G., Atherton, K., Power, C. & Jones, G. T. The influence of socioeconomic status on the reporting of regional and widespread musculoskeletal pain: results from the 1958 British Birth Cohort Study. *Ann. Rheum. Dis.* **68**, 1591–1595 (2009).
11. Jay, M. A., Bendayan, R., Cooper, R. & Muthuri, S. G. Lifetime socioeconomic circumstances and chronic pain in later adulthood: findings from a British birth cohort study. *BMJ Open* **9**, 1–10 (2019).
12. Mustard, C. A., Kalcevic, C., Frank, J. W. & Boyle, M. Childhood and early adult predictors of risk of incident back pain: Ontario child health study 2001 follow-up. *Am. J. Epidemiol.* **162**, 779–786 (2005).
13. Goosby, B. J. Early life course pathways of adult depression and chronic pain. *J. Health Soc. Behav.* **54**, 75–91 (2013).
14. Raphael, K. G., Widom, C. S. & Lange, G. Childhood victimization and pain in adulthood: a prospective investigation. *Pain* **92**, 283–293 (2001).
15. You, D. S., Albu, S., Lisenbardt, H. & Meagher, M. W. Cumulative childhood adversity as a risk factor for common chronic pain conditions in young adults. *Pain. Med.* **20**, 486–494 (2019).
16. Stickley, A., Koyanagi, A. & Kawakami, N. Childhood adversities and adult-onset chronic pain: results from the World Mental Health Survey, Japan. *Eur. J. Pain* **19**, 1418–1427 (2015).
17. Johnson, S. B., Riley, A. W., Granger, D. A. & Riis, J. The science of early life toxic stress for pediatric practice and advocacy. *Pediatrics* **131**, 319–327 (2013).
18. Bellis, M. A., Lowey, H., Leckenby, N., Hughes, K. & Harrison, D. Adverse childhood experiences: retrospective study to determine their impact on adult health behaviours and health outcomes in a UK population. *J. Public Health* **36**, 81–91 (2014).
19. Bowlby, J. *Attachment and Loss* (Hogarth Press and the Institute of Psycho-Analysis, 1969).
20. Brown, D. W. et al. Adverse childhood experiences and the risk of premature mortality. *Am. J. Prev. Med.* **37**, 389–396 (2009).
21. Bronfenbrenner, U. *The Ecology of Human Development* (Harvard University Press, 1979).
22. VanderWeele, T. J. et al. The global flourishing study: study profile and initial results on flourishing. *Nat. Ment. Health* **3**, 636–653 (2025).
23. Crabtree, S., English, C., Johnson, B. R., Ritter, Z. & VanderWeele, T. J. Global flourishing study: questionnaire development report. *Gallup Inc.* (2021). Available at: <https://osf.io/y3t6m>.
24. Johnson, B. R. et al. The global flourishing study. Available at: <https://doi.org/10.17605/OSF.IO/3J TZ8> (2024).
25. Padgett, R. N. et al. Analytic methodology for childhood predictor analyses for Wave 1 of the Global Flourishing Study. *BMC Glob. Public Health* <https://doi.org/10.1186/s44263-025-00142-0> (2024).
26. Padgett, R. N. et al. Global flourishing study statistical analyses code. Available at: <https://doi.org/10.17605/osf.io/vbype> (2024).
27. Ritter, Z. et al. Global flourishing study methodology. *Gallup Inc.* Available at: <https://osf.io/k2s7u> (2024).
28. Lomas, T. et al. The development of the global flourishing study questionnaire: charting the evolution of a new 109-item inventory of human flourishing. *BMC Glob. Public Health* **3**, 30 (2025).
29. Padgett, R. N. et al. Survey sampling design in wave 1 of the global flourishing study. *Eur. J. Epidemiol.* **40**, 391–406 (2024).
30. Vanderweele, T. J. On the promotion of human flourishing. *Proc. Natl. Acad. Sci. USA* **114**, 8148–8156 (2017).

31. Johnson, K. A., Moon, J. W., VanderWeele, T., Schnitker, S. & Johnson, B. R. Assessing religion and spirituality in a cross-cultural sample: development of religion and spirituality items for the Global Flourishing Study. *Relig. Brain Behav.* **0**, 1–14 (2023).
32. Borenstein, M., Hedges, L. V., Higgins, J. P. T. & Rothstein, H. R. A basic introduction to fixed-effect and random-effects models for meta-analysis. *Res. Synth. Methods* **1**, 97–111 (2010).
33. Hunter, J. E. & Schmidt, F. L. Fixed effects vs. random effects meta-analysis models: Implications for cumulative research knowledge. *Int. J. Sel. Assess.* **8**, 275–292 (2000).
34. Mathur, M. B. & VanderWeele, T. J. Sensitivity analysis for publication bias in meta-analyses. *J. R. Stat. Soc. Ser. C Appl. Stat.* **69**, 1091–1119 (2020).
35. R. Core Team. A language and environment for statistical computing. *R Foundation for Statistical Computing* (2024).
36. Viechtbauer, W. Conducting meta-analyses in R with the metafor. *J. Stat. Softw.* **36**, 1–48 (2010).
37. Wilson, D. J. The harmonic mean p-value for combining dependent tests. *Proc. Natl. Acad. Sci. USA* **116**, 1195–1200 (2019).
38. Abdi, H. Bonferroni and Šidák corrections for multiple comparisons. *Encycl. Meas. Stat.* **3**, 2007 (2007).
39. VanderWeele, T. J. & Mathur, M. B. Some desirable properties of the Bonferroni Correction: Is the Bonferroni Correction really so bad? *Am. J. Epidemiol.* **188**, 617–618 (2019).
40. VanderWeele, T. J. & Ding, P. Sensitivity analysis in observational research: introducing the E-Value. *Ann. Intern. Med.* **167**, 268–274 (2017).
41. Rubin, D. B. Multiple imputation after 18+ years. *J. Am. Stat. Assoc.* **91**, 473–489 (1996).
42. Acock, A. C. Working with missing values. *J. Marriage Fam.* **67**, 1012–1028 (2005).
43. Sterne, J. A. C. et al. Multiple imputation for missing data in epidemiological and clinical research: potential and pitfalls. *BMJ* **338**, b2393 (2009).
44. van Buren, S. *Flexible Imputation of Missing Data* 2nd edn (Chapman and Hall/CRC, 2023).
45. Lut, I. et al. Health outcomes, healthcare use and development in children born into or growing up in single-parent households: a systematic review study protocol. *BMJ Open* **11**, 1–5 (2021).
46. Brown, T. T. et al. Discrimination hurts: the effect of discrimination on the development of chronic pain. *Soc. Sci. Med.* **204**, 1–8 (2018).
47. Case, A., Fertig, A. & Paxson, C. The lasting impact of childhood health and circumstance. *J. Health Econ.* **24**, 365–389 (2005).
48. Repetti, R. L., Taylor, S. E. & Seeman, T. E. Risky families: family social environments and the mental and physical health of offspring. *Psychol. Bull.* **128**, 330–366 (2002).
49. Wiech, K. & Tracey, I. The influence of negative emotions on pain: Behavioral effects and neural mechanisms. *NeuroImage* **47**, 987–994 (2009).
50. Sheffler, J. L., Piazza, J. R., Quinn, J. M., Sachs-Ericsson, N. J. & Stanley, I. H. Adverse childhood experiences and coping strategies: identifying pathways to resiliency in adulthood. *Anxiety Stress Coping* **32**, 594–609 (2019).
51. Sege, R. D. & Harper Browne, C. Responding to ACEs with HOPE: health outcomes from positive experiences. *Acad. Pediatr.* **17**, S79–S85 (2017).
52. Koenig, H. G., VanderWeele, T. & Peteet, J. R. *Handbook of Religion and Health* (Oxford University Press, 2023).
53. VanderWeele, T. J. & Li, Y. Simple sensitivity analysis for differential measurement error. *Am. J. Epidemiol.* **188**, 1823–1829 (2019).
54. Macchia, L. Pain trends and pain growth disparities, 2009–2021. *Econ. Hum. Biol.* **47**, 101200 (2022).
55. Kilburn, R. & Karoly, L. A. The economics of early childhood policy: What the dismal science has to say about investing in children. RAND Corporation. <https://doi.org/10.4324/9781315665825-91> (2008).

Acknowledgements

The Global Flourishing Study was supported by funding from the John Templeton Foundation (grant #61665), Templeton Religion Trust (#1308), Templeton World Charity Foundation (#0605), Well-Being for Planet Earth Foundation, Fetzer Institute (#4354), Well Being Trust, Paul L. Foster Family Foundation, and the David and Carol Myers Foundation.

Author contributions

L.M. conducted the data analysis, wrote the paper. C.N.O. provided helpful comments to the written drafts. T.B. provided helpful comments to the written drafts. K.S. provided code for data analysis. A.P. provided helpful comments to the written drafts. B.R.J. coordinated data collection, participated in survey design, coordinated the creation of code for analysis, and provided helpful comments to the written drafts. T.J.V. coordinated data collection, participated in survey design, coordinated the creation of code for analysis, and provided helpful comments to the written drafts.

Competing interests

Tyler VanderWeele reports consulting fees from Gloo Inc., along with shared revenue received by Harvard University in its license agreement with Gloo according to the University IP policy. All other authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s43856-025-00997-2>.

Correspondence and requests for materials should be addressed to Lucia Macchia.

Peer review information *Communications Medicine* thanks Annamaria Guolo, Peyton Presto, and the other anonymous reviewer(s) for their contribution to the peer review of this work. [Peer reviewer reports are available].

Reprints and permissions information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

© The Author(s) 2025