



City Research Online

City St George's, University of London

Citation: Bussa, M., Ambrogi, F., Edefonti, V., O'Flaherty, M., Chavez-Ugalde, Y. & Colombet, Z. (2025). Identifying the patterns of ultra-processed food consumption and their characteristics in the UK adults using the UK National Diet and Nutritional Surveys 2008/09 to 2018/19. *Public Health Nutrition*, 28(1), e149. doi: 10.1017/s1368980025100840

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/35721/>

Link to published version: <https://doi.org/10.1017/s1368980025100840>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

Identifying the patterns of ultra-processed food consumption and their characteristics in the UK adults using the UK National Diet and Nutritional Surveys 2008/09 to 2018/19

Martino Bussa^{1,2}, Federico Ambrogi², Valeria Edefonti^{2,3}, Martin O'Flaherty⁴, Yanaina Chavez-Ugalde⁵, Zoè Colombet⁴

¹Bicocca Bioinformatics, Biostatistics and Bioimaging Centre, School of Medicine and Surgery, University of Milano-Bicocca, Milan, Italy

²Department of Clinical Sciences and Community Health, University of Milan - La Statale, Milan, Italy

³Fondazione IRCCS, Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy

⁴Department of Public Health, Policy and Systems, University of Liverpool, Liverpool, United Kingdom

⁵City St George's, University of London, London, UK

Corresponding author: Zoé Colombet, Department of Public Health and Policy, University of Liverpool, Liverpool, United Kingdom. Phone number: 0151 795 4129, E-mail: zoe.colombet@liverpool.ac.uk

Short title: Ultra-Processed Food Dietary Patterns in UK Adults



This is an Accepted Manuscript for Public Health Nutrition. This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI 10.1017/S1368980025100840

Public Health Nutrition is published by Cambridge University Press on behalf of The Nutrition Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

Acknowledgements: The data supporting the results of this study were provided by the National Diet and Nutrition Survey database. We thank all investigators and participants in the study for their contribution.

Financial Support: YCU collaborated to this research whilst being a postdoctoral research associate at the Medical Research Council (MRC) to the MRC Epidemiology Unit, University of Cambridge [grant number MC_UU_00006/5]. She is now at Research Fellow at City St George's, University of London and received no funding for this study.

Conflict of Interest: None.

Authorship: The authors' responsibilities were as follows: MB, MOF and ZC designed the study, MB drafted the manuscript and performed the statistical analysis, FA, VE, ZC checked the code and analysis, all authors contributed to the data interpretation and revised each draft for important intellectual content. All authors read and approved the final manuscript.

Ethical Standards Disclosure: This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving research study participants for collection of NDNS data were provided by the Oxfordshire A Research Ethics Committee. All participants provided written informed consent to take part in NDNS. Additional ethical approval for this secondary analysis of anonymised data was not required.

Data availability: The data used in this study are derived from the National Diet and Nutrition Survey (NDNS), conducted by Public Health England (PHE) and the Food Standards Agency (FSA) in the United Kingdom. Access to NDNS data is managed by the UK Data Service (UKDS) under license from PHE and FSA. Researchers interested in accessing NDNS data should consult the UK Data Service website (<https://www.ukdataservice.ac.uk/>) for information on data access procedures and conditions.

Abstract

Objective: To identify the dietary patterns of ultra-processed food (UPF) consumption in the UK adults, and to explore their nutritional characteristics and associated demographic and socioeconomic factors.

Design: UPF-based dietary patterns were identified using weighted principal component analysis and k-means cluster analysis on UPFs intakes (identified using Nova classification) from the cross-sectional NDNS data (2008-2019). Weighted multivariable logistic regression models were employed to identify the demographic and socioeconomic factors associated with the patterns.

Setting: United Kingdom.

Participants: 8,347 adults (≥ 18 y).

Results: UPFs accounted for 54% of total energy intake in the UK adult diet. Three distinct UPF-clusters were identified, labelled as "Sweet Foods," "Fast Foods," and "Traditional Foods" based on their predominant food intakes. Older participants (>68 years) were more likely to adhere to the "Sweet Foods" pattern (OR: 2.39; 95% CI: 1.99-2.87) and less likely to be part of the "Fast Foods" pattern (OR: 0.47; 95% CI: 0.40-0.55) compared to younger individuals (< 29). Participants in lower occupations were less likely to adhere to the "Fast Foods" pattern than participants in the higher occupations (OR: 0.82; 95% CI: 0.72-0.94) while being more likely to adhere to the "Traditional Foods" pattern (OR: 1.23; 95% CI: 1.06-1.43).

Conclusions: The UK diet was dominated by UPF products. Our analysis identified three distinct UPF dietary patterns with varying nutritional quality, influenced by key demographic and social factors. These findings provide valuable insights into the determinants of UPF consumption and highlight which population groups are more likely to consume certain types of UPFs.

Abbreviations: UPF: Ultra Processed Food, NCDs: Non-Communicable Diseases, PCA: Principal Component Analysis, SEM: Standard Error of Mean, CI: Confidence Interval, OR: Odds Ratio

Introduction

The prevalence of overweight and obesity has been steadily increasing worldwide since 1980, posing a significant global challenge⁽¹⁾. In 2021, 2 in 3 adults were either overweight or obese in England⁽²⁾. A large part of Non-Communicable Diseases (NCDs) deaths was attributable to poor diets, approximately 30% in 2019⁽³⁾, suggesting the improvement of diets as a well-established public health goal. The global rise in NCDs and obesity prevalence are partly attributed to rapid changes in global food systems, coupled with the development of increasingly sedentary lifestyles⁽⁴⁾. Recent social changes, such as technological evolution, urbanization, and increased per capita income, have profoundly influenced the global food system⁽⁵⁾, driving to the adoption of “Western” dietary patterns. These patterns are characterised by high levels of saturated fats, sugars, and refined foods, and low intakes of fibre-rich foods, primarily due to the high availability of cheap, energy-dense, nutrient-poor foods⁽⁶⁾. Within this nutrition transition, increased intakes of ultra-processed foods (UPFs) and ready-to-eat meals have been observed⁽⁷⁾. Nowadays, UPFs have become the primary source of energy in most high-income countries: 48% of the total dietary energy intake in Canada⁽⁸⁾, 57.9 % in the USA⁽⁹⁾, and 56.8% in the UK⁽¹⁰⁾.

Multiple classifications exist to identify UPFs, but the most frequently used is the Nova classification which categorises all products into four progressively processed groups, based on the extent of processing, with the highest category labelled as UPFs⁽¹¹⁾. UPFs are primarily industrial formulations made from substances extracted from foods, often chemically modified, and typically contain few 'natural' ingredients. Often, UPFs contain a variety of cosmetic additives, including preservatives, stabilizers, emulsifiers, solvents, binders, bulking agents, sweeteners, thickeners, sensory enhancers, carbonating, flavours, flavour enhancers, and colours^(11–13). UPFs are often described as energy-dense, containing high levels of sugars, sodium and saturated fats, and low levels of fibres, micronutrients, and phytochemicals⁽¹⁴⁾, even if considerable variation in these levels has been observed within each category of processing⁽¹⁵⁾. UPFs play a substantial role in nutrient intake⁽¹⁶⁾ and a study published in 2023 demonstrated how healthy dietary patterns with a high diet quality score, and adequate amounts of most macro- and micronutrients can be achieved using UPFs⁽¹⁷⁾. Indeed, the range of available UPFs is broad, from pre-packaged whole meals or non-sugared flavoured water as well as sugary soft drinks and highly processed snack foods like chips and candies.

While altering food from its natural state for safety, convenience, taste or palatability seems to have been a key contributor to human development, it also seems to be a substantial threat to health⁽¹⁸⁾. Especially, the consumption of UPFs seems to be associated with cancer, cardiovascular diseases, type 2 diabetes and other NCDs⁽¹⁹⁾.

Given the significant role of UPFs in modern diets and the growing evidence of their detrimental health effects, it is crucial to understand UPF consumption and its determinants to better inform effective public health policies. As UPFs encompass a wide range of products, identifying patterns in their consumption – such as which types are consumed together – can shed light on their impact on health. Further, examining whether these patterns correlate with individual characteristics like age or socioeconomic status can provide valuable insights for targeted interventions. Therefore, this study aims to identify the dietary patterns of UPF consumption among adults in the UK, exploring their nutritional characteristics, and the associated demographic and socioeconomic factors.

Subjects and Methods

Data source

We conducted a secondary analysis of individual-level data from the UK National Diet and Nutrition Survey (NDNS) using waves 1 to 11 (2008/09-2018/19). Details of the rationale, design and methods of the survey have been described elsewhere⁻. Briefly, the NDNS is an annual, cross-sectional survey collecting information on the nutrient intakes and status of individuals living in private households in the UK^(20–23). Between 2008 and 2019, the NDNS ran continuously as a rolling programme covering adults and children aged 1.5 years and over, using consistent data collection methods. A nationally representative sample was selected each year using a multi-stage random probability design, with one adult and one child chosen randomly from each household. Demographic and socioeconomic data were collected through questionnaires, and anthropometric data were measured^(20–23). Dietary data were collected through 4-day food diaries administered on random days of the week to ensure a balanced representation of dietary intakes. Once completed, interviewers reviewed diaries with respondents to add any missing details and enhance completeness. All those who completed three or four days of dietary recording were included in the dataset, giving a sample size of 6,828 participants for waves 1-4 combined, a sample size of 2,546 participants for years 5-6 combined, a sample size of 2,723 participants for waves 7-8 and a sample size of 3,558 participants for waves 9-11.

Inclusion criteria

Individuals who took part in NDNS waves 1 to 11, aged 18 years or older at data collection, and who had completed at least three food diary days (for the 4-day food diary) were included in these analyses.

Variables of interest

Socio-demographic characteristics

We used age in groups of 10-years, sex, country (England, Northern Ireland, Scotland, and Wales), and ethnicity (white and non-white), as demographic variables, and occupational categories (higher occupations; intermediate occupations; lower occupations; small employers and own account workers; routine/semi-routine occupations; and never worked) as socioeconomic variable in our analysis.

Ultra-processed food consumption

To identify the consumption of UPFs in the NDNS dietary data, we used the Nova classification developed by Monteiro and colleagues^(11,13). According to the Nova classification, all foods and drinks can be classified into one of these four groups according to the extent and purpose of the industrial food processing they undergo: “minimally processed foods”, “processed culinary ingredients”, “processed foods”, and “ultra-processed foods”. Each food and drink in the NDNS data was classified by two independent researchers (ZC and YCU), leading to a final 97% agreement after discussion, as detailed elsewhere⁽²³⁾. For the present study, UPFs intake was estimated using the percentage of daily total energy intake (without alcohol) provided by dietary items classified in the Nova group “ultra-processed foods”, while all other items (excluding alcohol) were classified non-UPFs.

Statistical analysis

To identify the UPF dietary patterns, we used a two-step procedure, detailed in the Supplemental Material. Firstly, a weighted principal component analysis (PCA) was applied to the covariance matrix of UPF intakes from 21 food groups (in % percentage of energy intake per day). Then, a clustering procedure was applied to the first retained principal components to group participants based on similarity in UPFs intakes operationalized via PCA, using a 50% explained variance threshold. Cluster analysis was performed using the HCPC algorithm⁽²⁴⁾, which combines hierarchical and partitional clustering. To determine the

optimal number of clusters, we examined the dendrogram and various cluster validation indices starting from either the 4-PCs or the 5-PCs solutions. Given that the inclusion of the fifth PC did not enhance cluster separation or compactness, the first four PCs were retained. Principal components loadings for the 4-PCs solution were reported in Supplementary Table 1 for completeness.

Cluster analysis identified distinct groups, interpreted as dietary patterns and labelled according to their predominant food intakes. Clusters were described according to their main nutritional characteristics. ANOVA was used to compare the means of nutritional characteristics (adjusted for daily energy intake) and food groups across the UPF-based dietary patterns; the Tukey test was used for pairwise comparisons. Weighted multivariable logistic regression models were employed to associate each cluster with selected demographic and socioeconomic characteristics. Adjusted odds ratios (ORs) were reported with 95% confidence intervals (CIs). A p-value of < 0.05 was considered statistically significant. Data management and statistical analyses were performed using R software, version 4.2.3, with the FactoMiner R package⁽²⁵⁾.

Results

Of the 15,655 participants across 11 survey waves, 8,347 were adults (aged 18 and above), and all these adults had completed at least three food diary days. Consequently, 8,347 participants were included in the analysis. The only missing data were for ethnicity (0.16%) and occupation (0.53%) (Supplementary Table 4). Overall, adults reported a mean total energy intake per day of 1,761.73 kcal (SEM 0.59) and 54% of these calories were provided by UPFs (Table 1). UPFs contributed highly to carbohydrates, fats, free sugars, and sodium intake.

Cluster analysis allowed to identify three dietary patterns. These were labelled according to their intakes as “Sweet Foods”, “Fast Foods” and “Traditional Foods”, and represented 23%, 47%, and 30% of the sample, respectively.

Differences in food groups and nutritional characteristics

Table 2 presents the UPFs intakes (in % of daily energy) within the 21 food groups included in the PCA across the three identified dietary patterns. The largest cluster (3,947 subjects), labelled the “Fast Foods” pattern, included adults with, on average, the highest daily energy intake of UPFs from pre-prepared meals, homemade dishes (e.g. meat pies and pastries),

manufactured poultry, pizza, hamburgers and kebabs, chips, breakfast cereals and soft drinks compared to the other two clusters. The second cluster (1,891 subjects), labelled the “Sweet Foods” pattern, consisted of adults with the highest daily energy intake of UPFs from biscuits and other sweet baked goods, industrial desserts and homemade desserts, as well as chocolate confectionery and yoghurts. Participants in this cluster had a lower daily energy intake of UPFs from meat products, pizza, beans, chips, crisps and savoury snacks, sauces and gravies, and notably, soft drinks, compared to the other clusters. The third cluster (2,509 subjects), labelled the “Traditional Foods” pattern, reported the highest average energy intake of UPFs from bread, processed meat, beans, margarine and other spreads. Additionally, the intake of UPFs from manufactured fish dishes, sugar confectionery, chocolate confectionery, and yoghurts was lower in this cluster than in the other two.

Table 3 describes the key nutritional characteristics of each UPF-based dietary pattern and provides an overview of the nutritional profile in the overall national sample study. Notably, the “Fast Foods” pattern exhibited the lowest average carbohydrate content (118 g/day) and the lowest energy intake from UPFs compared to the other two UPF patterns, as the lowest intake of fats and a low intake of protein. Conversely, the “Sweet Foods” pattern showed higher daily intakes of total sugars from UPFs, especially free sugars, and elevated levels of fats, saturated fats, and cholesterol. Furthermore, the “Traditional Foods” pattern showed the highest average sodium content from UPFs (1719 mg/day) and the lowest intake from free sugars compared to the other two. Supplementary Table 2 provides daily percentages of energy intake from UPFs in the food groups not included in the PCA across the identified UPF dietary patterns. Supplementary Table 3 details the daily intake (g/day) of selected food groups (included or not in the PCA) within the three identified dietary patterns.

Differences in demographic and socioeconomic characteristics

Supplemental Table 4 describes the demographic and socioeconomic characteristics of the participants and their distribution across the identified UPF dietary patterns. In our sample, 59% of participants were females, 60% lived in England, 92% were Whites, 15% were engaged in higher occupations, 33% in lower occupations, 26% in semi-routine/routine occupations, and 11% were small employers and self-employed workers. The sample exhibited a balanced distribution across all age groups.

Associations between each UPF dietary pattern and demographic/socioeconomic characteristics are presented in Table 4 as ORs and corresponding 95% CI. Older participants

(age > 68 years) were more likely to adhere to the “Sweet Foods” pattern (OR: 2.39; 95% CI: 1.99-2.87) and less likely to adhere to the “Fast Foods” pattern (OR: 0.47; 95% CI: 0.40-0.55) compared to younger individuals (age < 29 years). Women were also more likely to adhere to the “Sweet Foods” patterns than men (OR: 1.40; 95% CI: 1.26-1.56), while being less likely to adhere to the “Traditional Foods” (OR: 0.75; 95% CI: 0.69-0.83). Regarding the country, participants residing in Northern Ireland were more likely to adhere to the “Sweet Foods” (OR: 1.28; 95% CI: 1.10-1.50) and less likely to adhere to the “Fast Foods” pattern (OR: 0.76; 95% CI: 0.67-0.88) compared to those in England. Similarly, in Scotland, participants were less likely to adhere to the “Fast Foods” pattern (OR: 0.84; 95% CI: 0.74-0.96) than in England.

Regarding occupation, few associations were observed, but participants in lower and semiroutine/routine occupation categories were less likely to adhere to the “Fast Foods” pattern than participants in the higher occupation category (OR: 0.82; 95% CI: 0.72-0.94 and OR: 0.75; 95% CI: 0.65-0.86, respectively), while being more likely to adhere to the “Traditional Foods” pattern (OR: 1.23; 95% CI: 1.06-1.43 and OR: 1.56; 95% IC: 1.33-1.82, respectively). Interestingly, the participants who never worked and the retired ones were more likely to adhere to the “Traditional Foods” pattern than participants in the higher occupations category (OR: 1.40; 95% CI: 1.08-1.82).

Discussion

This study represents the first attempt to identify and describe dietary patterns of UPF intake among the adult population in the UK. Using a large, nationally representative sample, we identified three main UPF dietary patterns based on the type of ultra-processed products consumed: “Fast Foods”, “Sweet Foods”, and “Traditional Foods”, which differ by their nutritional intakes and the individual characteristics of their members.

In our sample, over half of the daily calories (54%) consumed by the UK adult population between 2008 and 2019, came from UPFs. These findings are consistent with previous analyses of the same NDNS data and other studies conducted in different countries^(9,15,26). For instance, Rauber and colleagues⁽¹⁰⁾ found that UPFs contributed to 56.8% of total energy intake and 64.7% of total free sugars in the UK diet based on NDNS 2008-2014 data. Additionally, in the NDNS 2008-2012, an average of 53% of total energy intake was provided by UPFs, while minimally processed foods contributed to an average of 28%⁽¹⁸⁾.

Moreover, household food purchase surveys conducted in 2008 revealed that UPFs constituted 63% of energy in the UK⁽¹²⁾.

But the number of products that fall into the category of UPFs according to the Nova classification system is high, and the type of ultra-processed products is broad. Within our sample, we have identified three UPF dietary patterns based on the main type of UPFs consumed. Almost half of the adults in our sample tend to eat UPFs we can classify as fast foods (e.g. prepared meals, pizza, hamburgers and kebabs, from the “Fast Foods” pattern). In contrast, 1 in 2 adults ate traditional UPFs (e.g. bread, processed meat, beans, margarine and other spreads, from the “Traditional Foods” pattern). Finally, the other adults tend to eat “sweet” UPFs, such as biscuits, desserts, chocolate confectionary and yoghurts, from the “Sweet Foods” pattern.

The high consumption of UPFs observed in the UK adult diet raises important nutritional concerns. Indeed, the diet observed in our sample exceeded recommended intake levels for free sugars, saturated fat, and sodium, while fiber intake fell below the recommended levels, according to WHO recommendations^(27–30). And UPFs seem to highly contribute to these unhealthy nutritional intakes. This is consistent with previous studies, which have found a high salt, free sugar and fat intakes associated with UPFs^(8,10,15). UPF products accounted for more than 70% of the daily sodium intake in our sample and 64% of the daily free sugar intake. UPF foods had almost 30% more fat than the diet fraction made up of non-UPF foods.

Interestingly, all the dietary patterns identified in our sample exhibited elevated fats, free sugars, and sodium levels, even if they varied in nutritional content. Specifically, the “Sweet Foods” patterns- demonstrated significantly higher levels of fats, free sugars, and energy content than the other two. In contrast, the “Traditional Foods” dietary pattern exhibited the highest daily sodium intake. These findings corroborate existing research indicating that the specific types of UPFs consumed can significantly impact overall nutritional intake. Some UPFs may offer a comparatively healthier nutritional profile than others, as previously demonstrated⁽¹⁷⁾. However, it is important to note that diets heavily reliant on UPFs, regardless of individual product choices, are still contributing to a high consumption of fat, salt, sugar and energy, which can be detrimental to health.

An interesting finding is the variation in UPF dietary patterns among adults in the UK, based on age, sex and occupation. Our results highlighted that older individuals were more likely to consume sweet UPFs and fewer ready-to-eat foods (“Fast foods”) than younger individuals.

This variation may be attributed to age-related declines in taste and smell sensitivity^(31–33), as well as a slower perception of sweetness in older compared to younger adults⁽³⁴⁾. These changes in sweet taste perception can influence food preferences⁽³⁵⁾. Additionally, for older adults, particularly those who are homebound, factors such as sensory appeal (or aesthetic appeal), convenience, and price were identified as playing a crucial role in food selection^(36,37). Conversely, younger participants showed a preference for “Fast Food” UPFs, a trend consistent with other national surveys that identified the youngest adult age group as having the highest consumption of junk foods^(18,26,38–40). Early adulthood is a period of life transition during which changes such as leaving the parental home or moving from school to further education or paid employment can lead to the disruption of pre-existing eating habits and dietary behaviours. Notably, leaving home and finishing school are associated with negative dietary changes⁽⁴¹⁾. These transitions may present critical opportunities for implementing effective interventions targeting diet and obesity. Similarly, women exhibited a preference for sweet UPFs compared to men, which is coherent with the literature, e.g. study showing women having a strong liking for the fat-and-sweet sensation⁽⁴²⁾, study reporting more craving for sweet foods⁽⁴³⁾ or reporting that women find sweet foods more pleasant⁽⁴⁴⁾.

Finally, occupation is also a determinant of food choice, and its association with higher UPF intake varies between countries⁽⁴⁵⁾. In our study, we observed an association between the type of UPFs chosen and occupation, with adults in low, routine and semi-routine occupations showing a reduced preference for the “Fast Foods” UPF pattern compared to those in higher-level occupations, instead favouring the “Traditional Foods” UPF pattern. These findings underline the importance of considering age, gender, and occupation differences when formulating dietary policies to promote healthier and more sustainable dietary choices, as well as understanding how other factors, such as marketing or price, influence choice in these groups.

Our study has some strengths and limitations. One key strength is the use of data from the NDNS, a large and nationally representative sample of the UK adult population, applying weighting to reduce sampling and non-response bias. Additionally, food diaries were used to collect dietary data, which are among the most comprehensive methods for assessing dietary intake⁽¹⁰⁾. Food diaries provide detailed accounts of consumed foods and are less prone to recall bias compared to other dietary assessment tools. However, there are also limitations to consider. The dietary data were self-reported, which may introduce self-reporting bias, including the under-reporting of certain items (notably, unhealthy foods). There is also a

potential for misclassification into the Nova classification due to the lack of detailed ingredient information in some foods in NDNS. Furthermore, multivariate methods come with inherent limitations, as they require subjective decisions regarding the number of components extracted, the choice of clusters, and their interpretation and naming. Such subjectivity can impact both the patterns identified and their subsequent interpretation, a limitation commonly acknowledged in studies employing these techniques^(46,47). It is also important to note that this is a cross-sectional study, so causal effects cannot be inferred. Also, as UPF consumption seems to have increased between the study period (2008-2018) and is likely to have increased since 2018, it will be interesting to evaluate if the UPF dietary patterns are changing with time. Additionally, the data were collected before the COVID-19 pandemic and recent inflation, a period which may have significantly altered dietary patterns in the UK.

Whether UPF-based diets are harmful to health simply due to poor nutritional quality, or if the nature and extent of the processing itself have health consequences, remains an ongoing debate that warrants continued research^(19,48,49). Based solely on their nutritional qualities, however, the three identified UPF patterns do not provide a sufficient causal basis to inform food policies. They do, however, provide good evidence into dietary patterns and socio-demographic and economic determinants of UPF consumption. Our findings suggest that policy efforts should focus on reducing overall UPF intake, while also improving the relative nutritional quality of UPFs - for instance, through reformulation - and promoting the intake of fresh and minimally processed foods.

Conclusion

This study confirms the continued dominance of UPFs in UK diets. Our analysis identified three distinct dietary patterns with varying nutritional quality, shaped by key demographic and socioeconomic factors. These findings provide valuable insights into the determinants of UPF consumption and highlight which population groups are more likely to consume certain types of UPFs.

References

1. Ng M, Fleming T, Robinson M, et al. (2014) Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* **384**, 766–781.
2. NHS (2022) Health Survey for England, 2021 part 1. <https://digital.nhs.uk/data-and-information/publications/statistical/health-survey-for-england/2021>.
3. Qiao J, Lin X, Wu Y, et al. (2022) Global burden of non-communicable diseases attributable to dietary risks in 1990-2019. *J Hum Nutr Diet* **35**, 202–213.
4. Popkin BM (2006) Global nutrition dynamics: the world is shifting rapidly toward a diet linked with noncommunicable diseases. *Am. J. Clin. Nutr.* **84**, 289–298.
5. Popkin BM (2015) Nutrition Transition and the Global Diabetes Epidemic. *Curr Diab Rep* **15**, 64.
6. Popkin BM, Adair LS & Ng SW (2012) Global nutrition transition and the pandemic of obesity in developing countries. *Nutr. Rev.* **70**, 3–21.
7. Tapsell LC, Neale EP, Satija A, et al. (2016) Foods, Nutrients, and Dietary Patterns: Interconnections and Implications for Dietary Guidelines. *Adv Nutr* **7**, 445–454.
8. Moubarac J-C, Batal M, Louzada ML, et al. (2017) Consumption of ultra-processed foods predicts diet quality in Canada. *Appetite* **108**, 512–520.
9. Martínez Steele E, Baraldi LG, Louzada ML da C, et al. (2016) Ultra-processed foods and added sugars in the US diet: evidence from a nationally representative cross-sectional study. *BMJ Open* **6**, e009892.
10. Rauber F, Louzada ML da C, Steele EM, et al. (2019) Ultra-processed foods and excessive free sugar intake in the UK: a nationally representative cross-sectional study. *BMJ Open* **9**, e027546. British Medical Journal Publishing Group.
11. Monteiro CA, Cannon G, Levy RB, et al. (2019) Ultra-processed foods: what they are and how to identify them. *Public Health Nutr* **22**, 936–941.

12. Moubarac J-C, Claro RM, Baraldi LG, et al. (2013) International differences in cost and consumption of ready-to-consume food and drink products: United Kingdom and Brazil, 2008–2009. *Global Public Health* **8**, 845–856. Taylor & Francis.
13. Monteiro CA, Cannon G, Moubarac J-C, et al. (2018) The UN Decade of Nutrition, the NOVA food classification and the trouble with ultra-processing. *Public Health Nutr* **21**, 5–17.
14. Moodie R, Stuckler D, Monteiro C, et al. (2013) Profits and pandemics: prevention of harmful effects of tobacco, alcohol, and ultra-processed food and drink industries. *Lancet* **381**, 670–679.
15. Poti JM, Mendez MA, Ng SW, et al. (2015) Is the degree of food processing and convenience linked with the nutritional quality of foods purchased by US households? *Am J Clin Nutr* **101**, 1251–1262.
16. Slimani N, Deharveng G, Southgate D a. T, et al. (2009) Contribution of highly industrially processed foods to the nutrient intakes and patterns of middle-aged populations in the European Prospective Investigation into Cancer and Nutrition study. *Eur J Clin Nutr* **63 Suppl 4**, S206-225.
17. Hess JM, Comeau ME, Casperson S, et al. (2023) Dietary Guidelines Meet NOVA: Developing a Menu for A Healthy Dietary Pattern Using Ultra-Processed Foods. *J Nutr* **153**, 2472–2481.
18. Adams J & White M (2015) Characterisation of UK diets according to degree of food processing and associations with socio-demographics and obesity: cross-sectional analysis of UK National Diet and Nutrition Survey (2008–12). *International Journal of Behavioral Nutrition and Physical Activity* **12**, 160.
19. Lane MM, Gamage E, Du S, et al. (2024) Ultra-processed food exposure and adverse health outcomes: umbrella review of epidemiological meta-analyses. *BMJ* **384**, e077310. British Medical Journal Publishing Group.
20. Bates B, Collins D, Jones K, et al. (2020) *National Diet and Nutrition Survey: rolling programme years 9 to 11 (2016/2017 to 2018/2019)*. 29.

21. Roberts C, Steer T, Maplethorpe N, et al. (2018) *National Diet and Nutrition Survey. Results from Years 7 and 8 (combined) of the Rolling Programme (2014/2015 to 2015/2016)*. .
22. Bates B, Lennox A, Prentice A, et al. (2014) *National Diet and Nutrition Survey. Results from Years 1, 2, 3 and 4 (combined) of the Rolling Programme (2008/2009 – 2011/2012)*. .
23. Chavez-Ugalde IY, de Vocht F, Jago R, et al. (2024) Ultra-processed food consumption in UK adolescents: distribution, trends, and sociodemographic correlates using the National Diet and Nutrition Survey 2008/09 to 2018/19. *Eur J Nutr*.
24. Husson F, Josse J & Pagès J (2010) Principal Component Methods Hierarchical Clustering Partitional Clustering: Why Would We Need to Choose for Visualizing Data? .
25. Lê S, Josse J & Husson F (2008) FactoMineR: An R Package for Multivariate Analysis. *Journal of Statistical Software* **25**, 1–18.
26. Nardocci M, Leclerc B-S, Louzada M-L, et al. (2019) Consumption of ultra-processed foods and obesity in Canada. *Can J Public Health* **110**, 4–14.
27. WHO (2023) *Total fat intake for the prevention of unhealthy weight gain in adults and children: WHO guideline*. Geneva, Switzerland: World Health Organization.
28. Schenker S (2012) UK recommendations for dietary fat: should they be reassessed in light of the recent joint FAO/WHO recommendations? *Nutrition Bulletin* **37**, 37–46.
29. WHO (2015) *Guideline: sugars intake for adults and children*. Geneva, Switzerland: World Health Organization.
30. WHO (2012) *Guideline: sodium intake for adults and children*. Geneva, Switzerland: World Health Organization.
31. Westenhoefer J (2005) Age and gender dependent profile of food choice. *Forum Nutr*, 44–51.
32. Schiffman SS (1997) Taste and smell losses in normal aging and disease. *JAMA* **278**, 1357–1362.

33. Methven L, Allen VJ, Withers CA, et al. (2012) Ageing and taste. *Proc Nutr Soc* **71**, 556–565.
34. Wada H, Matsumoto H, Takagiwa M, et al. (2024) Differences in time-intensity sensory profiles of sweet taste intensity of glucose between older and young adults. *Front Nutr* **11**, 1273055.
35. Sergi G, Bano G, Pizzato S, et al. (2017) Taste loss in the elderly: Possible implications for dietary habits. *Crit Rev Food Sci Nutr* **57**, 3684–3689.
36. Locher JL, Ritchie CS, Roth DL, et al. (2009) Food choice among homebound older adults: motivations and perceived barriers. *J Nutr Health Aging* **13**, 659–664.
37. Kamphuis CBM, de Bekker-Grob EW & van Lenthe FJ (2015) Factors affecting food choices of older adults from high and low socioeconomic groups: a discrete choice experiment. *Am J Clin Nutr* **101**, 768–774.
38. Marchese L, Livingstone KM, Woods JL, et al. (2021) Ultra-processed food consumption, socio-demographics and diet quality in Australian adults. *Public Health Nutr*, 1–11.
39. Khandpur N, Cediel G, Obando DA, et al. (2020) Sociodemographic factors associated with the consumption of ultra-processed foods in Colombia. *Rev Saude Publica* **54**, 19.
40. Ahmed F, Al-Radhwan L, Al-Azmi G, et al. (2014) Association between Stress and Dietary Behaviours among Undergraduate Students in Kuwait Gender Differences. *Journal of Nutrition and Health Sciences* **1**, 1. Annex Publishers.
41. Winpenny EM, van Sluijs EMF, White M, et al. (2018) Changes in diet through adolescence and early adulthood: longitudinal trajectories and association with key life transitions. *Int J Behav Nutr Phys Act* **15**, 86.
42. Lampuré A, Deglaire A, Schlich P, et al. (2014) Liking for fat is associated with sociodemographic, psychological, lifestyle and health characteristics. *Br. J. Nutr.* **112**, 1353–1363.

43. Hallam J, Boswell RG, DeVito EE, et al. (2016) Gender-related Differences in Food Craving and Obesity. *Yale J Biol Med* **89**, 161–173.
44. Grogan SC, Bell R & Conner M (1997) Eating sweet snacks: gender differences in attitudes and behaviour. *Appetite* **28**, 19–31.
45. Dicken SJ, Qamar S & Batterham RL (2024) Who consumes ultra-processed food? A systematic review of sociodemographic determinants of ultra-processed food consumption from nationally representative samples. *Nutr Res Rev* **37**, 416–456.
46. Hearty AP & Gibney MJ (2009) Comparison of cluster and principal component analysis techniques to derive dietary patterns in Irish adults. *Br J Nutr* **101**, 598–608.
47. Newby PK & Tucker KL (2004) Empirically derived eating patterns using factor or cluster analysis: a review. *Nutr. Rev.* **62**, 177–203.
48. Dicken SJ & Batterham RL (2021) The Role of Diet Quality in Mediating the Association between Ultra-Processed Food Intake, Obesity and Health-Related Outcomes: A Review of Prospective Cohort Studies. *Nutrients* **14**, 23.
49. Martini D, Godos J, Bonaccio M, et al. (2021) Ultra-Processed Foods and Nutritional Dietary Profile: A Meta-Analysis of Nationally Representative Samples. *Nutrients* **13**, 3390. Multidisciplinary Digital Publishing Institute.

Table 1. Daily intakes of selected nutritional characteristics in the adult population in the UK from NDNS 2009-19, and associated WHO recommendations (n=8347)

	Total				UPF _s				Non-UPF _s				WHO recommendations
	Mean	± SEM	% of total energy	of	Mean	± SEM	% of total energy	of	Mean	± SEM	% of total energy	of	
Total intake (g/day) ¹	2699.80	± 5.61	-		671.32	± 2.07	-		2028.48	± 3.60	-		-
Energy intake (kcal/day)	1761.73	± 0.59	-		947.98	± 0.73	53.81		813.75	± 0.89	46.19		-
Carbohydrates (g/day) ¹	213.30	± 0.68	48.43		130.48	± 0.47	29.63		82.82	± 0.23	18.80		40-70% of total energy intake
<i>Total sugars</i> (g/day) ¹	90.28	± 0.33	20.50		44.00	± 0.19	9.99		46.28	± 0.15	10.51		-
<i>Other sugars</i> (g/day) ¹	1.96	± 0.01	0.45		1.28	± 0.004	0.29		0.68	± 0.002	0.16		-
<i>Free sugars</i> (g/day) ¹	53.95	± 0.26	12.25		34.69	± 0.17	7.88		19.26	± 0.10	4.37		Less than 10% of total energy intake ³
Proteins (g/day) ¹	71.73	± 0.19	16.29		30.92	± 0.10	7.02		40.81	± 0.10	9.27		The Recommended Dietary Allowance of protein of 0.83 g per kg body weight per day ⁴
Fats (g/day) ¹	66.23	± 0.25	33.83		37.12	± 0.15	18.96		29.11	± 0.10	14.87		Not exceed 30% of total energy intake
<i>Saturated fatty acids</i> (g/day) ¹	24.69	± 0.09	12.61		12.73	± 0.05	6.50		11.96	± 0.04	6.11		Less than 10% of total energy intake
<i>Cis monounsaturated fatty acids</i>	24.16	± 0.09	12.34		14.28	± 0.06	7.30		9.88	± 0.03	5.04		15-20% of total energy intake

	<i>(g/day)</i> ¹						
<i>Cis-n3 fatty acids (g/day)</i> ¹	1.96 ± 0.01	1.00	1.10 ± 0.56	0.004	0.86 ± 0.44	0.002	0.5-2% of total energy intake
<i>Cis-n6 fatty acids (g/day)</i> ¹	9.34 ± 0.03	4.77	5.98 ± 3.05	0.02	3.36 ± 0.01	1.72	2.5-9% of total energy intake
<i>Trans fatty acids (g/day)</i> ¹	1.08 ± 0.55	0.0004	0.46 ± 0.23	0.002	0.62 ± 0.32	0.002	Less than 1% of total energy intake
	Total daily nutrient intake		UPF _s	% from UPFs	Non-UPF _s	% from non-UPFs	WHO
Cholesterol (mg/day) ¹	243.16 ± 0.69		63.74 ± 0.24	± 26.21	179.42 ± 0.46	± 73.79	-
AOAC fibres (g/day) ¹	18.09 ± 0.04		9.90 ± 0.03	± 54.73	8.19 ± 0.02	45.27	30 g per day
Sodium (g/day) ¹	2.06 ± 0.01		1.50 ± 0.01	± 72.82	0.56 ± 0.002	± 27.18	Less than 2 g per day
Water (g/day) ¹	2307.23 ± 4.37		460.60 ± 1.35	± 19.96	1846.63 ± 3.03	± 80.04	-

Abbreviation: SEM: standard error of mean; UPFs: Ultra-processed foods; WHO: World Health Organisation.

¹Adjusted for daily total energy intake

²AOAC Fiber: Dietary fibre measured using methods from the Association of Official Analytical Chemists (AOAC)

³Ideally less than 5% of total energy intake for additional health benefits

⁴For a healthy adult with minimal physical activity

Table 2. Daily intake of UPF in the 21 food groups included in the principal component analysis across UPF dietary patterns and in the overall national sample of the adult population in the UK from NDNS 2009-19 ($n=8347$)

	All	Sweet Foods	Fast Foods	Traditional Foods
n (%)		1891 (22.65)	3947 (47.29)	2509 (30.06)
Intake of UPF in the food groups (% of energy/day) ¹				
Industrial desserts	1.46 ± 0.003	1.81 ± 0.065 ^{a,b}	1.54 ± 0.044 ^c	1.05 ± 0.056
Dessert homemade	0.46 ± 0.001	0.53 ± 0.036 ^b	0.51 ± 0.025 ^c	0.33 ± 0.031
Pre-prepared meals	3.70 ± 0.003	3.34 ± 0.122 ^a	3.99 ± 0.084 ^c	3.53 ± 0.105
Homemade dishes	2.26 ± 0.002	2.15 ± 0.077 ^a	2.45 ± 0.053 ^c	2.03 ± 0.067
Biscuits and other sweet baked goods	6.04 ± 0.053	15.02 ± 0.091 ^{a,b}	3.14 ± 0.063 ^c	3.84 ± 0.079
Manufactured poultry	1.38 ± 0.002	1.12 ± 0.075 ^a	1.57 ± 0.052 ^c	1.28 ± 0.065
Processed meat	2.81 ± 0.005	2.24 ± 0.086 ^{a,b}	2.65 ± 0.059 ^c	3.47 ± 0.074
Bread	11.33 ± 0.055	9.00 ± 0.101 ^{a,b}	7.57 ± 0.070 ^c	18.98 ± 0.087
Manufactured fish dishes	0.41 ± 0.001	0.42 ± 0.035	0.45 ± 0.024 ^c	0.33 ± 0.031
Sauces, dressing and gravies	1.74 ± 0.002	1.41 ± 0.057 ^{a,b}	1.86 ± 0.039	1.79 ± 0.049
Pizza	1.56 ± 0.004	1.19 ± 0.097 ^a	1.93 ± 0.067 ^c	1.23 ± 0.084
Hamburgers and kebabs	0.68 ± 0.002	0.57 ± 0.052 ^a	0.82 ± 0.035 ^c	0.54 ± 0.044

Beans	0.78	±	0.64 ± 0.039 ^b	0.74 ± 0.027 ^c	0.96 ± 0.033
	0.001				
Breakfast cereals	3.38	±	3.40	±	3.82 ± 0.064 ^c
	0.005		0.093 ^{a,b}		2.68 ± 0.080
Chocolate confectionary	2.03	±	2.37 ± 0.081 ^b	2.23 ± 0.055 ^c	1.48 ± 0.069
	0.004				
Crisps and savoury snacks	1.75	±	1.58 ± 0.066 ^b	1.74 ± 0.045	1.89 ± 0.057
	0.001				
Soft drink	2.06	±	1.51 ± 0.091 ^a	2.55 ± 0.062 ^c	1.72 ± 0.078
	0.005				
Sugar confectionary	0.33	±	0.40 ± 0.033 ^b	0.35 ± 0.022 ^c	0.24 ± 0.028
	0.001				
Yoghourt	1.15	±	1.33 ± 0.054 ^b	1.28 ± 0.037 ^c	0.81 ± 0.047
	0.002				
Margarine and other spreads	1.86	±	1.69	±	1.24 ± 0.037 ^c
	0.008		0.054 ^{a,b}		2.98 ± 0.047
Chips	2.20	±	1.87 ± 0.085 ^a	2.50 ± 0.059 ^c	2.00 ± 0.074
	0.003				

Values are presented as mean ± standard error of the mean (SEM)

¹ 21 food groups used in the weighted principal component analysis. All *p* values < 0.01

P values were adjusted for multiple comparisons according to the False discovery rate (FDR) method

Contrasts (Tukey test): a: group Sweet Foods vs group Fast Foods *p* < 0.05; b: group Sweet Foods vs group Traditional Foods *p* < 0.05; c: group Fast Foods vs group Traditional Foods *p* < 0.05

Table 3. Selected nutritional characteristics across UPF dietary patterns and in the overall national sample of the adult population in the UK from NDNS 2009-19 (n= 8347)

	All	Sweet Foods	Fast Foods	Traditional Foods
n (%)	8347	1891 (22.65)	3947 (47.29)	2509 (30.06)
Total UPF intake (g/day) ¹	671.32 ± 2.07	648.00 ± 9.24 ^a	679.00 ± 6.34	677.00 ± 7.98
Energy intake from UPF (kcal/day)	947.98 ± 0.73	1057.00 ± 9.68 ^{a,b}	888.00 ± 6.65 ^c	961.00 ± 8.34
Carbohydrates (g/day) ¹	130.48 ± 0.48	143.00 ± 0.97 ^a	118.00 ± 0.67 ^c	141.00 ± 0.84
<i>Total sugar (g/day)¹</i>	44.00 ± 0.19	52.70 ± 0.61 ^{a,b}	43.40 ± 0.42 ^c	38.40 ± 0.53
<i>Other sugar (g/day)¹</i>	1.28 ± 0.004	1.49 ± 0.04 ^{a,b}	1.35 ± 0.03 ^c	1.03 ± 0.04
<i>Free sugar (g/day)¹</i>	34.69 ± 0.17	41.70 ± 0.57 ^{a,b}	34.60 ± 0.39 ^c	29.50 ± 0.50
Protein (g/day) ¹	30.92 ± 0.10	30.10 ± 0.25 ^b	29.40 ± 0.17 ^c	34.00 ± 0.22
Fat (g/day) ¹	37.12 ± 0.15	40.90 ± 0.31 ^{a,b}	34.80 ± 0.21 ^c	37.90 ± 0.27
<i>Saturated fatty acids (g/day)¹</i>	12.73 ± 0.05	15.20 ± 0.12 ^{a,b}	11.90 ± 0.08 ^c	12.30 ± 0.11
<i>Cis monounsaturated fatty acids (g/day)¹</i>	14.28 ± 0.06	15.30 ± 0.14 ^{a,b}	13.50 ± 0.09 ^c	14.70 ± 0.12
<i>Cis-n3 fatty acids (g/day)¹</i>	1.10 ± 0.004	1.11 ± 0.01 ^{a,b}	1.03 ± 0.01 ^c	1.19 ± 0.01
<i>Cis-n6 fatty acids (g/day)¹</i>	5.98 ± 0.02	6.03 ± 0.06 ^{a,b}	5.59 ± 0.04 ^c	6.55 ± 0.05
<i>Trans fatty acids (g/day)¹</i>	0.46 ± 0.002	0.52 ± 0.01 ^{a,b}	0.42 ± 0.01 ^c	0.46 ± 0.01
Cholesterol (mg/day) ¹	63.74 ± 0.24	69.80 ± 0.90 ^{a,b}	62.00 ± 0.62	62.00 ± 0.77
AOAC ² fiber (g/day) ¹	9.90 ± 0.03	10.24 ± 0.09 ^{a,b}	8.92 ± 0.06 ^c	11.20 ± 0.08
Sodium (mg/day) ¹	1503.22 ± 4.92	1441.00 ± 12.41 ^{a,b}	1396.00 ± 8.52 ^c	1719.00 ± 10.72
Water (g/day) ¹	460.60 ± 1.35	422.00 ± 8.52 ^{a,b}	485.00 ± 5.85 ^c	451.00 ± 7.36

Values are presented as mean ± standard error of mean (SEM)

¹Adjusted for total daily energy intake

² AOAC Fiber: Dietary fiber measured using methods from the Association of Official Analytical Chemists (AOAC)

All *p* values <0.01; *P* values were adjusted for multiple comparisons according to the False Discovery Rate (FDR) method

Contrasts (Tukey test): a: group Sweet Foods vs group Fast Foods *p* <0.05; b: group Sweet Foods vs group Traditional Foods *p* <0.05; c: group Fast Foods vs group Traditional Foods *p* <0.05

Table 4. Associations between demographic and socioeconomic characteristics and UPF dietary patterns in a sample of the adult population in the UK from NDNS 2009-2019 (n=8347)¹

	Sweet Foods vs other patterns			Fast Foods vs other patterns			Traditional Foods vs other patterns		
	OR	95%CI	p-value ²	O R	95%CI	p- value ²	OR	95%CI	p- value ²
Age (years)									
18-28	Re f.			Re f.			Re f.		
29-38	1.0 5	[0.86- 1.29]	0.690	0.8 5	[0.73- 0.99]	0.056	1.1 8	[1.0- 1.40]	0.108
39-48	1.2 9	[1.07- 1.56]	0.020	0.7 7	[0.66- 0.89]	0.001	1.1 4	[0.96- 1.34]	0.182
49-58	1.2 9	[1.06- 1.56]	0.020	0.8 2	[0.71- 0.96]	0.022	1.0 5	[0.89- 1.24]	0.589
59-68	1.3 0	[1.06- 1.58]	0.020	0.7 4	[0.64- 0.87]	0.001	1.1 7	[0.99- 1.39]	0.120
>68	2.3 9	[1.99- 2.87]	<0.001	0.4 7	[0.40- 0.55]	<0.001	1.1 3	[0.96- 1.33]	0.200
Sex									
Men	Re f.			Re f.			Re f.		
Women	1.4 0	[1.26- 1.56]	<0.001	1.0 1	[0.93- 1.11]	0.791	0.7 5	[0.69- 0.83]	<0.001
Country									
England	Re f.			Re f.			Re f.		
Northern Ireland	1.2 8	[1.10- 1.50]	0.008	0.7 6	[0.67- 0.88]	0.0004	1.1 1	[0.96- 1.28]	0.202
Scotland	1.1 7	[1.01- 1.37]	0.076	0.8 4	[0.74- 0.96]	0.022	1.0 7	[0.92- 1.23]	0.413
Wales	0.7 7	[0.65- 0.91]	0.009	0.9 8	[0.86- 1.12]	0.791	1.2 4	[1.08- 1.43]	0.010

Ethnicity	Re			Re			Re		
White	f.			f.			f.		
Non-white	0.9	[0.81-	0.954	0.9	[0.78-	0.442	1.1	[0.92-	0.341
	9	1.22]		2	1.09]		0	1.31]	
Occupational categories									
Higher occupations	Re			Re			Re		
	f.			f.			f.		
Intermediate occupation	0.9	[0.73-	0.417	0.9	[0.78-	0.501	1.1	[0.97—	0.142
	0	1.12]		3	1.11]		9	1.45]	
Lower occupation	1.0	[0.88-	0.700	0.8	[0.72-	0.011	1.2	[1.06-	0.024
	4	1.22]		2	0.94]		3	1.43]	
Small employers and own account workers	0.8	[0.71-	0.279	0.9	[0.77-	0.442	1.2	[1.03-	0.061
	7	1.08]		2	1.09]		4	1.50]	
Semiroutine/routine occupation	0.8	[0.75-	0.244	0.7	[0.65-	0.0002	1.5	[1.33-	<0.00
	9	1.05]		5	0.86]	4	6	1.82]	1
Never worked and other	0.7	[0.57-	0.173	0.8	[0.70-	0.442	1.4	[1.08-	0.033
	8	1.06]		9	1.14]		0	1.82]	

95% CI: 95% confidence interval; OR= Odds ratio

¹ Weighted multivariable logistic regression models adjusted for age, sex, country, ethnicity and occupational categories

² P values were adjusted for multiple comparison according to the False Discovery Rate method (FDR)