



City Research Online

City St George's, University of London

Citation: Bennett-Tennenhaus, R., Goodbody, I., Letz, J. C. & Shah, A. (2026). Tensor extriangulated categories. *Journal of Algebra*, 685, pp. 361-405. doi: 10.1016/j.jalgebra.2025.07.041

This is the published version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/35911/>

Link to published version: <https://doi.org/10.1016/j.jalgebra.2025.07.041>


Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).



Contents lists available at ScienceDirect

Journal of Algebra

journal homepage: www.elsevier.com/locate/jalgebra



Research Paper

Tensor extriangulated categories



Raphael Bennett-Tennenhaus^{a,*}, Isambard Goodbody^b,
Janina C. Letz^a, Amit Shah^c

^a Faculty of Mathematics, Bielefeld University, PO Box 100 131, 33501 Bielefeld, Germany

^b School of Mathematics and Statistics, University of Glasgow, University Place, Glasgow G12 8QQ, United Kingdom

^c Department of Mathematics, Aarhus University, Ny Munkegade 118, 8000 Aarhus, Denmark

ARTICLE INFO

Article history:

Received 18 March 2025
Available online 7 August 2025
Communicated by Volodymyr Mazorchuk

MSC:

primary 18M05
secondary 18E05, 18G80, 18A23, 18G99

Keywords:

Extriangulated category
Higher extensions
Cup product
Monoidal category
Tensor triangular geometry

ABSTRACT

A tensor extriangulated category is an extriangulated category with a symmetric monoidal structure that is compatible with the extriangulated structure. To this end we define a notion of a bixtriangulated functor $\mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$, with compatibility conditions between the components. We have two versions of compatibility conditions, the stronger depending on the higher extensions of the extriangulated categories. We give many examples of tensor extriangulated categories. Finally, we generalise Balmer’s classification of thick tensor ideals to tensor extriangulated categories.

© 2025 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Contents

1. Introduction 362

* Corresponding author.

E-mail addresses: raphaelbennetttennenhaus@gmail.com (R. Bennett-Tennenhaus), isambard.goodbody@glasgow.ac.uk (I. Goodbody), jletz@math.uni-bielefeld.de (J.C. Letz), a.shah1728@gmail.com (A. Shah).

2. Higher extensions in extriangulated categories	363
3. Bixtriangulated functors	371
4. Tensor extriangulated categories	385
5. Examples of tensor extriangulated categories	388
6. Classification of tensor ideals	395
Acknowledgment	403
Data availability	403
References	403

1. Introduction

Monoidal categories play an important role in algebra, representation theory and various other areas. The module category of a commutative ring and of a cocommutative Hopf algebra over a field naturally possess a monoidal structure. These give rise to tensor triangulated categories and finite tensor categories, which have become their own areas of study. The symmetric monoidal structure is at the heart of tensor triangular geometry, allowing one to attach a topological space to a tensor triangulated category; see [1]. Further, a monoidal category is the natural setting for the categorical notion of (co)algebra objects.

The aim of this paper is to define a compatibility condition of a monoidal structure with an extriangulated structure. Extriangulated categories were introduced by Nakaoka and Palu [51], and provide a generalisation of both exact and triangulated categories. An extriangulated structure is given by a class of extriangles, generalising exact sequences and exact triangles, and extension functors, generalising Ext^1 of exact categories and $\text{Hom}(-, \Sigma -)$ of triangulated categories.

We introduce *bixtriangulated* functors in Section 3.11 building on extriangulated functors introduced in [6]. Roughly speaking, a bixtriangulated functor is a functor $\mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$ that is extriangulated in each component, and where each associated natural transformation between extension functors is compatible with the other component. For many purposes this definition is adequate, though it does not recover bitriangulated functors, which involve $\text{Hom}(-, \Sigma^2 -)$. To recover the notion of a bitriangulated functor we introduce the notion of a *strong* bixtriangulated functor for which we use *higher extensions*.

For exact and triangulated categories the higher extensions are $\text{Ext}^n(-, -)$ and $\text{Hom}(-, \Sigma^n -)$. In Section 2 we recall the construction of higher extension functors: in the presence of enough injectives or projectives following [29]; and using coends following [26]. When any two of these constructions exist they are isomorphic. The higher extensions come equipped with a cup product. We show that whenever higher extensions exist, the corresponding cup product is compatible with any extriangulated functor, natural transformation or adjunction.

Finally, in Section 4 we define a category to be *monoidal extriangulated* if it is monoidal, extriangulated, the tensor product is bixtriangulated and the associator and

unitors are extriangulated natural transformations. If the monoidal structure is symmetric we say the category is *tensor extriangulated*. The higher extensions of the unit, together with the cup product, are the homogeneous components of a graded ring. This ring generalises the *Ext algebra* of finite tensor categories as well as the *graded central ring* (of the unit) from tensor triangular geometry; see for example [20] and [2]. In Lemma 4.5 we see this is a particularly nice ring when the monoidal structure is a strong bixtriangulated category:

Lemma A. *If $(\mathcal{A}, \mathbb{E}, \mathfrak{s}, \otimes, \mathbb{1})$ is a strong monoidal extriangulated category then the cup product makes $\mathbb{E}^*(\mathbb{1}, \mathbb{1}) = \{\mathbb{E}^n(\mathbb{1}, \mathbb{1})\}_{n \geq 0}$ into a graded-commutative graded ring.*

At the end of Section 3 and in Section 5 we present various examples of (strong) bixtriangulated functors and (strong) monoidal extriangulated categories:

- exact category of (bi)additive functors; see Examples 3.21 and 5.1;
- (stabilisation of the) category of matrix factorisations; see Example 3.26;
- category of flat modules over a commutative ring; see Example 5.2;
- (stabilisation of the) category of modules over a cocommutative Hopf algebra over a field; see Example 5.7; and
- extriangulated structure induced by pure-exact triangles; see Section 5.14.

We also look into the behaviour of (strong) bixtriangulated functors under stabilisation of extriangulated categories in Sections 3.22 and 5.4 and extriangulated substructures in Section 5.12.

In closing we generalise Balmer’s classification [1] of radical thick tensor ideals of a tensor triangulated category to tensor extriangulated categories. We use the approaches of Kock and Pitsch [38] and Buan, Krause and Solberg [9], and the proofs can be directly transferred to the extriangulated setting.

Theorem B. *Let $(\mathcal{A}, \mathbb{E}, \mathfrak{s}, \otimes, \mathbb{1})$ be a tensor extriangulated category. We assume the class of radical thick tensor ideals $\text{Rad}(\mathcal{A})$ is a set. Then there exists a spectral space $\text{BSpc}(\mathcal{A})$ and a bijection*

$$\text{Rad}(\mathcal{A}) \longleftrightarrow \{ \text{Thomason subsets of } \text{BSpc}(\mathcal{A}) \} .$$

This result is contained in Theorem 6.8. We compute the spectral space $\text{BSpc}(\mathcal{A})$ for the category of projective modules over various commutative rings in Section 6.11 and for stabilisations of extriangulated categories in Section 6.15.

2. Higher extensions in extriangulated categories

The notion of extriangulated categories generalises exact categories and triangulated categories. They were introduced by Nakaoka and Palu [51]. There exist notions of

higher extensions in exact categories (under some mild conditions) and in triangulated categories. These have been unified for extriangulated categories in [45,29,26]. In this section we recall the definitions and constructions and discuss some properties.

2.1. Extriangulated categories

An extriangulated category $\mathcal{A} = (\mathcal{A}, \mathbb{E}, \mathfrak{s})$ consists of an additive category \mathcal{A} , a functor $\mathbb{E}: \mathcal{A}^{\text{op}} \times \mathcal{A} \rightarrow \mathcal{A}b$, additive in each component, to the category $\mathcal{A}b$ of abelian groups, and a correspondence \mathfrak{s} associating an equivalence class $\mathfrak{s}(d)$ of a 3-term complex $X \rightarrow W \rightarrow Y$ to each $X, Y \in \mathcal{A}$ and $d \in \mathbb{E}(Y, X)$, satisfying some axioms; see [51, Definition 2.12]. By an *extriangle* we mean a representative of an equivalence class given by \mathfrak{s} , written

$$X \xrightarrow{f} W \xrightarrow{g} Y \dashrightarrow^d \quad \text{where } \mathfrak{s}(d) = [X \xrightarrow{f} W \xrightarrow{g} Y].$$

Let R be a commutative ring. An extriangulated category $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$ is *R-linear*, if the underlying category \mathcal{A} is R -linear, the functor \mathbb{E} factors through the forgetful functor $\text{Mod}(R) \rightarrow \mathcal{A}b$ and the induced functor $\mathcal{A}^{\text{op}} \times \mathcal{A} \rightarrow \text{Mod}(R)$ is R -bilinear; see [33, Section 3.1]. We denote the functor $\mathcal{A}^{\text{op}} \times \mathcal{A} \rightarrow \text{Mod}(R)$ also by \mathbb{E} . Every extriangulated category is trivially \mathbb{Z} -linear.

As for an exact or triangulated category, an extriangulated category is an additive category equipped with some additional extrinsic structure. Any exact or triangulated category has a natural extriangulated structure:

Example 2.2. Let \mathcal{E} be a (Quillen-)exact category. This means \mathcal{E} is equipped with a class of short exact sequences. Exact categories were introduced in [55, Section 2], for more details see [10]. When \mathcal{E} is an essentially small category or \mathcal{E} has enough \mathcal{E} -projectives or enough \mathcal{E} -injectives, then \mathcal{E} is an extriangulated category with $\mathbb{E}(Y, X) := \text{Ext}_{\mathcal{E}}^1(Y, X)$, the collection of Baer equivalence classes $[X \rightarrow W \rightarrow Y]$, and \mathfrak{s} the identity; for more details see [51, Example 2.13]. The assumptions on \mathcal{E} are sufficient to ensure that $\mathbb{E}(Y, X)$ is a set.

Example 2.3. Let \mathcal{T} be a triangulated category with suspension functor Σ . This means \mathcal{T} is equipped with a class of exact triangles. Triangulated categories were introduced by Verdier [59]. Then \mathcal{T} is an extriangulated category with

$$\mathbb{E}(Y, X) := \text{Hom}_{\mathcal{T}}(Y, \Sigma X) \quad \text{and} \quad \mathfrak{s}(d) := [X \rightarrow \Sigma^{-1} \text{cone}(d) \rightarrow Y]$$

for any $d \in \mathbb{E}(Y, X)$; for more details see [51, Proposition 3.22(1)].

Other examples of extriangulated categories include extension-closed full subcategories of triangulated categories [51, Remark 2.18], and certain ideal quotients of exact categories [51, Proposition 3.30]. We discuss the latter in more detail in Section 3.22.

In Examples 2.2 and 2.3 a notion of higher extensions exists. Motivated by higher extensions for exact categories, there are two constructions for higher extensions in extriangulated categories: via coends by [26] when the underlying category is essentially small, and via syzygies or cosyzygies by [45,29] when the extriangulated category has enough projectives or injectives, respectively. In the following sections we recall both constructions.

2.4. Higher extensions via coends

We briefly recall the definition of a coend. Let $F: \mathcal{A}^{\text{op}} \times \mathcal{A} \rightarrow \mathcal{B}$ be a functor where \mathcal{A} is essentially small and \mathcal{B} has set-indexed colimits. A *coend* of F is the coequaliser of the diagram

$$\coprod_{V, V' \in \text{sk}(\mathcal{A})} \coprod_{f \in \text{Hom}_{\mathcal{A}}(V, V')} F(V', V) \rightrightarrows \coprod_{W \in \text{sk}(\mathcal{A})} F(W, W),$$

defined by the morphisms $F(f, V)$ and $F(V', f)$ where $\text{sk}(\mathcal{A})$ denotes the skeleton of \mathcal{A} , ensuring the coproducts are set-indexed. In particular a coend is a colimit and, if it exists, is unique up to unique isomorphism. It is denoted by $\int^W F(W, W)$, and is equipped with morphisms $F(X, X) \rightarrow \int^W F(W, W)$ for every $X \in \mathcal{A}$.

Let R be a commutative ring and let $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$ be an essentially small R -linear extriangulated category. The *higher extensions* are functors

$$\begin{aligned} \mathbb{E}^n: \mathcal{A}^{\text{op}} \times \mathcal{A} &\rightarrow \text{Mod}(R) \quad \text{given by} \\ \mathbb{E}^0(Y, X) &:= \text{Hom}_{\mathcal{A}}(Y, X) \quad \text{and} \quad \mathbb{E}^n(Y, X) := \int^W \mathbb{E}^{n-1}(W, X) \otimes_R \mathbb{E}(Y, W) \end{aligned} \tag{2.4.1}$$

for $n \geq 1$; that is, the object $\mathbb{E}^n(Y, X)$ is the coend of the functor $\mathbb{E}^{n-1}(-, X) \otimes_R \mathbb{E}(Y, -)$. By [46, Proposition 2.2.1], we have $\mathbb{E}^1 = \mathbb{E}$. Using the universality of the colimit, it is straightforward to check that $\mathbb{E}^n(Y, X)$ is functorial in X and Y , and using also that tensoring over R is right exact and commutes with colimits, one has natural isomorphisms

$$\int^W \mathbb{E}^i(W, X) \otimes_R \mathbb{E}^j(Z, W) \xrightarrow{\cong} \mathbb{E}^n(Z, X) \tag{2.4.2}$$

for any integers $i, j \geq 0$ with $i + j = n$. This immediately yields the *cup product*

$$\smile: \mathbb{E}^i(Y, X) \otimes_R \mathbb{E}^j(Z, Y) \rightarrow \mathbb{E}^n(Z, X) \tag{2.4.3}$$

given by composing the morphism in (2.4.2) with the canonical morphism

$$\mathbb{E}^i(Y, X) \otimes_R \mathbb{E}^j(Z, Y) \rightarrow \int^W \mathbb{E}^i(W, X) \otimes_R \mathbb{E}^j(Z, W).$$

This is motivated by [26, Definition 3.10] where the cup product is introduced for $i+j = 1$. We have

$$f \smile d = \mathbb{E}^n(Z, f)(d) \quad \text{and} \quad e \smile g = \mathbb{E}^n(g, X)(e) \tag{2.4.4}$$

for morphisms $f: Y \rightarrow X$ and $g: Z \rightarrow Y$ in \mathcal{A} and $d \in \mathbb{E}^n(Z, Y)$ and $e \in \mathbb{E}^n(Y, X)$.


This construction of higher extension groups for exact categories goes back to [62, Section 4]. For extriangulated categories this was foreshadowed in [51, Remark 5.10] for the case where $i = j = 1$ in (2.4.3). It then was extended to any $i, j \geq 0$ in [26].

Remark 2.5. The construction in (2.4.1) can be done more generally to obtain a “folding” of R -bilinear functors. This defines a tensor product of R -bilinear functors. The tensor product already appears in [62, Section 4]; for a short survey, see [17]. We discuss this later in Example 3.21.

Example 2.6. Let \mathcal{E} be an exact category that is essentially small. We consider \mathcal{E} as an extriangulated category, as in Example 2.2. Roughly speaking, the coend acts as gluing exact sequences. This means $\mathbb{E}^n(Y, X)$ is the set of equivalence classes

$$[X = X_0 \rightarrow X_1 \rightarrow \cdots \rightarrow X_{n+1} = Y],$$

where $X_1, \dots, X_n \in \mathcal{E}$ and there is a commutative diagram of the form



such that $W_i \rightarrow X_{i+1} \rightarrow W_{i+1}$ is exact for each $i = 0, \dots, n - 1$; see [61, Section 3.4] and [62, Section 4.3]. The cup product is given by

$$\begin{aligned}
 [X = X_0 \rightarrow \cdots \rightarrow X_{n+1} = Y] \smile [Y = Y_0 \rightarrow \cdots \rightarrow Y_{m+1} = Z] \\
 = [X = X_0 \rightarrow \cdots \rightarrow X_n \rightarrow Y_1 \rightarrow \cdots \rightarrow Y_{m+1} = Z]
 \end{aligned}$$

where the morphism $X_n \rightarrow Y_1$ is the composition $X_n \rightarrow Y \rightarrow Y_1$.

Example 2.7. Let \mathcal{T} be a triangulated category with suspension functor Σ . We consider \mathcal{T} as an extriangulated category, as in Example 2.3. By [26, Corollary 3.23] we have

$$\mathbb{E}^n(Y, X) = \text{Hom}_{\mathcal{T}}(Y, \Sigma^n X),$$

and for $d \in \mathbb{E}^i(Y, X)$ and $e \in \mathbb{E}^j(Z, Y)$ the cup product is

$$d \smile e = (\Sigma^j d) \circ e \in \text{Hom}_{\mathcal{T}}(Z, \Sigma^{i+j} X) = \mathbb{E}^{i+j}(Z, X).$$

2.8. Higher extensions via syzygies or cosyzygies

We recall from [51, Section 3.4] the definition of an extriangulated category with enough projectives. Let $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$ be an extriangulated category. An object $P \in \mathcal{A}$ is \mathbb{E} -projective, if $\mathbb{E}(P, X) = 0$ for any $X \in \mathcal{A}$. We say $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$ has enough \mathbb{E} -projectives if for any $X \in \mathcal{A}$ there exists an \mathbb{E} -projective object P and an extriangle $W \rightarrow P \rightarrow X \dashrightarrow$. The object W is called a syzygy of X . The syzygy of an object X need not be unique. When \mathcal{A} has enough projectives, we fix a choice of syzygy, and denote it by ΩX ; cf. [29, Assumption 3.3]. We set

$$\Omega^0 X := X \quad \text{and} \quad \Omega^n X := \Omega(\Omega^{n-1} X)$$

for $n \geq 1$.

Let R be a commutative ring. In an R -linear extriangulated category $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$ with enough projectives, the higher extensions are functors

$$\begin{aligned} \mathbb{E}_\Omega^n : \mathcal{A}^{\text{op}} \times \mathcal{A} &\rightarrow \text{Mod}(R) \quad \text{given by} \\ \mathbb{E}_\Omega^0(Y, X) &:= \text{Hom}_{\mathcal{A}}(Y, X) \quad \text{and} \quad \mathbb{E}_\Omega^n(Y, X) := \mathbb{E}(\Omega^{n-1} Y, X) \end{aligned} \tag{2.8.1}$$

for $n \geq 1$; these functors are well-defined and independent of the choice of syzygy by [29, Proposition 3.4, Definition 3.6].

Dually, in an extriangulated category $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$ an object $I \in \mathcal{A}$ is \mathbb{E} -injective, if $\mathbb{E}(Y, I) = 0$ for any $Y \in \mathcal{A}$. We say $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$ has enough \mathbb{E} -injectives if for any $X \in \mathcal{A}$ there exists an \mathbb{E} -injective object I and an extriangle $X \rightarrow I \rightarrow W \dashrightarrow$. The object W is a cosyzygy of X . In an R -linear extriangulated category $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$ with enough injectives we fix a choice of cosyzygy ΣX for every $X \in \mathcal{T}$. The higher extensions are the functors

$$\begin{aligned} \mathbb{E}_\Sigma^n : \mathcal{A}^{\text{op}} \times \mathcal{A} &\rightarrow \text{Mod}(R) \quad \text{given by} \\ \mathbb{E}_\Sigma^0(Y, X) &:= \text{Hom}_{\mathcal{A}}(Y, X) \quad \text{and} \quad \mathbb{E}_\Sigma^n(Y, X) := \mathbb{E}(Y, \Sigma^{n-1} X) \end{aligned} \tag{2.8.2}$$

for $n \geq 1$; these functors are well-defined and independent of the choice of cosyzygy by [29, Proposition 3.4].

When the extriangulated category $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$ has enough \mathbb{E} -injectives and enough \mathbb{E} -projectives, the definitions of higher extensions via syzygies and cosyzygies are naturally isomorphic by [45, Lemma 5.1].

Example 2.9. Let \mathcal{T} be a triangulated category with suspension Σ . We consider \mathcal{T} as an extriangulated category as in Example 2.3. The zero object is the only \mathbb{E} -projective object, and for any object X there exists an exact triangle $\Sigma^{-1} X \rightarrow 0 \rightarrow X \xrightarrow{\text{id}_X} X$. Hence \mathcal{T} has enough \mathbb{E} -projectives and $\Sigma^{-1} X$ is a syzygy of X . Dually, any triangulated category has enough injectives and the suspension ΣX is a cosyzygy of X . It is straightforward to see that the higher extensions of any triangulated category \mathcal{T} are given by

$$\mathbb{E}_{\Sigma}^n(Y, X) = \text{Hom}_{\mathcal{T}}(Y, \Sigma^n X) \cong \text{Hom}_{\mathcal{T}}(\Sigma^{-n}Y, X) = \mathbb{E}_{\Omega}^n(Y, X).$$

Next we discuss the cup product on \mathbb{E}_{Ω} . The construction for \mathbb{E}_{Σ} works analogously. Let R be a commutative ring and let $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$ be an R -linear extriangulated category with enough \mathbb{E} -projectives. For an object $W \in \mathcal{A}$ there exists an \mathbb{E} -projective object P and an extriangle $\Omega W \rightarrow P \rightarrow W \dashrightarrow$. For any object Y this extriangle induces a surjective map of R -modules

$$\text{Hom}_{\mathcal{A}}(\Omega W, Y) \rightarrow \mathbb{E}(W, Y) \rightarrow 0$$

which is natural in Y ; see [51, Corollary 3.12]. Then the cup product for \mathbb{E}_{Ω} and $i, j > 0$ is defined as

$$\begin{aligned} \mathbb{E}_{\Omega}^i(Y, X) \otimes_R \mathbb{E}_{\Omega}^j(Z, Y) &\rightarrow \mathbb{E}_{\Omega}^i(\Omega^j Z, X) \cong \mathbb{E}_{\Omega}^{i+j}(Z, X) \\ d \otimes e &\mapsto d \smile e := \mathbb{E}_{\Omega}^i(f_e, X)(d), \end{aligned} \tag{2.9.1}$$

where f_e is a pre-image of e under the surjective map above for $W = \Omega^{j-1}Z$. By [29, Section 3.1] the cup product is independent of the pre-image, and there are natural isomorphisms $\mathbb{E}_{\Omega}^k(\Omega^{\ell}-, -) \cong \mathbb{E}_{\Omega}^{k+\ell}(-, -)$.

2.10. Comparison

When $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$ is essentially small and has enough \mathbb{E} -injectives or \mathbb{E} -projectives, then the definitions of higher extensions are naturally isomorphic by [26, Corollary 3.21]. The proof in [26] involves proving that the higher extensions and cup products are universal among bifunctors satisfying similar properties; see [26, Proposition 3.20]. We give an alternative proof, as well as a proof that the cup products are compatible with these natural isomorphisms. The main argument for both statements is contained in the following technical lemma.

Lemma 2.11. *Let R be a commutative ring and $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$ be an essentially small R -linear extriangulated category with enough \mathbb{E} -projectives. There exist R -module homomorphisms $\varphi_{i,j}(Y, X): \mathbb{E}^i(\Omega^j Y, X) \rightarrow \mathbb{E}^{i+j}(Y, X)$ for $i, j \geq 0$ and objects $X, Y \in \mathcal{A}$, such that:*

- (1) $\varphi_{i,j}(Y, X)$ is surjective when $i = 0$ and $j > 0$;
- (2) $\varphi_{i,j}(Y, X)$ is an isomorphism when $i \geq 1$ or $j = 0$;
- (3) $\varphi_{i,j}(Y, X)$ is natural in X ;
- (4) $\varphi_{i,j}(Y, X)$ is natural in Y in the sense that the associated diagram commutes for any lift $\Omega^j Y \rightarrow \Omega^j Y'$ of a morphism $Y \rightarrow Y'$; and
- (5) the morphisms are compatible with the cup product (2.4.3) in the sense that the following diagram commutes

$$\begin{array}{ccc}
 \mathbb{E}^i(Y, X) \otimes_R \mathbb{E}^j(\Omega^k Z, Y) & \xrightarrow{\sim} & \mathbb{E}^{i+j}(\Omega^k Z, X) \\
 \text{id} \otimes \varphi_{j,k}(Z, Y) \downarrow & & \downarrow \varphi_{i+j,k}(Z, X) \\
 \mathbb{E}^i(Y, X) \otimes_R \mathbb{E}^{j+k}(Z, Y) & \xrightarrow{\sim} & \mathbb{E}^{i+j+k}(Z, X)
 \end{array}$$

for all integers $i, j, k \geq 0$.

Proof. We use induction on j . For $j = 0$ we let $\varphi_{i,0}(Y, X)$ be the identity morphism. For $j = 1$ we have an extriangle $\Omega Y \rightarrow P \rightarrow Y \dashrightarrow$ with P an \mathbb{E} -projective object. Then there is a right exact sequence

$$\text{Hom}_{\mathcal{A}}(P, W) \rightarrow \text{Hom}_{\mathcal{A}}(\Omega Y, W) \rightarrow \mathbb{E}(Y, W) \rightarrow 0$$

in $\text{Mod}(R)$ that is natural in $W \in \mathcal{A}$. We apply $\int^W \mathbb{E}^i(W, X) \otimes_R -$ to the right exact sequence. Since coends preserve right exact sequences, the sequence

$$\mathbb{E}^i(P, X) \rightarrow \mathbb{E}^i(\Omega Y, X) \rightarrow \mathbb{E}^{i+1}(Y, X) \rightarrow 0$$

is exact; see [46, Proposition 2.2.1]. We let $\varphi_{i,1}(Y, X)$ be the second morphism. Finally, for $j > 1$ we define $\varphi_{i,j}(Y, X)$ as the composition

$$\mathbb{E}^i(\Omega^j Y, X) \xrightarrow{\varphi_{i,1}(\Omega^{j-1} Y, X)} \mathbb{E}^{i+1}(\Omega^{j-1} Y, X) \xrightarrow{\varphi_{i+1,j-1}(Y, X)} \mathbb{E}^{i+j}(Y, X).$$

The properties (1–3) hold by construction. Property (4) is straightforward to check from the construction, and (5) can be shown using induction on k . \square

Corollary 2.12. *Let R be a commutative ring and $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$ be an essentially small R -linear extriangulated category with enough \mathbb{E} -projectives. There are natural isomorphisms $\mathbb{E}_{\Omega}^i \implies \mathbb{E}^i$ for $i \geq 0$ that are compatible with the cup products defined in (2.4.3) and (2.9.1); that is the following diagram commutes*

$$\begin{array}{ccc}
 \mathbb{E}_{\Omega}^i(Y, X) \otimes_R \mathbb{E}_{\Omega}^j(Z, Y) & \xrightarrow[\text{(2.9.1)}]{\sim} & \mathbb{E}_{\Omega}^{i+j}(Z, X) \\
 \downarrow \cong & & \downarrow \cong \\
 \mathbb{E}^i(Y, X) \otimes_R \mathbb{E}^j(Z, Y) & \xrightarrow[\text{(2.4.3)}]{\sim} & \mathbb{E}^{i+j}(Z, X).
 \end{array}$$

Proof. For $i = 0$ take the identity on $\text{Hom}_{\mathcal{A}}(Y, X)$. For $i > 0$ we use the maps $\varphi_{1,i-1}(Y, X): \mathbb{E}_{\Omega}^i(Y, X) = \mathbb{E}(\Omega^{i-1} Y, X) \rightarrow \mathbb{E}^i(Y, X)$ from Lemma 2.11 which are isomorphisms by (2). We prove the square commutes by induction on j .

Let $d \in \mathbb{E}_\Omega^i(Y, X)$ and $e \in \mathbb{E}_\Omega^j(Z, Y)$. For the case $j = 0$ we obtain

$$\begin{aligned} \varphi_{1,i-1}(Z, X)(d \smile e) &= \varphi_{1,i-1}(Z, X)(\mathbb{E}_\Omega^i(e, X)(d)) \\ &= \varphi_{1,i-1}(Z, X)(\mathbb{E}(\Omega^{i-1}e, X)(d)) \\ &= \mathbb{E}^i(e, X)(\varphi_{1,i-1}(Y, X)(d)) \\ &= \varphi_{1,i-1}(Y, X)(d) \smile e; \end{aligned}$$

using property (4) for the third identification. The remaining identifications hold by the definitions (2.9.1), (2.8.1) and (2.4.3), respectively.

We assume that the desired square commutes for some $j \geq 0$. We now claim that the following diagram commutes:

$$\begin{array}{ccc} \mathbb{E}_\Omega^i(Y, X) \otimes_R \mathbb{E}_\Omega^{j+1}(Z, Y) & \xrightarrow[\text{(2.9.1)}]{\smile} & \mathbb{E}_\Omega^{i+j+1}(Z, X) \\ \cong \downarrow & & \downarrow \cong \\ \mathbb{E}_\Omega^i(Y, X) \otimes_R \mathbb{E}_\Omega^j(\Omega Z, Y) & \xrightarrow[\text{(2.9.1)}]{\smile} & \mathbb{E}_\Omega^{i+j}(\Omega Z, X) \\ \varphi_{1,i-1}(Y, X) \otimes \varphi_{1,j-1}(\Omega Z, Y) \downarrow & & \downarrow \varphi_{1,i+j-1}(\Omega Z, X) \\ \mathbb{E}^i(Y, X) \otimes_R \mathbb{E}^j(\Omega Z, Y) & \xrightarrow[\text{(2.4.3)}]{\smile} & \mathbb{E}^{i+j}(\Omega Z, X) \\ \text{id}_{\mathbb{E}^i(Y, X)} \otimes \varphi_{j,1}(Z, Y) \downarrow & & \downarrow \varphi_{i+j,1}(Z, X) \\ \mathbb{E}^i(Y, X) \otimes_R \mathbb{E}^{j+1}(Z, Y) & \xrightarrow[\text{(2.4.3)}]{\smile} & \mathbb{E}^{i+j+1}(Z, X). \end{array}$$

The upper square commutes by the definition of \mathbb{E}_Ω and its cup product. The middle square commutes by replacing Z with ΩZ in the inductive hypothesis. The lower square commutes by (5). It remains to observe that the outside square is precisely the claim. \square

Remark 2.13. Let $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$ be an extriangulated category. Assume that higher extensions exist. Then $\mathbb{E}^*(X, X) = \{\mathbb{E}^n(X, X)\}_{n \geq 0}$ together with the cup product is a graded ring for any $X \in \mathcal{A}$. Moreover, for any $X, Y \in \mathcal{A}$ the cup product induces a $\mathbb{E}^*(X, X)$ - $\mathbb{E}^*(Y, Y)$ -bimodule structure on $\mathbb{E}^*(Y, X) = \{\mathbb{E}^n(Y, X)\}_{n \geq 0}$.

Note that there is a category where the objects are the objects of \mathcal{A} , where $\mathbb{E}^*(X, Y)$ is the set of morphisms $X \rightarrow Y$, and where the cup product defines composition. This is a special case of the *additive tensor category* in [56].

Remark 2.14. In connection with the cup product it is common to introduce a Koszul sign convention; see [47, Theorem III.9.1]. This means, for $d \in \mathbb{E}^i(Y, X)$ we define morphisms

$$\begin{aligned} d_* : \mathbb{E}^j(W, Y) &\rightarrow \mathbb{E}^{i+j}(W, X), & e &\mapsto d \smile e \quad \text{and} \\ d^* : \mathbb{E}^j(X, Z) &\rightarrow \mathbb{E}^{i+j}(Y, Z), & c &\mapsto (-1)^{ij}c \smile d \end{aligned}$$

that are natural in W and Z , respectively. For $i = 0$ one has $d_* = \mathbb{E}^j(W, d)$ and $d^* = \mathbb{E}^j(d, Z)$. Further, the natural transformations for $i = 1$ and $j = 0$ extend those in [51, Definition 3.1]. The Koszul sign convention does not appear in any of the previous works on the cup product of an extriangulated category.

For a triangulated category \mathcal{T} the sign seems artificial, but when $\mathcal{T} = D(\mathcal{E})$ is the derived category of an exact category it is standard: An element in $\mathbb{E}^n(Y, X)$ is considered a degree n morphism $Y \rightarrow X$.

3. Bixtriangulated functors

We now discuss functors of the form $\mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$ of extriangulated categories. The notion of a bixtriangulated functor should generalise the notions for exact and triangulated categories; see for example [60, II.7.4] and [36, Definition 10.3.6], respectively.

3.1. Extriangulated functors

An *extriangulated functor*

$$F = (F, \beta): (\mathcal{A}, \mathbb{E}, \mathfrak{s}) \rightarrow (\mathcal{B}, \mathbb{F}, \mathfrak{t})$$

of extriangulated categories consists of an additive functor $F: \mathcal{A} \rightarrow \mathcal{B}$ and a natural transformation $\beta: \mathbb{E}(-, -) \implies \mathbb{F}(F(-), F(-))$ such that for any extriangle $X \xrightarrow{f} W \xrightarrow{g} Y \xrightarrow{d} X$ in \mathcal{A} there is an extriangle

$$F(X) \xrightarrow{F(f)} F(W) \xrightarrow{F(g)} F(Y) \xrightarrow{\beta_{Y,X}(d)} F(X)$$

in \mathcal{B} ; see [6, Definition 2.32] and also [50, Definition 2.11].

The composition of extriangulated functors $(F, \beta): (\mathcal{A}, \mathbb{E}, \mathfrak{s}) \rightarrow (\mathcal{B}, \mathbb{F}, \mathfrak{t})$ and $(G, \gamma): (\mathcal{B}, \mathbb{F}, \mathfrak{t}) \rightarrow (\mathcal{C}, \mathbb{G}, \mathfrak{u})$ is defined by (GF, α) where $\alpha_{Y,X} = \gamma_{FY,FX} \beta_{YX}$ for all $X, Y \in \mathcal{A}$; see [5, Definition 3.18(ii)].

Let $(F, \beta), (G, \gamma): (\mathcal{A}, \mathbb{E}, \mathfrak{s}) \rightarrow (\mathcal{B}, \mathbb{F}, \mathfrak{t})$ be extriangulated functors. An *extriangulated natural transformation* $\eta: (F, \beta) \implies (G, \gamma)$ is a natural transformation $\eta: F \implies G$ of additive functors such that, for all $X, Y \in \mathcal{A}$, the diagram

$$\begin{CD} \mathbb{E}(Y, X) @>\beta_{Y,X}>> \mathbb{F}(F(Y), F(X)) \\ @V\gamma_{Y,X}VV @VV(\eta_X)_*V \\ \mathbb{E}(G(Y), G(X)) @>(\eta_Y)_*>> \mathbb{F}(F(Y), G(X)) \end{CD}$$

commutes; see [50, Definition 2.11] and also [5, Definition 4.1].

Remark 3.2. The definition of an extriangulated natural transformation can be phrased using the cup product. Explicitly, a natural transformation $\eta: (\mathbf{F}, \beta) \implies (\mathbf{G}, \gamma)$ is extriangulated if and only if

$$\eta_X \smile \beta_{Y,X}(d) = \gamma_{Y,X}(d) \smile \eta_Y$$

for any $d \in \mathbb{E}(Y, X)$. This means the natural transformation is compatible with the connecting map from Hom to \mathbb{E} .

Specialising to exact or triangulated categories, these notions recover the classical definitions:

Example 3.3. Let $\mathbf{F}: \mathcal{E} \rightarrow \mathcal{F}$ be an exact functor of exact categories; this means any exact sequence is mapped to an exact sequence. The functor \mathbf{F} induces a map $\beta_{Y,X}: \text{Ext}_{\mathcal{E}}^1(Y, X) \rightarrow \text{Ext}_{\mathcal{F}}^1(\mathbf{F}(Y), \mathbf{F}(X))$, which is natural in X and Y since \mathbf{F} preserves pushout and pullback squares; for the latter see [10, Proposition 5.2]. If we view \mathcal{E} and \mathcal{F} as extriangulated categories as in Example 2.2, then \mathbf{F} , together with this natural transformation β , is an extriangulated functor.

Conversely, it is clear that any extriangulated functor between exact categories is exact; see [6, Theorem 2.34].

Lastly, we note that any natural transformation of exact functors is automatically extriangulated [50, p. 349]; see also [5, Example 5.4] for details.

Example 3.4. Let $(\mathbf{F}, \tau): \mathcal{S} \rightarrow \mathcal{T}$ be a triangulated functor of triangulated categories; this means $\tau: \mathbf{F}\Sigma \rightarrow \Sigma\mathbf{F}$ is a natural transformation and, for any exact triangle $X \xrightarrow{f} X \xrightarrow{g} Y \xrightarrow{h} \Sigma X$ in \mathcal{S} , there is an exact triangle

$$\mathbf{F}(X) \xrightarrow{\mathbf{F}(f)} \mathbf{F}(W) \xrightarrow{\mathbf{F}(g)} \mathbf{F}(Y) \xrightarrow{\tau_X \mathbf{F}(h)} \Sigma \mathbf{F}(X)$$

in \mathcal{T} . Then $(\mathbf{F}, \beta): \mathcal{S} \rightarrow \mathcal{T}$ is an extriangulated functor of triangulated categories with $\beta_{Y,X}$ given by the composition

$$\text{Hom}_{\mathcal{S}}(Y, \Sigma X) \xrightarrow{\mathbf{F}} \text{Hom}_{\mathcal{T}}(\mathbf{F}(Y), \mathbf{F}(\Sigma X)) \xrightarrow{\text{Hom}_{\mathcal{T}}(\mathbf{F}(Y), \tau_X)} \text{Hom}_{\mathcal{T}}(\mathbf{F}(Y), \Sigma \mathbf{F}(X))$$

for all $X, Y \in \mathcal{S}$; see [6, Theorem 2.33].

Conversely, let $(\mathbf{F}, \beta): \mathcal{S} \rightarrow \mathcal{T}$ be an extriangulated functor between triangulated categories. We set

$$\tau_X := \beta_{\Sigma X, X}(\text{id}_{\Sigma X}): \mathbf{F}(\Sigma X) \rightarrow \Sigma \mathbf{F}(X).$$

It is straightforward to check that this map is natural in X . It is an isomorphism, since it witnesses the mapping of the (ex)triangles

$$(X \rightarrow 0 \rightarrow \Sigma X \xrightarrow{\text{id}_{\Sigma X}} \Sigma X) \mapsto (F(X) \rightarrow 0 \rightarrow F(\Sigma X) \xrightarrow{\tau_X} \Sigma F(X)).$$

Hence (F, τ) is a triangulated functor.

Unlike the exact case, a natural transformation of triangulated functors is not necessarily extriangulated. It is extriangulated if and only if it is a morphism of triangulated functors in the sense of [36, Definition 10.1.9(ii)]; see [50, p. 349] and also [5, Example 5.3].

The next Lemma is a generalisation of the fact that the higher extensions, defined using syzygies, are functorial from [29, Proposition 3.4].

Lemma 3.5. *Let $(F, \beta): (\mathcal{A}, \mathbb{E}, \mathfrak{s}) \rightarrow (\mathcal{B}, \mathbb{F}, \mathfrak{t})$ be an extriangulated functor of extriangulated categories. For any morphism $f: Y \rightarrow Y'$ there exists a morphism $f_\omega: \Omega F(Y) \rightarrow F(\Omega Y')$ such that there is a commutative diagram*

$$\begin{array}{ccccccc} \Omega F(Y) & \longrightarrow & Q & \longrightarrow & F(Y) & \dashrightarrow & \\ \downarrow f_\omega & & \downarrow & & \downarrow f & & \\ F(\Omega Y') & \longrightarrow & F(P) & \longrightarrow & F(Y') & \dashrightarrow & \end{array}$$

where each row is an extriangle in \mathcal{B} and P and Q are \mathbb{E} -projective and \mathbb{F} -projective, respectively. Moreover, the morphism

$$\mathbb{F}((\text{id}_Y)_\omega, \text{id}_W): \mathbb{F}(F(\Omega Y), W) \rightarrow \mathbb{F}(\Omega F(Y), W)$$

is natural in $W \in \mathcal{B}$ and $Y \in \mathcal{A}$.

Proof. Let $\Omega Y' \rightarrow P \rightarrow Y' \dashrightarrow$ be an extriangle with P an \mathbb{E} -projective object. Further, let $\Omega F(Y) \rightarrow Q \rightarrow F(Y) \dashrightarrow$ be an extriangle with Q a \mathbb{F} -projective object. As $\mathbb{F}(Q, -) = 0$, applying $\text{Hom}(Q, F(-))$ to the first extriangle yields a short exact sequence by [51, Corollary 3.12]. That is we obtain a commutative diagram

$$\begin{array}{ccccccc} \Omega F(Y) & \longrightarrow & Q & \longrightarrow & F(Y) & \dashrightarrow & \\ & & \downarrow & & \downarrow f & & \\ F(\Omega Y') & \longrightarrow & F(P) & \longrightarrow & F(Y') & \dashrightarrow & . \end{array}$$

By the axiom $(\text{ET3})^{\text{op}}$ for an extriangulated category in [51, Definition 2.12], there exists a morphism $f_\omega: \Omega F(Y) \rightarrow F(\Omega Y)$ such that the diagram commutes. While this morphism need not be unique, the induced map

$$\mathbb{F}(f_\omega, \text{id}_W): \mathbb{F}(F(\Omega Y), W) \rightarrow \mathbb{F}(\Omega F(Y'), W)$$

is independent of the choice of f_ω . In particular, the two compositions

$$\Omega F(Y) \rightarrow \Omega F(Y') \rightarrow F(\Omega Y') \quad \text{and} \quad \Omega F(Y) \rightarrow F(\Omega Y) \rightarrow F(\Omega Y')$$

need not coincide, however after applying $\mathbb{F}(-, W)$ they will be the same. That means there is a commutative diagram

$$\begin{CD} \mathbb{F}(\Omega F(Y), W) @>{\mathbb{F}((\text{id}_Y)_\omega, \text{id}_W)}>> \mathbb{F}(F(\Omega Y), W) \\ @V{\mathbb{F}(\Omega F(f), \text{id}_W)}VV @VV{\mathbb{F}(F\Omega f, \text{id}_W)}V \\ \mathbb{F}(\Omega F(Y'), W) @>{\mathbb{F}((\text{id}_{Y'})_\omega, \text{id}_W)}>> \mathbb{F}(F(\Omega Y'), W) \end{CD}$$

where the vertical morphisms are well-defined by [29, Proposition 3.4]. \square

We record the interplay of extriangulated functors and the higher extensions:

Proposition 3.6. *Let $(F, \beta): (\mathcal{A}, \mathbb{E}, \mathfrak{s}) \rightarrow (\mathcal{B}, \mathbb{F}, \mathfrak{t})$ be an extriangulated functor of extriangulated categories. Assume higher extensions are well-defined in $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$ and $(\mathcal{B}, \mathbb{F}, \mathfrak{t})$ through the same construction, i.e. by means of:*

- (1) *coends following (2.4.1), assuming \mathcal{A} and \mathcal{B} are essentially small;*
- (2) *syzygies following (2.8.1), assuming \mathcal{A} and \mathcal{B} have enough projectives; or*
- (3) *cosyzygies following (2.8.2), assuming \mathcal{A} and \mathcal{B} have enough injectives.*

Then for each $n \geq 0$ there are induced natural transformations

$$\beta^n: \mathbb{E}^n(-, -) \implies \mathbb{F}^n(F(-), F(-))$$

such that $\beta_{Y,X}^i(d) \smile \beta_{Z,Y}^j(e) = \beta_{Z,X}^{i+j}(d \smile e)$ for $d \in \mathbb{E}^i(Y, X)$ and $e \in \mathbb{E}^j(Z, Y)$.

Proof. (1) Defining higher extensions via coends, we construct the natural transformation β^n by induction on n . For $n = 0$ it is the map given by F and for $n = 1$ we have $\beta^1 = \beta$. For $n \geq 2$ and any $X, Y, W \in \mathcal{A}$ we obtain the composition

$$\begin{CD} \mathbb{E}^{n-1}(W, X) \otimes_R \mathbb{E}(Y, W) @>>> \mathbb{F}^n(F(Y), F(X)) \\ @V{\beta_{W,X}^{n-1} \otimes \beta_{Y,W}}VV @AA{\smile}A \\ \mathbb{F}^{n-1}(F(W), F(X)) \otimes_R \mathbb{F}(F(Y), F(W)) @>>> \mathbb{F}^n(F(Y), F(X)) \end{CD}$$

The colimit property yields a unique morphism $\beta_{Y,X}^n: \mathbb{E}^n(Y, X) \rightarrow \mathbb{F}^n(F(Y), F(X))$, and it is straightforward to check that this map is natural in X and Y . By construction the cup product is compatible with these natural transformations.

(2, 3) We define higher extensions via syzygies. The argument for cosyzygies is analogous. Using Lemma 3.5 $(n - 1)$ -times we define

$$\begin{array}{ccc} \mathbb{E}(\Omega^{n-1}Y, X) & \xrightarrow{\beta_{\Omega^{n-1}Y, X}} & \mathbb{F}(\mathbb{F}(\Omega^{n-1}Y), \mathbb{F}(X)) \longrightarrow \mathbb{F}(\Omega^{n-1}\mathbb{F}(Y), \mathbb{F}(X)) \\ \parallel & & \parallel \\ \mathbb{E}_{\Omega}^n(Y, X) & \xrightarrow{\beta_{Y, X}^n} & \mathbb{F}_{\Omega}^n(\mathbb{F}(Y), \mathbb{F}(X)) \end{array}$$

for any $n \geq 2$. By Lemma 3.5 this morphism is natural in X and Y . It remains to check the compatibility of these natural transformations with the cup product. There is a canonical and commutative diagram of the form

$$\begin{array}{ccccc} \text{Hom}_{\mathcal{A}}(\Omega^n Y, X) & \longrightarrow & \mathbb{E}(\Omega^{n-1}Y, X) & = & \mathbb{E}_{\Omega}^n(Y, X) \\ \downarrow & & \downarrow & & \downarrow \beta_{Y, X}^n \\ \text{Hom}_{\mathcal{B}}(\mathbb{F}(\Omega^n Y), \mathbb{F}(X)) & \longrightarrow & \mathbb{F}(\mathbb{F}(\Omega^{n-1}Y), \mathbb{F}(X)) & & \\ \downarrow & & \downarrow & & \\ \text{Hom}_{\mathcal{B}}(\Omega^n \mathbb{F}(Y), \mathbb{F}(X)) & \longrightarrow & \mathbb{F}(\Omega^{n-1}\mathbb{F}(Y), \mathbb{F}(X)) & = & \mathbb{F}_{\Omega}^n(\mathbb{F}(Y), \mathbb{F}(X)). \end{array}$$

Hence from here it is straightforward to see that the natural transformation is compatible with the cup product. \square

3.7. Extriangulated adjunction

Let

$$(\mathbb{F}, \beta): (\mathcal{A}, \mathbb{E}, \mathfrak{s}) \rightarrow (\mathcal{B}, \mathbb{F}, \mathfrak{t}) \quad \text{and} \quad (\mathbb{G}, \gamma): (\mathcal{B}, \mathbb{F}, \mathfrak{t}) \rightarrow (\mathcal{A}, \mathbb{E}, \mathfrak{s})$$

be extriangulated functors. Following [5, Definition 4.9(i)], we say (\mathbb{G}, γ) is an *extriangulated right adjoint* of (\mathbb{F}, β) and (\mathbb{F}, β) is an *extriangulated left adjoint*, if (\mathbb{F}, \mathbb{G}) is a pair of adjoint functors and the unit and counit witnessing the adjunction are extriangulated natural transformations. In this situation we say $((\mathbb{F}, \beta), (\mathbb{G}, \gamma))$ is a *pair of extriangulated adjoint functors*.

Requiring that the unit and counit are extriangulated is superfluous for triangulated category. However, for exact categories it is necessary: For example, if R is a commutative ring and X is a flat non-projective R -module, then in the adjunction $-\otimes_R X \dashv \text{Hom}_R(X, -)$ the functor $-\otimes_R$ is exact while $\text{Hom}_R(X, -)$ is not.

A consequence of this definition is that we obtain an adjunction relation on the higher extensions.

Lemma 3.8. *Let $(\mathbb{F}, \beta): (\mathcal{A}, \mathbb{E}, \mathfrak{s}) \rightarrow (\mathcal{B}, \mathbb{F}, \mathfrak{t})$ be an extriangulated left adjoint of $(\mathbb{G}, \gamma): (\mathcal{B}, \mathbb{F}, \mathfrak{t}) \rightarrow (\mathcal{A}, \mathbb{E}, \mathfrak{s})$ and η and ε the associated unit and counit, respectively. Then the maps*

$$\begin{aligned} \mathbb{F}(\mathbb{F}(Y), X) &\xrightarrow{\gamma_{\mathbb{F}(Y), X}} \mathbb{E}(\mathbb{G}(\mathbb{F}(Y)), \mathbb{G}(X)) \xrightarrow{(\eta_Y)^*} \mathbb{E}(Y, \mathbb{G}(X)) \quad \text{and} \\ \mathbb{E}(Y, \mathbb{G}(X)) &\xrightarrow{\beta_{Y, \mathbb{G}(X)}} \mathbb{F}(\mathbb{F}(Y), \mathbb{F}(\mathbb{G}(X))) \xrightarrow{(\varepsilon_X)_*} \mathbb{F}(\mathbb{F}(Y), X) \end{aligned}$$

are inverse to each other for any $Y \in \mathcal{A}$ and $X \in \mathcal{B}$. In particular, this yields an isomorphism $\mathbb{F}(\mathbb{F}(Y), X) \cong \mathbb{E}(Y, \mathbb{G}(X))$ for all $Y \in \mathcal{A}$ and $X \in \mathcal{B}$ that is compatible with the adjunction of (\mathbb{F}, β) and (\mathbb{G}, γ) .

Proof. We show that the composition $\mathbb{F}(\mathbb{F}(Y), X) \rightarrow \mathbb{E}(Y, \mathbb{G}(X)) \rightarrow \mathbb{F}(\mathbb{F}(Y), X)$ is the identity map. The proof that the composition $\mathbb{E}(Y, \mathbb{G}(X)) \rightarrow \mathbb{F}(\mathbb{F}(Y), X) \rightarrow \mathbb{E}(Y, \mathbb{G}(X))$ is the identity is similar and omitted. Consider the following diagram:

$$\begin{array}{ccccc}
 \mathbb{F}(\mathbb{F}(Y), X) & \xrightarrow{\gamma_{\mathbb{F}(Y), X}} & \mathbb{E}(\mathbb{G}(\mathbb{F}(Y)), \mathbb{G}(X)) & & \\
 \downarrow (\varepsilon_{\mathbb{F}(Y)})^* & & \downarrow \beta_{\mathbb{G}(\mathbb{F}(Y)), \mathbb{G}(X)} & \searrow (\eta_Y)^* & \\
 \mathbb{F}(\mathbb{F}(\mathbb{G}(\mathbb{F}(Y))), X) & \xleftarrow{(\varepsilon_X)^*} & \mathbb{F}(\mathbb{F}(\mathbb{G}(\mathbb{F}(Y))), \mathbb{F}(\mathbb{G}(X))) & & \mathbb{E}(Y, \mathbb{G}(X)). \\
 \downarrow \mathbb{F}(\eta_Y)^* & & \downarrow \mathbb{F}(\eta_Y)^* & \swarrow \beta_{Y, \mathbb{G}(X)} & \\
 \mathbb{F}(\mathbb{F}(Y), X) & \xleftarrow{(\varepsilon_X)^*} & \mathbb{F}(\mathbb{F}(Y), \mathbb{F}(\mathbb{G}(X))) & &
 \end{array}$$

The top left square commutes since ε is an extriangulated natural transformation. The bottom left square commutes by virtue of \mathbb{F} being a bifunctor. The triangle on the right commutes as all the maps are natural. It remains to observe that the vertical composition on the left is the identity. \square

Corollary 3.9. Let $(\mathbb{F}, \beta): (\mathcal{A}, \mathbb{E}, \mathfrak{s}) \rightarrow (\mathcal{B}, \mathbb{F}, \mathfrak{t})$ be an extriangulated left adjoint of $(\mathbb{G}, \gamma): (\mathcal{B}, \mathbb{F}, \mathfrak{t}) \rightarrow (\mathcal{A}, \mathbb{E}, \mathfrak{s})$. With the isomorphisms from Lemma 3.8 the following diagram commutes

$$\begin{array}{ccccc}
 \mathbb{F}(\mathbb{F}(X), Y') & \xleftarrow{e_*} & \text{Hom}_{\mathcal{B}}(\mathbb{F}(X), Y) & \xrightarrow{(\beta_{X', X}(d))^*} & \mathbb{F}(\mathbb{F}(X'), Y) \\
 \updownarrow \cong & & \updownarrow \cong & & \updownarrow \cong \\
 \mathbb{E}(X, \mathbb{G}(Y')) & \xleftarrow{(\gamma_{Y, Y'}(e))^*} & \text{Hom}_{\mathcal{A}}(X, \mathbb{G}(Y')) & \xrightarrow{d^*} & \mathbb{E}(X', \mathbb{G}(Y))
 \end{array}$$

for any $d \in \mathbb{E}(X', X)$ and $e \in \mathbb{F}(Y, Y')$.

Proof. As the vertical maps are compositions of γ and the unit, or β and the counit, the claim follows from the definition of an extriangulated functor and Remark 3.2. \square

From the construction of the higher extensions and Proposition 3.6 we immediately obtain:

Corollary 3.10. Let $(\mathbb{F}, \beta): (\mathcal{A}, \mathbb{E}, \mathfrak{s}) \rightarrow (\mathcal{B}, \mathbb{F}, \mathfrak{t})$ be an extriangulated left adjoint of $(\mathbb{G}, \gamma): (\mathcal{B}, \mathbb{F}, \mathfrak{t}) \rightarrow (\mathcal{A}, \mathbb{E}, \mathfrak{s})$. We assume that the higher extensions are well-defined in $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$ and $(\mathcal{B}, \mathbb{F}, \mathfrak{t})$ through the same construction, in the sense of Proposition 3.6. Then $\mathbb{F}^n(\mathbb{F}(Y), X) \cong \mathbb{E}^n(Y, \mathbb{G}(X))$ for all $Y \in \mathcal{A}$ and $X \in \mathcal{B}$. Moreover, these isomorphisms are compatible with the cup product similar to Corollary 3.9. \square

3.11. Biextriangulated functors

Loosely speaking, a biextriangulated functor is a bifunctor that is extriangulated functor in each component, satisfying an additional compatibility condition between the components. Definition 3.12 below only requires an expected compatibility condition; later we discuss a stronger notion.

Definition 3.12. A *biextriangulated functor*

$$(\mathbb{F}, \varphi, \psi): (\mathcal{A}, \mathbb{E}, \mathfrak{s}) \times (\mathcal{B}, \mathbb{F}, \mathfrak{t}) \rightarrow (\mathcal{C}, \mathbb{G}, \mathfrak{u})$$

of extriangulated categories consists of:

- (1) a functor $\mathbb{F}: \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$;
- (2) for each $Y \in \mathcal{B}$, a natural transformation

$$\varphi^Y: \mathbb{E}(-, -) \rightarrow \mathbb{G}(\mathbb{F}(-, Y), \mathbb{F}(-, Y));$$

- (3) for each $X \in \mathcal{A}$, a natural transformation

$$\psi^X: \mathbb{F}(-, -) \rightarrow \mathbb{G}(\mathbb{F}(X, -), \mathbb{F}(X, -));$$

such that

- (a) for each $Y \in \mathcal{B}$, the functor $(\mathbb{F}(-, Y), \varphi^Y): \mathcal{A} \rightarrow \mathcal{C}$ is extriangulated;
- (b) for each $X \in \mathcal{A}$, the functor $(\mathbb{F}(X, -), \psi^X): \mathcal{B} \rightarrow \mathcal{C}$ is extriangulated;
- (c) for $X, X' \in \mathcal{A}$ and $g: Y \rightarrow Y'$ in \mathcal{B} , the following diagram commutes

$$\begin{CD} \mathbb{E}(X', X) @>\varphi_{X', X}^Y>> \mathbb{G}(\mathbb{F}(X', Y), \mathbb{F}(X, Y)) \\ @V\varphi_{X', X}^{Y'}VV @VV\mathbb{F}(X, g)_*V \\ \mathbb{G}(\mathbb{F}(X', Y'), \mathbb{F}(X, Y')) @>\mathbb{F}(X', g)^*>> \mathbb{G}(\mathbb{F}(X', Y), \mathbb{F}(X, Y')) \end{CD}$$

- (d) for $f: X \rightarrow X'$ in \mathcal{A} and $Y, Y' \in \mathcal{B}$, the following diagram commutes

$$\begin{CD} \mathbb{F}(Y', Y) @>\psi_{Y', Y}^X>> \mathbb{G}(\mathbb{F}(X, Y'), \mathbb{F}(X, Y)) \\ @V\psi_{Y', Y}^{X'}VV @VV\mathbb{F}(f, Y)_*V \\ \mathbb{G}(\mathbb{F}(X', Y'), \mathbb{F}(X', Y)) @>\mathbb{F}(f, Y')^*>> \mathbb{G}(\mathbb{F}(X, Y'), \mathbb{F}(X', Y)). \end{CD}$$

Remark 3.13. Conditions (c) and (d) mean that φ and ψ are *extranatural* in \mathcal{B} and \mathcal{A} , respectively; see [13].

Example 3.14. Let $F: \mathcal{E} \times \mathcal{F} \rightarrow \mathcal{G}$ be a *biexact* functor of exact categories; this means that $F(X, -)$ and $F(-, Y)$ are exact functors for every $X \in \mathcal{E}$ and $Y \in \mathcal{F}$; cf. [60, II.7.4]. We let φ^Y and ψ^X be the natural transformations obtained by applying $F(-, Y)$ and $F(X, -)$ to a short exact sequence in \mathcal{E} and \mathcal{F} , respectively; see Example 3.3. Hence, (a) and (b) hold. We show that (c) holds without any further assumptions. Let $X \rightarrow W \rightarrow X'$ be a short exact sequence in \mathcal{E} and $g: Y \rightarrow Y'$ a morphism in \mathcal{F} . By [10, Proposition 3.1], the morphism of short exact sequences induced by $F(-, g)$ factors as follows

$$\begin{array}{ccccc}
 F(X, Y) & \longrightarrow & F(W, Y) & \longrightarrow & F(X', Y) \\
 \downarrow F(X, g) & & \downarrow & & \downarrow = \\
 F(X, Y') & \longrightarrow & Z & \longrightarrow & F(X', Y) \\
 \downarrow = & & \downarrow & & \downarrow F(X', g) \\
 F(X, Y') & \longrightarrow & F(W, Y') & \longrightarrow & F(X', Y')
 \end{array}$$

where the top left and bottom right squares are bicartesian. The top row is precisely $\varphi^Y_{X', X}([X \rightarrow W \rightarrow X'])$ and the bottom row is $\varphi^{Y'}_{X', X}([X \rightarrow W \rightarrow X'])$. Hence the middle row can be obtained by applying $\text{Ext}^1_{\mathcal{G}}(F(X', Y), F(X, g))$ to the top row, or by applying $\text{Ext}^1_{\mathcal{G}}(F(X', g), F(X, Y'))$ to the bottom row. In particular, (c) holds. An analogous argument shows (d). So (F, φ, ψ) is a biexactriangulated functor.

Conversely, from Example 3.3 it is clear that any biexactriangulated functor of exact categories is a biexact functor.

Example 3.15. Let $(F, \vartheta, \zeta): \mathcal{S} \times \mathcal{T} \rightarrow \mathcal{U}$ be a *bitriangulated* functor; this means $\vartheta: F(\Sigma-, -) \rightarrow \Sigma F(-, -)$ and $\zeta: F(-, \Sigma-) \rightarrow \Sigma F(-, -)$ are natural transformations such that $(F(-, Y), \vartheta_{-, Y})$ and $(F(X, -), \zeta_{X, -})$ are triangulated functors for all $X \in \mathcal{S}$ and $Y \in \mathcal{T}$, and the following diagram anticommutes.

$$\begin{array}{ccc}
 F(\Sigma X, \Sigma Y) & \xrightarrow{\vartheta_{X, \Sigma Y}} & \Sigma F(X, \Sigma Y) \\
 \zeta_{\Sigma X, Y} \downarrow & (-1) & \downarrow \Sigma \zeta_{X, Y} \\
 \Sigma F(\Sigma X, Y) & \xrightarrow{\Sigma \vartheta_{X, Y}} & \Sigma^2 F(X, Y)
 \end{array} \tag{3.15.1}$$

We define the natural transformations φ^Y and ψ^X using $\vartheta_{-, Y}$ and $\zeta_{X, -}$, respectively, similarly to how β was defined in Example 3.4. Hence (a) and (b) hold. In this setting, condition (c) is the identification

$$(\Sigma F(X, g)) \circ \vartheta_{X, Y} \circ F(f, Y) = \vartheta_{X, Y'} \circ F(f, Y') \circ F(X', g)$$

for all $f: X' \rightarrow \Sigma X$ in \mathcal{S} and $g: Y \rightarrow Y'$ in \mathcal{T} . This holds as $\vartheta, F(f, -)$ and $F(-, g)$ are natural transformations. An analogous argument shows (d). So (F, φ, ψ) is a biexactriangulated functor.

The above argument does not use (3.15.1). In fact, a biextriangulated functor of triangulated categories need not induce a bitriangulated functor. We resolve this discrepancy using ‘strong’ biextriangulated functors in the next Section.

3.16. Strong biextriangulated functors

We now consider a stronger compatibility condition between the components of a biextriangulated functor. This will resolve the mismatch between biextriangulated and bitriangulated functors of triangulated categories, while not affecting the match between biextriangulated and biexact functors of exact categories. A possible disadvantage of this definition is that we need to assume that higher extensions exist, however these always exist for triangulated categories.

Definition 3.17. Let $F = (F, \varphi, \psi) : (\mathcal{A}, \mathbb{E}, \mathfrak{s}) \times (\mathcal{B}, \mathbb{F}, \mathfrak{t}) \rightarrow (\mathcal{C}, \mathbb{G}, \mathfrak{u})$ be a biextriangulated functor. We assume that higher extensions are well-defined in $(\mathcal{C}, \mathbb{G}, \mathfrak{u})$. We say F is *strong*, if for every $d \in \mathbb{E}(X, X')$ and $e \in \mathbb{F}(Y, Y')$ one has

$$\psi_{Y, Y'}^{X'}(e) \smile \varphi_{X, X'}^Y(d) = -\varphi_{X, X'}^{Y'}(d) \smile \psi_{Y, Y'}^X(e) \tag{3.17.1}$$

in $\mathbb{G}^2(F(X, Y), F(X', Y'))$.

Lemma 3.18. Let $F = (F, \varphi, \psi) : (\mathcal{A}, \mathbb{E}, \mathfrak{s}) \times (\mathcal{B}, \mathbb{F}, \mathfrak{t}) \rightarrow (\mathcal{C}, \mathbb{G}, \mathfrak{u})$ be a strong biextriangulated functor and assume that higher extensions are well-defined in each of the categories through the same construction, as in Proposition 3.6. Then

$$(\psi^{X'})_{Y, Y'}^j(e) \smile (\varphi^Y)_{X, X'}^i(d) = (-1)^{ij} (\varphi^{Y'})_{X, X'}^i(d) \smile (\psi^X)_{Y, Y'}^j(e)$$

for every $d \in \mathbb{E}^i(X, X')$ and $e \in \mathbb{F}^j(Y, Y')$.

Proof. By the construction in Proposition 3.6, the natural transformation $(\varphi^Y)^i$ is fully determined by $(\varphi^Y)^{\otimes i}$ on

$$\mathbb{E}(W_1, X) \otimes_R \mathbb{E}(W_2, W_1) \otimes_R \cdots \otimes_R \mathbb{E}(Y, W_{i-1}).$$

We obtain the claim by interchanging the \mathbb{E} and \mathbb{F} factors of $\mathbb{E}^i(X, X') \otimes_R \mathbb{F}^j(Y, Y')$. \square

For exact categories condition (3.17.1) is superfluous:

Proposition 3.19. Let $F : \mathcal{E} \times \mathcal{F} \rightarrow \mathcal{G}$ be a biexact functor. Then F is strong viewed as a biextriangulated functor. In particular, there is a one-to-one correspondence between biexact functors and strong biextriangulated functors of exact categories.

Proof. Let $d = [X \xrightarrow{f} Y \xrightarrow{g} Z] \in \text{Ext}_{\mathcal{E}}^1(X, Z)$ and $e = [X' \xrightarrow{f'} Y' \xrightarrow{g'} Z'] \in \text{Ext}_{\mathcal{F}}^1(X', Z')$. Then $F(d, X') \smile F(Z, e)$ is

$$[F(X, X') \xrightarrow{F(f, X')} F(Y, X') \xrightarrow{F(g, f')} F(Z, Y') \xrightarrow{F(Z, g')} F(Z, Z')]$$

and $F(X, e) \smile F(d, Z')$ is

$$[F(X, X') \xrightarrow{F(X, f')} F(X, Y') \xrightarrow{F(f, g')} F(Y, Z') \xrightarrow{F(g, Z')} F(Z, Z')];$$

see Example 2.6. We take the pushout of the span $F(X, Y') \leftarrow F(X, X') \rightarrow F(Y, X')$ and obtain a morphism of exact sequences

$$\begin{array}{ccccc} F(X, X') & \xrightarrow{F(f, X')} & F(Y, X') & \xrightarrow{F(g, X')} & F(Z, X') \\ \downarrow F(X, f') & & \downarrow & & \downarrow = \\ F(X, Y') & \xrightarrow{a} & W & \xrightarrow{b} & F(Z, X') \end{array}$$

The universal property of the pushout yields the commutative diagram

$$\begin{array}{ccccc} F(X, Y') & \xrightarrow{a} & W & \xrightarrow{c} & F(Y, Y') \\ \downarrow & & \downarrow b & & \downarrow F(g, Y') \\ 0 & \longrightarrow & F(Z, X') & \xrightarrow{F(Z, f')} & F(Z, Y') \\ & & \downarrow & & \downarrow F(Z, g') \\ & & 0 & \longrightarrow & F(Z, Z') \end{array}$$

such that $ca = F(f, Y')$, and so the rectangle defined by the top two squares is a pushout square. Since (a, b) is a kernel-cokernel pair the top left square is a pushout square. By the pasting lemma for pushout squares, every square is a pushout square; see [48, p. 72, Exercise 8]. So we obtain an extension

$$[F(X, X') \xrightarrow{\begin{pmatrix} F(X, f') \\ -F(f, X') \end{pmatrix}} F(X, Y') \oplus F(Y, X') \rightarrow F(Y, Y') \rightarrow F(Z, Z')],$$

and a commutative diagram

$$\begin{array}{ccccccc} F(X, X') & \xrightarrow{-F(f, X')} & F(Y, X') & \xrightarrow{F(g, f')} & F(Z, Y') & \xrightarrow{F(Z, g')} & F(Z, Z') \\ \uparrow = & & \uparrow & & \uparrow & & \uparrow = \\ F(X, X') & \xrightarrow{\begin{pmatrix} F(X, f') \\ -F(f, X') \end{pmatrix}} & F(X, Y') \oplus F(Y, X') & \longrightarrow & F(Y, Y') & \longrightarrow & F(Z, Z') \\ \downarrow = & & \downarrow & & \downarrow & & \downarrow = \\ F(X, X') & \xrightarrow{F(X, f')} & F(X, Y') & \xrightarrow{F(f, g')} & F(Y, Z') & \xrightarrow{F(g, Z')} & F(Z, Z') \end{array}$$

which exhibits the first claim. The second assertion follows from combining this with Example 3.14. \square

Proposition 3.20. *Let $(F, \varphi, \psi): \mathcal{S} \times \mathcal{T} \rightarrow \mathcal{U}$ be a strong bixtriangulated functor of triangulated categories. With*

$$\begin{aligned} \vartheta_{X,Y} &:= \varphi_{\Sigma X, X}^Y(\text{id}_{\Sigma X}): F(\Sigma X, Y) \rightarrow \Sigma F(X, Y) \quad \text{and} \\ \zeta_{X,Y} &:= \psi_{\Sigma Y, Y}^X(\text{id}_{\Sigma Y}): F(X, \Sigma Y) \rightarrow \Sigma F(X, Y) \end{aligned}$$

the functor (F, ϑ, ζ) is a bitriangulated functor.

Moreover, there is a one-to-one correspondence between bitriangulated functors and strong bixtriangulated functors on triangulated categories.

Proof. We assume that $(F, \varphi, \psi): \mathcal{S} \times \mathcal{T} \rightarrow \mathcal{U}$ is a bixtriangulated functor of triangulated categories, and that ϑ and ζ are given as above. From Example 3.4 we know that F , together with ϑ and ζ , is a triangulated functor in each component. It remains to show (3.15.1). By Example 2.7 this is equivalent to

$$\zeta_{X,Y} \smile \vartheta_{X,\Sigma Y} = -\vartheta_{X,Y} \smile \zeta_{\Sigma X,Y}.$$

This is just a specialisation of the definition of a strong bixtriangulated functor.

For the second part, it remains to show that the bixtriangulated functor that is induced by a bitriangulated functor is strong. This is straightforward to check using Examples 2.7 and 3.4. \square

We end this section with examples of bixtriangulated functors. For derived categories of module categories, the derived tensor product and derived Hom-functor are bixtriangulated. However, for module categories with the abelian exact structure the tensor product and the Hom-functor need not be bixtriangulated. This can be rectified by restricting to flat modules.

Example 3.21. Let R be a commutative ring. Let \mathcal{X} be an essentially small R -linear category. Let $\text{Mod}(\mathcal{X})$ and $\text{Mod}(\mathcal{X}^{\text{op}})$ be the categories of R -linear covariant and contravariant functors, respectively, of the form $\mathcal{X} \rightarrow \text{Mod}(R)$. For $F \in \text{Mod}(\mathcal{X}^{\text{op}})$ and $G \in \text{Mod}(\mathcal{X})$ define an R -bilinear functor $\mathcal{X}^{\text{op}} \times \mathcal{X} \rightarrow \text{Mod}(R)$ by $(X, Y) \mapsto F(X) \otimes_R G(Y)$. Taking coends, discussed in Section 2.4, gives a functor

$$\otimes_{\mathcal{X}}: \text{Mod}(\mathcal{X}^{\text{op}}) \times \text{Mod}(\mathcal{X}) \rightarrow \text{Mod}(R), \quad (F, G) \mapsto \int^W F(W) \otimes_R G(W);$$

see [21, Section 2]. When \mathcal{X} is a category with one object, the endomorphism ring is an R -algebra A and $\otimes_{\mathcal{X}}$ is the classical tensor product of modules over A .

Let $\text{Flat}(\mathcal{X})$ be the full subcategory of $\text{Mod}(\mathcal{X})$ consisting of functors F such that $F \otimes_{\mathcal{X}} - : \text{Mod}(\mathcal{X}^{\text{op}}) \rightarrow \text{Mod}(R)$ is exact, meaning F is *flat* in the sense of [53, Theorem 3.2(1)]. The subcategory $\text{Flat}(\mathcal{X})$ (respectively, $\text{Flat}(\mathcal{X}^{\text{op}})$) is extension-closed in $\text{Mod}(\mathcal{X})$ (respectively, $\text{Mod}(\mathcal{X}^{\text{op}})$). With the abelian exact structure on $\text{Mod}(R)$, one can check that the restriction of $\otimes_{\mathcal{X}}$ defines a biexact functor

$$\text{Flat}(\mathcal{X}^{\text{op}}) \times \text{Flat}(\mathcal{X}) \rightarrow \text{Mod}(R).$$

Thus this restriction is a strong biextriangulated functor by Example 3.14.

The functor categories and their subcategories of flat functors are strongly connected to the category of modules over a non-unital R -algebra. For a small R -linear category \mathcal{X} the *functor ring* is $A := \bigoplus_{X, Y \in \text{sk}(\mathcal{X})} \text{Hom}_{\mathcal{X}}(X, Y)$; this ring was studied in [23, Chapter II], also see [24]. This is a possibly non-unital ring with enough idempotents. By [23, II, Proposition 2], there is an R -linear equivalence between $\text{Mod}(\mathcal{X})$ and the category of unital left A -modules; one says M is *unital* if $AM = M$. This equivalence restricts to an equivalence between $\text{Flat}(\mathcal{X})$ and the category of flat unital left A -modules. Similarly, the category $\text{Mod}(\mathcal{X}^{\text{op}})$ (respectively $\text{Flat}(\mathcal{X}^{\text{op}})$) is equivalent to the category of (respectively, flat) unital right A -modules. Under this equivalence, the functor $\otimes_{\mathcal{X}}$ corresponds to the classical tensor product \otimes_A ; see for example [17, Lemma 2].

3.22. Stabilisation of extriangulated categories

There are a few ways to construct extriangulated categories. One is as a stabilisation of an exact category. We will see that any biextriangulated functor induced on the stabilisation from a biexact functor is strong.

Let $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$ be an extriangulated category. Let \mathcal{I} be a full subcategory of \mathcal{A} that is closed under finite direct sums and consists of objects that are \mathbb{E} -projective and \mathbb{E} -injective; for the definition of the latter see Section 2.8. The *stabilisation of \mathcal{A} along \mathcal{I}* , denoted by $\underline{\mathcal{A}}_{\mathcal{I}}$, has the same objects as \mathcal{A} and the morphisms are defined by

$$\text{Hom}_{\underline{\mathcal{A}}_{\mathcal{I}}}(X, Y) := \text{Hom}_{\mathcal{A}}(X, Y) / [\mathcal{I}](X, Y),$$

where $[\mathcal{I}](X, Y)$ is the two-sided ideal of morphisms $X \rightarrow Y$ in \mathcal{A} factoring through objects in \mathcal{I} .

By [51, Proposition 3.30], the extriangulated structure on \mathcal{A} descends to the stabilisation $\underline{\mathcal{A}}_{\mathcal{I}}$. The extriangulated structure $(\underline{\mathbb{E}}, \underline{\mathfrak{s}})$ on $\underline{\mathcal{A}}_{\mathcal{I}}$ is as follows: For objects $X, Y \in \underline{\mathcal{A}}_{\mathcal{I}}$, we have

$$\underline{\mathbb{E}}(Y, X) = \mathbb{E}(Y, X) \quad \text{and} \quad \underline{\mathfrak{s}}(d) = [X \xrightarrow{[f]} W \xrightarrow{[g]} Y]$$

where $\mathfrak{s}(d) = [X \xrightarrow{f} W \xrightarrow{g} Y]$.

Lemma 3.23. *Let $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$ be an extriangulated category and \mathcal{I} a full subcategory closed under finite direct sums that consists of objects that are \mathbb{E} -projective and \mathbb{E} -injective. If \mathcal{A} has enough \mathbb{E} -projectives or \mathbb{E} -injectives, then so does $\underline{\mathcal{A}}_{\mathcal{I}}$.*

Furthermore, whenever higher extensions are well-defined in $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$, then they are well-defined in $(\underline{\mathcal{A}}_{\mathcal{I}}, \underline{\mathbb{E}}, \underline{\mathfrak{s}})$ through the same construction, and

$$\underline{\mathbb{E}}^n(X, Y) = \mathbb{E}^n(X, Y)$$

for all $X, Y \in \mathcal{A}$ and $n \geq 1$.

Proof. The first claim holds, as under stabilisation \mathbb{E} -projective and \mathbb{E} -injective objects become $\underline{\mathbb{E}}$ -projective and $\underline{\mathbb{E}}$ -injective objects, respectively, and stabilisation preserves extriangles. The second claim is straightforward to check as the construction of the higher extensions commutes with the stabilisation. \square

Lemma 3.24 below recovers the $n = 1$ case of [5, Example 5.9] where (F, β) is the identity, $\mathcal{I} = 0$ and \mathcal{J} consists of all the projective-injectives.

Lemma 3.24. *Let $(F, \beta): (\mathcal{A}, \mathbb{E}, \mathfrak{s}) \rightarrow (\mathcal{B}, \mathbb{F}, \mathfrak{t})$ be an extriangulated functor. Let \mathcal{I} and \mathcal{J} be full subcategories of \mathcal{A} and \mathcal{B} , respectively, that are closed under finite direct sums and consist of objects that are projective and injective with respect to the respective extriangulated structure. If $F(\mathcal{I}) \subseteq \mathcal{J}$, then (F, β) induces an extriangulated functor*

$$(F, \beta): (\underline{\mathcal{A}}_{\mathcal{I}}, \underline{\mathbb{E}}, \underline{\mathfrak{s}}) \rightarrow (\underline{\mathcal{B}}_{\mathcal{J}}, \underline{\mathbb{F}}, \underline{\mathfrak{t}}).$$

Proof. By assumption we have $F([\mathcal{I}](X, Y)) \subseteq [\mathcal{J}](F(X), F(Y))$. Hence F induces an additive functor $\underline{F}: \underline{\mathcal{A}}_{\mathcal{I}} \rightarrow \underline{\mathcal{B}}_{\mathcal{J}}$. The natural transformation $\underline{\beta}$ is well-defined since \mathcal{I} and \mathcal{J} consist of projective and injective objects in the respective extriangulated structure. It now follows from the definition of $\underline{\mathfrak{s}}$ and $\underline{\mathfrak{t}}$ that $(\underline{F}, \underline{\beta})$ is an extriangulated functor. \square

Proposition 3.25. *Let $(F, \varphi, \psi): (\mathcal{A}, \mathbb{E}, \mathfrak{s}) \times (\mathcal{B}, \mathbb{F}, \mathfrak{t}) \rightarrow (\mathcal{C}, \mathbb{G}, \mathfrak{u})$ be a bixtriangulated functor. Let \mathcal{I}, \mathcal{J} and \mathcal{K} be full subcategories of \mathcal{A}, \mathcal{B} and \mathcal{C} , respectively, that are closed under finite direct sums and consist of objects that are projective and injective with respect to the respective extriangulated structure.*

If $F(\mathcal{I}, \mathcal{B}) \subseteq \mathcal{K}$ and $F(\mathcal{A}, \mathcal{J}) \subseteq \mathcal{K}$, then (F, φ, ψ) induces a bixtriangulated functor $(\underline{F}, \underline{\varphi}, \underline{\psi}): (\underline{\mathcal{A}}_{\mathcal{I}}, \underline{\mathbb{E}}, \underline{\mathfrak{s}}) \times (\underline{\mathcal{B}}_{\mathcal{J}}, \underline{\mathbb{F}}, \underline{\mathfrak{t}}) \rightarrow (\underline{\mathcal{C}}_{\mathcal{K}}, \underline{\mathbb{G}}, \underline{\mathfrak{u}})$. Moreover, if (F, φ, ψ) is strong, then so is the induced functor $(\underline{F}, \underline{\varphi}, \underline{\psi})$.

Proof. The first claim follows from Lemma 3.24, and the second claim follows from Lemma 3.23. \square

Stabilisation of an extriangulated category is motivated by the stabilisation of Frobenius exact categories. An exact category \mathcal{E} is *Frobenius*, if it has enough projectives and

enough injectives and the projectives and injectives coincide. In this situation, the stabilisation $\underline{\mathcal{E}} = \underline{\mathcal{E}}_{\mathcal{I}}$ with respect to the subcategory of projective-injective objects \mathcal{I} is triangulated; see [27, I.2]. As every biexact functor is strong when viewed as a bixtriangulated functor, such a functor always induces a strong bixtriangulated functor on any of its stabilisations by Proposition 3.25.

Example 3.26. Let S be a regular local ring. A *matrix factorisation* of $f \in S$ is a pair of maps of free modules $(\Phi: F \rightarrow G, \Psi: G \rightarrow F)$ such that

$$\Phi\Psi = f \operatorname{id}_G \quad \text{and} \quad \Psi\Phi = f \operatorname{id}_F .$$

The free modules F and G have the same rank, and when they have rank n , we say (Φ, Ψ) has *size* n . Matrix factorisations were introduced in [15, Section 5]; also see [63, Chapter 7]. We denote by $\operatorname{mf}_S(f)$ the category of matrix factorisations of f . Then $\operatorname{mf}_S(f)$ is Frobenius and the only indecomposable projective-injective objects are $(\operatorname{id}_S, f \operatorname{id}_S)$ and $(f \operatorname{id}_S, \operatorname{id}_S)$. The stabilisation $\underline{\operatorname{mf}}_S(f)$ is equivalent as a triangulated category to the stabilisation of the category of maximal Cohen–Macaulay modules over $S/(f)$; see [15, Section 6] and [63, Theorem 7.4].

Suppose K is a field and that $S = K[[x_1, \dots, x_s]]$ and $T = K[[y_1, \dots, y_t]]$ are power series rings. We fix $f \in S$ and $g \in T$. The tensor product of $(\Phi, \Psi) \in \operatorname{mf}_S(f)$ and $(\Phi', \Psi') \in \operatorname{mf}_T(g)$ is defined as

$$(\Phi, \Psi) \hat{\otimes} (\Phi', \Psi') := \left(\left(\begin{array}{cc} \Phi \otimes \operatorname{id} & \operatorname{id} \otimes \Phi' \\ -\operatorname{id} \otimes \Psi' & \Psi \otimes \operatorname{id} \end{array} \right), \left(\begin{array}{cc} \Psi \otimes \operatorname{id} & -\operatorname{id} \otimes \Phi' \\ \operatorname{id} \otimes \Psi' & \Phi \otimes \operatorname{id} \end{array} \right) \right) ;$$

see [64, Definition 2.1]. The tensor product is a matrix factorisation of $f + g$ over $S \otimes_K T$, and this yields a functor

$$\hat{\otimes} : \operatorname{mf}_S(f) \times \operatorname{mf}_T(g) \rightarrow \operatorname{mf}_{S \otimes_K T}(f + g) .$$

By [64, Lemmas 2.2, 2.8], this functor is biexact, and, by [64, Lemma 2.3], it induces a bixtriangulated functor on the stabilisations. However, as

$$(\Phi, \Psi) \hat{\otimes} (\operatorname{id}_T, g \operatorname{id}_T) \cong (\operatorname{id}_{S \otimes_K T}, (f + g) \operatorname{id}_{S \otimes_K T}) \oplus ((f + g) \operatorname{id}_{S \otimes_K T}, \operatorname{id}_{S \otimes_K T}) ,$$

the tensor product does not induce a functor on the intermediate stabilisations. In particular, it does not induce a tensor product on the category $\underline{\operatorname{mf}}_S(f)_{\mathcal{I}}$ for $\mathcal{I} = \{(\operatorname{id}, f \operatorname{id})\}$, which is equivalent to the category of maximal Cohen–Macaulay modules over $S/(f)$; see [63, Theorem 7.4].

There is another tensor product, called the multiplicative tensor product, defined in [22, Definition 2.6]. While it is a biexact functor, it does not respect the projective-injective objects, and hence does not induce a functor on the stabilisation.

4. Tensor extriangulated categories

In this section, we introduce tensor extriangulated categories, generalising the notion of a tensor triangulated category; cf. [32, Appendix A.2].

4.1. Monoidal categories

Recall that a *monoidal category* $(\mathcal{A}, \otimes, \mathbb{1})$ is a category \mathcal{A} together with a functor

$$\otimes : \mathcal{A} \times \mathcal{A} \rightarrow \mathcal{A},$$

called the *tensor product*, and a *unit* object $\mathbb{1} \in \mathcal{A}$ equipped with isomorphisms

$$\begin{aligned} \alpha_{X,Y,Z} : X \otimes (Y \otimes Z) &\xrightarrow{\cong} (X \otimes Y) \otimes Z, \\ \lambda_X : \mathbb{1} \otimes X &\xrightarrow{\cong} X \quad \text{and} \quad \rho_X : X \otimes \mathbb{1} \xrightarrow{\cong} X \end{aligned}$$

that are natural in each argument, satisfying some coherence axioms. A monoidal category is *symmetric* if, in addition, there exists an isomorphism

$$\sigma_{X,Y} : X \otimes Y \xrightarrow{\cong} Y \otimes X,$$

also natural in each argument, satisfying some coherence axioms. For a detailed definition see [48, XI.1], and [37] for minimally sufficient coherent axioms.

4.2. Tensor extriangulated categories

We now have all the ingredients to define a monoidal extriangulated category. Roughly speaking, it is an extriangulated category with a monoidal structure that respects the extriangulated structure. Explicitly, this means:

Definition 4.3. We call a tuple $(\mathcal{A}, \mathbb{E}, \mathfrak{s}, \otimes, \mathbb{1})$ a *monoidal extriangulated category* if:

- (1) $(\mathcal{A}, \otimes, \mathbb{1})$ is a monoidal category;
- (2) $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$ is an extriangulated category;
- (3) $\otimes = (- \otimes -, \varphi, \psi) : (\mathcal{A}, \mathbb{E}, \mathfrak{s}) \times (\mathcal{A}, \mathbb{E}, \mathfrak{s}) \rightarrow (\mathcal{A}, \mathbb{E}, \mathfrak{s})$ is a biextriangulated functor; and
- (4) the natural isomorphisms α, λ, ρ from the monoidal structure are extriangulated natural transformations in each variable.

We say a monoidal extriangulated category $(\mathcal{A}, \mathbb{E}, \mathfrak{s}, \otimes, \mathbb{1})$ is *strong*, if the biextriangulated functor $(- \otimes -, \varphi, \psi)$ is strong (see Definition 3.17).

A *tensor extriangulated category* is a monoidal extriangulated category where the monoidal structure is symmetric and the corresponding natural transformation σ is extriangulated in each variable.

Condition (4) has various consequences. For example, it yields

$$\varphi_{X,Y}^{\mathbb{1}} = (\rho_Y^{-1})_* \circ (\rho_X)^* \quad \text{and} \quad \psi_{X,Y}^{\mathbb{1}} = (\lambda_Y^{-1})_* \circ (\lambda_X)^*. \tag{4.3.1}$$

In particular, $\varphi^{\mathbb{1}}$ and $\psi^{\mathbb{1}}$ are natural isomorphisms. Further, in a tensor extriangulated category, the natural transformations φ and ψ from the tensor product determine each other; explicitly

$$(\sigma_{X',Y})_* \circ \varphi_{X,X'}^Y = (\sigma_{X,Y})^* \circ \psi_{X,X'}^Y$$

as σ is an extriangulated natural isomorphism in each variable.

4.4. Ring of higher extensions

Recall from Remark 2.13 that $\mathbb{E}^*(X, X)$ with the cup product is a graded ring. Similarly as for triangulated categories, a tensor structure makes the graded endomorphism ring graded-commutative.

Lemma 4.5. *Let $(\mathcal{A}, \mathbb{E}, \mathfrak{s}, \otimes, \mathbb{1})$ be a strong monoidal extriangulated category. We assume that higher extensions are well-defined in $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$. Then $\mathbb{E}^*(\mathbb{1}, \mathbb{1})$ with the cup product is a graded-commutative graded ring.*

Proof. Let $d \in \mathbb{E}^i(\mathbb{1}, \mathbb{1})$ and $e \in \mathbb{E}^j(\mathbb{1}, \mathbb{1})$. Then

$$\begin{aligned} d \smile e &= (\varphi^{\mathbb{1}})_{\mathbb{1},\mathbb{1}}^i(d) \smile (\psi^{\mathbb{1}})_{\mathbb{1},\mathbb{1}}^j(e) \\ &= (-1)^{ij} (\psi^{\mathbb{1}})_{\mathbb{1},\mathbb{1}}^j(e) \smile (\varphi^{\mathbb{1}})_{\mathbb{1},\mathbb{1}}^i(d) \\ &= (-1)^{ij} e \smile d. \end{aligned}$$

For the first and third equalities we implicitly use a higher version of (4.3.1). The second equality follows from Lemma 3.18. \square

Example 4.6. In Example 2.3 we explained that any triangulated category is naturally an extriangulated category. By Proposition 3.19 a strong bixtriangulated functor on a triangulated category is a bitriangulated functor. Hence the definition of a strong tensor extriangulated category recovers the notion of a tensor triangulated category in the sense of [32, A.2]. Note that other sources use a weaker definition of a tensor triangulated category, they might not assume that the monoidal structure is strong or the natural transformations α , ρ , λ and σ are extriangulated.

The extension ring $\mathbb{E}^*(\mathbb{1}, \mathbb{1})$ of a tensor triangulated category is precisely the non-negatively graded part of the endomorphism ring $\text{End}_{\mathcal{T}}^*(\mathbb{1})$.

Remark 4.7. In an essentially small extriangulated category negative extensions were defined in [26, Definition 5.1]. For a triangulated category these yield the negative part of the endomorphism ring $\text{End}_{\mathcal{T}}^*(\mathbb{1})$; see [26, Proposition 5.4]. We expect that Lemma 4.5 extends to $\mathbb{E}^*(X, X) = \{\mathbb{E}^n(X, X)\}_{n \in \mathbb{Z}}$.

We will see more examples of (strong) tensor extriangulated categories in Section 5.

4.8. Closed tensor extriangulated categories

A symmetric monoidal category $(\mathcal{A}, \otimes, \mathbb{1})$ is *closed* in the sense of [44, p. 119] if there is an *internal hom*

$$\text{hom}(-, -): \mathcal{A}^{\text{op}} \times \mathcal{A} \rightarrow \mathcal{A},$$

meaning a bifunctor such that, for every $X \in \mathcal{A}$, the functor $\text{hom}(X, -)$ is right adjoint to $- \otimes X$.

Definition 4.9. We say a tensor extriangulated category $(\mathcal{A}, \mathbb{E}, \mathfrak{s}, \otimes, \mathbb{1})$ is *closed* if there is a biextriangulated functor $\text{hom}(-, -): \mathcal{A}^{\text{op}} \times \mathcal{A} \rightarrow \mathcal{A}$ such that, for every $X \in \mathcal{A}$, the functor $\text{hom}(X, -)$ is an extriangulated right adjoint to $- \otimes X$.

We do not know whether the internal hom functor $\text{hom}(-, -)$ of a closed and strong tensor extriangulated category is strong.

4.10. Tensor extriangulated functors

The following definition generalises the notion of a tensor triangulated functor from [3, Definition 3].

Definition 4.11. A *tensor extriangulated functor*

$$(F, \beta): (\mathcal{A}, \otimes_{\mathcal{A}}, \mathbb{1}_{\mathcal{A}}) \rightarrow (\mathcal{B}, \otimes_{\mathcal{B}}, \mathbb{1}_{\mathcal{B}})$$

of tensor extriangulated categories consists of:

- (1) an extriangulated functor $(F, \beta): (\mathcal{A}, \mathbb{E}, \mathfrak{s}) \rightarrow (\mathcal{B}, \mathbb{F}, \mathfrak{t})$;
- (2) an isomorphism $u: F(\mathbb{1}_{\mathcal{A}}) \rightarrow \mathbb{1}_{\mathcal{B}}$; and
- (3) a natural isomorphism $\mu_{X, Y}: F(X \otimes_{\mathcal{A}} Y) \rightarrow F(X) \otimes_{\mathcal{B}} F(Y)$

such that the following diagrams commute:

$$\begin{array}{ccc}
 \mathbb{F}(\mathbb{1}_A \otimes_A X) & \xrightarrow{\mu_{\mathbb{1}_A, X}} & \mathbb{F}(\mathbb{1}_A) \otimes_B \mathbb{F}(X) \\
 \downarrow \mathbb{F}(\lambda_X) & & \downarrow u \otimes_B \mathbb{F}(X) \\
 \mathbb{F}(X) & \xleftarrow{\lambda_{\mathbb{F}(X)}} & \mathbb{1}_B \otimes_B \mathbb{F}(X)
 \end{array} \quad \text{and}$$

$$\begin{array}{ccc}
 \mathbb{E}(X, X') & \xrightarrow{\beta_{X, X'}} & \mathbb{F}(\mathbb{F}(X), \mathbb{F}(X')) \\
 \downarrow \varphi_{X, X'}^Y & & \downarrow \varphi_{\mathbb{F}(X), \mathbb{F}(X')}^{\mathbb{F}(Y)} \\
 \mathbb{E}(X \otimes_A Y, X' \otimes_A Y) & & \mathbb{F}(\mathbb{F}(X) \otimes_B \mathbb{F}(Y), \mathbb{F}(X') \otimes_B \mathbb{F}(Y)) \\
 \downarrow \beta_{X \otimes_A Y, X' \otimes_A Y} & & \downarrow (\mu_{X, Y})^* \\
 \mathbb{F}(\mathbb{F}(X \otimes_A Y), \mathbb{F}(X' \otimes_A Y)) & \xrightarrow{(\mu_{X', Y})^*} & \mathbb{F}(\mathbb{F}(X \otimes_A Y), \mathbb{F}(X') \otimes_B \mathbb{F}(Y)).
 \end{array}$$

5. Examples of tensor extriangulated categories

By construction the notion of a tensor extriangulated category generalises that of a tensor triangulated category. In comparison, there seem to be multiple different notions of a tensor exact category. Often the monoidal structure on an exact or abelian category is left exact, but not necessarily exact. In Example 3.21 we defined a biexact functor by restricting the tensor product to flat functors. In Example 5.1 we obtain a monoidal structure, which is not symmetric, by extending the tensor product to bimodules.

Example 5.1. Let R be a commutative ring and \mathcal{X} a small R -linear category. Let $\text{Bimod}(\mathcal{X}, \mathcal{X})$ be the category of R -bilinear functors $\mathcal{X}^{\text{op}} \times \mathcal{X} \rightarrow \text{Mod}(R)$. We extend the functor $\otimes_{\mathcal{X}}$ of Example 3.21 to a functor

$$\begin{aligned}
 \otimes_{\mathcal{X}} : \text{Bimod}(\mathcal{X}, \mathcal{X}) \times \text{Bimod}(\mathcal{X}, \mathcal{X}) &\rightarrow \text{Bimod}(\mathcal{X}, \mathcal{X}), \\
 (\mathbb{F}, \mathbb{G}) &\mapsto \left((X, Y) \mapsto \int^W \mathbb{F}(W, Y) \otimes_R \mathbb{G}(X, W) \right).
 \end{aligned}$$

It is straightforward to check that $\text{Bimod}(\mathcal{X}, \mathcal{X})$ equipped with $\otimes_{\mathcal{X}}$ is a monoidal category and $\text{Hom}_{\mathcal{X}}(-, -)$ serves as the monoidal unit; see [17, Proposition 1]. Now consider the subcategory $\text{Flat}(\mathcal{X}, \mathcal{X})$ of $\text{Bimod}(\mathcal{X}, \mathcal{X})$ consisting of R -bilinear functors \mathbb{F} such that $-\otimes_{\mathcal{X}} \mathbb{F}$ and $\mathbb{F} \otimes_{\mathcal{X}} -$ are exact. Then $\text{Flat}(\mathcal{X}, \mathcal{X})$ equipped with $\otimes_{\mathcal{X}}$ and $\text{Hom}_{\mathcal{X}}(-, -)$ is a monoidal extriangulated category.

When A is the functor ring of \mathcal{X} (see Example 3.21), then there is an equivalence of monoidal categories $\text{Bimod}(\mathcal{X}, \mathcal{X}) \rightarrow \text{Mod}(A^{\text{op}} \otimes_R A)$ under which $\otimes_{\mathcal{X}}$ corresponds to \otimes_A ; see [17, Proposition 2].

The monoidal structure in the previous Example is generally not symmetric. One obtains a symmetric monoidal structure for commutative rings:

Example 5.2. Let R be a commutative ring with local units; this means that for any $r_1, \dots, r_n \in R$, there exists an idempotent $e \in R$ such that $er_i = r_i = r_ie$ for each i . The standard tensor product \otimes_R is a symmetric monoidal structure on the category of unital R -modules $\text{Mod}(R)$; an R -module M is unital provided $M = RM$. However, the tensor product is generally only right exact. An R -module F is flat, if $F \otimes_R -$ is exact. The subcategory of flat R -modules, denoted $\text{Flat}(R)$, is an extension-closed subcategory of $\text{Mod}(R)$, and hence inherits an exact structure. Hence $(\text{Flat}(R), \otimes_R, R, \text{Ext}_R^1)$ is a tensor extriangulated category. The ring of extensions of the unit R is $\mathbb{E}^*(R, R) = \text{Ext}_R^*(R, R) = R$.

A big class of extriangulated categories, that are neither triangulated or exact, are the extension-closed subcategories of triangulated categories; see [51, Remark 2.18]. In general, extension-closed subcategories of a tensor triangulated category need not inherit the monoidal structure. However, in some situations we obtain a tensor extriangulated category.

Example 5.3. Let R be a commutative ring. The derived category $D(\text{Mod}(R))$ of R -modules is a tensor triangulated category with tensor product $\otimes_R^{\mathbf{L}}$ and unit R . The standard t-structure $(\mathcal{U}, \mathcal{V})$ on $D(\text{Mod}(R))$ is given by

$$\begin{aligned} \mathcal{U} &:= \{ X \in D(\text{Mod}(R)) \mid H_{<0}(X) = 0 \} \quad \text{and} \\ \mathcal{V} &:= \{ X \in D(\text{Mod}(R)) \mid H_{>0}(X) = 0 \} , \end{aligned}$$

the subcategories of complexes with homology concentrated in non-negative and non-positive degrees, respectively. As \mathcal{U} and \mathcal{V} are extension-closed subcategories of $D(\text{Mod}(R))$, they inherit an extriangulated structure. Further, the aisle \mathcal{U} is closed under the tensor product; that is $\mathcal{U} \otimes_R^{\mathbf{L}} \mathcal{U} \subseteq \mathcal{U}$ and $R \in \mathcal{U}$. Hence $(\mathcal{U}, \otimes_R^{\mathbf{L}}, R)$ is a tensor extriangulated category.

One gets further examples by taking the intersection of \mathcal{U} with the full subcategory of right-bounded complexes. When R is noetherian, then the intersection of \mathcal{U} with the right-bounded derived category of finitely generated modules yields a tensor extriangulated category as well.

5.4. Stabilisation of extriangulated categories

In Section 3.22 we gave conditions when a (strong) biextriangulated bifunctor induces a (strong) biextriangulated bifunctor on the stabilisation with respect to a subcategory of projective-injective objects. We now focus on monoidal extriangulated categories. We say an extension-closed subcategory \mathcal{I} of a monoidal extriangulated category $(\mathcal{A}, \otimes, \mathbb{1})$ is a two-sided tensor ideal, if $\mathcal{A} \otimes \mathcal{I} \subseteq \mathcal{I}$ and $\mathcal{I} \otimes \mathcal{A} \subseteq \mathcal{I}$. Then the following result is a direct consequence of Proposition 3.25:

Lemma 5.5. *Let $(\mathcal{A}, \otimes, \mathbb{1})$ be a monoidal extriangulated category. If \mathcal{I} is a two-sided tensor ideal consisting of objects that are \mathbb{E} -projective and \mathbb{E} -injective, then $\underline{\mathcal{A}}_{\mathcal{I}}$ inherits the monoidal structure from \mathcal{A} . In particular, $(\underline{\mathcal{A}}_{\mathcal{I}}, \otimes, \mathbb{1})$ is a monoidal extriangulated category. When $(\mathcal{A}, \otimes, \mathbb{1})$ is tensor extriangulated then so is $(\underline{\mathcal{A}}_{\mathcal{I}}, \otimes, \mathbb{1})$. When $(\mathcal{A}, \otimes, \mathbb{1})$ is strong, then so is $(\underline{\mathcal{A}}_{\mathcal{I}}, \otimes, \mathbb{1})$. \square*

Lemma 5.6. *Let $(\mathcal{A}, \mathbb{E}, \mathfrak{s}, \otimes, \mathbb{1})$ be a closed tensor extriangulated category. Then the \mathbb{E} -projectives form a two-sided tensor ideal.*

Proof. By [51, Corollary 3.12], the \mathbb{E} -projectives are extension-closed. Let $X, Y \in \mathcal{A}$. We are assuming $-\otimes X$ has an extriangulated right adjoint $\text{hom}(X, -)$. By Lemma 3.8 this means $\mathbb{E}(Y \otimes X, -) \cong \mathbb{E}(Y, \text{hom}(X, -))$. To say that an object Z is \mathbb{E} -projective is equivalent to saying that $\mathbb{E}(Z, -) = 0$. Altogether we have shown that if Y is \mathbb{E} -projective then so is $Y \otimes X$. Since we are working in a tensor extriangulated category, the tensor product is symmetric, and so we have shown that the \mathbb{E} -projectives form a two-sided tensor ideal. \square

Example 5.7. Let H be a cocommutative Hopf algebra over a field k . For H -modules M and N the comultiplication induces an H -module structure on $M \otimes_k N$. By [49, Chapter 12, Proposition 3], the tensor product is an exact, symmetric monoidal structure on $\text{Mod}(H)$. Hence $(\text{Mod}(H), \otimes_k, k, \text{Ext}_H^1)$ is a tensor extriangulated category.

Further, a finite-dimensional cocommutative Hopf algebra H over a field k is a Frobenius algebra by [43, p. 85, Remark]. Hence $\text{Mod}(H)$, as well as the full subcategory of finite-dimensional H -modules $\text{mod}(H)$, is Frobenius exact. It is well-known that $H \otimes_k M$ is free for every H -module M ; see for example [49, Chapter 12, Proposition 4]. Hence, every non-trivial two-sided tensor ideal of $\text{Mod}(H)$ contains all projective objects. In particular, there are exactly two distinct two-sided tensor ideals of $\text{Mod}(H)$ consisting of projective-injective objects, namely the zero subcategory and the full subcategory of projective-injective objects. Hence, the strong tensor extriangulated category $\text{Mod}(H)$ induces one other strong tensor extriangulated category via stabilisation, namely the tensor triangulated category $\underline{\text{Mod}}(H)$.

A special case of a finite-dimensional cocommutative Hopf algebra is a group algebra kG for a finite group G .

5.8. Dualisable objects

Example 5.7 is an instance of an abelian *rigid* monoidal category, that is an abelian monoidal category in which all objects are dualisable; see [58, p. 254, Examples]. We show that the existence of dualisable objects makes the category Frobenius.

Let $(\mathcal{A}, \otimes, \mathbb{1})$ be a closed monoidal category. The *dualisation functor* is given by

$$(-)^\vee := \text{hom}(-, \mathbb{1}): \mathcal{A}^{\text{op}} \rightarrow \mathcal{A}.$$

An object X in \mathcal{A} is called *strongly dualisable*, if the natural transformation

$$X^\vee \otimes Y \rightarrow \text{hom}(X, Y)$$

is an isomorphism; see [44, Chapter III, §1]. When the monoidal product is symmetric, then there exists a natural transformation $X \rightarrow (X^\vee)^\vee$. This is an isomorphism if X is strongly dualisable by [44, Proposition 1.3]. Note that if $(\mathcal{A}, \mathbb{E}, \mathfrak{s}, \otimes, \mathbb{1})$ is closed tensor extriangulated and if X is strongly dualisable then the natural isomorphisms $X^\vee \otimes - \rightarrow \text{hom}(X, -)$ and $X \otimes - \rightarrow (X^\vee)^\vee \otimes -$ are compositions of extriangulated natural transformations, and hence extriangulated.

Proposition 5.9. *Let $(\mathcal{A}, \mathbb{E}, \mathfrak{s}, \otimes, \mathbb{1})$ be a closed tensor extriangulated category. If each object is strongly dualisable, then \mathbb{E} -projectives are the same as \mathbb{E} -injectives.*

Proof. Let $X, Y \in \mathcal{A}$. Since X is strongly dualisable and \mathcal{A} is symmetric we have

$$\mathbb{E}(Y, X) \cong \mathbb{E}(Y, \text{hom}(X^\vee, \mathbb{1})) \cong \mathbb{E}(Y \otimes X^\vee, \mathbb{1}) \cong \mathbb{E}(X^\vee \otimes Y, \mathbb{1}) \cong \mathbb{E}(X^\vee, Y^\vee);$$

the second and fourth isomorphisms hold by Lemma 3.8. Hence X is \mathbb{E} -injective if and only if X^\vee is \mathbb{E} -projective, and Y is \mathbb{E} -projective if and only if Y^\vee is \mathbb{E} -injective.

Let $U \rightarrow V \rightarrow W \dashrightarrow$ be an extriangle in \mathcal{A} . Since X is strongly dualisable, the functor $\text{hom}(X^\vee, -) \cong (X^\vee)^\vee \otimes - \cong X \otimes -$ is an extriangulated right adjoint to $- \otimes X^\vee$. Combining the adjunction with the unitor we obtain a diagram

$$\begin{array}{ccccccc} \text{Hom}_{\mathcal{A}}(X^\vee, U) & \longrightarrow & \text{Hom}_{\mathcal{A}}(X^\vee, V) & \longrightarrow & \text{Hom}_{\mathcal{A}}(X^\vee, W) & \longrightarrow & \mathbb{E}(X^\vee, U) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \text{Hom}_{\mathcal{A}}(\mathbb{1}, X \otimes U) & \longrightarrow & \text{Hom}_{\mathcal{A}}(\mathbb{1}, X \otimes V) & \longrightarrow & \text{Hom}_{\mathcal{A}}(\mathbb{1}, X \otimes W) & \longrightarrow & \mathbb{E}(\mathbb{1}, X \otimes U) \end{array}$$

which commutes and in which every row is exact and every vertical map is an isomorphism; see Corollary 3.9 and [51, Proposition (2-ii)].

Suppose X is \mathbb{E} -projective. By Lemma 5.6 we have that $X \otimes W$ is \mathbb{E} -projective, and so the morphism $\text{Hom}_{\mathcal{A}}(\mathbb{1}, X \otimes W) \rightarrow \mathbb{E}(\mathbb{1}, X \otimes U)$ in the diagram above is zero, meaning that the map $\text{Hom}_{\mathcal{A}}(X^\vee, V) \rightarrow \text{Hom}_{\mathcal{A}}(X^\vee, W)$ is surjective. Altogether, for any extriangle $U \rightarrow V \rightarrow W \dashrightarrow$ every morphism $X^\vee \rightarrow W$ factors through the deflation $V \rightarrow W$. Thus X^\vee is \mathbb{E} -projective by [51, Proposition 3.24], and hence X is \mathbb{E} -injective.

If instead X is \mathbb{E} -injective, then X^\vee is \mathbb{E} -projective by the first argument above, and so X^\vee is \mathbb{E} -injective by the second, and so $(X^\vee)^\vee$ is \mathbb{E} -projective by the first, and $X \cong (X^\vee)^\vee$ since \otimes is symmetric and X is strongly dualisable. \square

Corollary 5.10. *Let $(\mathcal{A}, \mathbb{E}, \mathfrak{s}, \otimes, \mathbb{1})$ be a closed tensor extriangulated category in which each object is strongly dualisable. If $\mathbb{1}$ has an \mathbb{E} -projective cover or an \mathbb{E} -injective envelope, then \mathcal{A} is Frobenius.*

Proof. Thanks to Proposition 5.9 it is enough to check that \mathcal{A} has enough \mathbb{E} -projectives and \mathbb{E} -injectives. If $P \rightarrow \mathbb{1}$ is an \mathbb{E} -projective cover of $\mathbb{1}$ then $P \otimes X \rightarrow \mathbb{1} \otimes X \cong X$ is an \mathbb{E} -projective cover of X . Hence \mathcal{A} has enough \mathbb{E} -projectives. Further, the dual $\mathbb{1}^\vee \cong \mathbb{1} \rightarrow P^\vee$ of the \mathbb{E} -projective cover $P \rightarrow \mathbb{1}$ is an \mathbb{E} -injective cover. Hence by a similar argument as above \mathcal{A} has enough \mathbb{E} -injectives. \square

Remark 5.11. In a closed symmetric monoidal category the dual X^\vee of a strongly dualisable object X is a right and left dual of X . In this setting, assuming that all objects are strongly dualisable is equivalent to assuming the category is rigid. However, for an extriangulated tensor category that is rigid, it is not clear whether the functor given by the left or right duals is an extriangulated functor. In the case of abelian categories this can be resolved using the universal properties of kernels and cokernels; see [19, Proposition 4.2.9].

5.12. *Extriangulated substructure*

Let $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$ be an extriangulated category and \mathcal{I} a full subcategory of \mathcal{A} . Then there is a closed subfunctor of \mathbb{E} defined by

$$\mathbb{E}_{\mathcal{I}}(Y, X) := \{d \in \mathbb{E}(Y, X) \mid f^*(d) = 0 \text{ for all } W \in \mathcal{I} \text{ and } f \in \text{Hom}_{\mathcal{A}}(W, Y)\};$$

see [28, Proposition 3.16]. So, $(\mathcal{A}, \mathbb{E}_{\mathcal{I}}, \mathfrak{s}_{\mathcal{I}})$ is an extriangulated category by [28, Proposition 3.19] where $\mathfrak{s}_{\mathcal{I}}$ is the restriction of \mathfrak{s} .

Lemma 5.13. *Let $(\mathcal{A}, \mathbb{E}, \mathfrak{s}, \otimes, \mathbb{1})$ be a monoidal extriangulated category and let \mathcal{I} be a full subcategory of \mathcal{A} . Suppose that, for the bixtriangulated functor $(-\otimes -, \varphi, \psi)$,*

$$\varphi_{X', X}^Y(d) \in \mathbb{E}_{\mathcal{I}}(X' \otimes Y, X \otimes Y) \quad \text{and} \quad \psi_{Y', Y}^X(e) \in \mathbb{E}_{\mathcal{I}}(X \otimes Y', X \otimes Y)$$

for each $X, X', Y, Y' \in \mathcal{A}$, and each $d \in \mathbb{E}_{\mathcal{I}}(X', X)$ and $e \in \mathbb{E}_{\mathcal{I}}(Y', Y)$. Then $(\mathcal{A}, \mathbb{E}_{\mathcal{I}}, \mathfrak{s}_{\mathcal{I}}, \otimes, \mathbb{1})$ is a monoidal extriangulated category.

When $(\mathcal{A}, \mathbb{E}, \mathfrak{s}, \otimes, \mathbb{1})$ is symmetric or strong, then so is $(\mathcal{A}, \mathbb{E}_{\mathcal{I}}, \mathfrak{s}_{\mathcal{I}}, \otimes, \mathbb{1})$.

Proof. By [28, Proposition 3.16] we have that $\mathbb{E}_{\mathcal{I}}$ is a subfunctor of \mathbb{E} . By our assumption the restriction of φ^Y to $\mathbb{E}_{\mathcal{I}}(-, -)$ defines a subtransformation in the sense of [14, p. 261]. We call this restriction $\varphi_{\mathcal{I}}^Y$. We also obtain a restriction $\psi_{\mathcal{I}}^X$ of ψ^X . We need to check conditions (1–4) from Definition 4.3 hold. Condition (1) holds by assumption. Condition (2) holds by the discussion before the statement of the lemma. In what remains we check conditions (3) and (4).

We start with (3) and we check (a) from Definition 3.12 holds; that is that the functor $(-\otimes Y, \varphi_{\mathcal{I}}^Y)$ is extriangulated. Suppose $X \rightarrow X'' \rightarrow X' \xrightarrow{-d}$ is an extriangle in $(\mathcal{A}, \mathbb{E}_{\mathcal{I}}, \mathfrak{s}_{\mathcal{I}})$ with $d \in \mathbb{E}(X', X)$. Then it is an extriangle in $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$ and, as $(-\otimes Y, \varphi^Y)$ is an extriangulated functor,

$$X \otimes Y \rightarrow X'' \otimes Y \rightarrow X' \otimes Y \overset{\varphi_{\mathcal{I}}^Y(d)}{\dashrightarrow}$$

is an extriangle. As $\mathfrak{s}_{\mathcal{I}}$ is the restriction of \mathfrak{s} to $\mathbb{E}_{\mathcal{I}}$ the functor $(- \otimes Y, \varphi_{\mathcal{I}}^Y)$ is extriangulated. Using a similar argument shows that (b) from Definition 3.12 holds; that is $(X \otimes -, \psi_{\mathcal{I}}^X)$ is extriangulated.

To finish the proof we need to show that certain diagrams in $\text{Mod}(R)$ commute; namely the compatibility diagrams of $\varphi_{\mathcal{I}}$ and $\psi_{\mathcal{I}}$ from (c) and (d) of Definition 3.12, and their compatibility with α, ρ, λ of the monoidal structure. Each diagram is the restriction of the corresponding diagram for $(\mathcal{A}, \otimes, \mathbb{1})$. Hence they commute.

When $(\mathcal{A}, \otimes, \mathbb{1})$ is symmetric, then so is $(\mathcal{A}, \otimes, \mathbb{1})$ by the same argument as above. For strongness, note that the cup product is compatible with restriction. \square

5.14. Pure-exact extriangulated structure

In the next example we discuss a special case of extriangulated subfunctors for compactly generated triangulated categories. The extriangles arising are precisely the pure-exact triangles.

We recall some basic definitions concerning triangulated categories. Let \mathcal{T} be a triangulated category with suspension Σ . An object X is *compact* if the functor $\text{Hom}_{\mathcal{T}}(X, -): \mathcal{T} \rightarrow \mathcal{A}b$ preserves coproducts, and we write \mathcal{T}^c for the full subcategory of compact objects. The triangulated category \mathcal{T} is *compactly generated*, if it has small coproducts, \mathcal{T}^c is essentially small and for any non-zero object Y in \mathcal{T} there exists an object $X \in \mathcal{T}^c$ such that $\text{Hom}_{\mathcal{T}}(X, Y) \neq 0$.

Proposition 5.15. *Let $(\mathcal{T}, \otimes, \mathbb{1})$ be a compactly generated closed tensor triangulated category, considered as a tensor extriangulated category $(\mathcal{T}, \mathbb{E}, \mathfrak{s}, \otimes, \mathbb{1})$. In the notation of Section 5.12, we have that $(\mathcal{T}, \mathbb{E}_{\mathcal{T}^c}, \mathfrak{s}_{\mathcal{T}^c}, \otimes, \mathbb{1})$ is a tensor extriangulated category.*

We first recall the definition and some basic properties of phantom maps from [39] as they are closely connected to the extriangulated substructure $\mathbb{E}_{\mathcal{T}^c}$.

Let $\text{Mod}((\mathcal{T}^c)^{\text{op}})$ be the category of additive functors $(\mathcal{T}^c)^{\text{op}} \rightarrow \mathcal{A}b$. This is an abelian AB5 category. Let $\mathbf{Y}: \mathcal{T} \rightarrow \text{Mod}((\mathcal{T}^c)^{\text{op}})$ be the *restricted Yoneda functor*, defined on an object X by the restriction of $\text{Hom}_{\mathcal{T}}(-, X)$ to \mathcal{T}^c and likewise on morphisms. An additive functor $F: \mathcal{T} \rightarrow \text{Mod}((\mathcal{T}^c)^{\text{op}})$ is *cohomological* if any exact triangle

$$X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} \Sigma X \tag{5.15.1}$$

induces an exact sequence $F(X) \rightarrow F(Y) \rightarrow F(Z) \rightarrow F(\Sigma X)$ in $\mathcal{A}b$. In particular, the restricted Yoneda functor \mathbf{Y} is cohomological.

A morphism f in \mathcal{T} is called *phantom* if $\mathbf{Y}(f) = 0$. The composition of a phantom morphism with any morphism is phantom. A triangle (5.15.1) is said to be *pure-exact* if $0 \rightarrow \mathbf{Y}(X) \rightarrow \mathbf{Y}(Y) \rightarrow \mathbf{Y}(Z) \rightarrow 0$ is exact. So (5.15.1) is pure-exact if and only if h is phantom; see [39, Lemma 1.3].

Lemma 5.16. *Let $(\mathcal{T}, \otimes, \mathbb{1})$ be a closed tensor triangulated category. Let f and g be morphisms in \mathcal{T} . If f or g is phantom, then so is $f \otimes g$.*

Proof. As phantom morphisms are closed under composition with any other morphism, it is enough to show $f \otimes \text{id}$ is phantom when f is phantom and $\text{id} \otimes g$ is phantom when g is phantom. We give a proof for the former, the latter holds by symmetry.

The restricted Yoneda functor \mathbf{Y} preserves coproducts. Since $- \otimes X$ is a left adjoint for any object $X \in \mathcal{T}$, it also preserves coproducts. It follows that the composition $\mathbf{F} := \mathbf{Y} \circ (- \otimes X)$ is a cohomological functor $\mathcal{T} \rightarrow \text{Mod}((\mathcal{T}^c)^{\text{op}})$ that preserves coproducts.

Let f be a phantom morphism. By [39, Corollary 2.5] we have that $\mathbf{Y}(f \otimes \text{id}_X) = \mathbf{F}(f) = 0$ as \mathbf{F} is cohomological and preserves coproducts. \square

We can now give an explicit description of the extriangulated substructure $\mathbb{E}_{\mathcal{T}^c}$ from Proposition 5.15. If $d \in \mathbb{E}(Y, X) = \text{Hom}_{\mathcal{T}}(Y, \Sigma X)$, then $f^*(d) = d \circ f$ for every $f: W \rightarrow Y$ in \mathcal{T} . Hence $d \in \mathbb{E}_{\mathcal{T}^c}(Y, X)$ if and only if $d \circ f = 0$ for every map $f: C \rightarrow Y$ with $C \in \mathcal{T}^c$. That means $\mathbb{E}_{\mathcal{T}^c}(Y, X)$ is the subset of phantom maps in $\text{Hom}_{\mathcal{T}}(Y, \Sigma X)$, and $X \rightarrow Y \rightarrow Z \dashrightarrow$ is an extriangle if and only if $X \rightarrow Y \rightarrow Z \rightarrow \Sigma X$ is a pure-exact triangle.

Example 5.17. Let R be a non-artinian ring. The derived category $\text{D}(\text{Mod}(R))$ is compactly generated triangulated where compact objects are perfect complexes. We claim that the extriangulated structure on $\text{D}(\text{Mod}(R))$, given by [28, Proposition 3.19] taking \mathcal{I} to be the compact objects, is neither triangulated nor exact.

Using that R is not artinian, it cannot be pure-semisimple by the implication (i) \Rightarrow (a) in [54, Theorem 2.1]. By [25, Corollary 7.2], it follows that $\text{D}(\text{Mod}(R))$ is not pure-semisimple, and so there exists an object X that is not pure-injective. By [39, Definition 1.1], there exists a non-split pure-exact triangle $X \rightarrow Y \rightarrow Z \rightarrow \Sigma X$. Since categorical monics split in triangulated categories, $X \rightarrow Y$ cannot be monic. If the extriangulated structure on $\text{D}(\text{Mod}(R))$ were exact, the inflation $X \rightarrow Y$ would be a kernel, which is false.

It suffices to find a contradiction assuming the extriangulated structure is triangulated. Choose a morphism $X \rightarrow Y$ in $\text{D}(\text{Mod}(R))$ by concentrating an arbitrary homomorphism f in $\text{Mod}(R)$ in degree 0. By assumption we can complete $X \rightarrow Y$ to a pure-exact triangle $X \rightarrow Y \rightarrow Z \rightarrow \Sigma X$. By [25, Lemma 2.4] we have that $0 \rightarrow H_n(X) \rightarrow H_n(Y) \rightarrow H_n(Z) \rightarrow 0$ is pure-exact in $\text{Mod}(R)$ for each $n \in \mathbb{Z}$. Taking $n = 0$ implies that f is a pure-monomorphism. Since pure-monomorphisms in $\text{Mod}(R)$ are injective, a contradiction is found by choosing f not injective.

Finally we prove of Proposition 5.15 using the observations in Section 5.12.

Proof of Proposition 5.15. We check that the assumptions of Lemma 5.13 are satisfied. By Examples 3.4 and 3.15 we have

$$\varphi_{X',X}^Y(d) = (X' \otimes Y \xrightarrow{d \otimes \text{id}_Y} (\Sigma X) \otimes Y \xrightarrow{\cong} \Sigma(X \otimes Y))$$

for $d \in \mathbb{E}(X', X)$. If d is phantom, then so is $\varphi_{X',X}^Y(d)$ by Lemma 5.16. The analogous condition for ψ can be shown similarly. Hence $(\mathcal{T}, \mathbb{E}_{\mathcal{T}^c}, \mathfrak{s}_{\mathcal{T}^c}, \otimes, \mathbb{1})$ is an extriangulated tensor category by Lemma 5.13. \square

6. Classification of tensor ideals

The goal of this Section is the classification of the radical thick tensor ideals in an essentially small tensor extriangulated category. In [9, Sections 7 and 8] lattices of thick tensor ideals were considered for both abelian and triangulated categories simultaneously. The arguments for the classification of thick tensor ideals in a triangulated category of [1] and [38] generalise straight away to the extriangulated case. For convenience we give the details of some important parts.

6.1. Tensor ideals

Let $(\mathcal{A}, \otimes, \mathbb{1})$ be a symmetric monoidal category. A non-empty strictly full subcategory \mathcal{I} of \mathcal{A} is a *tensor ideal*, if for each $X \in \mathcal{A}$ and $Y \in \mathcal{I}$, we have $X \otimes Y \in \mathcal{I}$.

For an object $X \in \mathcal{A}$ and a positive integer n , we set $X^{\otimes n} := X \otimes \cdots \otimes X$, the n -fold tensor product of X . For convenience we also set $X^{\otimes 0} := \mathbb{1}$.

We say a tensor ideal \mathcal{I} is:

- (1) *radical* if, for all $X \in \mathcal{A}$, we have that $X^{\otimes n} \in \mathcal{I}$ for some $n \geq 1$ implies $X \in \mathcal{I}$; and
- (2) *prime* if, for all $X, Y \in \mathcal{A}$, we have that $X \otimes Y \in \mathcal{I}$ implies $X \in \mathcal{I}$ or $Y \in \mathcal{I}$.

Note that any prime tensor ideal is radical.

Let $(\mathcal{A}, \mathbb{E}, \mathfrak{s})$ be an extriangulated category. We say a non-empty full subcategory \mathcal{I} is *thick*, if it is closed under direct summands and has the *2-out-of-3 property*; that is, for any extriangle $X \rightarrow Y \rightarrow Z \dashrightarrow$, if any two of X, Y , or Z lie in \mathcal{I} , then so does the third.

Let \mathcal{C} be a collection of objects of \mathcal{A} . We denote by $\text{thick}^{\otimes}(\mathcal{C})$ the smallest thick tensor ideal containing \mathcal{C} , and by $\text{rad}(\mathcal{C})$ the smallest radical thick tensor ideal containing \mathcal{C} . These categories exist as the intersection of (radical) thick tensor ideals is again a (radical) thick tensor ideal. The radical thick closure can be described as

$$\text{rad}(\mathcal{C}) = \{ X \in \mathcal{A} \mid X^{\otimes n} \in \text{thick}^{\otimes}(\mathcal{C}) \text{ for some } n \geq 1 \}, \tag{6.1.1}$$

using the arguments from the proof of [38, Lemma 3.1.6] to see the right-hand side is thick. For a tensor extriangulated category $(\mathcal{A}, \otimes, \mathbb{1})$ we denote by $\text{Rad}(\mathcal{A})$ the class of all radical thick tensor ideals of \mathcal{A} . When \mathcal{A} is essentially small, then $\text{Rad}(\mathcal{A})$ is a set. In Proposition 6.5 we show that, if $\text{Rad}(\mathcal{A})$ is a set, it has a rich poset structure.

6.2. Lattices and frames

For completeness we recall the terminology and fix notation; we loosely follow [34]. Recall, in a poset L the *join* (respectively, *meet*) of a subset $S \subseteq L$ is the smallest (respectively, largest) element that contains (respectively, is contained in) every element of S . We denote the join of S by $\bigvee S$ and the meet of S by $\bigwedge S$. For elements $a, b \in L$ we write $a \vee b = \bigvee\{a, b\}$ and $a \wedge b = \bigwedge\{a, b\}$.

A *lattice* is a poset L in which every finite, non-empty subset has a join and a meet. A lattice L is *distributive* if L has a *greatest element* $1 = \bigwedge \emptyset$, a *least element* $0 = \bigvee \emptyset$ and it satisfies the *distributive law*; that is

$$a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c), \quad \text{or equivalently} \quad a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$$

for any $a, b, c \in L$. A *morphism of distributive lattices* is a map that respects the order, finite meets, finite joins, and the greatest and least element.

A lattice F is a *frame* if the join of any subset exist and it satisfies the *infinite join-distributivity law*; that is

$$a \wedge \bigvee S = \bigvee \{a \wedge s \mid s \in S\}$$

for any $a \in F$ and any subset $S \subseteq F$. A *morphism of frames* is a morphism of distributive lattices that preserves arbitrary joins.

6.3. Stone and Hochster duality

The prototypical example of a frame is the set of all open subsets of a topological space. In fact, there is a functor

$$\Omega: \mathcal{Top} \rightarrow \mathcal{Frm}^{\text{op}}$$

from the category of topological spaces with continuous maps to the category of frames. This functor has a right adjoint, given by the prime spectrum of a frame (F, \leq) : An element $a \in F$ is (*meet-*)*prime*, if $b \wedge c \leq a$ implies that either $a \leq b$ or $a \leq c$. The *prime spectrum of F* is the set of all prime elements in F endowed with a topology given by the closed sets $V(a) = \{p \mid a \leq p\}$. We denote the right adjoint by pt , due to the fact that prime elements in a frame correspond to points; for more details see [34, II.1.3].

The adjoint functors (Ω, pt) restrict to an equivalence between the category of sober spaces and the category of spatial frames. This is called *Stone duality*; for more details see [34, II.1.7]. This equivalence restricts further to an equivalence of the category of spectral spaces and the category of coherent frames. We recall the latter two notions.

An element a in a frame F is *compact* if for each subset S of F with $a \leq \bigvee S$ there is a finite subset $T \subseteq S$ such that $a \leq \bigvee T$. A frame F is *coherent* if the greatest element

1 is compact, $a \wedge b$ is compact for any compact elements $a, b \in F$, and for each $a \in F$ there exists a subset S of compact elements satisfying $a = \bigvee S$. A *morphism of coherent frames* is a morphism of frames that preserves compact elements.

In a coherent frame F the set of compact elements is a distributive sublattice of F , and the frame F can be recovered from the distributive lattice of compact elements. In fact, there is an equivalence $\mathcal{C}oh \rightarrow \mathcal{DLat}$ from the category of coherent frames to the category of distributive lattices; see [34, II.3.2].

On the other hand, a topological space X is *spectral*, if

- (1) X is quasi-compact;
- (2) the quasi-compact open subsets of X are closed under finite intersections and form a basis of open sets of the topology on X ; and
- (3) X is *sober*; that is every non-empty irreducible closed subset has a generic point;

see [30]. A *morphism of spectral spaces* is a continuous map such that the preimage of every quasi-compact open set is again quasi-compact. We denote the category of spectral spaces by $\mathcal{S}p$. The spectral spaces are precisely those topological spaces that occur as the Zariski spectrum of a commutative ring; see [30].

Putting everything together, we obtain a sequence of equivalences:

$$\mathcal{S}p \xrightleftharpoons[\text{pt}]{\Omega} \mathcal{C}oh^{\text{op}} \xrightarrow{(-)^c} \mathcal{DLat}^{\text{op}} \xleftarrow{(-)^{\text{op}}} \mathcal{DLat}^{\text{op}} \xleftarrow{(-)^c} \mathcal{C}oh^{\text{op}} \xrightleftharpoons[\text{pt}]{\Omega} \mathcal{S}p.$$

Here, the functor $(-)^{\text{op}}$ sends a lattice to its opposite lattice, which has the same underlying set and the reversed order. Note, that this is not a contravariant functor. The composition of these equivalences yields a functor $\mathcal{S}p \rightarrow \mathcal{S}p$, called *Hochster duality*. This equivalence is usually denoted by $(-)^{\vee}$, and $(X^{\vee})^{\vee} = X$ for a spectral space X . An open subset in X^{\vee} is precisely a *Thomason subset* in X , that is, a union $\bigcup X_i$ where each X_i is closed and has a quasi-compact complement in X .

6.4. Lattice of tensor ideals

Let $(\mathcal{A}, \otimes, \mathbb{1})$ be a tensor extriangulated category. The meet of two radical thick tensor ideals is given as their intersection. The join of a set of radical thick tensor ideals S is

$$\bigvee S = \text{rad}\left(\bigcup_{\mathcal{I} \in S} \mathcal{I}\right).$$

The least element is the nilradical ideal $\text{rad}(\{0\})$ and the greatest element is the whole category \mathcal{A} . Using the same arguments as for the triangulated case in [38, Theorem 3.1.9], we show the following:

Proposition 6.5. *Suppose $(\mathcal{A}, \otimes, \mathbb{1})$ is a tensor extriangulated category. If $\text{Rad}(\mathcal{A})$ is a set, then it is a coherent frame.*

Proof. We first show the infinite join-distributive law. Let $\mathcal{I} \in \text{Rad}(\mathcal{A})$ and $S \subseteq \text{Rad}(\mathcal{A})$. We show $\mathcal{I} \wedge \bigvee S \leq \bigvee_{\mathcal{J} \in S} (\mathcal{I} \wedge \mathcal{J})$, as the opposite inclusion always hold. For $X \in \mathcal{I} \wedge \bigvee S$ we define

$$\mathcal{J}_X := \{ Y \in \bigvee S \mid X \otimes Y \in \bigvee_{\mathcal{J} \in S} (\mathcal{I} \wedge \mathcal{J}) \}.$$

It is straightforward to check that \mathcal{J}_X is a radical thick tensor ideal. As $\mathcal{J}_X \geq \mathcal{J}$ for all $\mathcal{J} \in S$ we get

$$\mathcal{J}_X \geq \bigvee S \ni X.$$

Hence $X \otimes X \in \bigvee_{\mathcal{J} \in S} (\mathcal{I} \wedge \mathcal{J})$, and as $\bigvee_{\mathcal{J} \in S} (\mathcal{I} \wedge \mathcal{J})$ is radical, this means $X \in \bigvee_{\mathcal{J} \in S} (\mathcal{I} \wedge \mathcal{J})$. This shows $\text{Rad}(\mathcal{A})$ is a frame. Note that the compact objects in $\text{Rad}(\mathcal{A})$ are all of the form $\text{rad}(X)$ for an object $X \in \mathcal{A}$.

Let $X \in \mathcal{A}$ and assume $\text{rad}(X) \subseteq \bigvee S$ for some $S \subseteq \text{Rad}(\mathcal{A})$. Then $X^{\otimes n} \in \text{thick}^{\otimes}(\bigcup_{\mathcal{J} \in S} \mathcal{J})$. Taking the thick tensor closure is a finite process, hence there exists a finite subset $T \subseteq S$ such that $X^{\otimes n} \in \text{thick}^{\otimes}(\bigcup_{\mathcal{J} \in T} \mathcal{J})$. Hence $X \in \bigvee T$.

For coherence it remains to show that the meet of $\text{rad}(X)$ and $\text{rad}(Y)$ for any $X, Y \in \mathcal{A}$ is $\text{rad}(X \otimes Y)$. Since $\text{rad}(X)$ is a radical thick tensor ideal containing X it also contains $X \otimes Y$, and so $\text{rad}(X) \geq \text{rad}(X \otimes Y)$. Similarly $\text{rad}(Y) \geq \text{rad}(X \otimes Y)$. Thus $\text{rad}(X) \wedge \text{rad}(Y) \geq \text{rad}(X \otimes Y)$, and we now require the reverse inclusion. Let $U \in \text{rad}(X) \wedge \text{rad}(Y)$. Then $U^{\otimes 2} \in \text{rad}(X \otimes Y)$ and $U \in \text{rad}(X \otimes Y)$ by (6.1.1). This finishes the proof. \square

6.6. Classification

For the classification it remains to apply Stone and Hochster duality to Proposition 6.5. For tensor triangulated categories this result is due to Balmer [1]. The lattice-theoretic proof relies on work by [38] and [9]. However, first we describe the prime elements of $\text{Rad}(\mathcal{A})$.

Lemma 6.7. *Let $(\mathcal{A}, \otimes, \mathbb{1})$ be a tensor extriangulated category such that $\text{Rad}(\mathcal{A})$ is a set. A radical thick tensor ideal \mathcal{P} is prime in $\text{Rad}(\mathcal{A})$ if and only if \mathcal{P} is a prime thick tensor ideal.*

Proof. We assume \mathcal{P} is prime as an element in the lattice $\text{Rad}(\mathcal{A})$. Suppose $X \otimes Y \in \mathcal{P}$ for some objects $X, Y \in \mathcal{A}$. Then

$$\mathcal{P} \geq \text{rad}(X \otimes Y) = \text{rad}(X) \wedge \text{rad}(Y)$$

using an intermediate claim proven in Proposition 6.5. By assumption, we have that $\text{rad}(X) \leq \mathcal{P}$ or $\text{rad}(Y) \leq \mathcal{P}$. Hence $X \in \mathcal{P}$ or $Y \in \mathcal{P}$.

Conversely, we assume that \mathcal{P} is a prime thick tensor ideal. Let $\mathcal{I}, \mathcal{J} \in \text{Rad}(\mathcal{A})$ such that $\mathcal{I} \cap \mathcal{J} \leq \mathcal{P}$ and $\mathcal{I} \not\leq \mathcal{P}$. We need to show $\mathcal{J} \leq \mathcal{P}$. Let $X \in \mathcal{I}$ with $X \notin \mathcal{P}$. Then

$X \otimes Y \in \mathcal{P}$ for every $Y \in \mathcal{J}$. As \mathcal{P} is a prime thick tensor ideal, this yields $Y \in \mathcal{P}$ for every $Y \in \mathcal{J}$. Hence $\mathcal{J} \leq \mathcal{P}$. \square

Theorem 6.8. *Let $(\mathcal{A}, \otimes, \mathbb{1})$ be a tensor extriangulated category such that $\text{Rad}(\mathcal{A})$ is a set. We let $\text{BSpc}(\mathcal{A})$ be the set of all prime thick tensor ideals in \mathcal{A} and endow it with a topology generated by the closed sets*

$$\text{supp}(X) := \{ \mathcal{P} \in \text{BSpc}(\mathcal{A}) \mid X \notin \mathcal{P} \}$$

for any $X \in \mathcal{A}$. Then there is a bijection

$$\begin{aligned} \text{Rad}(\mathcal{A}) &\leftrightarrow \{ \text{Thomason subsets of } \text{BSpc}(\mathcal{A}) \}, \\ \mathcal{I} \mapsto \text{supp}(\mathcal{I}) &= \bigcup_{X \in \mathcal{I}} \text{supp}(X) \quad \text{and} \quad \{ X \in \mathcal{A} \mid \text{supp}(X) \subseteq V \} \leftarrow V. \end{aligned}$$

Proof. As the prime thick tensor ideals are precisely the prime elements of $\text{Rad}(\mathcal{A})$, we have

$$\text{BSpc}(\mathcal{A}) = (\text{pt}(\text{Rad}(\mathcal{A})))^\vee$$

as topological spaces. The classification follows from the description of the functors pt and $(-)^{\vee}$. \square

The topological space $\text{BSpc}(\mathcal{A})$ is called the *Balmer spectrum* of \mathcal{A} .

6.9. Functoriality

Let $(F, \beta): (\mathcal{A}, \otimes_{\mathcal{A}}, \mathbb{1}_{\mathcal{A}}) \rightarrow (\mathcal{B}, \otimes_{\mathcal{B}}, \mathbb{1}_{\mathcal{B}})$ be a tensor extriangulated functor. Then

$$\begin{aligned} \text{rad}(F(\bigvee S)) &= \text{rad}\left(\bigcup_{X \in S} F(X)\right) = \bigvee_{X \in S} \text{rad}(F(X)) \quad \text{and} \\ \text{rad}(F(\text{rad}(X) \wedge \text{rad}(Y))) &= \text{rad}(F(X \otimes Y)) = \text{rad}(F(X)) \wedge \text{rad}(F(Y)) \end{aligned}$$

for any set of objects S in \mathcal{A} and objects X and Y in \mathcal{A} . Hence

$$\text{rad}(F(-)): \text{Rad}(\mathcal{A}) \rightarrow \text{Rad}(\mathcal{B})$$

is a morphism of coherent frames. Applying Stone and Hochster duality we obtain a continuous map

$$\text{BSpc}(F): \text{BSpc}(\mathcal{B}) \rightarrow \text{BSpc}(\mathcal{A}) \quad \text{given by} \quad \mathcal{Q} \mapsto F^{-1}(\mathcal{Q}).$$

As for triangulated categories in [1, Corollary 3.8] and [4, Proposition 8.7], essentially surjective functors yield embeddings on the Balmer spectrum.

Lemma 6.10. *Let $(F, \beta): (\mathcal{A}, \otimes_{\mathcal{A}}, \mathbb{1}_{\mathcal{A}}) \rightarrow (\mathcal{B}, \otimes_{\mathcal{B}}, \mathbb{1}_{\mathcal{B}})$ be an essentially surjective tensor extriangulated functor. Then $\text{BSpc}(F): \text{BSpc}(\mathcal{B}) \rightarrow \text{BSpc}(\mathcal{A})$ is an embedding.*

Proof. The injectivity is clear from the definition of the induced map.

For the second claim, we need to show that $\text{BSpc}(F)$ is a closed map onto its image. By [12, Theorem 5.3.3], it is enough to show that $\text{BSpc}(F)(\text{supp}(Y))$ is closed in the image $\text{im}(\text{BSpc}(F))$ for any $Y \in \mathcal{B}$. Let $Y \in \mathcal{B}$. As F is essentially surjective, there exists $X \in \mathcal{A}$ with $F(X) \cong Y$. Then $Y \notin \mathcal{Q}$ if and only if $X \notin F^{-1}(\mathcal{Q})$ for any $\mathcal{Q} \in \text{BSpc}(\mathcal{B})$. Hence we have

$$\text{BSpc}(F)(\text{supp}(Y)) = \text{supp}(X) \cap \text{im}(\text{BSpc}(F)).$$

Hence the map $\text{BSpc}(F)$ is closed and thus an embedding. \square

The Balmer spectrum has been computed for many tensor triangulated categories, like the perfect complexes of a commutative ring [31,52,57] and the stable module category of a group algebra [7]. In the following we describe the Balmer spectrum of some of the examples from Section 5.

6.11. Projective modules

Let R be a commutative ring with local units. In Example 5.2 we saw that the category of flat modules $\text{Flat}(R)$ is a tensor extriangulated category. This category need not be essentially small. However, the subcategory of finitely presented flat modules is essentially small and so we may consider its Balmer spectrum. In fact, a module is finitely presented and flat if and only if it is finitely presented and projective. In the following we compute the Balmer spectrum of $\text{proj}(R)$ for some classes of commutative rings R .

Note that the exact structure on $\text{proj}(R)$ is the split exact structure, so the radical thick tensor ideals are precisely the radical tensor ideals that are closed under finite coproducts and direct summands.

Lemma 6.12. *Let R be a commutative ring that satisfies one of the following conditions:*

- (1) $R = S[x_1, \dots, x_n]$, the polynomial ring over a principal ideal domain S ;
- (2) R is a local ring; or
- (3) R is a Dedekind domain.

Then $\text{BSpc}(\text{proj}(R)) = \{\text{rad}(0)\}$; that is the Balmer spectrum consists of one point.

Proof. In the cases (1) and (2) every finitely generated projective R -module is free; for (1) this follows from the Quillen–Suslin theorem and for (2) from Kaplansky’s theorem; see for example [42, Chapter XXI, Theorem 3.7] and [35, Theorem 2], respectively. Hence

there are exactly two radical thick tensor ideals: $\text{rad}(0)$ and $\text{proj}(R)$, and $\text{rad}(0)$ is the only prime thick tensor ideal.

In the case (3) the finitely projective R -modules are precisely the torsion-free modules. In particular, any ideal is projective. Let I and J be non-zero ideals of R . By [16, Exercise 19.5], we obtain $J \oplus I^2 \cong JI \oplus I$. Since I is projective, we have $JI \cong J \otimes I$. This means $J \in \text{thick}(I)$. Furthermore, any finitely generated projective R -module is of the form $R^n \oplus I$ for some ideal I ; see [16, Exercise 19.6]. So $\text{proj}(R)$ has two radical thick tensor ideals, namely $\text{rad}(0)$ and $\text{proj}(R)$, and $\text{rad}(0)$ is the only prime thick tensor ideal. \square

Lemma 6.13. *Let $R = R_1 \times \dots \times R_n$ be a product of commutative rings. Then*

$$\text{BSpc}(\text{proj}(R)) = \text{BSpc}(\text{proj}(R_1)) \times \dots \times \text{BSpc}(\text{proj}(R_n))$$

with the product topology.

Proof. By induction it is enough to show the claim for $n = 2$. Let $R = S \times T$. As an R -module $R \cong S \oplus T$ and every R -module decomposes as $M \cong (S \otimes_R M) \oplus (T \otimes_R M)$. In particular, M is a finitely presented projective R -module if and only if $S \otimes_R M$ and $T \otimes_R M$ are finitely presented projective over S and T , respectively. The map

$$\begin{aligned} \text{Rad}(\text{proj}(R)) &\rightarrow \text{Rad}(\text{proj}(S)) \times \text{Rad}(\text{proj}(T)), \\ \mathcal{I} &\mapsto \{S \otimes_R M \mid M \in \mathcal{I}\} \times \{T \otimes_R M \mid M \in \mathcal{I}\} \end{aligned}$$

has an inverse given by

$$\text{rad}(\mathcal{I} \cup \mathcal{J}) \leftrightarrow \mathcal{I} \times \mathcal{J}$$

where we view the objects in \mathcal{I} and \mathcal{J} as R -modules using restriction. This bijection is an isomorphism of coherent frames and hence it yields the desired isomorphism of the Balmer spectra. \square

Note that a commutative noetherian ring is hereditary if and only if it is regular of Krull dimension 1; see for example [18, Corollary 3]. For such rings Corollary 6.14 describes the Balmer spectrum.

Corollary 6.14. *If R is a commutative hereditary noetherian ring, then the Balmer spectrum $\text{BSpc}(\text{proj}(R))$ is the finite set of connected components of the Zariski spectrum $\text{ZSpec}(R)$ equipped with the discrete topology, and $|\text{BSpc}(\text{proj}(R))|$ is equal to the number of indecomposable factors of R .*

Proof. Note that R cannot contain an infinite set of primitive idempotents since it is noetherian; see for example [41, Proposition 22.2]. By [8, Theorem 4.4], see also [11,

p. 44, Corollary 1], the ring R is a finite product of Dedekind domains and fields. Hence the claim follows from Lemmas 6.12 and 6.13. \square

6.15. Stabilisation of extriangulated categories

We discussed the stabilisation of a tensor extriangulated category in Section 5.4. In the stabilisation $\underline{\mathcal{A}}_{\mathcal{I}}$ the objects in \mathcal{I} are isomorphic to 0. Hence there are fewer radical thick tensor ideals in $\underline{\mathcal{A}}_{\mathcal{I}}$ than in \mathcal{A} . In fact, the identity induces a tensor extriangulated functor $\mathcal{A} \rightarrow \underline{\mathcal{A}}_{\mathcal{I}}$, and as a direct consequence of Lemma 6.10 we obtain:

Lemma 6.16. *Let $(\mathcal{A}, \otimes, \mathbb{1})$ be a tensor extriangulated category and \mathcal{I} a tensor ideal consisting of objects that are \mathbb{E} -projective and \mathbb{E} -injective. Then*

$$\text{BSpc}(\underline{\mathcal{A}}_{\mathcal{I}}) = \{ \mathcal{P} \in \text{BSpc}(\mathcal{A}) \mid \mathcal{P} \supseteq \mathcal{I} \}$$

with the subspace topology. \square

Example 6.17. Let H be a cocommutative Hopf algebra over a field k . We assume that H is finite-dimensional over k . Then the category finite-dimensional H -modules is essentially small, and by Example 5.7 it becomes a tensor extriangulated category with (\otimes_k, k) . Further, $\text{mod}(H)$ is a Frobenius exact category, and there exists exactly one non-trivial thick tensor ideal consisting of objects that are projective and injective. We denote by $\underline{\text{mod}}(H)$ the stabilisation of $\text{mod}(H)$ with respect to this tensor ideal. Hence under the induced morphism of coherent frames

$$\text{Rad}(\text{mod}(H)) \rightarrow \text{Rad}(\underline{\text{mod}}(H))$$

every ideal has precisely one pre-image except the zero ideal in $\underline{\text{mod}}(H)$, which has exactly two pre-images: the zero ideal and the ideal of all projective objects.

Let G be a finite group. By [7],

$$\text{BSpc}(\underline{\text{mod}}(kG)) = \text{Proj}(\text{H}^*(G, k)),$$

where $\text{H}^*(G, k)$ is the group cohomology of G and $\text{Proj}(\text{H}^*(G, k))$ the set of homogenous primes without the irrelevant ideal with the Zariski topology. Then

$$\text{BSpc}(\text{mod}(kG)) = \text{Spec}(\text{H}^*(G, k))$$

where the latter is the set of all homogenous prime ideals; also see [40, Example 17].

Example 6.18. Let $(\mathcal{A}, \otimes, \mathbb{1})$ be an additive monoidal category and $\text{Ch}(\mathcal{A})$ the category of chain complexes. The degree-wise split exact sequences endow $\text{Ch}(\mathcal{A})$ with an exact structure. It is straightforward to check that $\text{Ch}(\mathcal{A})$ is a tensor extriangulated category

with the tensor product induced by \otimes and the unit the complex with $\mathbb{1}$ concentrated in degree zero. Moreover, the exact structure on $\text{Ch}(\mathcal{A})$ is Frobenius and the projective objects are precisely the complexes of the form $\text{cone}(\text{id}_X)$ for any complex X ; these are precisely the contractible complexes.

We endow \mathcal{A} with the split exact structure. Then there is a bijection

$$\begin{aligned} \text{Rad}(\mathcal{A}) &\rightarrow \{ \mathcal{J} \in \text{Rad}(\text{Ch}(\mathcal{A})) \mid \mathcal{J} \subseteq \text{Proj}(\text{Ch}(\mathcal{A})) \} \\ &\text{given by } \mathcal{I} \mapsto \text{rad}(\{ \text{cone}(\text{id}_M) \mid M \in \mathcal{I} \}). \end{aligned}$$

Hence there may exist intermediate extriangulated stabilisations between the category of chain complexes $\text{Ch}(\mathcal{A})$ and its homotopy category

$$\text{K}(\mathcal{A}) = \text{Ch}(\mathcal{A}) / \text{Proj}(\text{Ch}(\mathcal{A}));$$

compare with Lemma 6.13 and Corollary 6.14.

Acknowledgment

R. B-T. is grateful to Isaac Bird for helpful discussions concerning Section 5.14. R. B-T. and A. S. were partly funded by the Danish National Research Foundation (DNRF156); the Independent Research Fund Denmark (1026-00050B); the Aarhus University Research Foundation (AUFF-F-2020-7-16). R. B-T. and J. C. L. were partly funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation)—Project-ID 491392403—TRR 358. J. C. L. was partly funded by the Alexander von Humboldt Foundation in the framework of a Feodor Lynen research fellowship endowed by the German Federal Ministry of Education and Research. I. G. was funded by a PhD scholarship from the Carnegie Institute for the Universities of Scotland.

Data availability

No data was used for the research described in the article.

References

- [1] Paul Balmer, The spectrum of prime ideals in tensor triangulated categories, *J. Reine Angew. Math.* 588 (2005) 149–168.
- [2] Paul Balmer, Spectra, spectra, spectra—tensor triangular spectra versus Zariski spectra of endomorphism rings, *Algebr. Geom. Topol.* 10 (3) (2010) 1521–1563.
- [3] Paul Balmer, Tensor triangular geometry, in: Rajendra Bhatia, Arup Pal, Govindan Rangarajan, Vasudevan Srinivas, Muthusamy Vanninathan (Eds.), *Proceedings of the Intl. Congr. of Mathematicians. Vol. II*, Hindustan Book Agency, New Delhi, 2010, pp. 85–112.
- [4] Paul Balmer, Beren Sanders, The Tate intermediate value theorem, arXiv e-prints, arXiv:2503.11719v1, 2025.
- [5] Raphael Bennett-Tennenhaus, Johanne Haugland, Mads Hustad Sandøy, Amit Shah, The category of extensions and a characterisation of n -exangulated functors, *Math. Z.* 305 (3) (2023) 44.

- [6] Raphael Bennett-Tennenhaus, Amit Shah, Transport of structure in higher homological algebra, *J. Algebra* 574 (2021) 514–549.
- [7] David J. Benson, Jon F. Carlson, Jeremy Rickard, Thick subcategories of the stable module category, *Fundam. Math.* 153 (1) (1997) 59–80.
- [8] George M. Bergman, Hereditary commutative rings and centres of hereditary rings, *Proc. Lond. Math. Soc.* 3 (23) (1971) 214–236.
- [9] Aslak Bakke Buan, Henning Krause, Øyvind Solberg, Support varieties: an ideal approach, *Homol. Homotopy Appl.* 9 (2005) 45–74.
- [10] Theo Bühler, Exact categories, *Expo. Math.* 28 (1) (2010) 1–69.
- [11] A.W. Chatters, Søren Jøndrup, Hereditary finitely generated P.I. algebras, *J. Algebra* 82 (1) (1983) 40–52.
- [12] Max Dickmann, Niels Schwartz, Marcus Tressl, *Spectral Spaces*, New Mathematical Monographs, vol. 35, Cambridge University Press, Cambridge, 2019.
- [13] Samuel Eilenberg, G. Maxwell Kelly, A generalization of the functorial calculus, *J. Algebra* 3 (1966) 366–375.
- [14] Samuel Eilenberg, Saunders MacLane, General theory of natural equivalences, *Trans. Am. Math. Soc.* 58 (1945) 231–294.
- [15] David Eisenbud, Homological algebra on a complete intersection, with an application to group representations, *Trans. Am. Math. Soc.* 260 (1) (1980) 35–64.
- [16] David Eisenbud, *Commutative Algebra with a View Toward Algebraic Geometry*, Grad. Texts in Math., vol. 150, Springer-Verlag, New York, 1995.
- [17] Laiachi El Kaoutit, BOCSES over small linear categories and corings, in: Mercedes Siles Molina, Laiachi El Kaoutit, Mohamed Louzari, L’Moufadal Ben Yakoub, Mohamed Benslimane (Eds.), *Associative and Non-associative Algebras and Applications*, Proc. of the 3rd Moroccan Andalusian Meeting on Algebras and Their Applications, in: Springer Proc. Math. Stat., vol. 311, Springer, Cham, 2020, pp. 273–286.
- [18] Shizuo Endo, On semi-hereditary rings, *J. Math. Soc. Jpn.* 13 (1961) 109–119.
- [19] Pavel Etingof, Shlomo Gelaki, Dmitri Nikshych, Victor Ostrik, *Tensor Categories*, Math. Surveys Monogr., vol. 205, American Mathematical Society, Providence, RI, 2015.
- [20] Pavel Etingof, Viktor Ostrik, Finite tensor categories, *Mosc. Math. J.* 4 (3) (2004) 627–654, 782–783.
- [21] Janet L. Fisher, The tensor product of functors; satellites; and derived functors, *J. Algebra* 8 (1968) 277–294.
- [22] Yves Baudelaire Fomatati, On tensor products of matrix factorizations, *J. Algebra* 609 (2022) 180–216.
- [23] Pierre Gabriel, Des catégories abéliennes, *Bull. Soc. Math. Fr.* 90 (1962) 323–448.
- [24] José L. García, Juan Martínez Hernández, Purity through Gabriel’s functor rings, *Bull. Soc. Math. Belg., Sér. A* 45 (1–2) (1993) 137–152.
- [25] Grigory Garkusha, Mike Prest, Triangulated categories and the Ziegler spectrum, *Algebr. Represent. Theory* 8 (2005) 499–523.
- [26] Mikhail Gorsky, Hiroyuki Nakaoka, Yann Palu, Positive and negative extensions in extriangulated categories, arXiv e-prints, arXiv:2103.12482v1, 2021.
- [27] Dieter Happel, *Triangulated Categories in the Representation Theory of Finite-Dimensional Algebras*, London Math. Soc. Lecture Note Ser., vol. 119, Cambridge University Press, Cambridge, 1988.
- [28] Martin Herschend, Yu Liu, Hiroyuki Nakaoka, n -exangulated categories (I): definitions and fundamental properties, *J. Algebra* 570 (2021) 531–586.
- [29] Martin Herschend, Yu Liu, Hiroyuki Nakaoka, n -exangulated categories (II): constructions from n -cluster tilting subcategories, *J. Algebra* 594 (2022) 636–684.
- [30] Melvin Hochster, Prime ideal structure in commutative rings, *Trans. Am. Math. Soc.* 142 (1969) 43–60.
- [31] Michael J. Hopkins, Global methods in homotopy theory, in: *Homotopy Theory*, Durham, 1985, in: London Math. Soc. Lecture Note Ser., vol. 117, Cambridge Univ. Press, Cambridge, 1987, pp. 73–96.
- [32] Mark Hovey, John H. Palmieri, Neil P. Strickland, *Axiomatic Stable Homotopy Theory*, Mem., vol. 128, Amer. Math. Soc. Amer.Math. Soc., Providence, RI, 1997.
- [33] Osamu Iyama, Hiroyuki Nakaoka, Yann Palu, Auslander-Reiten theory in extriangulated categories, *Trans. Am. Math. Soc. Ser. B* 11 (2024) 248–305.
- [34] Peter T. Johnstone, *Stone Spaces*, Cambridge Studies in Advanced Mathematics, vol. 3, Cambridge University Press, Cambridge, 1982.
- [35] Irving Kaplansky, Projective modules, *Ann. Math.* (2) 68 (1958) 372–377.

- [36] Masaki Kashiwara, Pierre Schapira, *Categories and Sheaves*, Grundlehren Math. Wiss., vol. 332, Springer-Verlag, Berlin, 2006.
- [37] G. Maxwell Kelly, On MacLane's conditions for coherence of natural associativities, commutativities, etc., *J. Algebra* 1 (1964) 397–402.
- [38] Joachim Kock, Wolfgang Pitsch, Hochster duality in derived categories and point-free reconstruction of schemes, *Trans. Am. Math. Soc.* 369 (1) (2017) 223–261.
- [39] Henning Krause, Smashing subcategories and the telescope conjecture—an algebraic approach, *Invent. Math.* 139 (1) (2000) 99–133.
- [40] Henning Krause, An analogue of Stone duality via support, *J. Pure Appl. Algebra* 228 (6) (2024) 107621.
- [41] Tsit Yuen Lam, *A First Course in Noncommutative Rings*, second edition, Grad. Texts in Math., vol. 131, Springer-Verlag, New York, 2001.
- [42] Serge Lang, *Algebra*, third edition, Grad. Texts in Math., vol. 211, Springer-Verlag, New York, 2002.
- [43] Richard Gustavus Larson, Moss Eisenberg Sweedler, An associative orthogonal bilinear form for Hopf algebras, *Am. J. Math.* 91 (1969) 75–94.
- [44] L. Gaunce Lewis Jr., J. Peter May, Mark Steinberger, James E. McClure, *Equivariant Stable Homotopy Theory*, Lecture Notes in Math., vol. 1213, Springer-Verlag, Berlin, 1986, With contributions by J.E. McClure.
- [45] Yu Liu, Hiroyuki Nakaoka, Hearts of twin cotorsion pairs on extriangulated categories, *J. Algebra* 528 (2019) 96–149.
- [46] Fosco Loregian, *(Co)end Calculus*, London Math. Soc. Lecture Note Ser., vol. 468, Cambridge University Press, Cambridge, 2021.
- [47] Saunders Mac Lane *Homology*, Classics in Mathematics, Springer-Verlag, Berlin, 1995, Reprint of the 1975 edition.
- [48] Saunders Mac Lane, *Categories for the Working Mathematician*, second edition, Grad. Texts in Math., vol. 5, Springer-Verlag, New York, 1998.
- [49] Harvey R. Margolis, *Spectra and the Steenrod Algebra*, North-Holland Math. Library, vol. 29, North-Holland Publishing Co., Amsterdam, 1983.
- [50] Hiroyuki Nakaoka, Yasuaki Ogawa, Arashi Sakai, Localization of extriangulated categories, *J. Algebra* 611 (2022) 341–398.
- [51] Hiroyuki Nakaoka, Yann Palu, Extriangulated categories, Hovey twin cotorsion pairs and model structures, *Cah. Topol. Géom. Différ. Catég.* 60 (2) (2019) 117–193.
- [52] Amnon Neeman, The chromatic tower for $D(R)$, *Topology* 31 (3) (1992) 519–532, With an appendix by Marcel Bökstedt.
- [53] Ulrich Oberst, Helmut Röhrl, Flat and coherent functors, *J. Algebra* 14 (1970) 91–105.
- [54] Mike Prest, Rings of finite representation type and modules of finite Morley rank, *J. Algebra* 88 (2) (1984) 502–533.
- [55] Daniel Quillen, Higher algebraic K -theory. I, in: Hyman Bass (Ed.), *Algebraic K -Theory, I: Higher K -Theories*, Proc. of the Conf., Seattle, 1972, in: *Lecture Notes in Math.*, vol. 341, Springer, Berlin, 1973, pp. 85–147.
- [56] Daniel Simson, Categories of representations of species, *J. Pure Appl. Algebra* 14 (1) (1979) 101–114.
- [57] Robert W. Thomason, The classification of triangulated subcategories, *Compos. Math.* 105 (1) (1997) 1–27.
- [58] Karl-Heinz Ulbrich, On Hopf algebras and rigid monoidal categories, in: *Hopf Algebras*, *Isr. J. Math.* 72 (1–2) (1990) 252–256.
- [59] Jean-Louis Verdier, Des catégories dérivées des catégories abéliennes. Number 239 in *Astérisque*, Soc. Math. France, Paris, 1996, With a preface by Luc Illusie, Edited and with a note by Georges Maltsiniotis.
- [60] Charles A. Weibel, *The K -Book, an Introduction to Algebraic K -Theory*, Graduate Studies in Mathematics, vol. 145, American Mathematical Society, Providence, RI, 2013.
- [61] Nobuo Yoneda, On the homology theory of modules, *J. Fac. Sci. Univ. Tokyo Sect. I* 7 (1954) 193–227.
- [62] Nobuo Yoneda, On Ext and exact sequences, *J. Fac. Sci. Univ. Tokyo Sect. I* 8 (1960) 507–576.
- [63] Yuji Yoshino, *Cohen–Macaulay Modules over Cohen–Macaulay Rings*, London Math. Soc. Lecture Note Ser., vol. 146, Cambridge Univ. Press, Cambridge, 1990.
- [64] Yuji Yoshino, Tensor products of matrix factorizations, *Nagoya Math. J.* 152 (1998) 39–56.