



## City Research Online

### City, University of London Institutional Repository

---

**Citation:** Szypula, J., Jarvstad, A., Jones, L. & Tapper, K. (2025). Personalizing a Weight Loss Program With Cognitive-Behavioral Phenotypes to Increase Engagement and Weight Loss in Adults With Overweight or Obesity: Quasi-Experimental Study (Preprint). JMIR Formative Research, doi: 10.2196/72645

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

---

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/36019/>

**Link to published version:** <https://doi.org/10.2196/72645>

**Copyright:** City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

**Reuse:** Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

---

City Research Online:

<http://openaccess.city.ac.uk/>

[publications@city.ac.uk](mailto:publications@city.ac.uk)

---

# **Personalizing a Weight Loss Program With Cognitive-Behavioral Phenotypes to Increase Engagement and Weight Loss in Adults With Overweight or Obesity: Quasi-Experimental Study**

Joanna Szygula, Andreas Jarvstad, Lucy Jones, Katy Tapper

Submitted to: JMIR Formative Research  
on: February 14, 2025

**Disclaimer:** © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

## *Table of Contents*

---

<b>Original Manuscript</b> .....	<b>5</b>
<b>Supplementary Files</b> .....	<b>26</b>
Figures .....	27
Figure 1.....	28
Figure 2.....	29
Figure 3.....	30
Multimedia Appendixes .....	31
Multimedia Appendix 0.....	32
Multimedia Appendix 1.....	32

Preprint  
JMIR Publications

# Personalizing a Weight Loss Program With Cognitive-Behavioral Phenotypes to Increase Engagement and Weight Loss in Adults With Overweight or Obesity: Quasi-Experimental Study

Joanna Szypula<sup>1</sup> PhD; Andreas Jarvstad<sup>1</sup> PhD; Lucy Jones<sup>2</sup> RD; Katy Tapper<sup>3</sup> Prof Dr

<sup>1</sup> School of Health and Psychological Science City St George's, University of London London GB

<sup>2</sup> Oviva London GB

<sup>3</sup> City St George's, University of London London GB

## Corresponding Author:

Joanna Szypula PhD

School of Health and Psychological Science  
City St George's, University of London  
Rhind Building, St John Street  
London  
GB

## Abstract

**Background:** Obesity remains a critical public health challenge, requiring innovative strategies to improve weight loss. Although preliminary evidence indicates that tailoring interventions to both cognitive and behavioral factors may enhance engagement and effectiveness, most research has focused on these elements in isolation. With the increasing scalability and convenience of digital weight loss programs, developing reliable methods for automated personalization has become essential. To address this, we constructed a questionnaire that assigns patients to cognitive-behavioral phenotypes, to enable tailored advice. This study evaluates the effectiveness of this approach in improving patient engagement and weight loss within a digital intervention.

**Objective:** To assess whether sending personalized weight loss advice can increase program engagement and lead to greater weight loss.

**Methods:** A quasi-experimental design was employed. Patients on a weight loss program were sent a 17-item questionnaire which matched them to one of four cognitive-behavioral profiles (phenotypes). Those who completed the questionnaire were sent phenotype-tailored weight loss advice once per week. As part of their weight loss program, patients used an app to track their meals and activities. Number of in-app events (i.e. engagements) was used as a proxy measure of program engagement. Self-reported weight was submitted as part of the structured weight loss program. Outcomes were compared to a historical cohort of patients who participated in the same weight loss program one year earlier, and those who were sent an invitation to complete the questionnaire but did not (i.e. non-responders).

**Results:** Those who received phenotype-tailored advice generated significantly more in-app engagements ( $M=257$ ,  $SD=232$ ), than those in the historical cohort ( $M=159$ ,  $SD=187$ ;  $P<.001$ ), and compared to non-responders ( $M=135$ ,  $SD=198$ ;  $P<.001$ ). There was a trend towards greater weight loss for those who received the tailored advice ( $M=-2.23$ kg,  $SD=7.97$ ), compared to the historical cohort ( $M=-1.6$ kg,  $SD=5.39$ ), and the non-responders ( $M=-0.69$ kg,  $SD=13.23$ ) but these differences were not significant, ( $P=.29$  and  $P=.23$ , respectively).

**Conclusions:** This study provides preliminary evidence that receiving weight loss advice tailored to an individual's cognitive-behavioral phenotype can increase engagement on a structured weight loss program, which could in turn support greater body mass reduction.

(JMIR Preprints 14/02/2025:72645)

DOI: <https://doi.org/10.2196/preprints.72645>

## Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

✓ **Please make my preprint PDF available to anyone at any time (recommended).**

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?


✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in [JMIR Publications](#)

No. Please do not make my accepted manuscript PDF available to anyone. I understand that if I later pay to participate in [JMIR Publications](#)

**Original Manuscript**



# Personalizing a Weight Loss Program With Cognitive-Behavioral Phenotypes to Increase Engagement and Weight Loss in Adults With Overweight or Obesity: Quasi-Experimental Study

Joanna Szypula<sup>1\*</sup>, Andreas Jarvstad<sup>1</sup>, Lucy Jones<sup>2</sup> & Katy Tapper<sup>1</sup>

<sup>1</sup> School of Health and Psychological Science, City St George's, University of London

<sup>2</sup> Oviva, 20 St Thomas Street, London SE1 9RS

\*Corresponding author, email: [joanna.szypula@city.ac.uk](mailto:joanna.szypula@city.ac.uk)

## Abstract

**Background:** Obesity affects more than one-quarter of adults in the United Kingdom and is a leading cause of preventable disease and health care costs. Digital behavior change programs can provide scalable weight management support, but maintaining engagement is challenging, and engagement is strongly associated with weight loss success. Tailoring interventions to cognitive-behavioral phenotypes, distinct patterns of thinking and behavior, offers one strategy to improve adherence. Although such approaches show promise in controlled settings, evidence from real-world digital programs is limited.

**Objective:** This study evaluated whether phenotype-tailored weekly advice improved engagement and weight loss in a national digital weight management program. Secondary aims were to assess correlations between advice use and outcomes, explore moderation by socioeconomic status, and capture participant perceptions of the advice.

**Methods:** We conducted a quasi-experimental study among UK adults enrolled in a free, 12-week program commissioned by the National Health Service. Eligible participants were aged 18–80 years with body mass index (BMI) >25 kg/m<sup>2</sup>. The phenotype group (n=148; mean age 48 years, 86% female, mean BMI 39 kg/m<sup>2</sup>) completed a 17-item questionnaire, were classified into one of four phenotypes, and received weekly tailored advice for 7 weeks. Comparators included a historical cohort enrolled one year earlier without phenotype advice (n=241; mean age 44 years, 71% female) and non-responders who did not complete the questionnaire (n=394; mean age 44 years, 76% female). Primary outcomes were program engagement (any in-app activity such as meal logging, activity tracking, content reading, or coach messaging) and self-reported weight.

**Results:** The phenotype group recorded a mean of 257 engagements (SD 232) over 7 weeks, significantly higher than both the historical cohort (159, SD 187; P<.001) and non-responders (135, SD 198; P<.001), representing 62–90% greater activity. All engagement types were significantly elevated (all P<.001). Mean weight loss was –2.23 kg (SD 7.97) in the phenotype group, compared with –1.60 kg (SD 5.39) in the historical cohort (P=.29) and –0.69 kg (SD 13.23) in non-responders (P=.23). The number of phenotype-specific advice documents opened correlated with engagement (r=0.48, P<.001) but not with weight loss (P=.42). Socioeconomic status did not moderate outcomes. Post-trial interviews (n=16) provided mixed feedback: many participants described the advice as clear, relevant, and motivating, while others considered it too general or poorly matched.

**Conclusions:** Phenotype-tailored weekly advice was associated with substantially higher engagement in a real-world digital program, although short-term weight differences were not statistically significant. While limited by non-randomized design, short follow-up, and reliance on self-reported weight, this study suggests phenotype-based tailoring may be a scalable strategy to strengthen adherence in digital weight loss interventions. Larger, randomized trials with longer follow-up are warranted to determine whether increased engagement translates into clinically meaningful weight loss.

**Registration:** <https://doi.org/10.17605/OSF.IO/UGWXB>

**Key words:** weight management; intervention adherence; personalization

## Introduction

Global obesity rates have doubled since 1990 [1], with around 1 in 4 adults in the United Kingdom now living with the condition [2]. While early interventions focused on telling individuals to ‘eat less and move more’, research over the past two decades highlights obesity as a complex, multifactorial issue requiring multicomponent, evidence-based treatment [3].

Many weight loss interventions are generic, offering identical advice to all participants [4,5]. This may partly explain the wide variability in individual responses, even to intensive treatments such as bariatric surgery [6]. Recent efforts have focused on personalizing interventions, often through genetic, anthropometric, or microbiota-based approaches [7–9]. There is growing evidence that personalized nutrition and exercise advice can enhance outcomes [8–10], but comparatively little attention has been paid to tailoring weight loss interventions based on cognitive-behavioral characteristics. Some research has used tools like the Three Factor Eating Questionnaire (TFEQ), Dutch Eating Behavior Questionnaire (DEBQ), or 6-Factor Questionnaire (6-FQ) to generate behavioral profiles [11–13], but the effects of tailoring advice based on these profiles remain largely untested.

Digital, app-based weight loss interventions are increasingly popular due to their scalability, accessibility, and low cost [14]. They can be as effective, or more so, than in-person programs [15–17], and allow for large-scale, real-world data collection to assess adherence and outcomes [18]. Individual differences such as personality may influence in-app engagement and outcomes in digital weight management programmes [19]. Tailoring interventions has been shown to increase engagement, which may lead to better outcomes [20–22], possibly because personalized advice feels more relevant and persuasive [23–26]. Improving engagement is particularly important in digital settings, where high attrition can limit effectiveness [26].

In-app engagement is commonly used as a proxy for adherence, with more frequent interaction indicating greater engagement [20]. A recent analysis by Lehmann et al. [18] found a significant relationship between Oviva app engagement and weight loss at 3 months across all countries analyzed, but not specifically within the United Kingdom. This non-significant relationship in the United Kingdom may be explained by the fact that this data came from more complex patients living with obesity, who often face greater challenges such as previous unsuccessful weight loss attempts, reliance on prescribed treatments (eg, GLP-1s or total dietary replacement), higher emotional eating, loss of control, significant comorbidities, and psychological barriers [27,28]. These factors could account for the weaker association between app engagement and weight loss in this group. Nevertheless, the broader international findings support the interpretation that app engagement reflects meaningful behavioral activity. App features prompt users to engage in evidence-based techniques such as self-monitoring, reflection, and problem-solving, all known to facilitate weight loss [29]. Therefore, increases in app engagement are likely to reflect active participation in self-management strategies, which may contribute to improved weight outcomes.

For this study, we collaborated with Oviva, a provider of digital weight loss services commissioned by the United Kingdom’s National Health Service (NHS). In a previous study, we developed a 17-item questionnaire to assign users to one of four cognitive-behavioral phenotypes [30]. In this paper, we aimed to test whether receiving phenotype-tailored advice would improve app engagement and weight loss in a real-world setting. Using a quasi-experimental design, patients enrolled in the Oviva program were invited via email to complete a ‘quiz’ that provided personalized advice. Those who responded received weekly, tailored emails for seven weeks. Two comparison groups were included: a historical cohort of patients enrolled a year earlier, and a group of non-responders who received the invitation but did not click the ‘quiz’ link.

As socioeconomic status is known to be linked to obesity prevalence [31–35], but its role in digital program outcomes remains unclear, we additionally explored this in the present study. Some evidence suggests eHealth interventions can support weight loss in low-income adults [36], while others report no effect [37–39]. However, many studies may be underpowered to detect interactions or may not recruit representative samples from lower socioeconomic groups [40–44].

We hypothesized that patients receiving phenotype-tailored advice would show greater number of in-app engagements and would lose more weight, than the comparison groups. We also predicted that these effects would be moderated by socioeconomic status (using Index of Multiple Deprivation [IMD] as proxy), with less benefit among more deprived participants. Finally, we expected greater app engagement and weight loss among those who accessed more of their phenotype-matched content.

## Method

### Ethical Considerations

Ethical approval for this study was granted by the City, University of London Psychology Department Research Ethics Committee (ETH2324-1944; amendments: ETH2324-2135 and ETH2324-2449). Participants did not sign a separate consent form, as patients enrolling in the Oviva program provide broad consent to be included in trials and for their anonymized data to be used for research, including secondary analyses. Eligible patients received an email inviting them to complete the phenotype questionnaire, with a clear notice that accessing the survey would enroll them into a 7-week automated email sequence. Participation was voluntary, and recipients could unsubscribe at any time. Historical comparison data were fully anonymized and covered under the original consent. Email addresses were used only to match individuals to anonymized patient user IDs, after which all identifiers were replaced. App engagement data were analyzed in anonymized form. Only authorized internal teams accessed identifiable data for email delivery and reporting, and no personal information was included in the research dataset. Participants were not financially compensated but received weekly educational materials as part of the intervention.

### Participants

Participants were recruited from Oviva, an NHS-contracted weight loss service provider, which is free for patients. They were sent an invitation email if they had started their weight management program with Oviva within the three-week recruitment period in May 2024. The weight management program used for recruitment is a 12-week digital behavioral and lifestyle intervention offered to NHS-referred patients. It includes support from specialist weight management dietitians and health coaches. Participants gain access to the Oviva app, a class IIa medical device, which features lessons on weight management and healthy lifestyle, goal setting with reminders, behavior tracking (eg., food, fluid intake, physical activity), and communication with their coach to monitor progress. Any activity on the app, such as logging, messaging the coach, or reading a lesson, is recorded as an "engagement". As this was a pragmatic, quasi-experimental study using convenience sampling over a fixed three-week recruitment period, enrolment was determined by natural sign-up rates.

### Program Eligibility

To join the weight management program at Oviva, patients must meet the following eligibility criteria: be aged 18-80, reside in or be registered with a GP practice in England, have a BMI  $\geq 30$ kg/m<sup>2</sup> (or  $\geq 27.5$ kg/m<sup>2</sup> for individuals of black African, African-Caribbean, and Asian origin), have a diagnosis of diabetes and/or hypertension, own a smartphone, and be able to speak English. Patients are ineligible if they have severe or moderate frailty as recorded on a frailty register, are pregnant, have been diagnosed with an eating disorder, have a significant unmanaged co-morbidity (eg,

chronic obstructive pulmonary disease not on medication), or have had bariatric surgery in the last two years.

## Design

This study employed a quasi-experimental design with post-test non-equivalent comparison groups. All Oviva weight management patients who started their program within the recruitment window were invited to complete the 'phenotypes quiz'. Those who clicked on the link and completed the questionnaire were part of the 'phenotypes group'. There were two comparison groups: i) patients who enrolled onto the same program 12 months before we recruited our sample (historical cohort), and ii) patients who were invited to complete the questionnaire but did not click on the link (non-responders). The historical cohort was randomly selected by the Oviva analytics team, and the researchers had no access or influence over which patients were included. The historical cohort was selected as a comparable reference group, as no substantial modifications were made to the app or program structure since their participation. While minor updates and iterative improvements were made, these were not expected to meaningfully impact engagement or outcomes.

The dependent variables were weekly app engagements and weight loss over a 7-week period. Any of the following activities counted as an app engagement: tracking an event (eg, meal, fluid, waist circumference, mood), sending a chat message to a coach, opening a lesson page (each page viewed increased the engagement count), opening a weekly article.

## Materials and Measures

### *Demographics*

Participant demographics were provided by Oviva's analytics team, typically sourced from GP practices, with weights self-reported. Socioeconomic status was indicated by IMD decile, a UK measure that estimates poverty based on geographical location, considering factors like income, employment, crime, and education (higher IMD scores indicate lower deprivation; [45]).

### *Phenotypes*

### *questionnaire*

The phenotypes questionnaire, or 'quiz', is a 17-item questionnaire developed in our earlier study to match patients to one of four profiles [30]. Briefly, we identified five cognitive and behavioral constructs which i) could affect weight or weight loss efforts, ii) have an existing validated measure, and iii) are potentially modifiable. The five constructs were: maladaptive eating, hedonic overeating, reward reactivity, self-regulation and psychological avoidance (see Supplementary Materials for full questionnaire). For further methodological details, please consult the development paper (currently under review) [46].

The quiz was created in Qualtrics, with each item rated on a scale from 1 (strongly disagree) to 5 (strongly agree). Participants were matched to a phenotype by calculating a percentage fit for each profile and assigning them to the one with the highest fit. They were shown their profile match along with a short 3-4 bullet point description of their phenotype. Participants could either continue with their matched phenotype or review all profile descriptions and choose the one that resonated most with them. If two or more fit scores were equal, participants were shown all profile descriptions and asked to select one.

### *The*

### *phenotypes*

There were four phenotypes a participant could get matched with. The profiles were named after colors to keep the assignment emotionally neutral, and each was paired with a spice name to create a distinctive cue for participants, helping them remember their profile name over the following weeks. 'Red Chilli' was characterized by high maladaptive and hedonic eating, low self-regulation, and high psychological avoidance. 'Yellow Saffron' was based on high hedonic eating and reward reactivity, and low maladaptive eating. 'Purple Lavender' represented low self-regulation and high psychological avoidance, as well as low maladaptive and hedonic eating. Lastly, Green Sage

reflected low maladaptive and hedonic eating, high self-regulation, and low psychological avoidance. The four cognitive-behavioral phenotypes are intended as practical, non-exclusive groupings that highlight dominant behavioral tendencies rather than rigid or mutually exclusive categories. Given the complexity of eating behavior, some overlap in traits and relevance of advice is expected. The profiles aim to guide individuals in identifying and addressing key obstacles to weight loss, while acknowledging that multiple factors may coexist.

### *Phenotype content*

Participants received weekly emails with advice tailored to their phenotype. The first document provided an overview and initial content, followed by weekly strategies addressing specific challenges (eg, overeating). Each document provided evidence-based advice, and a short exercise to help participants put this knowledge into practice.

The Red Chilli profile focused on reducing restriction and maladaptive eating using acceptance and commitment therapy, mindfulness, cognitive defusion, and self-compassion. The Yellow Saffron Profile addressed hedonic overeating with smaller portions of enjoyable foods, sensory eating techniques, craving management, and environmental architecture for healthier choices. The Purple Lavender profile improved goal setting and adherence by engaging with health goals daily, starting with small behaviors, and using routines, cognitive architecture, and defusion for challenging thoughts. Finally, the Green Sage profile enhanced diet quality and physical activity by adding vegetables, choosing foods with longer oral processing times, flexible eating (80% nutritious, 20% less nutritious), and cognitive architecture for activity goals.

### *Post-trial feedback*

After receiving all content emails, participants were sent a final email requesting feedback via a Qualtrics questionnaire, which included 11 Likert scale questions (1 = Strongly Disagree to 5 = Strongly Agree) assessing the helpfulness, effectiveness, and clarity of the advice. Participants were also given four optional open-ended questions to comment on the quiz, useful aspects, and suggestions for improving their profile or the advice.

### **Procedure**

Participants were sent an email inviting them to take part in the 'phenotypes quiz'. Those who completed the 'quiz' were automatically matched to a profile and asked to read through its description, based on which they could decide to accept their match or change it. Once their response was submitted, they received an automatic email reminding them which profile they were matched with and were added to a seven-week long campaign.

Participants received the first email with tailored advice a day after they took the quiz. This email contained links to four documents (one for each phenotype), with participants instructed to open the one matching their profile. Six more weekly emails followed, each containing four profile-specific documents. This means that throughout the trial, participants received links to 28 documents, but were instructed to only open seven of them. The final email contained a link to the post-trial feedback. The core program and care provided by dietitians and health coaches remained unchanged. Although the Tier 2 program lasts 12 weeks, the phenotype-specific advice was delivered over a 7-week period due to project timelines, which required all data collection and analysis to be completed within a fixed timeframe.

### **Analysis Plan**

The main analyses pertained to the effect of receiving phenotype-related advice on the average number of app engagements, and weight loss over a 7-week period. Hierarchical regressions were conducted to compare the average number of app engagements between the phenotype group, and the historical cohort group, as well as the non-responder group. We tested whether IMD moderated

this relationship by including it as step 2 of the regression, and the interaction term as step 3. The same approach was used to analyze the effect of condition on weight loss.

We further carried out equivalent Bayesian analyses using STAN (stan\_glm in rstanarm 2.32.1, brms 2.22.0, R4.4.1, RStudio 2024.12). We fit four models for each dependent variable (DV): 1)  $DV \sim 1$  (intercept only), 2)  $DV \sim \text{group}$ , 3)  $DV \sim \text{group} + \text{IMD}$ , and 4)  $DV \sim \text{group} * \text{IMD}$ . The engagement variable was square root transformed and the IMD variable standardized prior to fitting. The weight loss variable contained outliers and a preponderance of 0s. To maintain parity with the standard regression analyses, we did not further process these data in any way. We used weakly informative robust t-distribution priors ( $dF = 3$ ,  $M = 0$ ,  $SD = 10$ ), and a Gaussian link function. We ran 8 chains, each with 1000 discarded burn-in samples and 5000 retained samples, for a total of 20 000 retained samples per model. We verified model convergence using standard criteria (eg.,  $R_{hat} \sim 1$ , visually inspected chains for convergence). For each model we computed the log marginal likelihood and computed Bayes Factors (BF) first for target models against the null intercept-only model. For any models which provided BF in favor of the target models over the intercept-only model (ie,  $BF > 3$ ) we additionally computed pairwise BFs to establish which of the ‘surviving’ models best explain the data.

Change in the average number of engagements on a week-by-week basis was analyzed using a repeated-measures ANOVA.

Qualitative feedback from open-ended survey responses and semi-structured interviews was analyzed using a descriptive thematic approach. Responses were reviewed inductively to identify recurring sentiments and contrasting views across participants. Illustrative quotes were selected to reflect the diversity of opinions, and analysis focused on summarizing key perceived benefits, limitations, and personal relevance of the intervention.


The study protocol was uploaded to OSF prior to data analysis, and the analysis plan was pre-registered before obtaining engagement and weight loss data from Oviva [47]. Data were analyzed using SPSS v.28.

## Results

### Participant characteristics and study flow

Figure 1 presents the flow of participants through the study. There was a reasonable match in participant demographics across the three groups (Table 1) though phenotype group participants were slightly older, more likely to be female, and their average IMD score was higher than that of the comparison groups.

Figure 1. Flow of participants in a quasi-experimental study evaluating the effect of receiving phenotype-matched weight loss advice on app engagement and weight loss among patients actively participating in a UK weight loss program, recruited in May 2024.



Completed feedback survey (n=17)
-------------------------------------

Table 1. Demographic characteristics of participants in the phenotype group and two comparison groups (historical cohort and non-responders).

	Phenotypes group (n=148)	Historical cohort (n=241)	Non-responders (n=394)
Age in years	48 (12)	44 (10)	44 (11)
Sex (% female)	86%	71%	76%
Starting weight (kg)	108 (19.7)	111 (24.1)	109 (25.0)
Starting BMI (kg/m <sup>2</sup> )	39 (6)	39 (8)	39 (8)
Ethnicity			
- White	86%	69%	71%
- Black	5%	16%	10%
- Asian	5%	10%	8%
- Mixed	2%	1%	5%
- Other	1%	3%	3%
- Refused response	0%	1%	2%
- Missing data	1%	1%	1%
Mean IMD (SD) <sup>a</sup>	6 (3)	5 (3)	5 (3)
Median IMD	6	3	5
IQR <sup>b</sup> of IMD	5	6	4


<sup>a</sup>Note: scores range from 1-10, lower values indicate higher relative deprivation. Standard deviation shown in parentheses.

<sup>b</sup>Interquartile Range

## Phenotype assignment

Figure 2 shows how phenotype-group participants were assigned to their final profile. Note that 18 patients who received an invitation to the 'quiz' clicked on the link, but did not complete the questionnaire, meaning they received the weekly emails but were unaware which links they should click on. Data from these participants were excluded from all analyses.

Figure 2. Flow of participants in the phenotype group from starting the phenotype quiz to receiving a final profile match, which determined the tailored weight loss advice they received in the trial.



## Phenotype fidelity

Participants were asked to only match with one profile. They were matched with (eg, those matched with Red Chilli show documents for all profiles were multiple profiles. Fidelity was defined as opening at least one document tailored to the profile the participant was matched with, and not opening more than one document from a different profile. The fidelity rate was 73.6% throughout the trial. Among the remaining participants, 4.1% followed a different click pattern.

## Main analyses

### Effect of condition on total engagements

The average total number of engagements over a 7-week period was higher in the phenotype group, than in the historical cohort (see Table 2). A hierarchical linear regression was conducted to compare the average number of engagements between the two groups, and to examine whether this effect was moderated by IMD. Condition was entered at step 1, IMD at step 2, and the interaction term at step 3. The analysis revealed that participants in the phenotype group generated significantly more engagements, than those in the historical cohort ( $b = -93.61$ ,  $SE B = 22.39$ ,  $\beta = -0.22$ ,  $P < .001$ ), but the interaction was not significant ( $b = -4.44$ ,  $SE B = 7.54$ ,  $\beta = -0.12$ ,  $P = .56$ ). An equivalent Bayesian analysis confirmed that a model including only an intercept term ( $B = 11.91$ , 95% CDI = [11.05, 12.74]) and the main effect of phenotype group ( $B = 1.89$ , 95% CDI [1.05, 2.74]) was superior to the model that also included a main effect of IMD (BF = 144), as well as the interaction model (BF = 1429). All the models were superior to the baseline intercept-only model (all BFs > 14). Independent  $t$ -tests showed that all engagement subtypes (tracking activity, coach messaging, pages of educational modules viewed, and articles read) were significantly higher in the phenotype group than the historical cohort group, suggesting that the overall increase in engagement was not driven by any single type of app interaction (all  $P < .001$ ; see Supplementary Materials for all analyses).

### Effect of condition on weight loss

Average weight loss over a 7-week period was compared between the phenotype group, and the historical cohort (see Table 2). Another hierarchical regression was conducted to test the effect of

condition on weight loss, and whether IMD moderated this relationship. Condition was entered at step 1, IMD at step 2, and the interaction term at step 3. No significant differences in weight loss were found between the two groups ( $b = 0.85$ ,  $SE B = 0.80$ ,  $\beta = 0.06$ ,  $P = .29$ ), and the interaction was not significant ( $b = -0.26$ ,  $SE B = 0.27$ ,  $\beta = -0.22$ ,  $P = .33$ ). The equivalent Bayesian analysis showed that a base-line intercept model was substantially better than all the main effect and interaction models (all BFs  $\geq 567$ ), showing that neither group, or IMD, or their interaction explained variation in weight loss.

### *Relationship between documents opened and primary outcomes*

Among the phenotype cohort there was a significant positive correlation between the number of correct documents opened (ie, documents which matched the phenotype the participant was assigned to) and total engagements,  $r(146) = .481$ ,  $P < .001$ . There was no relationship between the number of correct documents opened and weight loss,  $r(119) = -.074$ ,  $P = .42$ .

### *Comparison with non-responders*

An independent groups *t*-test revealed participants in the phenotypes group generated significantly more engagements over the 7-week period, than non-responders (equal variances not assumed as Levene's Test  $F_{1,540} = 12.98$ ,  $P < .001$ ),  $t_{232.25} = 5.68$ ,  $P < .001$ ; see Table 2). Weight loss over the 7-week period was higher among the phenotype group, compared to non-responder, but this difference did not reach statistical significance,  $t_{445} = -1.21$ ,  $P = .23$  (see Table 2).

Table 2. Mean number of app engagements and mean weight loss in the phenotype group and two comparison groups (historical cohort and non-responders).

Group	Total app engagement count, mean (SD)	Weight loss (kg), mean (SD)
Phenotypes group (n=148)	257 (232)	2.23 (7.97)
Historical cohort (n=241)	159 (187)	1.60 (5.39)
Non-responders (n=394)	135 (198)	0.69 (13.23)

### *BMI and body mass percentage reduction*

On average, participants in the phenotype group reduced their BMI by  $0.79 \text{ kg/m}^2$  ( $SD = 2.90$ ), compared to a reduction of  $0.56 \text{ kg/m}^2$  ( $SD = 1.94$ ) in the historical cohort, over the 7-week period. This translated into a reduction of 1.93% ( $SD = 7.24$ ) body mass in the phenotype group, and 1.27% ( $SD = 4.14$ ) reduction in body mass in the historical cohort. Non-responders reduced their BMI by  $0.11 \text{ kg/m}^2$  ( $SD = 5.01$ ) and their body mass by 0.09% ( $SD = 13.01$ ).

## Exploratory analyses


### *Week-by-week engagements*

A repeated-measures ANOVA was conducted to assess the week by week change in app engagements, and the interactions between the phenotypes group, the historical cohort and the non-responders. As Mauchly's test of sphericity was significant ( $X^2(20) = 715.09$ ,  $P < .001$ , Mauchly's  $W = 0.40$ , Greenhouse-Geisser correction,  $\epsilon = .720$ ), Greenhouse-Geisser correction was applied. There was a significant effect of week number on app engagement,  $F_{4,32, 3367.68} = 81.85$ ,  $P < .001$ , and the interaction with condition was significant,  $F_{8,64, 3367.68} = 5.57$ ,  $P < .001$  (see Figure 3).

Figure 3. Mean weekly app engagements in the phenotype group (n=148) and the two comparison groups (historical cohort n=241; non-responders cohort n=394). Error bars denote 95% IC.

70

— Phenotypes group  
— Historical cohort  
— Non-responders



participant did not open  
analyzed. Overall, the

feedback on the phenotype-tailored advice was positive (see Table 3), as participants felt the advice they received was helpful and easy to understand. Participants were also willing to use the advice in the future.

Table 3. Post-trial participant ratings of phenotype-tailored advice (n=16), including perceived helpfulness, clarity, personalization, and intention to use the advice, scored on a 1–5 scale (higher ratings indicate greater endorsement).

Statement	Rating (SD)
The advice I received was helpful	3.75 (1.13)
The advice helped me make positive changes to my lifestyle	3.63 (1.26)
The advice helped me to lose weight	3.25 (1.24)
The length of the advice was just right (not too short and not too long)	3.50 (1.37)
The advice was clear and easy to understand	4.06 (1.00)
I learned something new by reading the advice in the resources	3.44 (1.31)
The advice I received felt personalized to me	3.13 (1.36)
The advice I received felt relevant to me	3.44 (1.31)
I followed the advice given to me in the resources	3.88 (0.72)
I completed the tasks in the 'Try it out' section in each of the resources	3.69 (1.01)
I will use the advice from the resources in the future	3.81 (0.98)

## Qualitative feedback

Qualitative feedback was collected in two ways: in the post-trial feedback questionnaire, and in post-trial interviews. The feedback was mixed, with some people feeling very positive about the phenotypes, and others feeling they were not as useful. One participant said the advice “was good” (Participant 9), and found it especially helpful to learn about “the reasons behind my eating when not hungry and the food I choose”. Another participant said “all advice was spot on (...) I looked forward to reading them and I will miss them. More would be great (...) I found this the most helpful even more than the learning modules (...) [understanding] how my mind works in relation to food certainly explained a lot” (Participant 10). Two more participants wrote “Excellent advice & profiles” (Participant 11) and “I found the information encouraging to read” (Participant 14). Two participants felt more neutral about their experience as one said “it was all useful, except I haven’t lost very much weight” (Participant 13) and “It was ok. To be honest I can’t recall much about it” (Participant 12). Others felt less positively about the phenotypes, saying the “quiz was not really helpful, advice was very vague and generalized (...) no specific tailored help” (Participant 2). Another participant said “Too much emphasis on how I feel. I hate being constantly asked to think

about how I feel. (...) I didn't really fit into any of the categories anyway" (Participant 4). Others said "it just feels generic and not very personal" (Participant 7) and "[the profile] didn't reflect me" (Participant 8).

Everyone who completed the post-trial survey was invited to participate in an interview, and 5 participants accepted. Participant 9 felt positively about the quiz and the Red Chilli profile she was matched with. She said the profile "was so accurate that it resonated" and "when I read the profile I was like (...) I know I'm an emotional eater (...) but actually, until I read that, it didn't make sense". She talked about how the profile "threw up things that you weren't aware of (...) you know what you should be doing. Why can't you do it? And I think that was the big question I really wanted to know the answer to (...) and then finding out that it's probably this and this (...) [that] was the most helpful thing". Participant 10, who was also matched with the Red Chilli profile, said: "honestly, every week, I've looked forward to reading them [the advice]. And I'm just sorry they've stopped now". She talked about how she used the learning from the profile to think "How do I feel? Am I being, you know, not very kind to myself? (...) it put things into perspective for me rather than just saying I've ate this, I've drank that". She mentioned how the advice was tailored to her needs, saying "I think I've lost well over a stone (...) I don't have a bad diet, I don't think I just choose fast, convenient food when I'm stressed (...) I've just put more thought into it and the thought that not being cruel to yourself, being actually like, oh, OK, you can do this". Moreover, she said "I found the language to be absolutely, you know, spot on for me that it had got me down to a T (...) I'm hoping that it [the advice] will make me stop and think (...) to actually keep the weight off because that's the second challenge, isn't it?"

Participant 6 was matched with the Yellow Saffron profile. She said: "quite often, diet or health things, they don't really allow for people who really love food in a foodie way (...) [the quiz] recognized that I was that sort of person (...) the advice was tailored to me (...) which actually I was really pleased with". She also said "I don't have to sort of feel defensive about it [being a foodie] (...) for dieting purposes, if you're in a particular profile, and they're not recognizing that in a positive way, so you get defensive and stressed (...) whereas this has recognized it in a positive way, and given me ways of thinking about food, how I like to live, support how I like to eat food (...) but at the same time, it said you could tweak this (...) I've probably been able to take a step back and view how I eat things in a slightly different way, but without feeling guilty about it".

In an interview with Participant 2, the participant said she did not feel her profile was the right fit and resonated with two profiles when shown all the profile descriptions. When asked for feedback, she was reviewing the core program, rather than the phenotype advice, and struggled to remember seeing the documents with the advice. Participant 7 opened four out of seven documents with advice for her profile, however, due to a system error, she was not receiving any coaching on the program. Thus, in the interview, she primarily talked about her dissatisfaction with the course in general, rather than about the phenotypes.

## Discussion

The aim of this quasi-experimental study was to assess the impact of providing tailored weight loss advice on app engagements and weight loss on a digital weight loss program. The participants were patients enrolled in a weight management program offered by Oviva, a service provider for the NHS. Advice was tailored based on responses to a 'phenotypes quiz', which matched patients to one of four cognitive-behavioral profiles. The evidence-based, tailored advice was delivered weekly over email, for seven weeks. App engagement (eg, logging meals, reading lessons) was used as a proxy for program adherence.

### *Effect on app engagements*

Patients who received phenotype-tailored advice generated significantly more app engagements over

a 7-week period, compared to a historical cohort of patients enrolled on the same program a year earlier, and compared to patients who received an invitation to take the 'quiz' but who did not click on the link (ie, non-responders). There was a 61.5% increase in app engagements between the phenotypes group and the historical cohort, and a 90.5% increase in engagements as compared to the non-responder group. There was a significant, positive correlation between the number of documents opened and the number of app engagements generated.

This finding aligns with our hypothesis that receiving tailored advice would boost program adherence. Additionally, the "try it out" section in the documents, which included exercises like logging hunger types in the app, likely contributed to increased engagement. This is unlikely to be driven purely by the instructions to use the app, as Oviva regularly encourages their users to log activities in the app, and such elevated engagement was not seen in the comparison groups. It could be that the coupling between tailored advice and instructions to log in the app was particularly effective. This is an important finding, as adherence to digital programs, often measured by app engagement, can be challenging. Enhancing adherence to digital interventions is crucial because it directly improves health-related outcomes [48]. Therefore, strategies that increase adherence are highly valuable for the success of eHealth interventions [49–51].

The increased engagement observed in the phenotype group may have been influenced by self-selection bias, given the study's quasi-experimental design. Since participants had to actively engage with multiple steps (opening an email, clicking the questionnaire link, and completing it) to receive their phenotype "match", it is possible that only the most motivated individuals chose to participate. This could, in turn, explain the higher engagement rates observed in this group. A randomized controlled trial is required to establish causality.

A randomized controlled trial could be feasibly integrated into digital weight management programs by randomizing eligible patients at the point of onboarding to either receive standard program content or phenotype-tailored advice in addition to the standard content. Randomization and content delivery could be automated within the app infrastructure, and outcomes (eg, app engagement, weight change) could be monitored via existing data pipelines. This design would allow for rigorous testing of the added benefit of phenotype-based personalization within a real-world setting, without changing or disrupting routine care.

Although engagement was higher in the phenotypes group, engagement levels declined week by week across all conditions. This pattern aligns with previous research on weight loss interventions [48–50] and Oviva's internal data. One possible explanation is habituation – as users become familiar with the app's features, their novelty wears off, leading to reduced interaction. Additionally, some users may have consistent meal routines, logging the same foods frequently in the early weeks but gradually reducing this behavior over time. Given that meal tracking is typically the primary contributor to in-app engagement, a decline in logging frequency could naturally result in lower overall engagement levels.

There was a slight imbalance in IMD scores between the phenotype group and the historical cohort, raising the question of whether relative deprivation status influenced engagement. However, the analysis showed that IMD was not associated with engagement, and adding IMD to the model did not improve its explanatory power beyond the effect of group alone. Furthermore, the interaction term was not informative due to the shared variance between predictor variables. While this may suggest that phenotype-based personalization supported engagement across socio-economic groups, the study was not powered to detect interaction effects, particularly given the unequal group sizes. Therefore, these null findings should be interpreted with caution.

### *Effect on weight loss*

**We also predicted that phenotype-tailored weight loss advice would lead to greater reduction in body mass over a 7-week period. While the phenotypes group showed more weight loss (-2.23kg) compared to the historical cohort (-1.60kg) and non-responders (-0.69kg), these differences were not statistically significant. IMD was not associated with weight loss, and the interaction between group and IMD was also non-significant. Bayesian analysis further confirmed that a baseline intercept model was substantially better than models including group, IMD, or their interaction, indicating that none of these variables explained variation in weight loss. This lack of statistically significant differences in weight loss between the groups may be explained by the fact that applying tailored advice takes time both to master and to impact weight loss, and our seven-week trial may have been too short to observe significant effects. A longer-term follow up on a larger sample may be worthwhile in future studies, to more clearly elucidate whether phenotype-tailored advice can increase weight loss.**

### *Post-trial feedback on the phenotypes*

Only a small proportion of participants (10.8%) provided post-trial feedback. While these results should be interpreted with caution, phenotypes and tailored advice were generally well received. The four most endorsed statements were that the advice was clear and easy to understand, helpful, followed by participants, and likely to be used in the future.

Interviews revealed mixed opinions. Some patients found the phenotypes useful, as they helped them understand the underlying reasons for their maladaptive eating behaviors. Others, however, felt the advice was too generic. In the survey, the statement with the lowest endorsement was that the advice felt personalized. One possible explanation is that the term "personalized" created expectations of a high level of individualization that were ultimately unmet. Additionally, some participants reported that none of the profiles fully resonated with them. This is not unexpected, as the four profiles are not exhaustive or mutually exclusive. They were designed to provide general guidance on the cognitive and behavioral patterns most likely to hinder progress, and some individuals may relate to more than one profile, or none at all.

A more hyper-personalized approach, where individual cognitive and behavioral dimensions are assessed separately and combined into tailored content, may improve both fit and impact. Such an approach is increasingly feasible through artificial intelligence-enabled systems. A recent study demonstrated that health advice can be tailored not only in terms of content but also in delivery, aligning with individuals' natural tendencies and cognitive styles [51]. This dual approach to personalization opens new possibilities for future work.

Recent advances in large language models (LLMs) are enabling digital health companies to deliver increasingly personalized messages, insights, and content [52–54]. Incorporating phenotypes into retrieval-augmented generation systems [55] offers a promising opportunity to enhance both relevance and precision of tailored interventions [56]. By aligning content with users' cognitive styles and behaviors, such systems may improve engagement and clinical outcomes [56–59]. While this study did not directly examine artificial intelligence-driven personalization, combining cognitive-behavioral profiling with LLMs represents a compelling avenue for future digital health innovation [59].

Another limitation relates to the delivery format. Some participants may have provided feedback on the core Oviva program rather than the phenotype-specific content (post-trial interviews confirmed at

least two instances), making it unclear whether low personalization ratings reflected dissatisfaction with the advice or broader program frustrations. Delivering tailored advice through a separate medium (email), while the main program was delivered in-app, may also have reduced coherence. Future studies would benefit from integrating tailored content directly into the digital intervention itself, to create a more seamless, unified experience that could further improve engagement and outcomes.

### *Strengths and limitations*

One strength of our study was testing phenotype-tailored advice on patients enrolled in an NHS-referred weight loss program, making our sample representative of UK patients seeking weight loss treatment and enhancing external validity. Additionally, by using an established program provider, we leveraged a well-developed app and core program, minimizing adherence issues due to app functionality, a common problem in prototype apps used in studies.

However, our study had limitations. The quasi-experimental design limits causal inferences. It is possible that because participants were not randomly assigned, only the most motivated individuals completed the 'phenotypes quiz', which could explain the higher engagement in this group. Without a randomized controlled trial or baseline motivation measurements, this alternative explanation remains possible.

Another limitation is that tailored content was delivered via email rather than integrated into the app, creating a disjointed experience that may have led participants to confuse the core program with the phenotype content. This separation could also have resulted in some participants missing the email content, leading to a fragmented experience. Finally, the phenotypes were based on data from four Oviva weight management programs, but the sample in this study was predominantly matched with the Red Chilli profile. Recruiting from a single program may have contributed to the observed differences in phenotype distribution. Such an uneven distribution might also indicate a need for further refinement of subgroups within profiles to better differentiate patient challenges.

**Finally, weight was self-reported, which may have introduced bias (eg, social desirability, systematic under-reporting), but self-reporting is standard practice in remotely delivered Tier 2 programs and accepted by the NHS for monitoring patient progress. On the program, participants track weight privately with their coach, with no penalties or incentives for weight change. In the study, phenotype-specific advice focused on cognition and behavior rather than weight outcomes, aligning with the app's broader behavioral framing. Participants were not specifically prompted to report weight or emphasize weight loss, and were unaware it would be assessed for the trial. Importantly, any bias in weight reporting is unlikely to have affected the study's primary aims, as analyses focused on differences across groups or time rather than absolute weight values.**

### *Conclusion*

In conclusion, our study provided preliminary evidence that advice tailored to cognitive-behavioral phenotypes may increase program adherence, as indicated by app engagements. Although the quasi-experimental design limits causal conclusions, the initial data trends are promising. Despite limited post-trial feedback, the positive ratings suggest further exploration of phenotype-tailored advice is warranted. To fully assess its impact on app engagement and weight loss, a randomized controlled trial with a larger sample and longer timeframe is needed.

## Acknowledgements

JS, KT, AJ and LJ contributed to the conceptualization and methodology of the study. JS, KT and AJ conducted the formal analysis, with AJ particularly focusing on the Bayesian approach. JS led the execution and coordination of the study, overseeing data collection and the provision of study materials. Additionally, JS prepared the original draft of the manuscript, which was subsequently reviewed and edited by KT, LJ and AJ. This work was conducted as part of a Knowledge Transfer Partnership (KTP number: 10029186), which was supported by UK Research and Innovation (grant number 13221). The authors attest that there was no use of generative artificial intelligence (AI) technology in the generation of text, figures, or other informational content of this manuscript.

## Data availability statement

**The datasets generated and/or analyzed during this study are not publicly available due to the potential risk of re-identification. While reasonable anonymization procedures were applied, there remains a theoretical risk that individuals could be re-identified if the data were combined with internal patient records held by Oviva. Although these records are not publicly accessible, this possibility means the study data cannot be shared.**

## Conflicts of interest

None declared.

## References

1. World Health Organization. Obesity and overweight. Accessed June 9, 2025. <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>
2. Obesity. nhs.uk. November 23, 2017. Accessed June 9, 2025. <https://www.nhs.uk/conditions/obesity/>
3. English S, Vallis M. Moving beyond eat less, move more using willpower: Reframing obesity as a chronic disease impact of the 2020 Canadian obesity guidelines reframed narrative on perceptions of self and the patient-provider relationship. *Clin Obes.* 2023;13(6):e12615. doi:10.1111/cob.12615
4. Annesi JJ, Whitaker AC. Psychological Factors Discriminating Between Successful and Unsuccessful Weight Loss in a Behavioral Exercise and Nutrition Education Treatment. *Int J Behav Med.* 2010;17(3):168-175. doi:10.1007/s12529-009-9056-2
5. Kwasnicka D, Luszczynska A, Hagger MS, et al. Theory-based digital intervention to promote weight loss and weight loss maintenance (Choosing Health): protocol for a randomised controlled trial. *BMJ Open.* 2020;10(11):e040183. doi:10.1136/bmjopen-2020-040183
6. Dent R, McPherson R, Harper ME. Factors affecting weight loss variability in obesity. *Metabolism.* 2020;113:154388. doi:10.1016/j.metabol.2020.154388
7. Holzapfel C, Dawczynski C, Henze A, Simon MC. Personalized dietary recommendations for weight loss. A scientific perspective from various angles. *Ernahrungs Umsch.* 2021;(68(2)):26-35. doi:10.4455/eu.2021.008
8. Ordovas JM, Ferguson LR, Tai ES, Mathers JC. Personalised nutrition and health. *BMJ.* Published online June 13, 2018:bmj.k2173. doi:10.1136/bmj.k2173

9. Livingstone KM, Ramos-Lopez O, Pérusse L, Kato H, Ordovas JM, Martínez JA. Reprint of: Precision nutrition: A review of current approaches and future endeavors. *Trends Food Sci Technol.* 2022;130:51-62. doi:10.1016/j.tifs.2022.10.010
10. Vandelanotte C, Trost S, Hodgetts D, et al. Increasing physical activity using an just-in-time adaptive digital assistant supported by machine learning: A novel approach for hyper-personalised mHealth interventions. *J Biomed Inform.* 2023;144:104435. doi:10.1016/j.jbi.2023.104435
11. Romain AJ, Avignon A, Macioce V, Boegner C, Attalin A, A S. Patterns of eating behavior in people with severe obesity seeking weight loss treatment: An exploratory study. *Appetite.* 2022;169:105797. doi:10.1016/j.appet.2021.105797
12. Pentikäinen S, Arvola A, Karhunen L, Pennanen K. Easy-going, rational, susceptible and struggling eaters: A segmentation study based on eating behaviour tendencies. *Appetite.* 2018;120:212-221. doi:10.1016/j.appet.2017.09.001
13. Kushner RF, Choi SW, Burns JL. Development of a six-factor questionnaire for use in weight management counseling. *Patient Educ Couns.* 2016;99(12):2018-2025. doi:10.1016/j.pec.2016.06.022
14. Wren GM, Koutoukidis DA, Scragg J, et al. Optimising a digitally delivered behavioural weight loss programme: study protocol for a factorial cluster randomised controlled trial. *Trials.* 2024;25(1):477. doi:10.1186/s13063-024-08320-5
15. Besson M, Gurviez P, Carins J. Using digital devices to help people lose weight: a systematic review. *J Soc Mark.* 2020;10(3):289-319. doi:10.1108/JSOCM-07-2019-0115
16. Kupila SKE, Joki A, Suojanen LU, Pietiläinen KH. The Effectiveness of eHealth Interventions for Weight Loss and Weight Loss Maintenance in Adults with Overweight or Obesity: A Systematic Review of Systematic Reviews. *Curr Obes Rep.* 2023;12(3):371-394. doi:10.1007/s13679-023-00515-2
17. Beleigoli AM, Andrade AQ, Cançado AG, Paulo MN, Diniz MDFH, Ribeiro AL. Web-Based Digital Health Interventions for Weight Loss and Lifestyle Habit Changes in Overweight and Obese Adults: Systematic Review and Meta-Analysis. *J Med Internet Res.* 2019;21(1):e298. doi:10.2196/jmir.9609
18. Lehmann M, Jones L, Schirmann F. App Engagement as a Predictor of Weight Loss in Blended-Care Interventions: Retrospective Observational Study Using Large-Scale Real-World Data. *J Med Internet Res.* 2024;26:e45469. doi:10.2196/45469
19. Hagiwara Y, Shimizu K, Adachi T, Kanai M, Miki T. Impact of personality prototypes on app engagement and weight loss in a mobile app-based lifestyle modification support program for individuals with obesity and non-communicable diseases. *Digit Health.* 2025;11:20552076251347891. doi:10.1177/20552076251347891
20. Jakob R, Harperink S, Rudolf AM, et al. Factors Influencing Adherence to mHealth Apps for Prevention or Management of Noncommunicable Diseases: Systematic Review. *J Med Internet Res.* 2022;24(5):e35371. doi:10.2196/35371

21. Ryan K, Dockray S, Linehan C. A systematic review of tailored eHealth interventions for weight loss. *Digit Health*. 2019;5:2055207619826685. doi:10.1177/2055207619826685
22. Lustria MLA, Noar SM, Cortese J, Van Stee SK, Glueckauf RL, Lee J. A Meta-Analysis of Web-Delivered Tailored Health Behavior Change Interventions. *J Health Commun*. 2013;18(9):1039-1069. doi:10.1080/10810730.2013.768727
23. Kreuter MW, Strecher VJ, Glassman B. One size does not fit all: The case for tailoring print materials. *Ann Behav Med*. 1999;21(4):276-283. doi:10.1007/BF02895958
24. Hunot-Alexander C, Croker H, Fildes A, Johnson F, Beeken RJ. Brief 'Appetitive Trait Tailored Intervention': Development in a Sample of Adults with Overweight and Obesity. *Behav Change*. 2022;39(2):106-122. doi:10.1017/bec.2021.22
25. Flores Mateo G, Granado-Font E, Ferré-Grau C, Montaña-Carreras X. Mobile Phone Apps to Promote Weight Loss and Increase Physical Activity: A Systematic Review and Meta-Analysis. *J Med Internet Res*. 2015;17(11):e253. doi:10.2196/jmir.4836
26. Burgess E, Hassmén P, Welvaert M, Pumpa KL. Behavioural treatment strategies improve adherence to lifestyle intervention programmes in adults with obesity: a systematic review and meta-analysis. *Clin Obes*. 2017;7(2):105-114. doi:10.1111/cob.12180
27. Arhi C, Karagianni C, Howse L, Siddiqui M, Borg CM. The Effect of Participation in Tier 3 Services on the Uptake of Bariatric Surgery. *Obes Surg*. 2021;31(6):2529-2536. doi:10.1007/s11695-021-05303-2
28. Hazlehurst JM, Logue J, Parretti HM, et al. Developing Integrated Clinical Pathways for the Management of Clinically Severe Adult Obesity: a Critique of NHS England Policy. *Curr Obes Rep*. 2020;9(4):530-543. doi:10.1007/s13679-020-00416-8
29. Berry R, Kassavou A, Sutton S. Does self-monitoring diet and physical activity behaviors using digital technology support adults with obesity or overweight to lose weight? A systematic literature review with meta-analysis. *Obes Rev*. 2021;22(10):e13306. doi:10.1111/obr.13306
30. Szypula J, Jones L, Barron A, Jarvstad A, Tapper K. Personalising weight loss strategies with cognitive-behavioural phenotypes. *Appetite*. 2024;199:107487. doi:10.1016/j.appet.2024.107487
31. Khanolkar AR, Patalay P. Socioeconomic inequalities in co-morbidity of overweight, obesity and mental ill-health from adolescence to mid-adulthood in two national birth cohort studies. *Lancet Reg Health - Eur*. 2021;6:100106. doi:10.1016/j.lanep.2021.100106
32. El-Sayed AM, Scarborough P, Galea S. Unevenly distributed: a systematic review of the health literature about socioeconomic inequalities in adult obesity in the United Kingdom. *BMC Public Health*. 2012;12(1):18. doi:10.1186/1471-2458-12-18
33. Newton S, Braithwaite D, Akinyemiju TF. Socio-economic status over the life course and obesity: Systematic review and meta-analysis. Xiao G, ed. *PLOS ONE*. 2017;12(5):e0177151. doi:10.1371/journal.pone.0177151
34. Szinay D, Forbes CC, Busse H, DeSmet A, Smit ES, König LM. Is the uptake, engagement, and

- effectiveness of exclusively mobile interventions for the promotion of weight-related behaviors equal for all? A systematic review. *Obes Rev.* 2023;24(3):e13542. doi:10.1111/obr.13542
35. Parkhurst A. *Fat Chance? Exploring the Evidence on Who Becomes Obese.* 2020health; 2015. <https://2020health.org/wp-content/uploads/2020/11/2020FATCHANCE-Final.pdf>
  36. Clark TL, Savin KL, Perez-Ramirez P, Valdez T, Toba G, Gallo LC. eHealth weight loss interventions for adults with low income: A systematic review. *Health Psychol.* 2023;42(6):353-367. doi:10.1037/hea0001278
  37. Richards R, Wren G, Whitman M. The Potential of a Digital Weight Management Program to Support Specialist Weight Management Services in the UK National Health Service: Retrospective Analysis. *JMIR Diabetes.* 2024;9:e52987. doi:10.2196/52987
  38. Toon J, Geneva M, Sharpe P, Lavin J, Bennett S, Avery A. Weight loss outcomes achieved by adults accessing an online programme offered as part of Public Health England's 'Better Health' campaign. *BMC Public Health.* 2022;22(1):1456. doi:10.1186/s12889-022-13847-w
  39. Birch JM, Jones RA, Mueller J, et al. A systematic review of inequalities in the uptake of, adherence to, and effectiveness of behavioral weight management interventions in adults. *Obes Rev.* 2022;23(6):e13438. doi:10.1111/obr.13438
  40. Coupe N, Cotterill S, Peters S. Enhancing community weight loss groups in a low socioeconomic status area: Application of the COM-B model and Behaviour Change Wheel. *Health Expect.* 2022;25(5):2043-2055. doi:10.1111/hex.13325
  41. Elfhag K, Rössner S. Who succeeds in maintaining weight loss? A conceptual review of factors associated with weight loss maintenance and weight regain. *Obes Rev.* 2005;6(1):67-85. doi:10.1111/j.1467-789X.2005.00170.x
  42. Rogers M, Lemstra M. The importance of community consultation and social support in adhering to an obesity reduction program: results from the Healthy Weights Initiative. *Patient Prefer Adherence.* Published online October 2015:1473. doi:10.2147/PPA.S91912
  43. Austin JL, Smith JE, Gianini L, Campos-Melady M. Attitudinal familism predicts weight management adherence in Mexican–American women. *J Behav Med.* 2013;36(3):259-269. doi:10.1007/s10865-012-9420-6
  44. Ahern AL, Boyland EJ, Jebb SA, Cohn SR. Participants' Explanatory Model of Being Overweight and Their Experiences of 2 Weight Loss Interventions. *Ann Fam Med.* 2013;11(3):251-257. doi:10.1370/afm.1446
  45. Noble S. English Indices of Deprivation 2019: research report.
  46. Szypula J, Jarvstad A, Jones L, Tapper K. Developing cognitive-behavioural phenotypes for weight management. Published online 2025. <https://osf.io/9nfx3>
  47. Tapper K, Szypula J, Jarvstad A. Testing the effectiveness of receiving phenotype-tailored content on engagement and weight loss. Published online 2024. doi:10.17605/OSF.IO/UGWXB
  48. Burke L, Acharya, Elci, et al. Adherence to a behavioral weight loss treatment program

- enhances weight loss and improvements in biomarkers. *Patient Prefer Adherence*. Published online June 2009:151. doi:10.2147/PPA.S5802
49. Lin PH, Grambow S, Intille S, et al. The Association Between Engagement and Weight Loss Through Personal Coaching and Cell Phone Interventions in Young Adults: Randomized Controlled Trial. *JMIR MHealth UHealth*. 2018;6(10):e10471. doi:10.2196/10471
  50. Power JM, Phelan S, Hatley K, et al. Engagement and Weight Loss in a Web and Mobile Program for Low-Income Postpartum Women: Fit Moms/ *Mamás Activas*. *Health Educ Behav*. 2019;46(2\_suppl):114S-123S. doi:10.1177/1090198119873915
  51. Briazu RA, Bell L, Dodd GF, et al. The effectiveness of personalised food choice advice tailored to an individual's socio-demographic, cognitive characteristics, and sensory preferences. *Appetite*. 2024;201:107600. doi:10.1016/j.appet.2024.107600
  52. Yarkoni T, Westfall J. Choosing Prediction Over Explanation in Psychology: Lessons From Machine Learning. *Perspect Psychol Sci*. 2017;12(6):1100-1122. doi:10.1177/1745691617693393
  53. Dao D, Teo JYC, Wang W, Nguyen HD. LLM-Powered Multimodal AI Conversations for Diabetes Prevention. In: *Proceedings of the 1st ACM Workshop on AI-Powered Q&A Systems for Multimedia*. ACM; 2024:1-6. doi:10.1145/3643479.3662049
  54. Gencer G, Gencer K. Large Language Models in Healthcare: A Bibliometric Analysis and Examination of Research Trends. *J Multidiscip Healthc*. 2025;Volume 18:223-238. doi:10.2147/JMDH.S502351
  55. Lewis P, Perez E, Piktus A, et al. Retrieval-Augmented Generation for Knowledge-Intensive NLP Tasks. In: Larochelle H, Ranzato M, Hadsell R, Balcan MF, Lin H, eds. *Advances in Neural Information Processing Systems*. Vol 33. Curran Associates, Inc.; 2020:9459-9474. [https://proceedings.neurips.cc/paper\\_files/paper/2020/file/6b493230205f780e1bc26945df7481e5-Paper.pdf](https://proceedings.neurips.cc/paper_files/paper/2020/file/6b493230205f780e1bc26945df7481e5-Paper.pdf)
  56. Kim DW, Park JS, Sharma K, et al. Qualitative evaluation of artificial intelligence-generated weight management diet plans. *Front Nutr*. 2024;11:1374834. doi:10.3389/fnut.2024.1374834
  57. Morgan-Bathke M, Raynor HA, Baxter SD, et al. Medical Nutrition Therapy Interventions Provided by Dietitians for Adult Overweight and Obesity Management: An Academy of Nutrition and Dietetics Evidence-Based Practice Guideline. *J Acad Nutr Diet*. 2023;123(3):520-545.e10. doi:10.1016/j.jand.2022.11.014
  58. Bush CL, Blumberg JB, El-Sohemy A, et al. Toward the Definition of Personalized Nutrition: A Proposal by The American Nutrition Association. *J Am Coll Nutr*. 2020;39(1):5-15. doi:10.1080/07315724.2019.1685332
  59. Zhang K, Meng X, Yan X, et al. Revolutionizing Health Care: The Transformative Impact of Large Language Models in Medicine. *J Med Internet Res*. 2025;27:e59069. doi:10.2196/59069

## Abbreviations

**DEBQ:** Dutch Eating Behavior Questionnaire

**GLP-1s:** Glucagon-like peptide-1 receptor agonists

**IMD:** Index of multiple deprivation

**LLMs:** Large language models


**NHS:** National Health Service

**TFEQ:** Three Factor Eating Questionnaire


**6-FQ:** 6 Factor Questionnaire

Preprint  
JMIR Publications


## Supplementary Files




## Figures




Flow of participants in a quasi-experimental study evaluating the effect of receiving phenotype-matched weight loss advice on app engagement and weight loss among patients actively participating in a UK weight loss program, recruited in May 2024.




Flow of participants in the phenotype group from starting the phenotype quiz to receiving a final profile match, which determined the tailored weight loss advice they received in the trial.



Mean weekly app engagements in the phenotype group (n=148) and the two comparison groups (historical cohort n=241; non-responders cohort n=394). Error bars denote 95% IC.



## Multimedia Appendixes



A completed TREND checklist.

URL: <http://asset.jmir.pub/assets/615c41e238f2bb51bfe5047ffc9f2d35.docx>

All items in the phenotypes questionnaire, and additional app engagement analyses.

URL: <http://asset.jmir.pub/assets/620934de135298c3c6c5c075d886a65a.doc>

