



City Research Online

City St George's, University of London

Citation: Meades, R., Dudeney, E., Sinesi, A., Cheyne, H., Maxwell, M., Best, C., McNicol, S., Shakespeare, J., Walker, J., Alderdice, F., et al (2026). Prevalence and risk factors for suicidal intent in pregnancy and after birth. *BJPsych International*, pp. 1-10. doi: 10.1192/bji.2026.10117

This is the published version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/38160/>

Link to published version: <https://doi.org/10.1192/bji.2026.10117>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

Prevalence and risk factors for suicidal intent in pregnancy and after birth

Rose Meades,¹ Elizabeth Dudeney,²  Andrea Sinesi,³  Helen Cheyne,⁴ Margaret Maxwell,⁵ Catherine Best,⁶ Stacey McNicol,³ Judy Shakespeare,⁷ James Walker,⁸ Fiona Alderdice,⁹ Julie Jomeen,¹⁰ Louise R. Williams,¹¹ Susan Ayers¹²  and the MAP Study Team[†]

¹Senior Research Fellow, Centre for Maternal and Child Health Research, City St George's, School of Health and Medical Sciences, City St George's University of London, UK

²Research Fellow, Centre for Maternal and Child Health Research, City St George's, School of Health and Medical Sciences, City St George's University of London, UK

³Research Fellow, Centre for Healthcare and Community Research, University of Stirling, UK

⁴Professor of Maternal and Child Health Research, Centre for Healthcare and Community Research, University of Stirling, UK

⁵Professor of Health Services and Mental Health Research, Centre for Healthcare and Community Research, University of Stirling, UK

⁶Associate Professor, Centre for Healthcare and Community Research, University of Stirling, UK

⁷Independent Scholar, Oxford, UK

⁸Professor of Obstetrics and Gynaecology, Faculty of Medicine and Health, St James University Hospital, University of Leeds, UK

⁹Senior Social Scientist and Co-Director NIHR Policy Research Unit Maternal and Neonatal Health and Care, National Perinatal Epidemiology Unit, University of Oxford, UK

¹⁰Professor of Midwifery, Faculty of Health, Southern Cross University, Australia

¹¹Research Fellow, School of Health and Medical Sciences, City St George's University of London, UK

¹²Professor of Maternal and Child Health, School of Health and Medical Sciences, City St George's University of London, UK. Email: susan.ayers.1@city.ac.uk

[†]For members of the MAP Study Team see Acknowledgements.

Keywords. Pregnancy; suicide; postpartum; evidence-based mental health; maternal mortality.

First received 7 Oct 2025 UTC
Final revision 6 May 2026 UTC
Accepted 28 May 2026 UTC

doi:[10.1192/bjpi.2026.10117](https://doi.org/10.1192/bjpi.2026.10117)

© The Author(s), 2026. Published by Cambridge University Press on behalf of Royal College of Psychiatrists. This is an Open

Background Suicide is one of the leading causes of maternal mortality in high-income countries. Suicide prevention policy calls for maternity care providers to actively explore potential risk factors and self-reported suicidal intent with perinatal women. However, longitudinal research establishing the prevalence of suicidal intent and associated risk factors with representative cohorts of perinatal women in the UK is lacking.

Aims This secondary analysis aimed to determine the prevalence of suicidal intent in pregnancy and postpartum, and examine factors associated with suicidal intent.

Method A total of 2210 women completed self-report assessments in early, mid- and late pregnancy and early postpartum. Suicidal intent was measured with the item 'I have made plans to end my life' (Clinical Outcomes in Routine Evaluation-10). Other variables included anxiety (Generalised Anxiety Disorder-7, Stirling Antenatal Anxiety Scale), depression (Whooley Questions), health status (EuroQoL EQ-5D-5L), mental health history, and sociodemographic and pregnancy variables.

Results Suicidal intent was indicated by 72 participants (3.26%, 95% CI 2.56–4.09) on at least one occasion. Suicidal intent was significantly associated with having negative feelings about pregnancy (odds ratio 3.63, 95% CI 1.99–6.63), unplanned pregnancy (odds ratio 2.03, 95% CI 1.25–3.30), previous and current mental health problems (odds ratio 3.70 95% CI 2.19–6.25), symptoms causing daily difficulties (odds ratio 5.94 95% CI 3.32–10.66) and being of Asian (odds ratio 2.62, 95% CI 1.44–4.77) or mixed ethnicity (odds ratio 3.43, 95% CI 1.49–7.93). Poor physical health in early pregnancy was also associated with suicidal intent ($t = 6.37$, d.f. = 2108, $p < 0.001$).

Conclusions These findings indicate that daily difficulties associated with mental health, and negative feelings about pregnancy are important and underexplored risk factors for experiencing suicidal intent. The findings also corroborate previous research indicating that previous and existing mental health problems and ethnicity are significant risk factors.

Suicide has been one of the leading direct causes of maternal death in the UK for over a decade,¹ and remains among the leading causes of maternal mortality in other high-income countries.² For example, between 2021 and 2023, a third (34%) of maternal deaths in the UK occurring between 6 weeks and 1 year after birth were attributable to mental health causes, with suicide the leading direct cause in this period.³ Globally, prevalence varies: suicide accounts for 5–20% of maternal deaths in high-income countries^{4–6} and 1–5% in low- and middle-income countries,⁷ although differences in reporting systems and classifications may underestimate true rates.^{5,8,9} Estimates of suicidal ideation in pregnancy and after birth are harder to establish because of variation in samples, designs and measures, but a recent review across 23 countries reported a pooled prevalence of 8%.¹⁰

Identifying maternal suicidality

Identifying suicidal ideation and behaviours could lead to preventative intervention and reductions in suicide. However, identifying suicidality is difficult. Suicide is a complex phenomenon that encompasses a spectrum of thoughts and behaviours, including suicidal ideation (thinking about ending one's life), suicidal planning or intent (formulating a method or plan) and suicide attempts (engaging in potentially self-lethal behaviour). Universal screening for common perinatal mental health problems is not currently recommended in the UK,¹¹ and stigma, shame, guilt and fear can inhibit disclosure.^{12,13} As a result, many women requiring specialist care go unidentified. Risk is shaped by multiple factors, including psychiatric history,¹⁴ comorbidity,⁵ substance misuse,¹⁵ intimate partner violence,¹⁶ adverse childhood experiences,¹⁷ pregnancy or infant loss,¹⁸ unplanned or unwanted pregnancy,¹⁹ sleep disturbance²⁰ and health complications.²¹ Sociodemographic risks include younger age, non-White ethnicity, deprivation, being unmarried²² and lack of social and partner support.²³ Nonetheless, meta-analytic evidence suggests that research on specific risk factors for perinatal suicide risk is lacking.²⁴

Although there is evidence that pregnancy and having a baby can protect against suicide,²⁵ the extent of these protective effect may be inflated,⁸ and the postnatal period carries higher risk of suicide deaths.³

Inequalities in maternal suicidality

Women from ethnic minority groups are disproportionately affected, facing barriers and inequalities in maternity mental health care that increase risks of untreated mental health problems and suicidality.^{26,27} Targeted, evidence-based approaches to identification, intervention and prevention are therefore needed. This priority is recognised in the UK's cross-government suicide prevention plan, which calls on maternity services to explore suicide risk factors at every contact.²⁸

Methodological considerations

Most studies on prevalence, correlates and risk factors for perinatal suicidality use cross-sectional designs,^{10,23,29} which limit inferences about directionality or temporal associations. Longitudinal approaches are needed to establish patterns, identify when risk is greatest and explore factors over time. This study therefore aimed to (a) determine the prevalence of suicidal intent in pregnancy and postpartum; and (b) examine associated risk factors for suicidal intent during pregnancy and the early postnatal periods – an understudied outcome. Improved understanding will help identify women at risk and inform prevention and intervention.

Method

Study design and population

Secondary analysis of the Methods of Assessing Perinatal Anxiety (MAP) longitudinal cohort of 2243 women recruited in the first trimester of pregnancy from November 2020 to November 2021 (research registry reference 5980).³⁰ Participants in the MAP study completed mental health assessment questionnaires at four time points: early pregnancy (mean gestation 11.4 weeks, s.d. 2.0, range 5–16), mid-pregnancy (mean gestation 23.0, s.d. 1.3, range 21–27), late pregnancy (mean gestation 31.9, s.d. 1.2, range 30–35) and postpartum (mean 7.9 weeks, s.d. 2.4, range 4–17). Women were eligible for the MAP study if they were aged 16 years or over, <15 weeks pregnant at the time of recruitment, able to provide written informed consent and had sufficient English to understand and complete questionnaires. A 10:1 ratio of participants from England and Scotland were recruited to reflect the number of annual births for the two nations.

Secondary analyses were based on 2210 women who completed the measure of suicidal intent on at least one time point. Participants were aged 16–50 years (mean 31.34, s.d. 5.18). Most were married or cohabiting ($n = 1749$, 87.84%), of White ethnicity ($n = 1548$, 77.25%) and educated to undergraduate degree level or above ($n = 1235$, 61.62%). Around a third ($n = 741$, 36.34%) reported a history of mental health problems (see Supplementary Table 1). For analysis of risk factors, participants were grouped into those who

expressed suicidal intent at any time point ($n = 72$) and those who did not ($n = 2138$).

Measures

Suicidal intent and general distress were measured with the Clinical Outcomes in Routine Evaluation (CORE-10).³¹ The CORE-10 is a ten-item measure of psychological distress derived from the larger CORE – Outcome Measure. It asks respondents about how they have been feeling over the past week. The ten items are scored on a 0–4 Likert scale. Suggested cut-off scores are as follows: 5–10 indicate low-level problems, 10–15 indicates mild psychological distress, 15–20 indicates moderate distress, 20–25 indicates moderately severe distress and 25–40 indicates severe psychological distress.³¹

Suicidal intent was measured by the item on the CORE-10 'I have made plans to end my life', with response options of 'not at all', 'only occasionally', 'sometimes', 'often' or 'most or all of the time'. Suicidal intent was coded as present for participants who made any response other than 'not at all'.

The CORE-10 has good psychometric properties with perinatal women and performs better than measures of anxiety (Generalised Anxiety Disorder-2) and depression (Whooley Questions) at identifying women worried about their psychological health.³⁰ The CORE-10 is recommended by the UK Royal College of Psychiatrists as a core outcome measure for perinatal services.³²

Anxiety was assessed with two measures. The Generalised Anxiety Disorder-7 (GAD-7)³³ consists of seven self-report items used to identify probable cases of generalised anxiety disorder. Items are scored on a 0–3 scale and higher scores reflect greater anxiety severity. Total scores range from 0 to 21, with a recommended cut-off score of 10.³⁰ The Stirling Antenatal Anxiety Scale (SAAS)³⁴ is a ten-item, clinically derived measure developed specifically for anxiety in pregnancy and subsequently validated for use postpartum. The SAAS includes general anxiety and pregnancy-specific anxiety items. The pregnancy-specific items are about the birth and baby, meaning the scale can be used postnatally. Items are scored on a 0–4 Likert scale, with a cut-off score of 9 and above giving optimal sensitivity and specificity in diagnosing anxiety disorders. The scale has good diagnostic accuracy and there is some evidence it performs better than the GAD-7 at identifying women with perinatal anxiety.³⁰

Depression was assessed with the Whooley Questions,³⁵ two yes/no questions widely used in UK maternity services.³⁶ Answering 'yes' to one or both of the questions indicates possible depression. The GAD-7, CORE-10 and Whooley Questions are included in UK clinical guidance on perinatal outcome measures and care pathways for perinatal mental health.^{36,37,38}

Difficulty with daily life was assessed by one item asking how difficult psychological symptoms made

it for participants to 'do your work, take care of things at home, or get along with other people?'. Answers were scored on a four-point scale (not at all difficult, somewhat difficult, very difficult, extremely difficult).

Participants were also asked in early pregnancy whether they had ever experienced psychological or mental health problems (yes/no/don't know) and, if so, to indicate which mental health problem(s) from a list of anxiety or anxiety related problems, depression, bipolar disorder, post-traumatic stress disorder, borderline/personality disorder, psychosis, dissociation/dissociative disorders, eating disorders, sleep or wake disorders, attention-deficit hyperactivity disorder, obsessive-compulsive disorder, schizophrenia and other.

Sociodemographic characteristics were measured using standard classifications for England³⁹ or Scotland,⁴⁰ and included age, ethnicity, education and relationship status.

General health was measured with the Euro-QoL EQ-5D-5L.⁴¹ The EQ-5D-5L is a preference-based measure of health status and quality of life widely used in health research. It includes a visual analogue scale asking participants to assess their overall health that day, on a scale from 0 to 100. In addition to the EQ-5D-5L, participants were asked at baseline to indicate whether they had existing health conditions (high blood pressure, stroke, diabetes, angina/heart attack, anxiety/depression, arthritis, back problems, thyroid problem, eczema/psoriasis, liver disease, kidney disease, asthma, chronic bronchitis, emphysema, cancer, irritable bowel syndrome, migraine, other).

Pregnancy experiences were measured with individual items which asked women whether their pregnancy was planned (yes/no), how they felt about becoming pregnant (five-point scale from very unhappy to very happy), whether women had a previous pregnancy (yes/no) how many children women had before the current pregnancy and whether they had previously experienced a miscarriage or stillbirth (yes/no/prefer not to say).

Procedure

All participants were recruited by research midwives/nurses or clinical midwives around the time of their antenatal booking appointment or pregnancy scan. Information about the study was given at this point, and participants could either provide informed consent and be given the first questionnaire at that point or consent for their contact details to be shared with the research team who would follow up with them, provide more information, answer questions and gain consent. Questionnaire measures were sent and completed online or by post, according to participants' preferences. The order of presentation of the assessment measures was counterbalanced to minimise risk of bias in response patterns.

Safeguarding protocols were in place for participants who reported suicidal intent. Participants who scored any response other than 'not at all' on the suicide item on the CORE-10 were contacted by telephone within 24 h of receiving their questionnaire and a suicide risk assessment was conducted. If participants did not answer they were messaged by email or text to explain the research team were trying to contact them. Three contact attempts were made and risk assessments discussed with the clinical leads for the MAP programme. Action was taken if appropriate (e.g. referral or contact with participant's local specialist mental health midwife or general practitioner).

Regardless of a participant's responses to the item on suicide, if they scored above clinical thresholds for anxiety and/or depression, they were contacted by post or email to encourage them to see a health professional and provide information about options such as helplines and self-referral to mental health services.

Ethical approval

The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2013. All procedures involving human patients were approved by the National Health Service West of Scotland Research Ethics Service (WoSREC 3. reference 20/WS/0065), Health Research Authority (reference IRAS 274901) and City St Georges, University of London (reference ETH1920-0572).

Statistical analysis

As women completed the questions at four time points, their responses were analysed in two ways: by observation and by participant.

Sociodemographic characteristics that were stable over time and/or were only assessed at baseline (highest level of education, age, ethnicity, previous experience of mental health problems, previous treatment, type of treatment, number of children, previous pregnancy loss, whether the pregnancy was planned, previous mental health problems, health conditions) were analysed at the level of participant. Results are presented as number and percentage of participants in each category, and are reported separately for those who ever reported suicidal intent during the study waves versus those who never reported suicidal intent. The association of suicidal intent with categorical independent variables was assessed with Firth logistic regression, which uses penalised maximum likelihood and is suitable for rare outcomes.⁴² For continuous independent variables, the association was assessed through independent samples *t*-tests.

For independent variables that were captured as repeated measures over multiple time points,

these were analysed as observations nested within participants. The association of mental health continuous independent variables (scores on the CORE-10, GAD-7, Whooley Questions and SAAS) with suicidal intent at the same time point was examined with Bayesian mixed-effects models. Bayesian mixed-effects models were employed because of the sparseness of the outcome variable. A weakly informative prior was used (time point and independent variables coefficients distributed normal (0,2.5), and constant distributed normal (0,10)). The outcome was coded as a binary variable of any suicidal intent at that time point versus none. A random effect of participant was included in the model to account for correlation of repeated measures on individuals, and time point was included as a covariate. The mental health continuous independent variables (scores on the CORE-10, GAD-7, Whooley Questions and SAAS) were converted to standard scores (mean 0, s.d. 1) to enable direct comparison of the odds ratios. Odds ratios are reported with their 95% confidence interval. Because of the low prevalence of suicidal intent, it was not possible to fit multivariable models including multiple predictors simultaneously. Therefore, each independent variable was examined in a separate model, with longitudinal analyses adjusted for time point and

clustering within individuals. Analysis was conducted with Stata version 16 or above for Windows (StataCorp LLC, College Station, TX, USA; <https://www.stata.com>).

Results

Prevalence of suicidal intent in pregnancy and after birth

The number of participants who reported suicidal intent is shown in Fig. 1, with the corresponding prevalence. In this sample, 72 participants (3.26%, 95% CI 2.56–4.09) reported some level of suicidal intent on at least one time point during pregnancy or after birth. Of these 72 participants, nearly all ($n = 69$) only reported suicidal intent at one time point, two reported suicidal intent at two time points and one participant reported suicidal intent at three time points. The prevalence of suicidal intent ranged from 1.65% (95% CI 1.16–2.28) in early pregnancy to 0.64% (95% CI 0.30–1.22) in late pregnancy (see Table 1). Women were more likely to report suicidal intent in early pregnancy compared with the late pregnancy time point (odds ratio 2.59, 95% CI 1.22–5.51). There were no other significant differences between time points in the probability of reporting suicidal intent.




Fig. 1

Number of participants and proportion of sample reporting suicidal intent.

Table 1

Prevalence of suicidal intent at different time points

	Early pregnancy	Mid-pregnancy	Late pregnancy	Postpartum
I have made plans to end my life	<i>n</i> (%) 95% CI	<i>n</i> (%) 95% CI	<i>n</i> (%) 95% CI	<i>n</i> (%) 95% CI
Not at all	2141 (98.35) 97.72–98.81	1469 (98.99) 98.33–99.39	1387 (99.36) 98.76–99.66	1345 (98.82) 98.09–99.28
Only occasionally	27 (1.24) 0.85–1.80	13 (0.88) 0.51–1.50	1 (0.07) 0.01–0.51	8 (0.59) 0.29–1.17
Sometimes	8 (0.37) 0.18–0.73	2 (0.13) 0.03–0.54	5 (0.36) 0.15–0.86	7 (0.51) 0.25–1.08
Often	0 (0)	0 (0)	2 (0.14) 0.04–0.57	1 (0.07) 0.01–0.52
Most or all of the time	1 (0.05) 0–0.33	0 (0)	1 (0.07) 0.01–0.51	0 (0)
Total suicidal intent	36 (1.65) 1.16–2.28	15 (1.01) 0.57–1.66	9 (0.644) 0.30–1.22	16 (1.18) 0.67–1.90
Total number	2177 (100)	1484 (100)	1396 (100)	1361 (100)

Factors associated with suicidal intent

Associations between suicidal intent and socio-demographic, pregnancy and health risk factors are shown in [Table 2](#).

Sociodemographic factors

Suicidal intent at any time point was significantly associated with being younger (mean age: no suicidal intent 31.40, s.d. 5.15 years; suicidal intent mean 29.54, s.d. 5.55 years; $t = 2.91$, d.f. = 2116, $p = 0.004$) and of Asian/British Asian ethnicity (see [Table 2](#)). Although most participants reporting suicidal intent were of White ethnicity ($n = 38$, 59.38%, 95% CI 46.78–70.85), Asian and Asian British participants comprised 12.89% (95% CI 11.48–14.42) of the sample, but 25% ($n = 16$, 95% CI 15.77–37.24) of those reporting suicidal intent. The odds ratio for suicidal intent in women from Asian backgrounds was 2.62 (95% CI 1.44–4.77) compared with participants from White ethnicity. The odds ratio was 3.43 (95% CI 1.49–7.93) for women from a mixed ethnic background compared with women from a White background. There was no significant association between suicidal intent and level of education or relationship status.

Pregnancy factors

Suicidal intent was significantly associated with unplanned pregnancy, negative feelings about pregnancy and having had a previous pregnancy (see [Table 2](#)). Unplanned pregnancy was reported by 23.81% ($n = 520$, 95% CI 22.07–25.64%) of the sample, but 38.03% ($n = 27$, 95% CI 27.37–49.98%) of participants who reported suicidal intent. Women who reported they were unhappy or ambivalent ('not happy or unhappy') about their pregnancy were more likely to report suicidal intent at some point in the perinatal period (odds ratio 3.63, 95% CI 1.99–6.63). Women who had a previous pregnancy were more

likely to report suicidal intent (odds ratio 1.94, 95% CI 1.12–3.35). However, there was no significant relationship between suicidal intent and previous pregnancy loss, having no children versus one or more children, or having two or less children versus three or more children.

Mental and physical health

Suicidal intent was also associated with previous mental health problems, which were reported by 67.19% of those reporting suicidal intent compared with 35.34% of the sample overall (see [Table 2](#)). Analysis of different types of previous mental health problems found nearly all were associated with increased risk of suicidal intent (anxiety, attention-deficit hyperactivity disorder, personality disorder, depression, obsessive-compulsive disorder, post-traumatic stress disorder, psychosis, sleep disorders, dissociative disorders). The exceptions were bipolar disorder, eating disorders and schizophrenia, but only a small number of participants reported these disorders, so our results should be interpreted with caution (see Supplementary Table 2).

[Table 3](#) shows the association between suicidal intent and concurrent or previously reported psychological symptoms of anxiety, depression and distress. Suicidal intent was significantly associated with current psychological symptoms on all measures. The odds of reporting suicidal intent increased by three times for every additional one standard deviation increase in a score on the CORE-10 or SAAS. Similarly, the increase in odds of reporting suicidal intent more than doubled for every standard deviation increase in score on the GAD-7 and the Whooley Questions. Mental health symptoms reported on questionnaires at the preceding time point were also significantly associated with suicidal intent. If participants rated their current mental health symptoms as making daily life difficult, the odds of them reporting suicidal intent were almost six times higher than

Table 2

Associations between suicidal intent and sociodemographic, pregnancy and health-related risk factors

	No suicidal intent	Suicidal intent	Odds ratio
	<i>n</i> (%)	<i>n</i> (%)	(95% CI)
Ethnicity	1940 (100)	64 (100)	
White	1510 (77.84)	38 (59.38)	Reference
Mixed/multiple ethnic groups	82 (4.18)	7 (10.94)	3.61 (1.60–8.14)*
Asian/Asian British	243 (12.53)	16 (25.00)	2.66 (1.47–4.81)*
Black/African/Caribbean/ Black British	85 (4.38)	3 (4.69)	1.61 (0.53–4.90)
Other ethnic group	21 (1.08)	0 (0)	0.91 (0.05–15.34)
Level of education	1940 (100)	64 (100)	
Less than a degree	744 (38.35)	25 (39.06)	Reference
Degree level or higher	1196 (61.65)	39 (60.94)	0.96 (0.58–1.60)
Relationship status	1927 (100)	64 (100)	
In a relationship, non- cohabiting	157 (8.15)	6 (9.38)	Reference
Cohabiting	662 (34.35)	19 (29.69)	0.71 (0.29–1.76)
Married/civil partnership	1033 (53.61)	35 (54.69)	0.83 (0.35–1.95)
Separated/divorced/single	75 (3.89)	4 (6.25)	1.44 (0.42–4.96)
Planned pregnancy	2102 (100)	71 (100)	
Yes	1612 (76.69)	44 (61.97)	Reference
No	490 (23.31)	27 (38.03)	2.03 (1.25–3.30)*
Had a previous pregnancy	2103 (100)	71 (100)	
No	807 (38.37)	17 (23.94)	Reference
Yes	1296 (61.63)	54 (76.06)	1.94 (1.12–3.35)*
Feelings about pregnancy	2103(100)	71 (100)	
Other response	11 (0.52)	1 (1.41)	Reference
Very unhappy/unhappy	8 (0.38)	2 (2.82)	3.63 (1.99–6.63)*
Had children already	2117 (100)	71 (100)	
No children	1058 (49.98)	35 (49.30)	Reference
One or more children	1059 (50.02)	36 (50.70)	1.02 (0.64–1.63)
Number of children	2106 (100)	71 (100)	
Two or fewer children	1962 (93.16)	65 (91.55)	Reference
More than two children	144 (6.84)	6 (8.45)	1.27 (0.59–2.75)
Previous pregnancy loss ^a	2080 (100)	69 (100)	
No	1473 (70.82)	44 (63.77)	Reference
Yes	607 (29.18)	25 (36.23)	1.39 (0.85–2.28)
History of mental health problems ^b	1975 (100)	64 (100)	
Yes	698 (35.34)	43 (67.19)	3.70 (2.19–6.25)*
No	1277 (64.66)	21 (32.81)	Reference
Pre-existing physical health condition	2041 (100)	66 (100)	
Yes	565 (27.68)	23 (34.85)	1.41 (0.85–2.35)
No	1476 (72.32)	43 (65.15)	Reference
How difficult (mental health symptoms) make daily life	2049 (100)	69 (100)	
Not at all	1234 (60.22)	13 (18.84)	Reference
Somewhat/very/extremely difficult	815 (39.78)	56 (81.16)	5.94 (3.32–10.66)*

a. Data for participants who responded 'prefer not to say' are not reported in the table and were excluded from this analysis.

b. Data for participants who responded 'don't know' were excluded from this analysis.

* $p < 0.05$.

respondents who did not (odds ratio 5.94, 95% CI 3.32–10.66) (Table 2).

Suicidal intent was also significantly associated with poor physical health in early pregnancy

($t = 6.37$, d.f. = 2108, $p < 0.001$). The mean self-rated health score on the visual analogue scale of the EQ-5D-5L at the early pregnancy time point was 80.05 (s.d. 16.58) in participants who never

Table 3

Odds of reporting suicidal intent by concurrent and previous psychological symptoms

	Concurrent ^a psychological symptoms and suicidal intent, odds ratio (95% credible interval)	Previous ^b psychological symptoms and suicidal intent, odds ratio (95% credible interval)
General distress (CORE-10)	3.67 (3.10–4.42)*	2.24 (1.76–2.79)*
Anxiety (GAD-7)	2.47 (2.10–2.93)*	2.30 (1.79–2.90)*
Anxiety (SAAS)	3.19 (2.65–3.88)*	2.56 (1.93–3.37)*
Depression (Whooley Questions)	2.62 (2.06–3.30)*	2.29 (1.70–3.08)*

CORE-10, Clinical Outcomes in Routine Evaluation; GAD-7, Generalised Anxiety Disorder-7; SAAS, Stirling Antenatal Anxiety Scale.

a. Concurrent psychological symptoms are scores on each of the listed measures at the same time point as suicidal intent.

b. Previous psychological symptoms are scores on the measures from the lagged time point relative to suicidal intent.

* $p < 0.05$.

experienced suicidal intent, compared with 66.84 (s.d. 21.18) in participants who had experienced suicidal intent. Mixed-effects logistic regression indicated that across all time points, better physical health scores were associated with lower odds of suicidal intent (odds ratio 0.96, 95% CI 0.95–0.97). There was no significant relationship between suicidal intent and having specific pre-existing physical health conditions such as arthritis, asthma, cancer, diabetes or other health conditions. However, there were relatively few women with pre-existing health conditions, so the lack of association may be attributable to the small sample size in the group with suicidal intent.

Discussion

This study aimed to determine the prevalence of suicidal intent in pregnancy and the early post-natal period, and examine factors associated with suicidal intent in a large UK sample. Results showed that suicidal intent was most strongly associated with experiencing difficulties with daily life, previous mental health problems and being unhappy about being pregnant. Suicidal intent was also associated with being younger, of mixed or Asian ethnicity, having poor current mental health, poor current physical health, unplanned pregnancy and previous pregnancies. Women were most likely to report suicidal intent in early pregnancy. This is an important finding because it suggests that although most suicides in the perinatal period happen after birth, assessment or discussion of suicidal ideation, behaviours and/or intent with perinatal women needs to be considered from the beginning of pregnancy.³ The observed associations between suicidal intent and both negative feelings about the pregnancy and unplanned pregnancy may reflect experiences of entrapment, a construct implicated in contemporary models of suicidality in which individuals perceive themselves as unable to escape an adverse or unwanted situation.^{43,44} Because these are pregnancy-specific factors routinely assessed in antenatal care, maternity practitioners are well placed to identify potential risk and act early, representing an important opportunity for prevention.

Ethnicity and age were the only sociodemographic variables with a significant relationship to suicidal intent in this study. In line with previous research, being younger and from a mixed ethnic background was associated with suicidal intent.^{22,45} There is less previous evidence that women from Asian backgrounds have an increased risk of suicide,⁴⁵ but, overall, evidence shows that Asian and Black women are two to three times more likely than White women to die from any cause during pregnancy.³ Research also shows that perinatal women from ethnic minority groups experience more inequality and barriers in maternity care services than White women, which can increase their risk of poor mental health and/or suicidality and affect their access to perinatal mental healthcare.^{26,27} These findings reflect the need for targeted approaches that consider ethnicity, religious and cultural beliefs, stigma, and attitudes and beliefs in mental health for identifying and supporting perinatal women from different ethnic backgrounds who may be experiencing suicidality.

An important finding from this study is the indication that mental and physical health have a relationship to suicidal intent. A higher proportion of women who reported suicidal intent had previous and current mental health problems, as well as reporting existing health conditions in early pregnancy. Having a diagnosis of severe physical illness is associated with suicide risk in the general population,⁴⁶ but research also suggests that the effect of physical ill health in terms of disruption to a person's daily activity is another important predictor for suicide.⁴⁷

In the perinatal literature, there is some evidence that having a chronic medical illness is significantly associated with suicidal behaviour during pregnancy,²¹ and other research has reported that physical pregnancy-related factors, such as severe hyperemesis gravidarum, pregnancy-induced hypertension, gestational diabetes mellitus,⁴⁸ perineal laceration, caesarean delivery⁴⁹ and obesity,⁵⁰ increase the risk of suicidal ideation. Although we did not ask women how their physical health affected daily functioning, our results showed that the perceived impact of mental health difficulties on daily functioning was associated with suicidal intent. Taken together, these findings

underscore the importance of the link between physical and mental health, and suggest that the impact on functioning in the perinatal period may be a possible focus for assessing need for support. Given that universal maternity service appointments largely focus on the mother's physical health, and that midwives are more used to asking about physical health, this provides an opportunity to explore the interface between mothers' physical and mental health and feelings about being pregnant.

Although multivariable modelling was not possible because of the low prevalence of suicidal intent, these findings provide an important foundation for future research in an underexplored area. Larger or pooled data-sets, including linkage to routine health records, are needed to enable adequately powered analyses of independent risk factors and extend these findings to more diverse populations.

Strengths and limitations

This is the first study to assess suicidal intent (rather than suicidal ideation or self-harm) at multiple time points over pregnancy and in the early postpartum period in a representative and ethnically diverse UK sample. The characteristics and prevalence of perinatal women who endorse the item 'I have made plans to end my life' on the CORE-10 have not been explored previously in a representative sample. This item is different to many other self-report items that only ask about suicidal or self-harm ideation. Although the sample was recruited through routine antenatal care and is broadly reflective of UK maternity populations, it was relatively socioeconomically advantaged and required sufficient English to complete questionnaires, which may limit applicability to more diverse and underserved groups. Also, although this is a large study, the low prevalence of suicidal intent in this population means that multivariate analysis was not possible. Nevertheless, this is the first study to have a sufficiently large sample to examine suicidal intent by socio-demographic characteristics in a perinatal population, so it makes an important contribution to this area. A further limitation is that because women received a follow-up telephone call as part of the safeguarding procedure if they reported suicidal intent, this may have affected their response to follow-up questionnaires.

Conclusions

Our findings suggest that difficulties in daily life caused by mental health symptoms, negative feelings about being pregnant and previous mental health problems were significantly associated with suicidal intent. Young age, Asian or mixed ethnicity, poor current mental health, poor current physical health, unplanned pregnancy and previous pregnancy were also risk factors. These findings emphasise the link between physical and mental health, suggesting that assessing

daily functioning in the perinatal period could identify support needs. Given the focus on physical health in maternity appointments, this offers an opportunity to explore the intersection of physical health, mental health and women's feelings toward their pregnancy.

Supplementary material

To view supplementary material for this article, please visit <https://doi.org/10.1192/bji.2026.10117>

Data availability

Individual participant-level data are currently not available, but authors can provide sample-level data and information on request, after publication. The study protocol is available at <https://njl-admi.nihr.ac.uk/document/download/2034506>.

Acknowledgements

We gratefully thank the women who took part in the MAP study for their time and generosity. We are also grateful for the input and advice of colleagues in the MAP Study Team and Steering Committee in conducting this research. The MAP Study Team includes the authors, Georgina Constantinou, Simon Gilbody, Debra Salmon, Agnes Hann, Una Hutton, Jennifer Holly, Nazihah Uddin and Cassandra Yuill.

Author contributions

A.S., C.B., F.A., H.C., J.J., J.S., L.R.W., M.M., R.M., S.A. and J.W. all contributed to the conceptualisation of the study, methodology and oversight. Additionally, R.M., A.S. and L.R.W. were involved in data collection, data validation and project administration. C.B. and S.M. provided statistical expertise and conducted the analyses. The manuscript was drafted by R.M. and E.D. All listed authors and the MAP study team (see Acknowledgements) reviewed and approved the manuscript.

Funding

This work was supported by the National Institute for Health Research (NIHR), Health Services & Delivery Research Programme (NIHR 17/105/16). The views expressed are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care.

Declaration of interest

A.S., H.C. and M.M. developed and published one of the measures used in this study, the Stirling Antenatal Anxiety Scale. All other authors have no conflicts of interest to declare.

References

- 1 Knight M, Nair M, Tuffnell D, Kenyon S, Shakespeare J, Brocklehurst P, et al. *Saving Lives, Improving Mothers' Care: Surveillance of Maternal Deaths in the UK 2012–14 and Lessons Learned to Inform Maternity Care from the UK and Ireland Confidential Enquiries into Maternal Deaths and Morbidity 2009–14*. National Perinatal Epidemiology Unit, 2016 (<https://www.npeu.ox.ac.uk/mbrace-uk/reports/maternal-reports/maternal-report-2012-2014>).
- 2 Diguisto C, Saucedo M, Kallianidis A, Bloemenkamp K, Bodker B, Buoncristiano M, et al. Maternal mortality in eight European countries with enhanced surveillance systems: descriptive population based study. *BMJ* 2022; 379: e070621.
- 3 Felker A, Patel R, Kotnis R, Kenyon S, Knight M. *Saving Lives, Improving Mothers' Care Compiled Report – Lessons Learned to Inform Maternity Care from the UK and Ireland Confidential Enquiries into Maternal Deaths and Morbidity 2021–23*. National Perinatal Epidemiology Unit, University of Oxford, 2025 (<https://www.npeu.ox.ac.uk/mbrace-uk/reports/maternal-reports/maternal-report-2021-2023>).

- 4 Grigoriadis S, Wilton AS, Kurdyak PA, Rhodes AE, Vonder Porten EH, Levitt A, et al. Perinatal suicide in Ontario, Canada: a 15-year population-based study. *Can Med Assoc J* 2017; 189: e1085–92.
- 5 Howard LM, Khalifeh H. Perinatal mental health: a review of progress and challenges. *World Psychiatry* 2020; 19: 313–27.
- 6 Trost SL, Beauregard J, Chandra G, Njie F, Berry J, Harvey A, et al. *Pregnancy-Related Deaths: Data from Maternal Mortality Review Committees in 36 US States, 2017–2019*. Centers for Disease Control and Prevention, US Department of Health and Human Services, 2022 (<https://www.cdc.gov/maternal-mortality/php/data-research/mmrc/index.html> [cited 3 Jul 2026]).
- 7 Fuhr DC, Calvert C, Ronsmans C, Chandra PS, Sikander S, De Silva MJ, et al. Contribution of suicide and injuries to pregnancy-related mortality in low-income and middle-income countries: a systematic review and meta-analysis. *Lancet Psychiatry* 2014; 1: 213–25.
- 8 Chin K, Wendt A, Bennett IM, Bhat A. Suicide and maternal mortality. *Curr Psychiatry Rep* 2022; 24: 239–75.
- 9 Simmons E, Gong J, Daskalopoulou Z, Quigley MA, Alderdice F, Harrison S, et al. Global contribution of suicide to maternal mortality: a systematic review protocol. *BMJ Open* 2024; 14: e087669.
- 10 Xiao M, Hu Y, Huang S, Wang G, Zhao J, Lei J. Prevalence of suicidal ideation in pregnancy and the postpartum: a systematic review and meta-analysis. *J Affect Disord* 2022; 296: 322–36.
- 11 UK National Screening Committee. *Antenatal and Postnatal Mental Health*. UK Government, 2019 (<https://view-health-screening-recommendations.service.gov.uk/antenatal-and-postnatal-mental-health/>).
- 12 Dudeney E, Coates R, Ayers S, McCabe R. Acceptability and content validity of suicidality screening items: a qualitative study with perinatal women. *Front Psychiatry* 2024; 15: 1359076.
- 13 Dudeney E, Meades R, Ayers S, McCabe R. Perinatal women's views and experiences of discussing suicide in maternity care settings: a qualitative study. *Women Birth* 2024; 37: 101662.
- 14 Martini J, Bauer M, Lewitzka U, Voss C, Pfennig A, Ritter D, et al. Predictors and outcomes of suicidal ideation during peripartum period. *J Affect Disorders* 2019; 257: 518–26.
- 15 Modini C, Leske S, Roberts S, Whelan N, Chitakis A, Crompton D, et al. Maternal deaths by suicide in Queensland, Australia, 2004–2017: an analysis of maternal demographic, psychosocial and clinical characteristics. *Arch Womens Ment Health* 2021; 24: 1019–25.
- 16 Alhusen JL, Frohman N, Purcell G. Intimate partner violence and suicidal ideation in pregnant women. *Arch Womens Ment Health* 2015; 18: 573–8.
- 17 Jasthi DL, Nagle-Yang S, Frank S, Masotya M, Huth-Bocks A. Associations between adverse childhood experiences and prenatal mental health and substance use among urban, low income women. *Community Ment Health* 2022; 58: 595–605.
- 18 Knight M, Bunch K, Tuffnell D, Patel R, Shakespeare J, Kotnis R, et al. *Saving Lives, Improving Mothers' Care: Lessons Learned to Inform Maternity Care from the UK and Ireland Confidential Enquiries into Maternal Deaths and Morbidity 2017–19*. National Perinatal Epidemiology Unit, 2021 (<https://www.npeu.ox.ac.uk/mbrace-uk/reports/maternal-reports/maternal-report-2017-2019>).
- 19 Fellmeth G, Nosten S, Khirikoekkong N, Oo MM, Gilder ME, Plugge E, et al. Suicidal ideation in the perinatal period: findings from the Thailand–Myanmar border. *J Public Health (Oxf)* 2021; 44: e514–8.
- 20 Palagini L, Cipriani E, Miniati M, Bramante A, Gemignani A, Geoffroy PA, et al. Insomnia, poor sleep quality and perinatal suicidal risk: a systematic review and meta-analysis. *J Sleep Res* 2023; 33: e14000.
- 21 Molla A, Nigussie J, Girma B. Prevalence and associated factors of suicidal behavior among pregnant mothers in southern Ethiopia: a cross-sectional study. *BMC Public Health* 2022; 22: 490.
- 22 Orsolini L, Valchera A, Vecchiotti R, Tomasetti C, Lasevoli F, Fornaro M, et al. Suicide during perinatal period: epidemiology, risk factors and clinical correlates. *Front Psychiatry* 2016; 7: 1–6.
- 23 Arditi-Arbel B, Hamdan S, Winterman M, Gvion Y. Suicidal ideation and behavior among perinatal women and their association with sleep disturbances, medical conditions, and known risk factors. *Front Psychiatry* 2023; 13: 987673.
- 24 Gelabert E, Plaza A, Roca-Lecumberri A, Bramante A, Brenna V, Garcia-Esteve L, et al. Suicide attempts during pregnancy and postpartum: a systematic review and meta-analysis. *Matern Child Health J* 2024; 28: 1443–53.
- 25 Khalifeh H, Hunt IM, Appleby L, Howard LM. Suicide in perinatal and non-perinatal women in contact with psychiatric services: 15 year findings from a UK national inquiry. *Lancet Psychiatry* 2016; 3: 233–42.
- 26 Conneely M, Packer KC, Bicknell S, Jankovic J, Kaur Sihre H, McCabe R, et al. Exploring Black and South Asian women's experiences of help-seeking and engagement in perinatal mental health services in the UK. *Front Psychiatry* 2023; 14: 1119998.
- 27 Jankovic J, Parsons J, Jovanovic N, Berrisford G, Copello A, Fazil Q, et al. Differences in access and utilisation of mental health services in the perinatal period for women from ethnic minorities – a population-based study. *BMC Med* 2020; 18: 245.
- 28 Department of Health and Social Care. *Suicide Prevention in England: 5-Year Cross-Sector Strategy*. Department of Health and Social Care, 2023 (<https://www.gov.uk/government/publications/suicide-prevention-strategy-for-england-2023-to-2028/suicide-prevention-in-england-5-year-cross-sector-strategy>).
- 29 Reid HE, Pratt D, Edge S, Wittkowski A. Maternal suicide ideation and behaviour during pregnancy and the first postpartum year: a systematic review of psychological and psychosocial risk factors. *Front Psychiatry* 2022; 13: 765118.
- 30 Ayers S, Coates R, Sinesi A, Cheyne H, Maxwell M, Best C, et al. and the MAP study team assessment of perinatal anxiety: diagnostic accuracy of five measures. *Br J Psychiatry* 2024; 25: 1–7.
- 31 Barkham M, Bewick B, Mullin T. The CORE-10: a short measure of psychological distress for routine use in the psychological therapies. *Counsell Psychother Res* 2013; 13: 3–13.
- 32 Royal College of Psychiatrists (RCPsych). *Framework for Routine Outcome Measurement in Perinatal Psychiatry: College Report CR216*. RCPsych, 2018 (https://www.rcpsych.ac.uk/docs/default-source/improving-care/better-mh-policy/college-reports/college-report-cr216.pdf?sfvrsn=12b1e81c_2).
- 33 Spitzer RL, Kroenke K, Williams JB, Lowe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med* 2006; 166: 1092–7.
- 34 Sinesi A, Cheyne H, Maxwell M, O'Carroll R. The Stirling Antenatal Anxiety Scale (SAAS): development and initial psychometric validation. *J Affect Disord* 2022; 8: 100333.
- 35 Whooley MA, Avins AL, Miranda J, Browner WS. Case-finding instruments for depression. Two questions are as good as many. *J Gen Intern Med* 1997; 12: 439–45.
- 36 National Collaborating Centre for Mental Health. *The Perinatal Mental Health Care Pathways*. National Collaborating Centre for Mental Health, 2018 (<https://www.england.nhs.uk/publication/the-perinatal-mental-health-care-pathways/>).
- 37 National Institute for Health and Care Excellence (NICE). *Antenatal and Postnatal Mental Health: Clinical Management and Service Guidance (CG192)*. NICE, 2020 (<https://www.nice.org.uk/guidance/cg192>).

- 38 Scottish Intercollegiate Guidelines Network (SIGN). *Management of Perinatal Mood Disorders (SIGN 127)*. SIGN, 2023 (https://www.sign.ac.uk/assets/sign127_update.pdf).
- 39 Office for National Statistics (ONS). *English and Wales Census*. ONS, 2011 (<https://www.ons.gov.uk/peoplepopulationandcommunity/culturalidentity/ethnicity/articles/ethnicityandnationalidentityinenglandandwales/2012-12-11>).
- 40 National Records of Scotland. *Scotland's Census*. National Records of Scotland, 2011 (<https://www.scotlandscensus.gov.uk/metadata/ethnic-group/#panel-2>).
- 41 Herdman M, Gudex C, Lloyd A, Janssen M, Kind P, Parkin D, et al. Development and preliminary testing of the new five-level version of EQ-5D (EQ-5D-5L). *Qual Life Res* 2011; 20: 1727–36.
- 42 Firth D. Bias reduction of maximum likelihood estimates. *Biometrika* 1993; 80: 27–38.
- 43 Reid HE, Pratt D, Edge D, Wittkowski A. What makes a perinatal woman suicidal? A grounded theory study. *BMC Psychiatry* 2022; 22: 386.
- 44 O'Connor RC, Kirtley OJ. The integrated motivational-volitional model of suicidal behaviour. *Philos Trans R Soc Lond B Biol Sci* 2018; 373: 20170268.
- 45 Office for National Statistics (ONS). *Sociodemographic Inequalities in Suicides in England and Wales 2011 to 2021: A Population Level Analysis Comparing the Risk of Dying by Suicide across Sociodemographic Groups in Adults in England and Wales*. ONS, 2023 (<https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/healthinequalities/bulletins/sociodemographicinequalitiesinsuicidesinenglandandwales/2011to2021>).
- 46 Nafilyan V, Morgan J, Mais D, Sleeman KE, Butt A, Ward I, et al. Risk of suicide after diagnosis of severe physical health conditions: a retrospective cohort study of 47 million people. *Lancet Reg Health Eur* 2022; 25: 100562.
- 47 Onyeka IN, Maguire A, Ross E, O'Reilly D. Does physical ill-health increase the risk of suicide? A census-based follow-up study of over 1 million people. *Epidemiol Psychiatr Sci* 2020; 29: e140.
- 48 Girardi P, Pompili M, Innamorati M, Serafini G, Berrettoni C, Angeletti G, et al. Temperament, postpartum depression, hopelessness, and suicide risk among women soon after delivering. *Women Health* 2011; 51: 511–24.
- 49 Kim JJ, La Porte LM, Saleh MP, Allweiss S, Adams MG, Zhou Y, et al. Suicide risk among perinatal women who report thoughts of self-harm on depression screens. *Obstet Gynecol* 2015; 125: 885–93.
- 50 Kalmbach DA, Cheng P, Ong JC, Ciesla JA, Kingsberg SA, Sangha R, et al. Depression and suicidal ideation in pregnancy: exploring relationships with insomnia, short sleep, and nocturnal rumination. *Sleep Med* 2020; 65: 62–73.