



## City Research Online

### City, University of London Institutional Repository

---

**Citation:** Haberman, S., Denuit, M. and Renshaw, A. E. (2013). Approximations for quantiles of life expectancy and annuity values using the parametric improvement rate approach for modelling and projecting mortality. *European Actuarial Journal*, 3(1), pp. 191-201. doi: 10.1007/s13385-013-0065-9

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

---

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/4051/>

**Link to published version:** <http://dx.doi.org/10.1007/s13385-013-0065-9>

**Copyright:** City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

**Reuse:** Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

---

---

---

City Research Online:

<http://openaccess.city.ac.uk/>

[publications@city.ac.uk](mailto:publications@city.ac.uk)

---

# Approximations for quantiles of life expectancy and annuity values using the parametric improvement rate approach to modelling and projecting mortality

## Abstract

In this paper, we develop accurate approximations for medians of life expectancy and life annuity pure premiums viewed as functions of future mortality trends as predicted by parametric models of the improvement rates in mortality. Numerical illustrations show that the comonotonic approximations perform well in this case, which suggests that they can be used in practice to evaluate the consequences of the uncertainty in future death rates. Prediction intervals based on 5% and 95% quantiles are also considered but appear to be wider compared to simulated ones. This provides the practitioner with a conservative shortcut, thereby avoiding the problem of simulations within simulations in, for instance, Solvency 2 calculations.

*Key words and phrases:* Life annuity, life expectancy, mortality projection, comonotonicity, simulation.

## 1 Introduction

Forecasting mortality in actuarial studies is generally based on extrapolation methods that capture the pattern in historical mortality rates by means of appropriate parametric predictor structures. Foremost among such structures is the Poisson log-bilinear specification proposed by Brouhns, Denuit and Vermunt (2002) and Renshaw and Haberman (2003) in line with the seminal paper by Lee and Carter (1992). Recently, Haberman and Renshaw (2012) have introduced and investigated parametric mortality projection methods based on mortality improvement rates (as opposed to mortality rates). This approach provides an efficient alternative to the direct parametric modelling and projecting of mortality rates.

In this paper, we consider present values of life annuity benefits as functions of the unknown life table applying in the future (as well as life expectancies, corresponding to zero interest rate). Deriving the exact distribution for this random variable requires extensive simulations of numerical evaluations. Therefore, we take the comonotonic approximations proposed by Denuit and Dhaene (2007) and Denuit (2007) in the random walk with drift case and extended to general *ARIMA* models by Denuit, Haberman and Renshaw (2010). Specifically, we adapt this approach to the parametric projection models targeting mortality improvement rates (rather than mortality rates) proposed by Haberman and Renshaw (2012). The approach developed in the present paper helps avoid the requirement to conduct simulations with simulations in, for instance, Solvency 2 reserving calculations. Numerical illustrations show that the comonotonic approximations perform well for medians (and other central quantiles), which suggests that they can be used in practice to evaluate the consequences of the uncertainty in future death rates. Prediction intervals based on 5% and 95% quantiles are also considered but appear to be wider compared to simulated ones. This provides the practitioner with a conservative shortcut avoiding simulations within simulations.

This paper is organised as follows. Section 2 describes the mortality projection method based on parametric improvement rates. The comonotonic approximations are derived in Section 3. Section 4 is devoted to numerical illustrations. The final Section 5 briefly discusses the results.

## 2 Mortality improvement rates

We consider a rectangular data array, partitioned into unit squares of size one year corresponding to ages  $x = x_1, x_2, \dots, x_k$  and periods  $t = t_1, t_2, \dots, t_n$ . Denote  $m_x(t) \equiv m_{x,t}$  the central rate of mortality (or death rate) at age  $x$  in period  $t$ .

Referring to Haberman and Renshaw (2012), under their *Route II* approach we consider the period-based mortality improvement rates (*MIR*) given by

$$Z_{x,t} = 2 \frac{1 - m_{x,t}/m_{x,t-1}}{1 + m_{x,t}/m_{x,t-1}}.$$

In this definition, we consider the ratio of the period one-step mortality improvements to the average of the two adjacent mortality rates. A more natural definition of improvement rate would involve the initial rate in the denominator i.e. the rate at time  $t-1$ . The definition here avoids the phase difference between the numerator and denominator that would otherwise be present and background calculations indicate that it leads to improved modelling results. Following modelling and extrapolation, the *MIR* are converted to mortality rates (*MR*) using the reverse relationship

$$m_{x,t_n+j} = m_{x,t_n+j-1} \frac{(2 - z_{x,t_n+j})}{(2 + z_{x,t_n+j})} = m_{x,t_n+j-1} g(z_{x,t_n+j})$$

where the function  $g$  is defined as

$$g(z) = \frac{2 - z}{2 + z}.$$

Given the nature of  $z$ , which typically take values well within the range  $(-0.5, 0.5)$  as can be seen for Figure 3 in Renshaw and Haberman (2012), we can safely restrict  $g$  to the domain  $z \in (-1, 2)$  where  $g$  is positive and decreasing.

In this paper, we consider

$$Z_{x+j,t_n+i} \equiv Z_{ji} = \beta_{x+j} \kappa_{t_n+i},$$

where  $Z_{ji}$  and  $\kappa_{t_n+i}$  are random variables and the  $\beta_{x+j}$  are considered as known constants. Henceforth, we assume that  $\kappa_{t_n+i}$  obeys some ARIMA time series model (and is therefore multivariate Normal). Then conditional on  $t_0$  we have

$$Z_{ji} \sim N(\mu_{ji}, \sigma_{ji}^2)$$

with

$$\mu_{ji} = \beta_{x+j} E(\kappa_{t_n+i}), \quad \sigma_{ji}^2 = (\beta_{x+j})^2 \text{Var}(\kappa_{t_n+i}).$$

### 3 Comonotonic approximations

In this section, we show that the theoretical arguments which formed the basis of Denuit, Haberman and Renshaw (2010) can be extended to provide approximations for quantiles of life expectancy and annuity predictions under parametric improvement rate modelling as defined in Section 2. There are, however some fundamental differences, as stressed below.

As in Section 2, we decompose the incremental mortality rate changes into

$$Z_{x+j,t_n+i} = \beta_{x+j} \kappa_{t_n+i} \sim N(\mu_{ji}, \sigma_{ji}^2).$$

The next result shows that assuming that the incremental mortality rate changes are perfectly correlated provides a conservative upper bound on future death rates. In this paper, we concentrate on u-type approximations for quantiles as the numerical study performed in Denuit, Haberman and Renshaw (2010) showed that they were more accurate than their l-type counterparts.

Before proceeding with this result, let us recall the definition of some useful stochastic order relations. For more details, we refer the interested reader to Denuit, Dhaene, Goovaerts and Kaas (2005). The increasing convex order, or stop-loss order (denoted as  $\leq_{ICX}$ ) is defined for random variables  $X$  and  $Y$  as follows:  $X \leq_{ICX} Y$  if  $E[h(X)] \leq E[h(Y)]$  for all the non-decreasing convex functions  $h$  for which the expectations exist. In words,  $X \leq_{ICX} Y$  means that  $X$  tends to be “smaller” and “less variable” than  $Y$ . The supermodular order (denoted as  $\leq_{SM}$ ) is defined for random vectors  $(X_1, \dots, X_n)$  and  $(Y_1, \dots, Y_n)$  as  $(X_1, \dots, X_n) \leq_{SM} (Y_1, \dots, Y_n)$  if  $E[h(X_1, \dots, X_n)] \leq E[h(Y_1, \dots, Y_n)]$  for all the supermodular functions  $h$  for which the expectations exist. Recall that a (regular) supermodular function has a non-negative mixed partial derivative with respect to each pair of distinct components. In words,  $(X_1, \dots, X_n) \leq_{SM} (Y_1, \dots, Y_n)$  means that the

components of  $(X_1, \dots, X_n)$  are “less positively dependent” than the components of  $(Y_1, \dots, Y_n)$ .

**Property.** Let  $Z \sim N(0,1)$ . We then have the following upper bound on the death rate at age  $x+j$  in calendar year  $t_n+j$ :

$$m_{x+j,t_n+j} \leq_{ICX} m_{x+j,t_n} \prod_{i=1}^j g(\mu_{ji} + \sigma_{ji}Z).$$

**Proof.** Whatever the dependent structure between the  $\kappa_{t_n+i}$ ,  $i=1,2,3,\dots$ , we have from Proposition 6.3.7 of Denuit, Dhaene, Goovaerts and Kaas (2005) that

$$(Z_{x+j,t_n+1}, Z_{x+j,t_n+2}, \dots, Z_{x+j,t_n+j}) \leq_{SM} (\mu_{j1} + \sigma_{j1}Z, \mu_{j2} + \sigma_{j2}Z, \dots, \mu_{jj} + \sigma_{jj}Z).$$

The more the  $\kappa_{t_n+i}$  are positively related, the closer is the incremental mortality rate random vector to the upper bound in the  $\leq_{SM}$  sense. Now, we get from Property 3.4.61(ii) of Denuit *et al.* (2005) that

$$(g(Z_{x+j,t_n+1}), \dots, g(Z_{x+j,t_n+j})) \leq_{SM} (g(\mu_{j1} + \sigma_{j1}Z), \dots, g(\mu_{jj} + \sigma_{jj}Z))$$

also holds. From Proposition 6.3.9 of Denuit *et al.* (2005), we finally see that

$$\prod_{i=1}^j g(Z_{x+j,t_n+i}) \leq_{ICX} \prod_{i=1}^j g(\mu_{ji} + \sigma_{ji}Z)$$

from which the announced result follows since a ranking in the  $\leq_{ICX}$  sense is not affected by scaling (i.e. multiplication by  $m_{x+j,m}$ ). This completes the proof.

Now, let us denote as  ${}_dP_x(t_n | \boldsymbol{\kappa})$  the random  $d$ -year survival probability for an individual aged  $x$  in calendar year  $t_n$ , that is, the conditional probability that this individual reaches age  $x+d$  in year  $t_n+d$ , given the vector  $\boldsymbol{\kappa}$  of the  $\kappa_t$ . It is formally defined as

$${}_dP_x(t_n | \boldsymbol{\kappa}) = \exp(-S_d)$$

where

$$S_d = m_{x,t_n} + \sum_{j=1}^{d-1} m_{x+j,t_n} \prod_{i=1}^j g(Z_{x+j,t_n+i}).$$

We know from Proposition 3.4.29 of Denuit *et al.* (2005) that

$$S_d \leq_{ICX} S_d^u = m_{x,t_n} + \sum_{j=1}^{d-1} m_{x+j,t_n} \prod_{i=1}^j g(\mu_{ji} + \sigma_{ji} Z).$$

Here, we take  $\exp(-S_d^u)$  as an approximation to the  $d$ -year survival probability  ${}_dP_x(t_n | \boldsymbol{\kappa}) = \exp(-S_d)$  and we investigate its accuracy in the next section, based on numerical illustrations.

As a final comment, let us mention that the approach developed in the present section also applies to alternative specifications for  $Z_{x,t}$ . For instance, the comonotonic approximations also hold for models with a cohort effect as long as the individual under interest belongs to a cohort whose effect can be estimated from the available historical data. Specifically, we can also consider

$$Z_{x+j,t_n+i} = \beta_{x+j} \kappa_{t_n+i} + \iota_{t_n-x+i-j} \sim N(\mu_{ji}, \sigma_{ji}^2).$$

with

$$\mu_{ji} = \beta_{x+j} E(\kappa_{t_n+i}) + \iota_{t_n-x+i-j}, \quad \sigma_{ji}^2 = (\beta_{x+j})^2 \text{Var}(\kappa_{t_n+i})$$

as long as the cohort effect  $\iota$  can be considered as constant (*i.e.* estimated from past data).

#### 4. Numerical illustrations

Let us consider a basic life annuity contract paying 1 unit of currency at the end of each year, as long as the annuitant survives. The random life annuity single premium, that is, the conditional expectation of the payments made to an annuitant aged  $x$  in the year  $t_n$  given  $\kappa_{t_n}, \kappa_{t_n+1}, \kappa_{t_n+2}, \dots$  is

$$a_x(t_n | \boldsymbol{\kappa}) = \sum_{d \geq 1} {}_dP_x(t_n | \boldsymbol{\kappa}) \nu(0, d),$$

where  $\nu(\dots)$  is the discount factor (precisely,  $\nu(s, t)$  is the present value at time  $s$  of a unit payment made at time  $t$ ). Note that  $a_x(t_n | \boldsymbol{\kappa})$  corresponds to the generation aged  $x$  in calendar year  $t_n$ , and accounts for future mortality improvements experienced by this particular cohort. Clearly,  $a_x(t_n | \boldsymbol{\kappa})$  is a random variable that depends on the future

trajectory  $\kappa_n, \kappa_{n+1}, \kappa_{n+2}, \dots$ . An analytical computation of the distribution function of  $a_x(t_n | \boldsymbol{\kappa})$  is out of reach.

From the approximation  $S_d^u$  assumed for  $S_d$ , we get the following approximation for the random survival probabilities

$${}_d P_x(t_n | \boldsymbol{\kappa}) \approx \exp\left(-F_{S_d^u}^{-1}(1-U)\right)$$

where  $U$  is uniformly distributed on the interval  $(0,1)$ . Note that the same random variable  $U$  is used for all of the values of  $d$ , making the approximations to the conditional survival probabilities comonotonic. Hence, we obtain the following approximation for  $a_x(t_n | \boldsymbol{\kappa})$

$$a_x(t_0 | \boldsymbol{\kappa}) \approx \sum_{d \geq 1} \exp\left(-F_{S_d^u}^{-1}(1-U)\right) v(0, d).$$

Since this approximation is a sum of comonotonic random variables, its quantile functions is additive. So, we obtain the following approximations for the quantile function  $F_{a_x(t_n | \boldsymbol{\kappa})}^{-1}(z)$  of  $a_x(t_n | \boldsymbol{\kappa})$

$$F_{a_x(t_0 | \boldsymbol{\kappa})}^{-1}(z) \approx \sum_{d \geq 1} \exp\left(-F_{S_d^u}^{-1}(1-z)\right) v(0, d)$$

where

$$F_{S_d^u}^{-1}(z) \approx m_{x, t_n} + \sum_{j=1}^{d-1} m_{x+j, t_n} \prod_{i=1}^j g\left(\mu_{ji} + \sigma_{ji} \Phi^{-1}(z)\right).$$

The random cohort life expectancy  $e_x^\square(t_n | \boldsymbol{\kappa})$  is the conditional expected remaining lifetime of an individual aged  $x$  in year  $t_n$ , given  $\kappa_n, \kappa_{n+1}, \kappa_{n+2}, \dots$ . Keeping the assumption that deaths are uniformly distributed over each calendar year, this demographic indicator is given by

$$e_x^\square(t_n | \boldsymbol{\kappa}) = \frac{1}{2} + \sum_{d \geq 1} {}_d P_x(t_n | \boldsymbol{\kappa}).$$

We use the  $\square$  superscript to indicate that we work along a diagonal band in the Lexis diagram. Except for the additive constant  $1/2$ ,  $e_x^\square(t_n | \boldsymbol{\kappa})$  coincides with  $a_x(t_n | \boldsymbol{\kappa})$  if we



let the interest rate tend to zero. As was the case for  $a_x(t_n | \boldsymbol{\kappa})$ , an analytic computation of the distribution function of  $e_x^\square(t_n | \boldsymbol{\kappa})$  is out of reach.

From the approximation  $S_d^u$  assumed for  $S_d$ , we get the approximation for  $e_x^\square(t_n | \boldsymbol{\kappa})$

$$e_x^\square(t_n | \boldsymbol{\kappa}) \approx \frac{1}{2} + \sum_{d \geq 1} \exp(-S_d^u).$$

Since the  $S_d^u$ 's are sums of comonotonic random variables, their quantile functions are additive. Moreover, the  $z$ th quantile of  $\exp(-S_d^u)$  is  $\exp(-F_{S_d^u}^{-1}(1-z))$ . This provides the following approximation for the quantile function  $F_{e_x^\square(t_n | \boldsymbol{\kappa})}^{-1}(z)$  of  $e_x^\square(t_n | \boldsymbol{\kappa})$

$$F_{e_x^\square(t_n | \boldsymbol{\kappa})}^{-1}(z) \approx \frac{1}{2} + \sum_{d \geq 1} \exp(-F_{S_d^u}^{-1}(1-z)).$$

For the numerical results which follow, we use the 1961-2009 USA male and female mortality experiences with deaths and matching exposures by individual calendar year for individual ages 20-104 (the full age range being 0-109) available through the Human Mortality Database (HMD). Preliminary analysis including an analysis of residuals (not reproduced) is supportive of the inclusion of the cohort effects terms  $\iota_{t-x}$  for males but not for females. Hence, we report the results for the respective  $H_1$  formulation (see below for a precise definition) for males and the  $LC$  formulation for females by depicting the fitted parameter values in Figure 1. Also included are the period component time series forecasts using the selected  $AR(1)$  process for males and the simple  $AR(0)$  process for females, fitted as Gaussian regression models. Thus, for males, we have chosen to model the  $MIR$  Gaussian structure using the so-called  $H_1$  formulation:

$$Z_{x+j,t_n+i} = \beta_{x+j} \kappa_{t_n+i} + \iota_{t_n-x+i-j} \sim N(\mu_{ji}, \sigma_{ji}^2)$$

whereas for females, we have modelled the structure based on the  $LC$  formulation:

$$Z_{x+j,t_n+i} = \beta_{x+j} \kappa_{t_n+i} \sim N(\mu_{ji}, \sigma_{ji}^2)$$

where

$$\mu_{ji} = \beta_{x+j} E(\kappa_{t_n+i}) + \iota_{t_n-x+i-j} \text{ or } \mu_{ji} = \beta_{x+j} E(\kappa_{t_n+i}) \text{ and } \sigma_{ji}^2 = (\beta_{x+j})^2 \text{Var}(\kappa_{t_n+i}).$$

Using these parameters estimates and forecasts we tabulate (Table 1: 1<sup>st</sup> and 3<sup>rd</sup> panels) details for life expectancy and 4% annuity predictions, computed by cohort trajectory for ages 40, 45, 50, ...75 focused on the year 2009. For convenience, we have not used the topping-out procedure advocated by Haberman and Renshaw (2012) for dealing with extrapolating the life table to the oldest ages. For comparison, we also tabulate (Table 1: 2<sup>nd</sup> and 4<sup>th</sup> panels) the respective equivalent life expectancies and 4% annuity predictions generated by the simulation method described in Haberman and Renshaw (2012), using a total of 10,000 simulations for each age. Referring to Table 1 and Figure 1 we note the following points:

- On comparing like for like, there is an exceptionally close agreement between the matching theoretical and simulated median predictions. However, the interval prediction widths in the theoretical cases are much wider when compared with the matching simulated cases.
- We note the narrowness of the simulated males prediction intervals, which are appreciably narrower than equivalent simulated intervals for the England & Wales male mortality experience depicted in Figure 8 of Haberman & Renshaw (2012), where topping-out by age has been applied but this seems to have little effect on increasing the interval widths.
- With the exception of a few isolated ages in the male experience, the beta parameters are positive over the full age range for males and females and therefore for both modelling structures. It would be possible to adapt the algorithms so that the beta parameters are constrained to be positive. We note further that the period index forecasts for mortality improvement rates are negative for males using  $H_1$  but positive for females using  $LC$ .
- In order to reach the 99.5% solvency probability required under Solvency 2, we assume that the policyholders are required to provide premiums adding up to the 75th quantile of the present value of annuity payments and the insurer pays for the difference between the 99.5th quantile of these payments and the aforementioned 75th quantile. Here, we make the assumption that the size of the portfolio is large enough to neglect diversifiable risk so that only the systematic risk matters. The latter is equal to the size of the portfolio multiplied by the expected present value of the annuity payments given future mortality. The difference in the 99.5<sup>th</sup> and 75<sup>th</sup> quantiles appears in the last column of each panel in Table 1. Comparing the differences based on the approximations derived in the present paper to the simulated ones, we see there that the amount of capital is over-estimated when the approximations are used.

In the standard model's module for longevity risk, the Value-at-Risk at probability level 99.5% is approximated by the change in net asset value due to a pre-specified longevity shock based on a 25% reduction in mortality rates at all ages. Let us now explore the accuracy of the approximations derived in the present paper in dealing with such a shock. Specifically, we compare the values obtained from the approximations derived in the present paper to those coming from the standard formula which consists of reducing death rates by 25%. Table 2 is the same as Table 1 subject to a reduction in all projected mortality rates by a factor of 25%. The close agreement between simulated and theoretical medians is preserved, as in Table 1. Also the medians in Table 2 are

consistently higher than their matching counter-parts in Table 1 as expected. The approximations to the 95<sup>th</sup> quantiles listed in Table 1 are reasonably close to their simulated counterparts in Table 2 based on a reduction of death rates by 25%. This shows that the approximations derived in the present paper for high quantiles may be used as an alternative to the standard approach which involves decreasing all death rates by, say, 25%.

## 5. Discussion

Combining a conservative shift with non necessary conservative ones, the approximations derived in the present paper appear to be very accurate in the centre of the distribution (around the median) but tend to over estimate the tails (left and right). Using the proposed easy-to-compute approximation may thus be a good strategy for the calculation of the percentiles in the centre of the distribution (for example, the valuation of the median) as it would considerably reduce the computational burden and save time.

## Acknowledgements

The authors would like to express their gratitude to an anonymous Referee whose comments have been extremely useful to revise a previous version of the present work. The financial support of PARC “Stochastic Modelling of Dependence” 2012-17 awarded by the *Communauté française de Belgique* is gratefully acknowledged by Michel Denuit.

## References

- Brouhns, N., Denuit, M., Vermunt, J.K. 2002. A Poisson log-bilinear approach to the construction of projected life tables. *Insurance: Mathematics and Economics* 31, 373-393.
- Denuit, M. 2007. Distribution of the random future life expectancies in log-bilinear mortality projection models. *Lifetime Data Analysis*, 13, 381-397.
- Denuit, M., Dhaene, J. 2007. Comonotonic bounds on the survival probabilities in the Lee-Carter model for mortality projection. *Computational and Applied Mathematics* 203, 169-176.
- Denuit, M., Dhaene, J., Goovaerts, M.J., Kaas, R. 2005. *Actuarial Theory for Dependent Risks: Measures, Orders and Models*. Wiley, New York.
- Denuit, M., Haberman, S., Renshaw A.E. 2010. Comonotonic approximations to quantiles of life annuity conditional expected present values: extensions to general ARIMA models and comparison with the bootstrap. *ASTIN Bulletin* 40, 331-349.
- Haberman, S., Renshaw A.E. 2009. On age-period-cohort parametric mortality rate projections. *Insurance: Mathematics and Economics* 45, 255-270.
- Haberman, S., Renshaw A.E. 2012. Parametric mortality improvement rate modelling and projecting. *Insurance: Mathematics and Economics* 50, 309-333.
- Lee, R.D., Carter, L., 1992. Modelling and forecasting the time series of US mortality. *Journal of the American Statistical Association* (with discussion) 87, 659-272.

- Renshaw, A.E., Haberman, S. 2003. Lee-Carter mortality forecasting with age-specific enhancement. *Insurance: Mathematics and Economics* 33, 255-272.
- Renshaw, A.E., Haberman, S. 2006. A cohort based extension to the Lee-Carter model for mortality reduction factors. *Insurance: Mathematics and Economics* 38, 556-570.

<i>age  quantile</i>	<i>life expectancy</i>				<i>4% annuity value</i>			
	<i>0.05</i>	<i>0.5</i>	<i>0.95</i>	<i>0.995-0.75</i>	<i>0.05</i>	<i>0.5</i>	<i>0.95</i>	<i>0.995-0.75</i>
40	36.64	45.31	56.21	8.929	18.38	19.92	21.40	1.217
45	32.70	40.27	50.22	8.687	17.34	18.91	20.52	1.397
50	28.86	35.37	44.28	8.295	16.17	17.76	19.48	1.577
55	25.11	30.61	38.40	7.720	14.88	16.44	18.23	1.740
60	21.46	25.98	32.55	6.910	13.43	14.92	16.72	1.843
65	17.97	21.59	26.91	5.927	11.85	13.24	14.97	1.871
70	14.68	17.46	21.56	4.797	10.17	11.41	12.98	1.792
75	11.64	13.67	16.62	3.587	8.43	9.46	10.80	1.579

**USA females- LC: theoretical predictions**

<i>age  quantile</i>	<i>life expectancy</i>				<i>4% annuity value</i>			
	<i>0.05</i>	<i>0.5</i>	<i>0.95</i>	<i>0.995-0.75</i>	<i>0.05</i>	<i>0.5</i>	<i>0.95</i>	<i>0.995-0.75</i>
40	43.56	45.22	46.94	2.002	19.64	19.91	20.17	0.288
45	38.70	40.21	41.76	1.746	18.62	18.90	19.18	0.310
50	33.95	35.32	36.74	1.635	17.44	17.75	18.06	0.352
55	29.36	30.57	31.85	1.448	16.11	16.43	16.76	0.373
60	24.87	25.95	27.06	1.328	14.58	14.91	15.25	0.396
65	20.65	21.56	22.50	1.132	12.89	13.23	13.57	0.395
70	16.70	17.44	18.22	0.914	11.07	11.39	11.73	0.375
75	13.04	13.64	14.28	0.749	9.15	9.45	9.76	0.356

**USA females- LC: simulated predictions (10,000 simulations)**

<i>age  quantile</i>	<i>life expectancy</i>				<i>4% annuity value</i>			
	<i>0.05</i>	<i>0.5</i>	<i>0.95</i>	<i>0.995-0.75</i>	<i>0.05</i>	<i>0.5</i>	<i>0.95</i>	<i>0.995-0.75</i>
40	40.38	44.61	49.14	4.909	18.87	19.62	20.33	0.760
45	34.71	38.13	41.97	4.330	17.51	18.24	18.97	0.803
50	29.84	32.58	35.72	3.636	16.12	16.80	17.51	0.801
55	25.46	27.61	30.12	2.941	14.65	15.28	15.95	0.765
60	22.05	23.77	25.76	2.348	13.35	13.92	14.53	0.707
65	18.66	19.99	21.52	1.811	11.90	12.41	12.95	0.633
70	15.10	16.05	17.16	1.305	10.19	10.60	11.06	0.531
75	11.68	12.31	13.04	0.869	8.31	8.63	8.97	0.411

**USA males-H1: theoretical predictions**

<i>age  quantile</i>	<i>life expectancy</i>				<i>4% annuity value</i>			
	<i>0.05</i>	<i>0.5</i>	<i>0.95</i>	<i>0.995-0.75</i>	<i>0.05</i>	<i>0.5</i>	<i>0.95</i>	<i>0.995-0.75</i>
40	43.88	44.54	45.18	0.729	19.49	19.61	19.73	0.131
45	37.52	38.10	38.66	0.618	18.11	18.24	18.36	0.133
50	32.06	32.54	33.03	0.566	16.67	16.80	16.92	0.145
55	27.18	27.59	28.00	0.444	15.15	15.28	15.40	0.137
60	23.39	23.74	24.09	0.418	13.79	13.91	14.03	0.140
65	19.67	19.96	20.25	0.335	12.28	12.40	12.51	0.131
70	15.81	16.03	16.25	0.267	10.49	10.59	10.69	0.122
75	12.13	12.29	12.46	0.193	8.53	8.62	8.70	0.100

**USA males- H1: simulated predictions (10,000 simulations)**

Table 1. USA female & male 2009 life expectancy and 4% annuity quantile predictions, ages 40(05)75: comparison of theoretical & simulated predictions.

<i>age  quantile</i>	<i>life expectancy</i>				<i>4% annuity value</i>			
	<i>0.05</i>	<i>0.5</i>	<i>0.95</i>	<i>0.995-0.75</i>	<i>0.05</i>	<i>0.5</i>	<i>0.95</i>	<i>0.995-0.75</i>
40	38.80	48.32	58.29	7.493	18.92	20.48	21.78	1.000
45	34.82	43.21	52.47	7.410	17.95	19.56	21.00	1.167
50	30.91	38.21	46.67	7.209	16.85	18.49	20.06	1.340
55	27.09	33.34	40.88	6.856	15.62	17.26	18.93	1.508
60	23.35	28.57	35.10	6.602	14.23	15.83	17.55	1.638
65	19.75	24.00	29.45	5.575	12.70	14.22	15.92	1.714
70	16.33	19.67	24.00	4.681	11.06	12.44	14.04	1.700
75	13.14	15.63	18.86	3.654	9.33	10.52	11.92	1.563

**USA females- LC: theoretical predictions**

<i>age  quantile</i>	<i>life expectancy</i>				<i>4% annuity value</i>			
	<i>0.05</i>	<i>0.5</i>	<i>0.95</i>	<i>0.995-0.75</i>	<i>0.05</i>	<i>0.5</i>	<i>0.95</i>	<i>0.995-0.75</i>
40	46.51	48.22	49.94	1.876	20.22	20.47	20.71	0.265
45	41.58	43.14	44.65	1.670	19.28	19.55	19.80	0.277
50	36.72	38.16	39.60	1.623	18.18	18.48	18.78	0.323
55	32.02	33.28	34.58	1.489	16.94	17.25	17.56	0.353
60	27.41	28.54	29.68	1.437	15.49	15.83	16.15	0.393
65	23.01	23.99	24.97	1.144	13.88	14.22	14.55	0.383
70	18.86	19.66	20.52	0.997	12.11	12.44	12.78	0.397
75	14.98	15.64	16.34	0.769	10.21	10.52	10.84	0.352

**USA females- LC: simulated predictions (10,000 simulations)**

<i>age  quantile</i>	<i>life expectancy</i>				<i>4% annuity value</i>			
	<i>0.05</i>	<i>0.5</i>	<i>0.95</i>	<i>0.995-0.75</i>	<i>0.05</i>	<i>0.5</i>	<i>0.95</i>	<i>0.995-0.75</i>
40	43.47	47.88	52.22	4.477	19.55	20.26	20.90	0.660
45	37.72	41.40	45.23	4.122	18.30	19.01	19.69	0.723
50	32.73	35.74	39.00	3.593	17.00	17.69	18.37	0.746
55	28.20	30.63	33.30	3.011	15.62	16.27	16.93	0.737
60	24.62	26.59	28.76	2.450	14.38	14.98	15.60	0.697
65	21.03	22.59	24.30	1.937	12.97	13.52	14.08	0.641
70	17.23	18.39	19.66	1.454	11.27	11.74	12.23	0.561
75	13.53	14.34	15.22	1.021	9.37	9.75	10.15	0.458

**USA males-H1: theoretical predictions**

<i>age  quantile</i>	<i>life expectancy</i>				<i>4% annuity value</i>			
	<i>0.05</i>	<i>0.5</i>	<i>0.95</i>	<i>0.995-0.75</i>	<i>0.05</i>	<i>0.5</i>	<i>0.95</i>	<i>0.995-0.75</i>
40	47.16	47.78	48.39	0.693	20.14	20.24	20.35	0.117
45	40.79	41.34	41.91	0.646	18.89	19.00	19.12	0.128
50	35.22	35.73	36.21	0.548	17.57	17.69	17.80	0.128
55	30.20	30.63	31.06	0.485	16.15	16.27	16.39	0.133
60	26.23	26.59	26.95	0.427	14.86	14.98	15.09	0.135
65	22.28	22.59	22.89	0.351	13.40	13.52	13.63	0.127
70	18.16	18.41	18.65	0.282	11.64	11.75	11.85	0.118
75	14.19	14.37	14.56	0.211	9.68	9.77	9.86	0.103

**USA males- H1: simulated predictions (10,000 simulations)**

Table 2. USA female & male 2009 life expectancy and 4% annuity quantile predictions, ages 40(05)75: comparison of theoretical & simulated predictions subject to a 25% reduction in projected mortality rates.