



City Research Online

City, University of London Institutional Repository

Citation: Khawaja, A. P., Crabb, D. P. & Jansonius, N. M. (2015). Time to abandon over-simplified surrogates of ocular perfusion pressure in glaucoma research. *Acta Ophthalmologica*, 93(1), e85-e86. doi: 10.1111/aos.12431

This is the published version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/4997/>

Link to published version: <https://doi.org/10.1111/aos.12431>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk

Letter to the Editor

Time to abandon over-simplified surrogates of ocular perfusion pressure in glaucoma research

Anthony P. Khawaja,¹ David P. Crabb² and Nomdo M. Jansonius^{3,4}

¹Department of Public Health and Primary Care, Institute of Public Health, University of Cambridge School of Clinical Medicine, Cambridge, UK;

²Department of Optometry and Visual Science, City University London, London, UK;

³Department of Ophthalmology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands;

⁴Department of Epidemiology, Erasmus Medical Center, Rotterdam, The Netherlands

doi: 10.1111/aos.12431

© 2014 The Authors. *Acta Ophthalmologica* published by John Wiley & Sons Ltd on behalf of *Acta Ophthalmologica Scandinavica Foundation*.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Editor,

The role of ocular perfusion pressure (OPP) in the pathogenesis of glaucoma has attracted a great deal of research interest, as highlighted in a

review article recently published in *Acta Ophthalmologica* (Costa et al. 2013). For the majority of studies in this field, blood pressure (BP) minus intraocular pressure (IOP) has been used as a simple surrogate measure of OPP. The authors of the review article correctly acknowledge that interpretation of these surrogate measures is problematic and that any crude association observed between OPP and glaucoma may be related solely to the IOP component, given the known strength of IOP as a risk factor for glaucoma. However, we strongly disagree with the authors' conclusion that statistically adjusting for IOP is a satisfactory solution to this problem. We have previously shown that it is quite impossible to untangle the effects of IOP and BP in a model containing OPP results (Khawaja et al. 2013). In short, adjusting for IOP in a model containing OPP will inevitably result in the situation that the coefficients for OPP actually represent the effect of BP only, and not OPP. This has been substantiated theoretically and demonstrated clearly using a simulated dataset; the coefficients for OPP in IOP adjusted regression models were exactly the same as those for BP in IOP adjusted models (Khawaja et al. 2013). Therefore, in studies that have found a significant association between OPP and glaucoma using regression models adjusted for IOP, it is actually a significant association between BP and glaucoma that has been demonstrated, and no conclusions can be drawn regarding perfusion pressure. OPP may well be

important in glaucoma pathogenesis, but unfortunately the surrogate approach of calculating BP minus IOP is over-simplified, and results have been consistently misinterpreted, which has not furthered our understanding of the disease process. Given the critical issues highlighted, we strongly suggest abandoning the approach of assuming that BP minus IOP reflects OPP and that research resources are directed towards new methods in this important field.

References

- Costa VP, Harris A, Anderson D et al. (2013): Ocular perfusion pressure in glaucoma. *Acta Ophthalmol* [Epub ahead of print].
Khawaja AP, Crabb DP & Jansonius NM (2013): The role of ocular perfusion pressure in glaucoma cannot be studied with multi-variable regression analysis applied to surrogates. *Invest Ophthalmol Vis Sci* **54**: 4619–4620.

Correspondence:

Anthony P. Khawaja
Strangeways Research Laboratory (EPIC)
2 Worts' Causeway
Cambridge CB1 8RN
UK
Tel: +44 (0) 1223 748682
Fax: +44 (0) 1223 748676
Email: anthony.khawaja@gmail.com

Anthony Khawaja is funded by a Wellcome Trust Clinical Research Fellowship.