



City Research Online

City, University of London Institutional Repository

Citation: Atkinson, L., Olander, E. K. and French, D. P. (2013). Why don't many obese pregnant and post-natal women engage with a weight management service?. *Journal of Reproductive and Infant Psychology*, 31(3), pp. 245-256. doi: 10.1080/02646838.2013.809518

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/5782/>

Link to published version: <http://dx.doi.org/10.1080/02646838.2013.809518>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk

1 Why don't many obese pregnant and post-natal women engage with a weight management
2 service?

3

4 Lou Atkinson, Applied Research Centre in Health and Lifestyle Interventions, Coventry
5 University, Coventry, UK l.atkinson@coventry.ac.uk

6 Ellinor K Olander, Applied Research Centre in Health and Lifestyle Interventions, Coventry
7 University, Coventry, UK e.olander@coventry.ac.uk

8 David P French, Applied Research Centre in Health and Lifestyle Interventions, Coventry
9 University, Coventry, UK david.french@manchester.ac.uk

10

11 Corresponding author at: Lou Atkinson, Applied Research Centre in Health and Lifestyle
12 Interventions, Coventry University, Priory Street, Coventry, CV1 5FB, UK tel. 02476
13 887470, fax. 02476795987, email. l.atkinson@coventry.ac.uk

14

15

16

1 Abstract

2 Objective: This study investigated the views and experiences of obese pregnant and post-
3 natal women who had declined or disengaged from an evidence-based weight management
4 service, and their reasons for doing so.

5 Background: Despite significant risks of maternal obesity to both mother and baby, the
6 majority of obese women offered tailored weight management support during or after
7 pregnancy declined to use it, and many women who accepted the service disengaged soon
8 after.

9 Methods: Semi-structured interviews regarding women's views and experiences were
10 conducted with obese pregnant and post-natal women who declined the service (N=7) and
11 women who disengaged from the service (N=11), and analysed thematically.

12 Results: Four main themes were identified. "First contact counts" related to inadequate
13 explanation of the service by the referrer, being offended by the referral, and negative
14 expectations of the service. "Missed opportunities for support" describes what support
15 declining women identified as desirable, such as regular weight monitoring. "No need for
16 help", and "Service not meeting needs" related to personal choices regarding weight
17 management, including not wanting support and preferring group-based services.

18 Conclusions: While some obese pregnant and post-natal women did not want any type of
19 weight management support, many did but failed to engage with the service on offer due to a
20 variety of barriers. A more sensitive and transparent referral process and further tailoring of
21 the service to meet individual needs may increase uptake and continued use of this service.
22 The inclusion of non-participants and non-completers formed a valuable element of service
23 evaluation.

24 Keywords: Maternal obesity, Qualitative research, Interviews, Service evaluation, Pregnancy,
25 Post-natal

26

1 Introduction

2

3 It has been estimated that 15.6% of women in England have a Body Mass Index (BMI)
4 $\geq 30\text{kg/m}^2$ when they become pregnant, with women from deprived backgrounds being more
5 at risk of maternal obesity (Heslehurst, Rankin, Wilkinson, & Summerbell, 2010). Being
6 obese during pregnancy carries serious health risks for both mother and child (Abenheim,
7 Kinch, Morin, Benjamin, & Usher, 2007), and significantly increases the risk of obesity in
8 the resulting infant (Griffiths, Hawkins, Cole, & Dezateux, 2010; Pirkola, et al., 2010). In
9 recognition of this, the National Institute for Health and Clinical Excellence (NICE) in
10 England and Wales recently issued new guidance on interventions for weight management
11 before, during and after pregnancy (National Institute for Health and Clinical Excellence,
12 2010). This guidance and associated commissioning guide (National Institute for Health and
13 Clinical Excellence, 2011) recommends early identification of pregnant women with a BMI
14 $\geq 30\text{kg/m}^2$, and provision of practical and tailored advice from health professionals regarding
15 healthy eating and physical activity during pregnancy. The guidance also suggests
16 encouragement and support to achieve a healthy weight up to two years after childbirth and
17 before the birth of a subsequent baby, including provision of tailored physical activity and
18 dietary advice, signposting to community services and encouragement to breastfeed.

19

20 The Maternal and Early Years Healthy Weight Service (MAEYS) is a care pathway which
21 has been cited as an exemplar of NICE's recommendations (National Institute for Health and
22 Clinical Excellence, 2011). MAEYS was introduced in June 2009, initially as a one year pilot
23 programme to assess feasibility in six Primary Care Trusts (PCTs) in the West Midlands area
24 of England where over 20% of women are obese at the start of pregnancy (Sheikh, Malik, &
25 Gardosi, 2010), and is one of the first evidence-based care pathways developed in the UK
26 specifically for obese women during and after pregnancy (Baker, 2011). MAEYS is delivered

1 on a one to one basis, in the client's home, by dedicated Healthy Weight Advisors who
2 received 12 days training to develop specific skills and competences in diet, physical activity,
3 infant feeding and behaviour change (see Baker, 2011). The service is free and available
4 from early pregnancy until the child is two years old. It provides tailored advice and support
5 with behaviour change on healthy eating, physical activity and infant feeding, along with
6 regular weight monitoring. Eligible women are referred by midwives and other health
7 professionals. The primary aim of MAEYS is to prevent childhood obesity through
8 minimizing gestational weight gain. Secondary aims are to reduce the significant health risks
9 associated with being obese during pregnancy, to support post-natal weight loss and to
10 establish healthy infant feeding habits.

11

12 Previous research has highlighted the need to more fully explore how to engage obese women
13 in weight management services (Heslehurst, Bell, & Rankin, 2011), and maternity-care
14 professionals have identified the importance of reaching all obese women, not just those who
15 are already motivated to improve their health behaviour (Heslehurst, Moore, et al., 2010). A
16 qualitative evaluation of MAEYS has shown good levels of acceptability among a sample of
17 service users (Atkinson, Olander & French, in review) and data collected from women
18 completing a whole phase of the service showed positive results for gestational weight gain
19 and breastfeeding rates (Baker, 2011). However, often this is where evaluation ends, without
20 investigating the views and experiences of people who declined or failed to complete
21 interventions (Sørensen, Skovgaard, & Puggaard, 2006). During the pilot of MAEYS half of
22 the women referred declined to use the service, and two thirds of those who started using the
23 service failed to complete at least three quarters of planned sessions. Other weight
24 management services in pregnancy have reported initiation of the service to be as low as
25 14.5% (Knight & Wyatt, 2010) and 35% (Davis et al, 2012) of the women who were invited

1 to participate, and one service reporting zero attendance from 76 women referred (West,
2 2010). Multiple barriers to recruitment for group-based maternal healthy weight services
3 have been reported by obese pregnant women, (Olander & Atkinson, 2013) and referring
4 health professionals (Davis et al, 2012; West, 2010) including inconvenient time and
5 location, work commitments, feeling unwell, not wanting to focus on weight management
6 during pregnancy, and health professionals' difficulties in raising the issue. Thus while the
7 individualised home-based design of MAEYS may have overcome some of the pragmatic
8 barriers related to more structured, group-based services, it is important when assessing the
9 acceptability of the care pathway to understand why the majority of eligible women did not
10 take advantage of MAEYS. Previous research has demonstrated that valuable insights into
11 acceptability and adherence can be gained by interviewing decliners and non-completers
12 when evaluating a community-based lifestyle intervention (Barter-Godfrey, Taket, &
13 Rowlands, 2007). Additionally, evidence regarding why women disengage from maternal
14 healthy weight services is lacking, as few such services have been delivered or evaluated.
15 Without this understanding, significant health inequalities could develop or worsen. In
16 addition, levels of uptake and adherence to health services are important contributors to their
17 cost-effectiveness. Therefore, the aim of the present study is to investigate, through
18 qualitative methods a) why some women who were referred to MAEYS chose not to use the
19 service, and b) why some women who began using MAEYS disengaged from the service.

20

21 Method

22

23 *Design*

24 Cross-sectional interview study, conducted one year after the service was introduced into the
25 antenatal and post-natal care pathways.

26

1 *Procedure*

2 Following ethical approval from the authors' University Ethics Committee, recruitment of
3 participants was initiated by MAEYS project leads within each of the pilot PCTs. Women
4 who had declined the service and women who had been clients of MAEYS but had
5 voluntarily ceased using the service were contacted, and asked if they would be willing to be
6 contacted regarding the evaluation. If women agreed to be contacted, or did not submit an
7 "opt-out" response, they were contacted by the research team and invited to participate in
8 either a face to face or telephone interview, according to their preference.

9 All participants were provided with an information sheet and gave either written or verbal
10 (for telephone interviews) consent prior to being interviewed. Interviews were conducted by
11 a researcher with the use of a topic guide. Topics for discussion included: Introduction to the
12 service, expectations for the service, reasons for declining or leaving the service, positive and
13 negative experiences of the service and suggestions for changes. However, participants were
14 encouraged to talk freely about their experiences of MAEYS, and their own weight
15 management. Interview duration ranged from 10 minutes to 32 minutes, with a median
16 duration of 16 minutes. Interviews were recorded and transcribed verbatim.

17

18 *Sample*

19 See Table 1 for details of sample breakdown. [Table 1 near here]

20 Seven women who had declined MAEYS agreed to be interviewed, all of whom described
21 themselves as White British. Ages ranged from 27 to 36 years, and parity ranged from first to
22 fifth pregnancy. Eleven women who had disengaged from the service agreed to be
23 interviewed, ten of whom described themselves as White British, and one woman described
24 herself as British Pakistani. Ages ranged from 21 to 35 years and parity ranged from first to

1 third pregnancy. Both groups contained at least one participant from each of the six PCTs.
2 All participants had a pre-pregnancy BMI ≥ 30 kg/m².

3

4 *Analysis*

5 Data were analysed separately for each sample via a process of inductive thematic analysis,
6 using a realist approach (Braun & Clarke, 2006). As such, the main goal of the analysis was
7 to provide a detailed description of the entire dataset, without being led by the questions
8 asked, in order to inform current and future health service design. Briefly, this process
9 involved; familiarisation with the data by repeated readings of the transcripts, coding of all
10 data relating to participants' experiences of the service, identification of patterns (themes)
11 within the data, checking identified themes against the coded data, and finally defining the
12 themes through examination of the precise content of groups of similar data. Two researchers
13 independently completed the analysis for each sample and resulting themes were compared.
14 There was a high degree of agreement between researchers concerning the nature of the
15 dominant themes and the final themes were produced following discussion. Finally the
16 analyses for each sample were compared to identify congruence and divergence of themes
17 between women who had declined the service and women who had disengaged from the
18 service.

19

20 *Results*

21

22 Four main themes emerged from the data: "First contact counts", and "Missed opportunities
23 for support" were aggregated themes relevant to both women who had declined the service
24 and women who had disengaged. "No need for help", and "Service not meeting needs" were

1 relevant to women who had declined and women who had disengaged respectively. Selected
2 quotes are included for illustration of key points. Pseudonyms are used to protect anonymity.

3

4 *First contact counts*

5

6 This theme related to aspects of the referral experience.

7

8 Many of the women reported that they were not told that they had been referred or were not
9 informed about the service, prior to being contacted. These women reported being shocked
10 and confused after unexpectedly receiving a phone call or letter.

11

12 *“I just received a letter in the post, and saying that I’d been put forward for it by my*
13 *midwife... And no-one had warned me about it, and I think if someone had warned me about*
14 *it, it would have been better, because I just opened the letter, read the letter and thought ‘Oh*
15 *God’ and just burst into tears. It’s a sensitive time anyway because your body’s changing and*
16 *you’re conscious of what you look like anyway.”*

17 Paula (declined)

18

19 Several women reported being upset by the referral. They often reported that they felt the
20 referral was a form of judgement, either of themselves as a person or as a parent.

21

22 *“At the end of the day I’ve brought up my first child and he’s fine....So it’s like he’s not*
23 *overweight, I know how to wean him, I know about healthy eating and all that, it’s just me*
24 *that’s overweight.”*

25 Tara (declined)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

Some women were referred to the service following discussions with a health professional regarding their weight, with the subject either raised by them or the health professional. In these cases the women reported being happy to be referred. This was only reported by women who had disengaged. Additionally, some women had seen leaflets or posters and had self-referred. Aside from the women who were unaware of the referral, several women reported being told about the service, but stated that they were given very little information about it by the referrer. Some women cited this lack of information as a contributing reason for not engaging with the service.

“The nurse said there’s a lady in here from some programme, she didn’t have any information on her whatsoever...I did feel maybe if I got a bit more information I’d have followed it up a bit more.”

Crystal (disengaged)

However, some women reported having the service explained to them in more depth and stated that they were happy with this.

Some women who had been told about the referral reported not being contacted by the service as promised, and this was then given as a reason for not engaging with the service.

“Nobody’s mentioned it again since...I never just, never thought to bring it up when you attend the appointments, you’re worried about other things.”

Shelly (declined)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

No need for help

Aside from issues related to the referral process, two further issues emerged within the women’s explanations for declining the service. Firstly, some reported feeling that the referral unjustly implied that they were not able to make healthy choices for themselves and their child.

“I just feel as if like you’re being, because they want to offer you help with the weaning, which is all very good if, like I say, it’s your first child, but I felt like they were going to come in and try to tell me how to raise my child, that’s how I felt. And I don’t like things like that, it makes me feel like they think I’m a bad mother.”

Tara (declined)

Secondly, some women reported already having a healthy lifestyle or using other services.

“Since I’ve had the little one I have joined Weightwatchers, which is something I have done before and it’s worked, or not worked, depending on what mood I’m in really.”

Shelly (declined)

“Generally speaking we have a very balanced diet, so I’ll probably just carry on like that. I don’t really know what they could do that would make a difference.”

Beatrice (declined)

Service not meeting needs

1 Aside from pragmatic barriers to continuing the service, such as moving to a different
2 borough, family commitments, and a lack of willpower, some women reported that the
3 service did not meet their needs in a variety of ways.

4

5 For example, some women reported needing more information and/or structure for the
6 service to work for them.

7

8 *“Maybe if it was writ [sic] out like in a certain way, like you can have these many calories a*
9 *day and then you’re given your list like, of all the foods and how many calories are in the*
10 *food.”*

11 Cathy (disengaged)

12

13 However, other women suggested that they would prefer a less intensive service, or perceived
14 the service to be too prescriptive.

15

16 *“She didn’t seem to grasp that I couldn’t physically write down everything that I’d eaten that*
17 *day, like times and portions sizes. She just kept saying ‘when she [baby] goes to sleep’, but*
18 *that’s when I go to sleep, not write down what I’ve eaten!”*

19 Monica (disengaged)

20

21 Some women expressed a preference for a group service, rather than the one to one home
22 visits, and some women reported wanting to wait until after the baby was born to start
23 thinking about weight management, often citing weight gain as an inevitable consequence of
24 pregnancy.

25

1 *“I thought, if she’s going to come and weigh me, and then, I mean, I’ve put weight on,*
2 *obviously, because the baby’s growing, and I would have put weight on and stuff. And then I*
3 *just thought, well I can’t focus like that.”*

4 Judy (disengaged)

5

6 *Missed opportunities for support*

7

8 Two notable missed opportunities emerged. Firstly, many of the women who had declined
9 the service reported wanting support with weight management. When questioned about what
10 help they would like, answers from many of these women reflected elements of the service
11 itself, including; forming a personalised action plan, goal-setting, healthy recipe ideas, and
12 for support to be flexible and inexpensive.

13

14 *“More recipes to try, things I could cook that were healthy, and help us all being healthy. I*
15 *want to lose weight, things I can eat more or less of, good to have goals, keep me on track,*
16 *rather than doing it on my own.”*

17 Alma (declined)

18

19 Secondly, across both participant groups, regular weight monitoring was mentioned
20 frequently as being desirable, and often cited as a motivational tool. Women expressed
21 confusion or disappointment that they were not weighed regularly by their midwife or even
22 sometimes by the healthy weight advisor.

23

1 *“When I was first pregnant I put on a lot of weight and this time I wanted to monitor my*
2 *weight but I wasn’t weighed...It’s a health issue, you need people to tell you. Without people*
3 *telling you, you just don’t see it.”*

4 Katherine (disengaged)

5

6 Discussion

7

8 Both women who had declined to access a personalised weight management service
9 (MAEYS) during and/or after pregnancy, and women who had initially engaged with
10 MAEYS but had ceased using the service, reported a variety of negative experiences relating
11 to their referral to MAEYS. These included being upset by the referral or initial contact from
12 the service, as well not receiving enough information about the service, and lack of contact
13 from the service after accepting the referral. However, a few women reported a more positive
14 referral experience and these women all reported being happy to be referred to MAEYS. The
15 primary reason given for declining to use MAEYS was feeling that health professionals
16 would make assumptions about referred women’s abilities to care for themselves and their
17 child. Some women preferred to use alternative weight management services or to self-
18 manage their weight. The dominant reason given for ceasing to use MAEYS was that the
19 service did not meet the woman’s needs in terms of format and content. Some of the women
20 who had declined the service reported wanting support with weight management. Many
21 women reported wanting to have their weight gain regularly monitored and some women
22 who had experienced MAEYS were disappointed not to be weighed by their healthy weight
23 advisor.

24

1 There are several strengths of the present study. Firstly, the use of qualitative methods
2 allowed participants to fully explain their experiences and feelings about various aspects of
3 MAEYS, and their decisions relating to whether or not to participate. Further, as most
4 qualitative evaluations of interventions or services solely include the views of service users
5 and providers (Adolfsson, Carlson, Undén, & Rössner, 2002; Morgan, et al., 2010; Stewart,
6 Chapple, Hughes, Poustie, & Reilly, 2008), our approach allowed the views of non-
7 participants and non-completers to be taken into account alongside those of the service users,
8 whose experiences are reported elsewhere (Atkinson et al, in review). For the acceptability
9 of an intervention to be fairly assessed, it is important to explore and understand which
10 aspects of the intervention potential recipients find unacceptable, why this may deter their
11 participation, and the reasons why the intervention fails to keep participants engaged. Finally,
12 while previous qualitative studies have examined the perceived barriers to accessing weight
13 management, healthy eating or physical activity support during pregnancy in the UK (Clarke
14 & Gross, 2004; Olander, Atkinson, Edmunds, & French, 2011) their findings relate to views
15 regarding an imagined service, whereas the present findings relate to actual experiences of an
16 intervention delivered as part of a routinely delivered service. While the pilot phase of
17 MAEYS provided an opportunity to conduct evaluation of the service, most of the PCTs
18 involved planned to continue to offer MAEYS as part of their antenatal and postnatal care
19 pathways in the long term. The evaluation of an ongoing service often has a different focus
20 than the evaluation of a quasi-experimental or controlled trial of a proposed intervention, as
21 described by the Medical Research Council's guidance on developing and evaluating
22 complex interventions (Craig et al, 2008). As such, the goal of the present evaluation was to
23 assess the acceptability of the intervention as routinely delivered, which is usually with lower
24 fidelity than when interventions are delivered in a comparatively controlled research setting.
25 The data gained from the present evaluation has highlighted a number of process and

1 operational issues which may not have been identified by an evaluation of an intervention
2 delivered as part of a research study. For example, specially trained research midwives are
3 often used to recruit women to intervention trials, (e.g. Smith et al, 2010), whereas MAEYS
4 relied on every midwife and health visitor within the PCTs to refer as part of their everyday
5 practice, regardless of their personal knowledge of or interest in weight management. As
6 such, the quality of referrals was variable and had a negative impact on recruitment. Settings
7 included PCTs with differences in population density, ethnicity and deprivation, as well as
8 different organisational structures. As such, the relevance of the experiences and decisions
9 reported in the present evaluation is not restricted to a single locale or PCT, and the findings
10 can be directly applied to future service provision based on the MAEYS pathway.

11

12 Regarding study limitations, our sample sizes were small, in common with much in-depth
13 qualitative research, and women self-selected to participate. Previous research has
14 demonstrated the difficulties in recruiting obese pregnant women to a qualitative study to
15 discuss their experiences (Tierney, et al., 2010). Additionally, it has been shown that
16 attendees of a health service are more likely to respond to a request to participate in research
17 than non-attendees (Sutton, Bickler, Sancho-Aldridge, & Saidi, 1994). Therefore it was not
18 unexpected that recruitment would be challenging, and multiple and varied attempts were
19 made to recruit participants. Nevertheless, the small sample does reduce the reliability of
20 conclusions drawn from the data. Participants were also predominantly White British. It was
21 not possible to purposively sample for the present evaluation as the ethnic background of
22 women was not known prior to contact from the research team. However, providers of
23 MAEYS during the pilot phase reported over-representation of White British women using
24 the service compared to the local population (Baker, 2011). There was, however, significant
25 diversity in ages and parity within the sample.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

The authors believe the present evaluation to be the first of an intervention which exemplifies the NICE guidance for interventions targeting weight management, during and after pregnancy (National Institute for Health and Clinical Excellence, 2010). There are a limited number of qualitative evaluations of the acceptability of maternal obesity interventions (Atkinson et al, in review; Heslehurst et al., 2010a), and those that have been conducted have almost exclusively relied upon data from participants who have been actively involved with the intervention (Claesson, et al., 2008). One recent study has reported the views of obese women who declined a healthy weight service in pregnancy (Olander & Atkinson, in review). The service offered was a free, short term (six week), structured, group-based service, held at community venues. As such the service differs significantly from MAEYS. As might be expected, pragmatic barriers such as an inability to travel to the venue or to take time away from work to attend were the most prevalent reasons for women to decline the service (Olander & Atkinson, in review). Similarly, women who had declined a post-partum healthy lifestyle programme reported doing so due to lack of time, prioritising one’s children over personal health goals and the logistics of attending (Carter-Edwards, et al., 2009). The programme in question was again a structured intervention which required participants to travel to a specific location at regular times. Although these pragmatic barriers were identified in the present study, they were less prevalent. This is likely because MAEYS was designed to be flexible enough to overcome some of these barriers, for example delivering the service in the home, and outside of women’s working hours. Decliners of a group based service also reported a lack of motivation towards weight management, mentioning that they did not want to focus on their weight during pregnancy (Olander & Atkinson, in review). In contrast, decliners of MAEYS mostly expressed wanting support to minimise pregnancy-related weight gain. Some women did state that they wanted to wait until after pregnancy to

1 focus on their weight. As MAEYS is accessible for up to two years postnatally, this service
2 may be more acceptable to some women than one only available during pregnancy. The
3 present study suggests that at least some of the women who declined MAEYS would have
4 found the service acceptable and valuable as they expressed wanting the type of support
5 MAEYS offered, but were deterred from trying the service due to the way the referral was
6 handled or the service was explained. Although participants did not highlight specific terms
7 or language that they found upsetting, rather that they were upset by perceived inferences of
8 the referral itself, the need for training to enable health professionals to communicate
9 sensitively in relation to obesity to encourage long term engagement has previously been
10 highlighted (Smith et al, 2012; Heslehurst et al, 2011). When comparing the present results
11 with the findings from an evaluation of MAEYS based on active service users (Atkinson et
12 al, in review), it is possible to identify differing experiences and preferences which may have
13 reduced uptake and usage of the service, e.g. a poorly handled referral, or a personal
14 preference for group-based interventions. It is also important to recognise that while health
15 promotion is now considered integral to the role of the midwife, it has been noted that by
16 attempting to address a wide range of public health issues as part of maternity care, midwives
17 run the risk of becoming “Jill of all trades and master of none” (Beldon & Crozier, 2005,
18 p.218). Having a service such as MAEYS to provide specialist support may relieve some of
19 the pressure on midwives to help women with weight management, but they still face the
20 challenge of finding the time to sensitively introduce the service to women amongst many
21 competing priorities during antenatal appointments,

22

23 Implications for practice

24

1 Findings suggest that modification of the referral process could increase uptake of MAEYS
2 and similar interventions based on the recommendations of NICE (National Institute for
3 Health and Clinical Excellence, 2010). Similarly, the present findings suggest that premature
4 withdrawal from the service may be reduced if women are given more information about the
5 service at the start, to manage their expectations. For example, fully informing the woman of
6 the referral and the reasons for it could reduce upset. Furthermore, a more detailed
7 explanation of what the service entails could reduce women's perception of the referral as
8 implying concern about their parenting abilities. Lack of confidence in raising the issue of
9 obesity has previously been cited by health professionals as a barrier to recruitment to
10 maternal weight management services (Knight & Wyatt, 2010) and further research is
11 required to understand how health professionals can approach referrals to maximise uptake.
12 However, initially measures could be taken to ensure that referring health professionals have
13 a detailed understanding of the aims of the service, and what is involved, in order that they
14 can effectively communicate this to their patients.

15 Early referral and reducing the time between the referral and first contact from the service
16 may also increase uptake, as some of the women in this study were willing to accept the
17 referral to the service, but when contact from the service was delayed they either forgot about
18 it or suggested it was too late for them to benefit. It also appears that regular weight
19 monitoring is an anticipated, and in many cases, desirable element of the weight management
20 service, and should be offered to all participating women. Further research is needed to
21 explore the healthy weight advisors' barriers to weighing women at their appointments, but
22 health professionals currently reluctant to offer weight monitoring should be encouraged that
23 the women in this study viewed monitoring as a useful tool within a weight management
24 service.

25

1 Importantly, the present results also suggest that while some of the issues identified could be
2 addressed with further tailoring of the service to meet individual needs, some of the reasons
3 given for non-participation or non-completion are not modifiable within the current
4 intervention design. For example, some women simply prefer to use a different format of
5 service, e.g. group-based sessions, or a more rigid diet plan. Service providers should
6 therefore consider increasing the flexibility of the intervention further and offering obese
7 women a choice of formats and services.

8

9 Conclusion

10 This qualitative evaluation of a tailored, home-based weight management intervention for
11 obese women during and after pregnancy which included women who had declined or
12 disengaged from the service has identified several aspects of the intervention which could be
13 modified to increase acceptability. Uptake and continued use of maternal obesity services
14 may be increased by a more sensitive referral process and more detailed explanation of the
15 aims and format of the service. Women who were interested in weight management support
16 during or after pregnancy valued the inclusion of regular weight monitoring and service
17 providers should consider how best to fulfil this need. However, it is also apparent that some
18 obese pregnant and post-natal women are unlikely to want weight management support
19 during this time.

20

21

22

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

References

- Abenhaim, H., Kinch, R., Morin, L., Benjamin, A., & Usher, R. (2007). Effect of prepregnancy body mass index categories on obstetrical and neonatal outcomes. *Archives of Gynecology and Obstetrics*, 275(1), 39-43.
- Adolfsson, B., Carlson, A., Undén, A.-L., & Rössner, S. (2002). Treating obesity: A qualitative evaluation of a lifestyle intervention for weight reduction. *Health Education Journal*, 61(3), 244-258.
- Atkinson, L., Olander, E.K., French, D.P. (in review). Acceptability of a weight management intervention for pregnant and post-natal women with BMI >30kg/m2: A qualitative evaluation of a service delivered in primary care settings which meets current UK public health guidelines. *BMC Pregnancy and Childbirth*, Under review.
- Baker, J. (2011). Developing a care pathway for obese women in pregnancy and beyond. *British Journal of Midwifery*, 19(10), 632-643.
- Barter-Godfrey, S., Taket, A., & Rowlands, G. (2007). Evaluating a community lifestyle intervention: adherence and the role of perceived support. *Primary Health Care Research & Development*, 8(04), 345-354.
- Beldon, A. & Crozier, S. (2005). Health promotion in pregnancy: the role of the midwife. *Perspectives in Public Health*, 125(5), 216-220.
- Braun, V., & Clarke, V. (2006). Using thematic analysis in psychology. [Qualitative Research in Psychology]. *Qualitative Research in Psychology*, 3(2), 77-101.
- Carter-Edwards, L., Ostbye, T., Bastian, L. A., Yarnall, K. S., Krause, K. M., & Simmons, T. J. (2009). Barriers to adopting a healthy lifestyle: insight from postpartum women. *BMC Research Notes*, 2, 161.
- Claesson, I.-M., Josefsson, A., Cedergren, M., Brynhildsen, J., Jeppsson, A., Nyström, F., et al. (2008). Consumer satisfaction with a weight-gain intervention programme for obese pregnant women. *Midwifery*, 24(2), 163-167.
- Clarke, P. E., & Gross, H. (2004). Women's behaviour, beliefs and information sources about physical exercise in pregnancy. *Midwifery*, 20(2), 133-141.
- Craig, P., Dieppe, P., Macintyre, S., Michie, S., Nazareth, I. & Petticrew, M. (2008). Developing and evaluating complex interventions: the new Medical Research Council guidance. *BMJ* 2008;337:a1655
- Davis, D.L., Raymond, J.E., Clements, V., Adams, C., Mollart, L.J., Teate, A.J. & Fourer, M.J. (2012). Addressing obesity in pregnancy: The design and feasibility of an innovative intervention in NSW, Australia. *Women and Birth*, 25, 174-180.
- Griffiths, L. J., Hawkins, S. S., Cole, T. J., & Dezaux, C. (2010). Risk factors for rapid weight gain in preschool children: findings from a UK-wide prospective study. *International Journal of Obesity*, 34(4), 624-632.
- Heslehurst, N., Bell, R., & Rankin, J. (2011). Tackling maternal obesity: The challenge for public health. *Perspectives in Public Health*, 131(4), 161-162.
- Heslehurst, N., Moore, H., Rankin, J., Ells, L. J., Wilkinson, J. R., & Summerbell, C. D. (2010a). How can maternity services be developed to effectively address maternal obesity? A qualitative study. *Midwifery*, doi:10.1016/j.midw.2010.1001.1007.
- Heslehurst, N., Rankin, J., Wilkinson, J. R., & Summerbell, C. D. (2010b). A nationally representative study of maternal obesity in England, UK: trends in incidence and demographic inequalities in 619 323 births, 1989-2007. *International Journal of Obesity*, 34(3), 420-428.
- Knight, B.A. & Wyatt, K. (2010). Barriers encountered when recruiting obese pregnant women to a dietary intervention. *Nursing Times*, 106(32), 20-22.

- 1 Morgan, P. J., Collins, C. E., Plotnikoff, R. C., McElduff, P., Burrows, T., Warren, J. M., et
2 al. (2010). The SHED-IT community trial study protocol: a randomised controlled
3 trial of weight loss programs for overweight and obese men. *BMC Public Health*,
4 10(1), 701.
- 5 National Institute for Health and Clinical Excellence. (2010). *NICE public health guidance*
6 *27: Dietary interventions and physical activity interventions for weight management*
7 *before, during and after pregnancy*. London.
- 8 National Institute for Health and Clinical Excellence. (2011). *NICE Weight management*
9 *before, during and after pregnancy (CMG36)* National Institute for Health and
10 Clinical Excellence.
- 11 Olander, E., Atkinson, L., Edmunds, J., & French, D. (2011). The views of pre- and post-
12 natal women and health professionals regarding gestational weight gain: An
13 exploratory study. *Sexual and Reproductive Healthcare*, 2, 43 - 48.
- 14 Olander, E.K. & Atkinson, L. Can't or won't: obese women's reasons for not attending a
15 weight management service during pregnancy. (*In review*).
- 16 Pirkola, J., Pouta, A., Bloigu, A., Hartikainen, A.-L., Laitinen, J., Jarvelin, M.-R., et al.
17 (2010). Risks of Overweight and Abdominal Obesity at Age 16 Years Associated
18 With Prenatal Exposures to Maternal Prepregnancy Overweight and Gestational
19 Diabetes Mellitus. *Diabetes Care*, 33(5), 1115-1121.
- 20 Sheikh, A. E., Malik, A., & Gardosi, J. (2010). *West Midlands Perinatal and Infant Mortality*
21 *2008-9*. Retrieved from http://www.perinatal.nhs.uk/pnm/KHD_2008-9.pdf.
- 22 Smith, D.M., Cooke, A., & Lavendar, T. (2012). Maternal obesity is the new challenge; a
23 qualitative study of health professionals' views towards suitable care for pregnant
24 women with a Body Mass Index (BMI) ≥ 30 kg/m². *BMC Pregnancy & Childbirth*,
25 12:157 doi:10.1186/1471-2393-12-157
- 26 Smith D.M., Whitworth M., Sibley C., Taylor W., Gething J., Chmiel C. & Lavender T.
27 (2010). The design of a community lifestyle programme to improve the physical and
28 psychological well-being of pregnant women with a BMI of 30 kg/m² or more. *BMC*
29 *Public Health*, 10:284 doi:10.1186/1471-2458-10-284
- 30 Sørensen, J. B., Skovgaard, T., & Puggaard, L. (2006). Exercise on prescription in general
31 practice: A systematic review. *Scandinavian Journal of Primary Health Care*, 24(2),
32 69-74.
- 33 Stewart, L., Chapple, J., Hughes, A. R., Poustie, V., & Reilly, J. J. (2008). The use of
34 behavioural change techniques in the treatment of paediatric obesity: qualitative
35 evaluation of parental perspectives on treatment. *Journal of Human Nutrition and*
36 *Dietetics*, 21(5), 464-473.
- 37 Sutton, S., Bickler, G., Sancho-Aldridge, J., & Saidi, G. (1994). Prospective study of
38 predictors of attendance for breast screening in inner London. *Journal of*
39 *Epidemiology and Community Health*, 48(1), 65-73.
- 40 Tierney, S., Butterfield, C., Stringer, E., Fox, J. R. E., McGowan, L., & Furber, C. (2010).
41 Difficulties in recruiting pregnant women with eating or weight issues. *British*
42 *Journal of Midwifery*, 18(11), 717-723.
- 43 West, C. (2010) Developing a support service for overweight women. *The Practising*
44 *Midwife*, 13(10), 19-21
- 45
46
47
48