



City Research Online

City, University of London Institutional Repository

Citation: MacFarlane, A.J., Blondel, B., Mohangoo, A.D., Cuttini, M., Nijhuis, J., Novak, Z., Olafsdottir, H.S., Zeitlin, J. & the Euro-Peristat Scientific Committee, . (2015). Wide differences in mode of delivery within Europe: risk-stratified analyses of aggregated routine data from the Euro-Peristat study. *BJOG: an International Journal of Obstetrics and Gynaecology*, 123(4), pp. 559-568. doi: 10.1111/1471-0528.13284

This is the published version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/6985/>

Link to published version: <https://doi.org/10.1111/1471-0528.13284>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Wide differences in mode of delivery within Europe: risk-stratified analyses of aggregated routine data from the Euro-Peristat study

AJ Macfarlane,^a B Blondel,^b AD Mohangoo,^c M Cuttini,^d J Nijhuis,^e Z Novak,^f HS Ólafsdóttir,^g J Zeitlin,^b the Euro-Peristat Scientific Committee

^a Centre for Maternal and Child Health Research, City University London, London, UK ^b INSERM, Obstetrical Perinatal and Paediatric Epidemiology Research Team, Centre for Epidemiology and Biostatistics (U1153), Paris-Descartes University, Paris, France ^c Department of Child Health, TNO, the Netherlands Organisation for Applied Scientific Research, Leiden, the Netherlands ^d Research Unit of Perinatal Epidemiology, Bambino Gesù Children's Hospital, Rome, Italy ^e Department of Obstetrics and Gynaecology, GROW School of Oncology and Developmental Biology, Maastricht University Medical Centre, Maastricht, the Netherlands ^f Perinatology Unit, University Medical Centre, Ljubljana University, Ljubljana, Slovenia ^g Department of Obstetrics and Gynaecology, Landspítali University Hospital, Landspítali v/ Hringbraut, Iceland

Correspondence: Professor A Macfarlane, School of Health Sciences, City University London, 1 Myddelton Street, London EC1R 1UW, UK. Email A.J.Macfarlane@city.ac.uk

Accepted 3 November 2014. Published Online 6 March 2015.

Objective To use data from routine sources to compare rates of obstetric intervention in Europe both overall and for subgroups at higher risk of intervention.

Design Retrospective analysis of aggregated routine data.

Setting Thirty-one European countries or regions contributing data on mode of delivery to the Euro-Peristat project.

Population Births in participating countries in 2010.

Methods Countries provided aggregated data about overall rates of obstetric intervention and about caesarean section rates for specified subgroups.

Main outcome measures Mode of delivery.

Results Rates of caesarean section ranged from 14.8% to 52.2% of all births and rates of instrumental vaginal delivery ranged from 0.5% to 16.4%. Overall, there was no association between rates of instrumental vaginal delivery and rates of caesarean section, but similarities were observed between some countries that are geographically close and may share common traditions of practice. Associations were observed between caesarean section rates for

women with breech and vertex births and with singleton and multiple births but patterns of association for women who had and had not had previous caesarean sections were more complex.

Conclusions The persisting wide variations in caesarean section and instrumental vaginal delivery rates point to a lack of consensus about practice and raise questions for further investigation. Further research is needed to explore the impact of differences in clinical guidelines, healthcare systems and their financing and parents' and professionals' attitudes to care at delivery.

Keywords Breech birth, caesarean section, Euro-Peristat, instrumental vaginal delivery, international comparisons, mode of delivery, parity, repeat caesarean section, twins.

Linked article This article is commented on by FG Mariona. To view this mini commentary visit <http://dx.doi.org/10.1111/1471-0528.13354>. This article is also commented on by M Mathai. To view this mini commentary visit <http://dx.doi.org/10.1111/1471-0528.13344>.

Please cite this paper as: Macfarlane AJ, Blondel B, Mohangoo AD, Cuttini M, Nijhuis J, Novak Z, Ólafsdóttir HS, Zeitlin J, the Euro-Peristat Scientific Committee. Wide differences in mode of delivery within Europe: risk-stratified analyses of aggregated routine data from the Euro-Peristat study. BJOG 2015; DOI: 10.1111/1471-0528.13284.

Introduction

The recommendation from a World Health Organization (WHO) conference in 1985 that 'Countries with some of the lowest perinatal mortality rates in the world have caesarean section rates of <10%. There is no justification for

any region to have a rate higher than 10–15%',¹ is still frequently cited even though rates of obstetric intervention have continued to rise substantially in both high- and middle-income countries.^{2–11} Data from the Organisation for European Co-operation and Development (OECD) show a continuing rise in caesarean section rates in most member

countries, despite signs of flattening off in a few countries with high rates.^{10,11} Consequences of the rise in caesarean section rates include associations with raised risks for mothers and babies, including placenta accreta, placenta praevia, placental abruption and stillbirth in subsequent pregnancies.^{12–15} A number of factors have been cited as possible explanations, including fear of litigation, financial incentives related to methods of payment,^{16–18} women's requests for caesarean section¹⁹ and the perception that caesarean section is a safe procedure.²⁰

Caesarean sections are clearly necessary in some high-risk situations, such as placenta praevia or fetal distress. For other situations, there is ongoing debate about the systematic use of caesarean section, for example for breech presentation,^{21–25} multiple births^{26–31} or women with previous caesarean section.^{20,32,33}

A succession of comparative analyses, from the 1980s onwards, have shown major variations between high-income countries in their rates of obstetric intervention.^{2,3,5,6,10,11,34} These include the first Euro-Peristat study, which compiled routinely collected data for the 15 members of the European Union in 2000. This found that rates of caesarean section varied from 11.7% to 30.8% and rates of instrumental vaginal delivery, using either forceps or vacuum extraction, ranged from 4.9% to 15.0%.³⁴

The subsequent Euro-Peristat project extended data collection to the 25 states that were members of the European Union in 2005, plus Norway.³⁵ It was based on data for 2004 and found that rates of caesarean section had increased and differences had widened, ranging from 14.4% to 37.8%. Induction rates ranged even more widely, from 5.9% to 37.9%.

It is well established that caesarean rates vary by a number of factors, notably parity, previous caesarean section, fetal presentation and multiplicity,^{36–41} but to date international comparisons based on aggregated data have not stratified methods of delivery by these factors.

This study aims to use aggregated population-based data from routine sources to explore differences between intervention rates at delivery for European countries, and the extent to which clinicians in countries with high overall caesarean rates were also more likely to intervene in specific situations.

Methods

Data sources

Data come from the Euro-Peristat project, a collaboration between 26 member states of the European Union and Norway, Iceland and Switzerland, to assess perinatal health in Europe using a common set of ten core and 20 recommended perinatal health indicators.^{42,43} A distinctive feature of the Euro-Peristat project is the collection of data by

prespecified subgroups to improve the comparability and usefulness of the indicators.

Each participant was requested to provide nationally aggregated population-based data from its routine data collection systems for the year 2010 and these were used to compile the full set of indicators, including data about women's demographic characteristics, care provided and outcome for mothers and babies. These are available on the Euro-Peristat web site.⁴³ In some member states, data collection is devolved to constituent regions or countries. In Belgium, Flanders, Brussels and Wallonia provided the data from separate regional sources and England, Wales, Scotland and Northern Ireland provided data for the countries of the UK. France provided data from the National Perinatal Survey, a nationally representative survey of a sample of 14 000 births in 2010, which was the most recent of its series of such surveys.⁴⁴

Participants were also asked to document the way data were recorded in their countries' routine systems and the definitions they used. The implications of the observed differences in these were discussed at collaborators' meetings, and are documented in Euro-Peristat publications.^{34,43,45}

This analysis includes 31 countries and regions that contributed data about mode of delivery. Of these, 27 contributed data about at least one specified subgroup for the mode of delivery indicator.

Definitions

The mode of delivery indicator was defined as the percentage distribution of total births, live and stillborn, by mode of delivery for all births. This was then subdivided by mother's parity, whether she had had a previous caesarean section, fetal presentation and plurality.

In Poland, Portugal, England and Wales, rates were reported per woman. This may result in slight underestimates of operative deliveries, as two or more multiple births to the same woman were counted only once, but the impact of this was minimal.

Mode of delivery was subdivided into spontaneous vaginal delivery, operative vaginal delivery and two categories of caesarean section. Countries differed in the ways that they classified caesarean section. Some countries subdivided them according to whether they were undertaken before or during labour. Others used the subdivision into elective caesarean section, which includes all those planned before the onset of labour and so was likely to include a few that took place after labour had started, and emergency or unplanned caesarean section, which could include a few caesareans in emergency situations before labour started. The definitions used in each country were documented but in the Euro-Peristat tables, data about elective caesareans were grouped with caesarean sections before labour and emergency sections were grouped with those undertaken in labour.

Some countries were excluded from specific subgroup analyses because of questions about the validity of the data. Malta was excluded from analyses of repeat caesarean sections because the percentages of previous caesareans were unlikely and Lithuania was excluded from analyses by presentation because only 0.4% of births were recorded as breech.

Statistical analysis

Associations between caesarean section rates and rates of instrumental vaginal delivery were examined, as were associations between the percentages of caesarean sections to women with and without previous caesarean section and associations between caesarean section rates for breech and vertex births and between rates for twins and singletons. Statistical associations were tested by calculating Spearman's ranked correlations, using IBM SPSS STATISTICS, version 21 (IBM, Armonk, NY, USA).

Results

Mode of delivery

The numbers of births ranged from under 5000 in Malta and Iceland to well over 500 000 in England, Italy and Germany (Table 1). There was wide variation in overall rates of caesarean section in Europe in 2010, from 14.8% in Iceland to 52.2% in Cyprus, with a median value of 25.2%.

By far the highest rates of caesarean section that were undertaken or planned before labour were 38.8% in Cyprus and 24.9% of births in Italy. Elsewhere, rates ranged from 6.6% in Finland to 17.9% in Luxembourg. In Romania, 33.1% of births were by caesarean section undertaken in an emergency or in labour. This was very much higher than that for any other country. Rates for other countries ranged less than for elective rates, from 8.6% in Sweden to 16.7% in Malta. Hence, if the extreme rates for Cyprus, Italy and Romania were disregarded, the range of rates of caesarean sections that were planned or undertaken before labour was wider than the range of rates of caesarean section that were classified as emergency or undertaken in labour.

Rates of instrumental vaginal delivery also varied widely, from 0.5% in Romania to 16.4% in Ireland, with a median value of 7.5%. There was no statistical association between rates of caesarean section and instrumental vaginal delivery (Figure 1), but the data showed considerable differences in practice. The countries with very low rates of instrumental vaginal delivery, $\leq 2\%$ had caesarean section rates ranging from 23.1% to 34.0%. When rates for these countries were excluded, there was still no statistical association between rates of caesarean section and spontaneous vaginal birth. In the 24 countries that could subdivide caesarean rates, there was no statistical association,

positive or negative, between instrumental vaginal delivery and rates of caesarean sections that were emergency or undertaken in labour. This was still the case when the countries that made minimal use of operative vaginal delivery were excluded (not shown).

Caesarean section rates by subgroup

Caesarean section rates for babies by parity, by whether or not women had had a previous caesarean section, as well as for babies in vertex and breech presentations and for singleton and twin babies, are shown in Table 2.

Parity and previous caesarean section

Countries with high caesarean section rates for births to primiparous women tended to have high rates among births to multiparous women.

Caesarean section rates among women who had a previous caesarean section were high overall but ranged from between 45% and 55% in The Netherlands, Norway, Finland and Iceland to 92.9% in Latvia and 93.5% in Cyprus (Table 2, Figure 2). Caesarean section rates among women without a previous caesarean section ranged from 11.3% in The Netherlands to 28.9% in Italy. This was highly correlated with rates among those with a previous caesarean section, but the degree of association varied.

Repeat section rates were high in Italy, Cyprus and Malta, which also had high rates among births to women without a previous caesarean section, but were also high in Latvia and Lithuania where primary caesarean section rates were lower. Among the countries that had lower rates for births to women without previous caesarean sections, repeat caesarean section rates ranged more widely, from <50% to 80% (Figure 2).

Breech presentation

Breech deliveries accounted for around 4% of all births. In the 21 countries that could contribute data, a high degree of correlation was found between rates of caesarean section for breech births and rates for all vertex births (Figure 2). More than 80% of breech babies were delivered by caesarean section in 16 countries. In nine of these, rates were over 90%. Nevertheless, caesarean section rates for breech births varied widely in countries that had low rates for vertex births.

Twin and singleton births

Caesarean section rates were high for twin births, but here again there were wide variations in practice. The strong association between caesarean section rates for twin births with rates for all singleton births is shown in Figure 2. It also shows that, on the other hand, caesarean section rates for twin births varied very widely between countries that had low rates for singletons.

Table 1. Births in participating countries by mode of onset of labour and delivery

Country/coverage	Number of total births	Number with mode of delivery stated	Percentage of total births with mode stated: Mode of delivery				
			Vaginal spontaneous	Vaginal instrumental	Caesarean elective/before labour	Caesarean emergency/during labour	Caesarean section, all
Belgium							
Brussels	25 098	25 009	71.5	8.3	9.7	10.4	20.2
Flanders	69 976	69 976	69.6	10.4	11.3	8.8	20.1
Wallonia	38 430	38 310	71.6	7.5	10.4	10.5	20.9
Czech Republic	114 406	113 917	75.1	1.8	12.7	10.4	23.1
Denmark	63 513	63 460	71.0	6.9	9.4	12.8	22.1
Germany	637 664	619 903	62.2	6.4	15.4	15.9	31.3
Estonia	15 884	15 884	74.0	4.9	7.8	13.4	21.2
Ireland	75 595	75 564	56.6	16.4			27.0
Greece							
Spain	478 037	478 037					26.3
France	14 903	14 731	66.9	12.1	11.3	9.7	21.0
Italy	547 568	546 133	58.6	3.4	24.9	13.1	38.0
Cyprus	8603	8591	45.3	2.5	38.8	13.4	52.2
Latvia	19 246	19 246	74.0	1.6	11.5	13.0	24.4
Lithuania	30 977	30 977	73.5	1.3	9.4	15.8	25.2
Luxembourg	6560	6560	59.9	10.2	17.9	12.1	30.0
Hungary							
Malta	4036	4036	63.0	3.9	16.4	16.7	33.1
The Netherlands	178 838	177 607	72.9	10.0	7.7	9.4	17.0
Austria	78 989	78 989	65.6	5.6			28.8
Poland	402 826	402 578	64.6	1.4			34.0
Portugal	100 280	100 130	48.8	14.9			36.3
Romania	174 692	174 692	62.5	0.5	3.8	33.1	36.9
Slovenia	22 416	22 404	77.5	3.5	8.3	10.8	19.1
Slovak Republic							
Finland	61 371	61 368	74.5	8.6	6.6	10.2	16.8
Sweden	114 955	114 955	74.9	7.6	8.9	8.6	17.5
United Kingdom							
England	622 303	661 926	62.8	12.6	9.9	14.7	24.6
Wales	32 649	32 523	61.3	12.6	11.1	15.0	26.1
Scotland	57 438	57 166	59.7	12.6	11.9	15.9	27.8
Northern Ireland	25 359	24 884	57.0	13.1	14.6	15.2	29.9
Iceland	4903	4903	78.6	6.5			14.8
Norway	62 591	62 591	73.0	9.9	6.6	10.5	17.1
Switzerland	80 276	79 565	55.8	11.0			33.1
Median	62 591	62 591	66.2	7.5	10.7	12.9	25.2
Minimum	4036	4036	45.3	0.5	3.8	8.6	14.8
Maximum	637 664	661 926	78.6	16.4	38.8	33.1	52.2
Number	31	31	30	30	24	24	31

Discussion

Main findings

This analysis confirms the wide variations in overall caesarean section rates reported elsewhere, as well as in specific subgroups. Caesarean section rates among women with a previous caesarean section were highly correlated

statistically with rates for births to women without a previous caesarean section. In addition, there were strong statistical associations between caesarean rates for breech and vertex births and singleton and twin births. The patterns of association are more complex, however, with a wide range of rates, despite the highly significant rank correlations. There was no inverse association between




Figure 1. Comparison between caesarean section and vaginal instrumental birth rates, 2010.

rates of operative vaginal delivery and overall caesarean section rates.

Strengths and limitations

These tabulations of mode of delivery by parity, previous caesarean section, presentation and multiplicity are not found in other international reports based on data that are collected routinely within member states. Routine data are collected every year for countries' own purposes and the data in the Euro-Peristat report came from 129 different systems. As they cover the whole population of countries and regions, they avoid the selection biases that can arise in studies based on samples of individual maternity units.^{4,8} With some well-documented exceptions, routine data are of good quality and virtually complete. Most of the gaps arise where particular data items are not included in specific countries' systems.^{33,43,45}

In the Euro-Peristat project, steps are taken to make comparisons as reliable as possible by using harmonised definitions, compiling numbers as well as rates to enable checking for inconsistencies and collecting information about missing data.⁴³ The clinicians and data experts on the Euro-Peristat Scientific Committee have been actively involved in discussing and interpreting the results.

On the other hand, with aggregated data, it is not possible to adjust operative delivery rates for known risk factors:

sociodemographic factors such as mothers' ages, individual or area-based measures of socio-economic status or clinical complications. These may not explain observed differences between countries, however. For example, an analysis of national age-specific caesarean rates for OECD countries found that adjusting caesarean section rates for age made little difference to the wide variation observed.⁴⁶

Risk-adjusted analyses to compare rates for maternity units in England have found that known risk factors explained only a small proportion of the variance, even though they explained a higher proportion of variation in emergency than elective caesarean section rates,^{37–40} but the authors pointed out that their conclusions may not necessarily apply elsewhere. An analysis of data about deliveries of low-risk women in France found differences in intervention rates for units after adjustment for mothers' and unit characteristics. It found higher rates of instrumental vaginal delivery in units with over 3000 births and higher rates of any intervention in private units compared with public units.¹⁷ Similar analyses within countries with different levels of intervention or with separate public and private systems of health care might yield further information about factors that can influence intervention rates.

Studies that collect data about population or unit-based samples of individual women and their babies^{4,8,47} can define their data items prospectively and thus consistently, but are expensive. This restricts either their sample size or frequency. The WHO surveys of practice included only small numbers of hospitals in each participating country.^{4,8} The National Caesarean Section Sentinel Audit collected more detailed data than routine systems,⁴⁷ but was expensive and has not been repeated. As with studies using routinely collected data, known risk factors explained only a small proportion of the variance between units and the cost of replicating it in every country in Europe would be prohibitive.

Interpretation

The differences observed raise questions about why there are such wide variations in clinical practice.

In some specific situations, the need for intervention is clear. For others there is ongoing debate, for example about the use of caesarean section for breech presentation,^{21–25, 48–50} multiple births^{26–31} and women with previous caesarean section.^{20,30,31,51,52} The data presented here suggest that for breech births and twin births, practice is associated with factors that influence the overall rate of caesarean section, although the extent of association can differ between countries.

The data do suggest that there are some groups of countries with common or similar traditions, and that identifying these could explain some similarities in practice, such as the low rates of operative vaginal delivery in some

Table 2. Caesarean section rates in participating countries by parity, fetal presentation and multiplicity

Country/ coverage	Caesarean sections as percentages of total births in category for which data are available							
	Nulliparous women	Multiparous women	Women with previous caesarean section	Women without previous caesarean section	Vertex presentation	Breech presentation	Singleton birth	Multiple birth
Belgium								
Brussels	22.0	18.8	64.0	15.1	15.8	85.0	18.2	59.7
Flanders	21.9	18.5	66.2	15.2	15.9	89.7	18.7	53.7
Wallonia	22.7	19.5	66.6	15.4	16.2	86.8	19.6	58.1
Czech Republic	26.0	20.4	72.1	19.0	19.4	94.0	21.9	78.4
Denmark	23.8	20.8	66.8	16.1	18.7	88.6	20.5	59.6
Germany	35.9	26.5	64.2	26.6	27.3	91.0	29.6	74.8
Estonia	23.4	19.5	70.8	16.1		89.4	19.8	65.9
Ireland	28.5	26.0					25.7	63.6
Greece								
Spain	29.4	24.8					25.4	69.4
France	23.2	19.0	65.2	15.6	17.7	75.2	19.9	54.8
Italy	37.3	36.6	89.7	28.9	35.1	93.3	36.5	85.6
Cyprus	57.0	47.3	93.5	28.5	49.1	94.5	49.9	91.6
Latvia	25.3	23.6	92.9	18.1	21.5	82.7	23.4	65.3
Lithuania	27.3	23.3	89.2	19.5	25.2		24.2	59.5
Luxembourg	32.7	27.5	71.5	24.1	25.8	91.8	28.2	77.1
Hungary								
Malta	33.3	32.9	78.8	26.4	31.0	96.1	30.3	98.6
The Netherlands	19.5	14.8	52.9	11.3	13.6	76.1	16.1	43.9
Austria	30.5	27.4					26.8	83.9
Poland								
Portugal								
Romania								
Slovenia	21.4	16.7	78.7	15.3	15.3	82.3	17.5	58.8
Slovak Republic								
Finland	15.8	13.8	47.6	13.8	14.1	73.2	15.8	49.4
Sweden	19.1	15.5			11.5	87.4	16.4	53.6
United Kingdom								
England	24.9	24.4	70.3	18.0			23.9	62.6
Wales								
Scotland	28.5	27.2	79.1	21.4	23.5	100.0	26.4	73.3
Northern Ireland	30.5	20.5					28.6	70.2
Iceland	15.4	14.4	47.6		12.6	91.2	14.3	31.1
Norway	19.1	15.6	54.2	13.6	14.4	69.0	16.0	47.4
Switzerland					27.8	95.4	31.4	77.5
Median	24.9	20.5	70.3	17.1	18.2	89.6	22.6	63.1
Minimum	15.4	13.8	47.6	11.3	11.5	69.0	14.3	31.1
Maximum	57.0	47.3	93.5	28.9	49.1	100.0	49.9	98.6
Number	27	27	21	20	22	22	28	28

countries in the east of Europe and the overall low levels of intervention in the Nordic countries and the Netherlands.

A number of approaches could be used to investigate the differences in practice documented here. A review of national policies and guidelines would be useful for comparing national policies with data about practice. It

would have to take account of the fact that, where within-country data are available, they show considerable variations between maternity units within the same healthcare system with common guidelines and policies.^{17,37–40} Where individual level data are available, constructing the ten groups defined by Robson,^{41,53} might be a useful approach




Figure 2. Comparison between caesarean rates for women with previous caesarean section and rates for women with no previous caesarean section, between rates for breech births and vertex births, and rates for twin births and singleton births

to identifying categories which differ, although many EU countries do not record all the data items required.

There is growing evidence from surveys and other research that health system issues, notably differences in the organisation of health care and methods of payment, can influence choice of mode of delivery, both in Europe^{16–18} and more widely,^{54–56} with higher rates in the private sector. Comparisons of the content and scope of obstetric and midwifery education and the roles and responsibilities of midwifery and obstetric staff would also be informative. Women's requests for caesarean section have been cited as a reason for the rising rates but only a minority of women in a wide range of countries have expressed a preference for caesarean section.^{57–59} Studies in a number of countries have shown that many women who ended up having caesarean section would have preferred vaginal delivery and are influenced by models of care and healthcare providers' preferences.^{54, 60–63} The results of the WHO global surveys of maternal and perinatal health in Latin America and Asia have pointed to the potential dangers and excessive costs associated with caesarean sections that are clinically unnecessary.^{4,8,9}

Any initiatives to counter trends towards higher rates of clinically unnecessary obstetric intervention need monitoring using routine data, so an ongoing European perinatal monitoring system is needed. Using the subgroups in Euro-Peristat to monitor trends over time in mode of delivery is more informative than simply monitoring over-

all trends in caesarean section. Future studies could add additional risk subgroups, such as the ten groups defined by Robson,^{41,63} but to do this, some countries would have to expand the range of data items that they record routinely.

Conclusions

The persisting wide differences in mode of delivery in Europe are a cause for concern because they point to a lack of consensus about best practice. Research is needed to investigate the reasons for these differences, including comparing the basis for formulation of and adherence to clinical guidelines, exploring the impact of differences in healthcare systems and their financing and parents' and professionals' attitudes to care at delivery.

Disclosure of interest

The authors declare that they have no competing interests to disclose.

Contribution to authorship

AJM and BB discussed the design of the paper, ADM compiled the data from data contributed by all members of the Euro-Peristat group, AJM drafted the paper. All the authors commented on drafts of the paper and approved the final version. JZ had overall scientific responsibility for the project.

Details of ethics approval

This study was based on aggregated routinely collected data, so ethics approval was not required.

Funding

The results from this study are based on data from the Euro-Peristat project, a European project for monitoring and evaluating perinatal outcomes at a European level. The Euro-Peristat project received funding from the European Union in the framework of the Health Programme (grant number: 20101301). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

This article drew on the work of the many people who contributed to the European Perinatal Health Report: Health and Care of Pregnant Women and Babies in Europe in 2010. They include statisticians, researchers, clinicians, administrators and others from each of the collaborating countries who compiled aggregated data for their countries and sent them to Euro-Peristat. They are too numerous to list here, but their names can be found online at Appendix S1 and in Appendix A1 of the European Health Report at www.europeristat.com. The authors would like to thank them for their contributions.

The Euro-Peristat scientific committee

Gerald Haidinger (Austria), Sophie Alexander (Belgium), Pavlos Pavlou (Cyprus), Petr Velebil (Czech Republic), Anne-Marie Nybo Andersen (Denmark), Luule Sakkeus (Estonia), Mika Gissler (Finland), Béatrice Blondel (France), Nicholas Lack (Germany), Aris Antsaklis (Greece), István Berik (Hungary), Helga Sól Ólafsdóttir (Iceland), Sheelagh Bonham (Ireland), Marina Cuttini (Italy), Janis Misins (Latvia), Jone Jaseliuniene (Lithuania), Yolande Wagener (Luxembourg), Miriam Gatt (Malta), Jan Nijhuis (Netherlands), Kari Klungsoyr (Norway), Katarzyna Szamotulska (Poland), Henrique Barros (Portugal), Mihai Horga (Romania), Jan Cap (Slovakia), Živa Novak-Antolic (Slovenia), Francisco Bolívar (Spain), Karin Gottvall (Sweden), Sylvie Berrut (Switzerland), Alison Macfarlane (United Kingdom). Project coordination: Jennifer Zeitlin, Marie Delnord, Ashna Mhangoo.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Appendix S1. Acknowledgements. ■

References

- 1 World Health Organisation. Appropriate technology for birth. *Lancet* 1985;326:436–7.
- 2 Notzon FC, Placcek PJ, Taffel SM. Comparisons of national caesarean section rates. *N Engl J Med* 1987;316:386–9.
- 3 Macfarlane AJ, Chamberlain G. What is happening to caesarean section rates? *Lancet* 1993;342:1005–6.
- 4 Villar J, Valladoes E, Wojdyla D, Zavaleta N, Carroli G, Velazco A, et al. Caesarean delivery rates and pregnancy outcomes: the 2005 WHO global survey on maternal and perinatal health in Latin America. *Lancet* 2006;367:1819–29.
- 5 Althabe FA, Sosa C, Belizan JM, Gibbons L, Jacquerioz F, Bergel E. Caesarean section rates and maternal and neonatal mortality in low, medium and high-income countries: an ecological study. *Birth* 2006;33:270–6.
- 6 Betrán AP, Merialdi M, Lauer JA, Bing-Shun W, Thomas J, Van Look LP, et al. Rates of caesarean section: analysis of global, regional and national estimates. *Paediatr Perinat Epidemiol* 2007;21:98–113.
- 7 Baldo MH. Caesarean section in countries of the Eastern Mediterranean Region. *East Mediterr Health J* 2008;14:470–88.
- 8 Lumpiganon P, Laopaiboon M, Gulmezoglu AM, Souza JP, Taneepanichskul S, Ruyan P, et al. Method of delivery and pregnancy outcomes in Asia: the WHO global survey on maternal and perinatal health 2007–08. *Lancet* 2010;375:490–9.
- 9 Belizan JM, Althabe F, Cafferata ML. Health consequences of the increasing caesarean section rates. Commentary. *Epidemiology* 2007;18:485–6.
- 10 OECD. Health data 2011. [http://dotstat.oecd.org/Index.aspx?DataSetCode=HEALTH_PROC] Accessed 16 April 2012.
- 11 Declercq E, Young R, Cabral H, Ecker J. Is a rising caesarean delivery rate inevitable? Trends in industrialised countries, 1987 to 2007. *Birth* 2011;38:99–104.
- 12 Yang Q, Wen S, Oppenheimer L, Chen X, Black D, Gao J, et al. Association of caesarean delivery for first birth with placenta praevia and placental abruption in second pregnancy. *BJOG* 2007;114:609–13.
- 13 Gray R, Quigley MA, Hockley C, Kurinczuk JJ, Goldacre M, Brocklehurst P. Caesarean delivery and risk of stillbirth in subsequent pregnancy: a retrospective cohort study in an English population. *BJOG* 2007;114:264–70.
- 14 Deneux-Tharoux C, Carmona E, Bouvier-Colle MH, Breart G. Postpartum maternal mortality and cesarean delivery. *Obstet Gynecol* 2006;108:541–8.
- 15 Jackson S, Fleege L, Fridman M, Gregory K, Zelop C, Olsen J. Morbidity following primary caesarean delivery in the Danish National Birth Cohort. *Am J Obstet Gynecol* 2012;206:139.e1–5.
- 16 Bertolini R, Di Lallo D, Spadea T, Perucci C. Caesarean section rates in Italy by payment mode: an analysis based on birth certificates. *Am J Public Health* 1992;82:257–61.
- 17 Coulm B, Le Ray C, Lelong N, Drewniak N, Zeitlin J. Obstetric interventions for low-risk pregnant women in France: do maternity unit characteristics make a difference? *Birth* 2012;39:183–91.
- 18 Grytten J, Monkerud L, Hagen TP, Sørensen R, Eskild A, Skau I. The impact of hospital revenue on the increase in Caesarean sections in Norway. A panel data analysis of hospitals 1976–2005. *BMC Health Serv Res* 2011;11:267.
- 19 Habiba M, Kaminski M, Da Frè M, Marsal K, Bleker O, Librero J, et al. Caesarean section on request: a comparison of obstetricians' attitudes in eight European countries. *BJOG* 2006;113:647–56.
- 20 American College of Obstetricians and Gynecologists. Vaginal birth after previous cesarean delivery. *Obstet Gynecol* 2010;1:450–63.

- 21 Hannah ME, Hannah WJ, Hewson SA, Hodnett ED, Saigal S, Willan AR. Planned caesarean section versus planned vaginal birth for breech presentation at term: a randomised multicentre trial: term breech trial collaborative group. *Lancet* 2000;21:1375–83.
- 22 Glezerman M. Five years to the term breech trial: the rise and fall of a randomised controlled trial. *Am J Obstet Gynecol* 2006;194:20–5.
- 23 Hannah ME, Whyte H, Hannah WJ, Hewson S, Amankwah K, Cheng M, et al. Maternal outcomes at 2 years after planned caesarean section versus planned vaginal birth for breech presentation at term: The international randomized Term Breech Trial. *Am J Obstet Gynecol* 2004;191:917–27.
- 24 Whyte H, Hannah ME, Saigal S, Hannah WJ, Hewson S, Amankwah K, et al. Outcomes of children at 2 years after planned caesarean birth versus planned vaginal birth for breech presentation at term: The International Randomized Term Breech Trial. *Am J Obstet Gynecol* 2004;191:864–71.
- 25 Kotaska A. Commentary: routine cesarean section for breech: the unmeasured cost. *Birth* 2011;38:162–4.
- 26 National Collaborating Centre for Women's and Children's Health. *Caesarean Section. Commissioned by the National Institute for Health and Clinical Excellence*. London: RCOG Press, 2011.
- 27 Vayssière C, Benoist G, Blondel B, Deruelle P, Favre R, Gallot D, et al. Twin pregnancies: guidelines for clinical practice from the French College of Gynaecologists and Obstetricians. *Eur J Obstet Gynecol Reprod Biol* 2011;156:12–7.
- 28 Hofmeyr JG, Barrett JF, Crowther CA. Planned caesarean section for women with a twin pregnancy. *Cochrane Database Syst Rev* 2, 2011. 00075320-100000000-05226.
- 29 Dodd JM, Deussen AR, Grivell RM, Crowther CA. Elective birth at 37 weeks' gestation for women with an uncomplicated twin pregnancy. *Cochrane Database of Systematic Reviews*. 2, 2014. 00075320-100000000-02641.
- 30 Rossi AC, Mullin PM, Chmait RH. Neonatal outcomes of twins according to birth order, presentation and mode of delivery: a systematic review and meta-analysis [Review]. *BJOG* 2011;118:523–32.
- 31 Barrett JF, Hannah ME, Hutton EK, Willan AR, Allen AC, Armson BA, et al. A randomized trial of planned cesarean or vaginal delivery for twin pregnancy. *N Engl J Med* 2013;369:1295–305.
- 32 Dodd JM, Crowther CA. Elective repeat caesarean section versus induction of labour for women with a previous caesarean birth. *Cochrane Database Syst Rev* 2006;4, Art. NO. CD004906.
- 33 Dodd JM, Crowther CA, Huertas E, Guise JM, Horey D. Planned elective caesarean section versus planned vaginal birth for women with a previous caesarean birth. *Cochrane Database Syst Rev* 2004; 4, Art. No. CD004224.
- 34 Wildman K, Blondel B, Nijhuis J, Defoort P, Bakoula C. European indicators of health care during pregnancy, delivery and the postpartum period. *Eur J Obstet Gynecol Reprod Biol* 2003;111: S53–65.
- 35 Euro-Peristat project in collaboration with SCPE, EUROCAT and EURONEOSTAT. Better statistics for better health for pregnant women and their babies in 2004. European Perinatal Health Report 2008.[www.europeristat.com] Accessed 10 February 2015.
- 36 Guihard P, Blondel B. Les facteurs associe a la pratique d'une césarienne en France. Resultats de l'enquete nationale perinatale de 1995. *J Gynecol Obstet Biol Reprod* 2001;30:444–543.
- 37 Bragg F, Cromwell D, Edozien LC, Guroi-Uganci I, Mahmood TA, Templeton A, et al. Variation in rates of caesarean section among English NHS trusts after accounting for maternal and clinical risk: cross sectional study. *BMJ* 2010;341:c5065.
- 38 Alves B. Risk factors and variation in caesarean section rates in England in the late 1990s. MSc dissertation. London School of Hygiene and Tropical Medicine, 2002.
- 39 Paranjothy S, Frost C, Thomas J. How much variation in CS rates can be explained by case mix differences. *BJOG* 2005;112:658–66.
- 40 RCOG, London School of Hygiene and Tropical Medicine. Patterns of maternity care in NHS hospitals 2011/12. London: RCOG. [http://www.rcog.org.uk/our-profession/research-and-audit/clinical-indicators-project]. Accessed 3 May 2013.
- 41 Robson M. Classification of caesarean sections. *Fetal Matern Med Rev* 2001;12:23–39.
- 42 Zeitlin J, Wildman K, Breart G, Alexander S, Barros H, Blondel B, et al. Selecting an indicator set for monitoring and evaluating perinatal health in Europe: criteria, methods and results from the PERISTAT project. *Eur J Obstet Gynecol Reprod Biol* 2003;111(Suppl 1):5–14.
- 43 Euro-Peristat project with SCPE and Eurocat. European Perinatal health report. The health of pregnant women and babies in Europe in 2010. May 2013. [www.europeristat.com]. Accessed 10 February 2015.
- 44 Blondel B, Lelong N, Kermarrec M, Goffinet F. Trends in perinatal health in France from 1995 to 2010. Results from the French National Perinatal Surveys. *J Gynecol Obstet Biol Reprod (Paris)* 2012;41:e1–15.
- 45 Gissler M, Mohangoo AD, Blondel B, Chalmers J, Macfarlane A, Gaizauskiene A, et al. Perinatal health monitoring in Europe: results from the EURO-PERISTAT project. *Inform Health Soc Care* 2010;35:64–79.
- 46 McPherson K, Gon G, Scott M. International variations in a selected number of surgical procedures. OECD Health Working Papers, No. 61. OECD Publishing. [http://dx.doi.org/10.1787/5k49h4p5g9mw-en]. Accessed 2 May 2014.
- 47 Thomas J, Paranjothy S. *The National Sentinel Caesarean Section Audit*. London: RCOG Press; 2001.
- 48 Carayol M, Blondel B, Zeitlin J, Breart G, Goffinet F. Changes in the rates of caesarean delivery before labour for breech presentation at term in France: 1972–2003. *Eur J Obstet Gynecol Reprod Biol* 2007;132:20–6.
- 49 Goffinet F, Carayol M, Foidart JM, Alexander S, Uzan S, Subtil D, et al. Is planned vaginal delivery for breech presentation at term still an option? Results of an observational prospective survey in France and Belgium *Am J Obstet Gynecol* 2006;194:1002–11.
- 50 Molkenboer JFM, Bouckaert PJXM, Roumen FJME. Recent trends in breech delivery in the Netherlands. *BJOG* 2003;110:948–51.
- 51 Royal College of Obstetricians and Gynaecologists. *Birth after Previous Caesarean Birth. Green-top Guideline No. 45*. London: RCOG; February 2007.
- 52 Guihard P, Blondel B. Trends in risk factors for caesarean sections in France between 1981 and 1995; lessons for reducing the rates in the future. *BJOG* 2001;108:48–55.
- 53 Brennan DJ, Robson MS, Muphy M, O'Herlihy C. Comparative analysis of international caesarean delivery rates using 10-group classification identifies significant variation in spontaneous labor. *Am J Obstet Gynaecol* 2009;201:308e1–8.
- 54 Domingues RMSM, Dias MAB, Pereira MN, Torres JA, d'Orsi E, Pereira APE, et al. Process of decision-making regarding the mode of birth in Brazil: from the initial preference of women to the final mode of birth. *Cad Saúde Pública* 2014; 30(suppl. 1) Rio de Janeiro Aug. 2014. [http://dx.doi.org/10.1590/0102-311X00105113]. Accessed 22 September 2014.
- 55 Hopkins K, de Lima Amaral EF, Mourão AN. The impact of payment source and hospital type on rising cesarean section rates in Brazil, 1998–2008. *Birth* 2014;41:169–77.
- 56 Shorten A, Shorten B. What happens when a private hospital comes to pact of the 'public' to 'private' hospital shift on regional birthing outcomes. *Women Birth* 2007;20:49–55.

- 57 Kingdon C, Neilson J, Singleton V, Gyte G, Hart A, Gabbay M, et al. Choice and birth method: mixed method study of caesarean delivery for maternal request. *BJOG* 2009;116:886–95.
- 58 McCourt C, Weaver J, Statham H, Beake S, Gamble J, Creedy D. Elective caesarean section and decision making: a critical review of the literature. *Birth* 2007;34:65–79.
- 59 Mazzoni A, Althabe F, Liu NH, Bonotti AM, Gibbons L, Sánchez AJ, et al. Women's preference for caesarean section: a systematic review and meta-analysis of observational studies. *BJOG* 2011;118:391–9.
- 60 Deng W, Klemetti R, Long Q, Wu Z, Duan C, Zhang WH, et al. Cesarean section in Shanghai: women's or healthcare provider's preferences. *BMC Pregnancy Childbirth* 2014;14:285.
- 61 Donati S, Grandolfo ME, Andreozzi S. Do Italian mothers prefer Cesarean Delivery? *Birth* 2003;30:89–93.
- 62 Dweik D, Girasek E, Mészáros G, Tőreki A, Keresztúri A, Pál A. Non-medical determinants of cesarean section in a medically dominated maternity system. *Acta Obstet Gynecol Scand* 2014;93:1025–33.
- 63 Coulm B, Blondel B, Alexander S, Bouvain M, Le Ray C. Potential avoidability of planned cesarean sections in a French national database. *Acta Obstet Gynecol Scand* 2014;93:905–12.