



City Research Online

City, University of London Institutional Repository

Citation: Sun, Z., Longmire, E. K. & Krizan, D. (2014). Interactions of copepods with fractal-grid generated turbulence based on Tomo-PIV and 3D-PTV. Paper presented at the 67th Annual Meeting of the APS Division of Fluid Dynamics, 23-11-2014 - 25-11-2014, San Francisco, USA.

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/7933/>

Link to published version:

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk

Abstract Submitted
for the DFD14 Meeting of
The American Physical Society

Interactions of Copepods with Fractal-Grid Generated Turbulence based on Tomo-PIV and 3D-PTV¹ ZHENGZHONG SUN, DANIEL KRIZAN, ELLEN LONGMIRE, University of Minnesota — A copepod escapes from predation by sensing fluid motion caused by the predator. It is thought that the escape reaction is elicited by a threshold value of the maximum principal strain rate (MPSR) in the flow. The present experimental work attempts to investigate and quantify the MPSR threshold value. In the experiment, copepods interact with turbulence generated by a fractal grid in a recirculating channel. The turbulent flow is measured by time-resolved Tomo-PIV, while the copepod motion is tracked simultaneously through 3D-PTV. Escape reactions are detected based on copepod trajectories and velocity vectors, while the surrounding hydrodynamic information is retrieved from the corresponding location in the 3D instantaneous flow field. Measurements are performed at three locations downstream of the fractal grid, such that various turbulence levels can be achieved. Preliminary results show that the number of escape reactions decreases at locations with reduced turbulence levels, where shorter jump distances and smaller change of swimming orientation are exhibited. Detailed quantitative results of MPSR threshold values and the dynamics of copepod escape will be presented.

¹Supported by NSF-IDBR Grant #0852875.

Zhengzhong Sun
University of Minnesota

Date submitted: 29 Jul 2014

Electronic form version 1.4